



**HAL**  
open science

## Hyperstructures associated to a connected graph

Gabriel Turinici

► **To cite this version:**

Gabriel Turinici. Hyperstructures associated to a connected graph. An. Stiint. Univ. Ovidius Constanta, Ser. Mat., 1994, 2, pp.179-187. hal-00536508

**HAL Id: hal-00536508**

**<https://hal.science/hal-00536508>**

Submitted on 27 Jun 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## HYPERSTRUCTURES ASSOCIATED TO A CONNECTED GRAPH

Gabriel Turinici

Seminarul Matematic, Univ. "Al. I. Cuza" Iasi

Bd. Copou 11, Ro-6600 Iasi

Romania

### Abstract

We are searching to associate, in a natural way, a hyperstructure to a connected graph showing that in many relevant cases it is a join-space (under the convention in [3]).

1. Let  $H$  be a nonempty set and  $\mathcal{P}^*(H) = \{A \mid A \subseteq H, A \neq \emptyset\}$  (where  $\subseteq$  stands for the non-strict inclusion). A *hyperoperation* on  $H$  is a function  $(x, y) \rightarrow xoy$  from  $H \times H$  to  $\mathcal{P}^*(H)$ . We will denote  $A \circ B = \cup\{a \circ b \mid a \in A, b \in B\}$  for  $A, B \in \mathcal{P}^*(H)$ .

A hyperoperation  $\circ$  is called *associative* if

$$(\forall)x, y, z \in H : (x \circ y) \circ z = x \circ (y \circ z).$$

In this case  $(H, \circ)$  will be referred to as a *semi-hypergroup*. The hyperoperation  $\circ$  is called *reproducible*, if  $xoH = H$  or  $x \circ H = H$  for each  $x \in H$ .

The semi-hypergroup  $(H, \circ)$  with " $\circ$ " reproducible is called a *hypergroup*. In this case we introduce for each  $a, b \in H$   $a/b = \{c \in H \mid a \in b \circ c\}$ . A commutative hypergroup  $(H, \circ)$  which satisfies:

$$(J) \quad (\forall)(a, b, c, d) \in H^4, a/b \cap c/d \neq \emptyset \implies a \circ d \cap b \circ c \neq \emptyset$$

is called a *join-space* (as in [3]).

If in addition the join-space  $(H, \circ)$  has the property (see [1], pag. 71):

$$x \circ x = \{x\} = x/x \text{ for all } x \in H$$

$(H, \circ)$  is said to be a *geometrical join-space*. Otherwise it is called a non-geometrical join-space.

2. Let  $G=(V, E)$  be a tree (from now on we suppose  $\text{card}(V) \geq 2$ ).

(a)  $\forall x \in V, x \circ x = \{x\}$ ;

(b)  $\forall x, y \in V, x \neq y, x \circ y = [xy] = \{z \in V \mid z \text{ is a vertex on the elementary path that joins } x \text{ and } y\}$ . Note that because  $G$  is a tree, (b) is consistent and

non-contradictory ( $G$  tree  $\implies \forall x, y \in V, x \neq y$ , there exists exactly one path joining  $x$  and  $y$ ).

**Theorem 1.** Let  $G=(V,E)$  be a tree and  $(V,o)$  be its associated hyperstructure by the method above. Then  $(V,o)$  is a non-geometrical join-space.

**Proof.** We notice for the beginning that  $o$  is commutative,  $x \circ x = x/x = \{x\}$  for each  $x \in V$ , and  $\{x, y\} \subseteq x \circ y$ , for any  $x, y \in V$  with  $x \neq y$ .

To prove the associativity we start by showing that:

$$(V)x, y, z \in V : (x \circ y) \circ z \subseteq x \circ (y \circ z). \tag{1}$$

If  $x = y$  then  $(x \circ y) \circ z = x \circ z, x \circ (y \circ z) = x \circ (x \circ z)$  and since  $x \circ z \subseteq x \circ (x \circ z)$ , (1) follows.

If  $x \neq y$ , denote by  $x = \alpha_1, \alpha_2, \dots, \alpha_n = y$  the path  $[xy]$  that joins  $x$  and  $y$ , and by  $y = \beta_1, \beta_2, \dots, \beta_k = z$  the path  $[yz]$ . Two situations occur:

i)  $x \circ y \cap y \circ z = \{y\}$ ; then one can easily see that

$$(x \circ y) \circ z = x \circ y \cup y \circ z = \{\alpha_1, \alpha_2, \dots, \alpha_n = y = \beta_1, \beta_2, \dots, \beta_k = z\}$$

and also

$$x \circ (y \circ z) = x \circ \{\beta_1, \beta_2, \dots, \beta_k = z\} = \{\alpha_1, \alpha_2, \dots, \alpha_n = y = \beta_1, \beta_2, \dots, \beta_k = z\};$$

so we have equality.

ii)  $x \circ y, y \circ z$  have common points different from  $y$ . As  $G$  is a tree, there exists  $p \in \{1, \dots, n\}, q \in \{1, \dots, k\}$  such that  $q = \max\{i \mid \beta_i \in x \circ y\}$   $\alpha_p = \beta_q$  and

$$\begin{aligned} x \circ y &= \{\alpha_1 = x, \alpha_2, \dots, \alpha_{p+1} = \beta_{q-1}, \dots, \alpha_p = \beta_q, \alpha_n = y = \beta_1\} \\ y \circ z &= \{\beta_1 = y = \alpha_n, \beta_2 = \alpha_{n-1}, \dots, \beta_q = \alpha_p, \beta_{q+1}, \dots, \beta_k = z\}. \end{aligned}$$

As a consequence of this,

$$x \circ z = \{x = \alpha_1, \alpha_2, \dots, \alpha_p = \beta_q, \beta_{q+1}, \dots, \beta_k = z\} \text{ (because } G \text{ has no cycles).}$$

Finally one obtains

$$(x \circ y) \circ z = x \circ (y \circ z) = \{\alpha_1, \dots, \alpha_n, \beta_{q+1}, \dots, \beta_k\}.$$

So far we proved (1), therefore we can write:

$$(x \circ y) \circ z \subseteq x \circ (y \circ z) = (y \circ z) \circ x \subseteq y \circ (z \circ x) = (z \circ x) \circ y \subseteq z \circ (x \circ y) = (x \circ y) \circ z.$$

Now all the signs  $\subseteq$  can be replaced by  $=$  in this last relation; and therefore, the associativity of  $o$  is proved.

We noticed at the beginning that  $(V)x, y \in V : y \in x \circ y$ . So  $(V)y \in V : y \in x \circ y \subseteq x \circ V$  where from  $(V)y \in V : y \in x \circ V = V \circ x$ . Hence  $o$  is reproducible.

To finish the proof we must show that (J) holds. Let  $(a, b, c, d) \in V^4$  such that  $a/b \cap c/d \neq \emptyset$ . This means  $(\exists)\alpha \in V, \alpha \in a/b \cap c/d$  (hence  $a \in b \circ \alpha, c \in d \circ \alpha$ ). We have two cases to analyze:

i)  $b \circ \alpha \cap d \circ \alpha = \{\alpha\}$ . As  $a \circ d \cap b \circ c$  contains at least  $a, c, \alpha$  (J) follows.

ii)  $b \circ \alpha \cap d \circ \alpha \neq \{\alpha\}$ . Let  $b \circ \alpha = \{\alpha_1 = b, \dots, \alpha_i = a, \dots, \alpha_n = \alpha\}, d \circ \alpha = \{\beta_1 = d, \dots, \beta_j = c, \dots, \beta_m = \alpha\}$  and  $\beta_{j_0} = \max\{k \mid \beta_k \in b \circ \alpha\}, \alpha_{i_0} = \beta_{j_0} = e$ .

There are two possibilities for  $a$ :  $a \in b \circ e$  or  $a \in e \circ \alpha$  and likewise for  $c$ :  $c \in d \circ e$  or  $c \in e \circ \alpha$ . As we will see, in all cases (J) is valid:

a)  $a \in b \circ e, c \in e \circ \alpha; b \circ c \cap a \circ d$  contains  $a, e$ .

b)  $a \in b \circ e, c \in d \circ e$ ; we find  $a, c, e$  are in  $a \circ c$  in  $b \circ c \cap a \circ d$ .

c)  $\{a \in e \circ \alpha, c \in e \circ \alpha\}$  or  $\{a \in b \circ e, c \in d \circ e\}$ ; then  $e \in b \circ c \cap a \circ d$ .

The proof is now complete.

**Remark 1.** Denote by  $H_G$  the join-space  $(V,o)$  associated to a tree  $G=(V,E)$  like above. If  $\Gamma$  is the set of isomorphism classes of trees and  $G$  the set of isomorphism classes of join-spaces, then the function  $f : \Gamma \rightarrow G$  defined by  $f(\bar{G}) = \overline{H_G}$  is injective (here  $\bar{x}$  is the class of  $x$ ). Indeed, the condition  $x \circ y = \{x, y\}$  in  $H_G$  is equivalent to  $(x, y)$  being an edge in the tree  $G$  and so we can construct in only one way the graph  $G$  starting from  $H_G$ .

3. Let  $G=(V,E)$  be a graph. We say that  $G$  is *2-connected* if  $\text{card}(V) > 2$ ,  $G$  is connected and by removing a vertex one obtains a connected graph too. A block in  $G$  is a maximal 2-connected subgraph of  $G$ .

Let  $G=(V,E)$  be a connected graph and denote by  $b_1, \dots, b_k$  its blocks. Construct the graph  $G^* = (V^*, E^*)$  where  $V^* = V \cup \{b_1, \dots, b_k\}$  and  $(x, y) \in E^*$  if and only if one of the following cases occurs:

i)  $x \in V, y = b_i$  (for some  $i \in \{1, \dots, k\}$ ) and  $x$  is a vertex of  $b_i$  in  $G$  (or vice versa  $y \in V, x = b_i$ .)

ii)  $(x, y) \in E$  and there is no elementary cycle passing through  $x$  and  $y$ .

**Example 1.** Let  $G$  as below (fig. 4),  $b_1 = \{6, 7, 8, 9\}, b_2 = \{2, 3, 4, 5\}$ . Then  $G^*$  is as in fig. 5.

**Remark 2.**  $G^*$  is a tree. Indeed,  $G$  connected implies  $G^*$  connected, and if in  $G^*$  there exists a cycle it contains at least two blocks  $b_i$  and  $b_j$  of  $G$  as points. Then in  $G^*$  are two paths from  $b_i$  to  $b_j$  and the same will be in  $G$ . But then  $b_i$  and  $b_j$  are contained in the same block of  $G$ , contradiction.

Let  $G=(V,E)$  be a connected graph and  $G^*$  its block-tree. We denote by  $H_G$  the join space  $H_{G^*}$  and we will shall say that  $H_G$  is the join-space associated to  $G$ .

**Remark 3.** i) If  $G$  is a tree then  $G$  is its own block-tree and so the definition above extends the one we gave in 2.

ii) If we denote by  $\Gamma$  the set of isomorphism classes of graphs then  $f^* : \Gamma \rightarrow G$  defined by  $f^*(\bar{G}) = \overline{H_G}$  is (unlike in the case of  $f$ ) not injective.

iii) When  $G$  is 2-connected its block-tree is a star-shaped graph (there exists a point  $b \in V$  such that any edge in  $G^*$  is of a form  $(b, a)$  or  $(a, b)$ ,  $a \in V$ ) and the join-space associated to it

$$(V^*, \circ) : V^* = V \cup \{b\}, x \circ y = \{x, y, b\} \text{ for any } x, y \in V, x \neq y$$

gives no information about the graph-structure of  $G$ . In fact  $H_G$  characterizes the tree-structure of a graph; so it is natural to ask if there exists another hyperstructure construction for 2-connected graphs. An answer is given in Section 5.

4. In Section 3 we associated a join-space to a connected graph using its path-structure. A question arises: given a join-space  $(J, \circ)$  with  $\text{card}(J) < \infty$ , under which conditions upon  $J$  there exists a graph  $G$  (in fact a tree) such that  $J = H_G$ ?

In this section we suppose  $J$  is a finite set having at least two elements and  $\circ$  is a hyperoperation on  $J$ . Let us consider the following conditions:

- (A<sub>1</sub>)  $\circ$  is associative;
- (A<sub>2</sub>)  $\circ$  is commutative;
- (A<sub>3</sub>)  $x \circ J = J \circ x = J$ ,  $\forall x \in J$ ;
- (A<sub>4</sub>)  $x \circ x = x$ ,  $\forall x \in J$ .

**Proposition 1.** *If (A<sub>1</sub>) - (A<sub>4</sub>) and (J) are valid for (J, \circ), then for any  $x, y, z \in J$  such that  $z \in x \circ y$  and  $y \in x \circ z$  one has  $y = z$ .*

**Proof.** Indeed,  $z \in x \circ y \Rightarrow x \in z/y$  and  $y \in x/y$  and  $y \in x \circ z \Rightarrow x \in y/z$ . Then  $z/y \cap y/z \neq \emptyset \Rightarrow y \circ y \cap z \circ z \neq \emptyset \Rightarrow y = z$ .

A join-space for which (A<sub>1</sub>) - (A<sub>4</sub>) and (J) hold, as well as (G)  $\forall x, y \in J$ ,  $x \neq y : \{x, y\} \subseteq x \circ y$  is called a (g) - join space.

**Proposition 2.** *Let (J, \circ) be a (g)-join-space. Then for each  $x \in J$  there exists  $y \in J$  such that  $x \circ y = \{x, y\}$ .*

**Proof.** Let  $y \in J$  be an element of  $J \setminus \{x\}$  for which  $\text{card}(x \circ y)$  is minimal. By hypothesis (J, \circ) is a (g)-join-space; so that  $\text{card}(x \circ y) \geq 2$ . If  $\text{card}(x \circ y) = 2$  than  $(x \circ y) = \{x, y\}$ . If  $\text{card}(x \circ y) \geq 3$  there exists  $z \in x \circ y \setminus \{x, y\}$ . Now,  $z \in x \circ y$  gives  $x \circ z \subseteq x \circ (x \circ y) = (x \circ x) \circ y = x \circ y$ . The case  $x \circ z = x \circ y$  yields  $y \in x \circ z \subseteq x \circ y$  (where  $\subseteq$  stands for the strict inclusion) implies  $\text{card}(x \circ z) < \text{card}(x \circ y)$ , contradiction. This completes the proof.

Let  $(J, \circ)$  be a (g)-join-space. We define a relation  $\sim$  over  $J$  by:

$$x \sim y \leftrightarrow \{x \neq y \text{ and } x \circ y = \{x, y\}\}.$$

0	a	b	c	d	e
a	a	a,b	a,b,c	a,d,e	a,e
b	a,b	b	b,c	b,c,d	a,b,e
c	a,b,c	b,c	c	c,d	c,d,e
d	a,d,e	b,c,d	c,d	d	d,e
e	a,e	a,b,e	c,d,e	d,e	e

One can easily see that  $\sim$  is symmetrical whenever  $\circ$  is commutative. Any (g)-join-space  $(J, \circ)$  which satisfies the condition:

$$(P) \forall k \in N, (\forall) \{x_1, x_2, \dots, x_k\} \subseteq J \text{ with } x_1 \sim x_2, \dots, x_{k-1} \sim x_k \text{ then } \{x_1, \dots, x_k\} \subseteq x_1 \circ x_k$$

is called a path join-space.

**Remark 4.** i) From  $\{x_1, \dots, x_k\}$  being a set it follows  $x_i \neq x_j$  for all  $i, j \in \{1, \dots, k\}$ ,  $i \neq j$ .

ii) The above definition is not trivial. This is shown from the following example: Let  $(J, \circ), J = \{a, b, c, d, e\}$  be a (g)-join-space as below:

Note that  $x \circ y$  is the minimal path that joins  $x$  and  $y$  in the graph from fig. 6. This (g)-join-space is not a path join-space.

**Theorem 2.** *Let (J, \circ) be a path join-space. Then there is a tree  $G = (V, E)$  such that  $J = H_G$ .*

**Proof.** Let  $V = J$ ,  $E = \{(x, y) \in J^2 \mid x \sim y\}$ ,  $G = (V, E)$ . We shall show that  $G$  is a tree and  $J = H_G$ . To prove there is no cycle in  $G$ , suppose by contradiction that there exists  $\{x_1, \dots, x_k\} \subseteq J$  such that  $x_1 \sim x_2, x_2 \sim x_3, \dots, x_{k-1} \sim x_k, x_k \sim x_1$  ( $k \geq 3$ ). Then:

$$x_2 \in x_1/x_k \text{ (because } x_k \sim x_1, x_1 \sim x_2 \Rightarrow x_1 \in x_2 \circ x_k)$$

$$x_2 \in x_k/x_1 \text{ (because } x_1 \sim x_k, x_k \sim x_{k-1}, \dots, x_3 \sim x_2 \Rightarrow x_k \in x_1 \circ x_2).$$

Hence  $x_1/x_k \cap x_k/x_1 \neq \emptyset$  where from  $x_1 \circ x_k \cap x_k \circ x_1 \neq \emptyset$ , which is impossible.

Now suppose that  $G$  is not connected, i.e. there exist at least two connected components in  $G$ . Let  $x_i$  and  $x_j$  be such that  $\text{card}(x_i \circ x_j)$  is minimal with the property that  $x_i$  and  $x_j$  are not in the same component. It follows that  $x_i \not\sim x_j$ ; so  $(\exists) z \in x_i \circ x_j \setminus \{x_i, x_j\}$ . We have seen that  $x_i \circ z \subseteq x_i \circ x_j, z \circ x_j \subseteq x_i \circ x_j$  implies  $\text{card}(x_i \circ z) < \text{card}(x_i \circ x_j), \text{card}(z \circ x_j) < \text{card}(x_i \circ x_j)$ . Since  $x_i, x_j$  are not in the same component at least one of the pairs  $(x_i, z), (x_j, z)$  has the same property, in contradiction with the minimality of  $\text{card}(x_i \circ x_j)$ . So far  $G$

is a tree. Let  $\square$  be the hyperoperation associated to the tree-structure of  $G$ . By condition (P),  $x \square y \subseteq x \circ y$  for each  $x, y \in J$ . Suppose that:

$$(*) \quad c \square y \subset x \circ y \text{ (obviously } x \neq y) \text{ for some } x, y \in J,$$

and take  $(x, y)$  with the property that  $\text{card}(x \circ y)$  has the minimum value among all the pairs which satisfy (\*). Let  $z \in x \circ y \setminus x \square y$  (of course  $z \notin \{x, y\}$ ). Since  $z \in x \circ y$  and  $z \notin \{x, y\}$ ,  $\text{card}(x \circ z) < \text{card}(x \circ y)$ ,  $\text{card}(z \circ y) < \text{card}(x \circ y)$  and, by minimality,  $x \square z = x \circ z$  and  $z \square y = z \circ y$ .

In  $G$ ,  $x \square z$  intersects  $z \square y$  at least in one point (for example  $z$ ). If  $x \square z \cap z \square y = \{z\}$  we obtain  $z \in x \square y$ , contradiction. It follows that  $\exists t \in (x \square z \cap z \square y) \setminus \{z\}$ . Let us choose  $t$  such that  $x \square t \cap y \square t = \{t\}$ . Then  $t \in x \square z \subseteq x \circ z$ , and this yields  $x \in t/z$ . Since  $z \in x \circ y \Rightarrow x \in z/y$  it follows that:

$$z/y \cap t/z \neq \emptyset \Rightarrow z \circ z \cap t \circ y \neq \emptyset \Rightarrow z \in y \circ t.$$

On the other hand we have in  $(J, \square)$   $y \square x = y \square t \cup t \square x$ . This yields  $y \square t \subseteq y \square x$  where from  $z \in y \square x$ , contradiction. Hence  $x \square y = x \circ y$ ,  $\forall x, y \in J$  and consequently,  $H = H_G$ .

**Remark 5.** The result above shows that - supposing we keep the set  $V$  and we make no difference among paths joining the same end-points - the construction in Section 2 is unique and is possible only for trees.

5. Let  $G=(V, E)$  be a 2-connected graph. We have seen in Section 3 that in this case a new construction is needed. This is founded on an equivalent definition of a 2-connected graph according to which for any two different elements  $x, y \in V$  there is a cycle passing through  $x, y$ . Define a hyperoperation  $\circ : V \times V \rightarrow \mathcal{P}^*(V)$  by:

- a.  $x \circ x = \{x\} \forall x \in V$ ;
  - b.  $x \circ y = \{z \in V \mid \text{there is a cycle through } x, y, z\} \forall x, y \in V, x \neq y$ .
- $G$  being 2-connected  $\circ$  is well defined. It is easy to see that  $\circ$  is commutative and

(\*\*) if  $\{x, y, z\} \subseteq V$  then  $x \in y \circ z \Leftrightarrow y \in x \circ z \Leftrightarrow z \in x \circ y$ .  
In general,  $\circ$  is not associative. This is shown by the following

**Example 2.** Let  $H = \{a_1, a_2, b_1, b_2, c_1, c_2, m_1, m_2, n_1, n_2, p_1, p_2\}$  and the graph (fig. 8)

We notice that  $(a_1 \circ a_1) \circ b_1 = a_1 \circ b_1 = H \setminus \{a_2, b_2\}$  and  $a_1 \circ (a_1 \circ b_1) = a_1 \circ H \setminus \{a_2, b_2\} = H$ , so  $\circ$  is not associative.

**Theorem 3.** Let  $G=(V, E)$  be a 2-connected graph and  $\circ$  be the hyperoperation just defined. Then  $\circ$  is associative if and only if  $x \circ y = V$  for each  $x, y \in V, x \neq y$ .



**Proof.** One can easily see that  $\circ$  is reproducible and so the implication  $\Rightarrow$  is obvious. Conversely, assume that  $\circ$  is associative and let  $a, b, c \in V$  with  $b \neq c$ . We will shall prove  $a \in b \circ c$ . Suppose on the contrary  $a \notin b \circ c$ . If there exists  $f \in b \circ c \setminus \{b, c\}$  with  $f \in a \circ b$  then  $a \neq f$  (because  $a \notin b \circ c$ ). But  $f \in a \circ b \Rightarrow a \in b \circ f$ ; hence  $a \in b \circ f \subseteq b \circ (b \circ c) = (b \circ b) \circ c = b \circ c$ , contradiction. It follows  $a \circ b \cap b \circ c = \{b\}$  ( $c \in a \circ b$  would imply  $a \in b \circ c$ ). Similarly, we may show that  $a \circ c \cap b \circ c = \{c\}$ . Suppose now that there exists  $x \in a \circ c \cap a \circ b, x \neq a$ . Of course, if  $x = b$  then  $b \in a \circ c$ , impossible. Hence  $x \neq b$  and in the same way  $x \neq c$ . Now  $x \in a \circ c, x \neq a, c \Rightarrow a \in c \circ x$  and  $x \in a \circ b, x \neq a, b \Rightarrow b \in a \circ x$ . So,  $b \in (c \circ x) \circ x$  where from  $b \in c \circ x$  which in turn gives  $x \in b \circ c$ , in contradiction with  $x \in a \circ c$ . We therefore proved that:

$$a \circ b \cap b \circ c = \{b\}; \quad a \circ b \cap a \circ c = \{a\}; \quad b \circ c \cap a \circ c = \{c\}. \tag{2}$$

In  $a \circ b$  there is a path  $P_{ab}$  joining  $a$  and  $b$ ; let also  $P_{ac}$  and  $P_{bc}$  be paths in  $a \circ c$  and  $b \circ c$  respectively. We obtain from (2) that  $P_{ab} \cup P_{ac} \cup P_{bc}$  is a cycle containing  $a, b, c$ ; hence  $a \in b \circ c$ , contradiction. Summarizing up,  $a \notin b \circ c$  leads to a contradiction; so  $a \in b \circ c$  for any  $b, c \in V, b \neq c$ , and the proof is complete.

**Remark 6.** i) The results above can be restated as:  $\circ$  is associative if and only if for any three points in  $V$  there is a cycle containing them.

ii) An interesting class of 2-connected graphs is the one consisting of all such objects that contain a hamiltonian cycle. In this case  $\circ$  is associative but this property is not sufficient for a 2-connected graph to have a hamiltonian cycle as one can see from the example in fig. 9.



References

1. Corsini, P., *Prolegomena of hypergroup theory*, Suppl. alla Riv. di mat. pura ed. appl., Aviani Editore, 1991.
2. Nieminen J., *Join spaces graphs*, J. of Geometry, 33 (1988).
3. Prenowitz W., Jantosciak J., *Geometries and Join Spaces*, J. Reine und Angewandte Math., vol. 257 (1972), pp 100-128.
4. Harary F., *Graph theory*, Addison-Wesley Publishing Company, 1972.

Received: September 1994




Fig. 4




Fig. 5



Fig. 7




Fig. 6




Fig. 8




Fig. 9