



HAL
open science

Shiga-Watanabe's time inversion property for self-similar diffusion processes

Juha Vuolle-Apiala

► **To cite this version:**

Juha Vuolle-Apiala. Shiga-Watanabe's time inversion property for self-similar diffusion processes. *Statistics and Probability Letters*, 2009, 77 (9), pp.920. 10.1016/j.spl.2007.01.001 . hal-00538013

HAL Id: hal-00538013

<https://hal.science/hal-00538013>

Submitted on 20 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

Shiga-Watanabe's time inversion property for self-similar diffusion processes

Juha Vuolle-Apiala


PII: S0167-7152(07)00031-4
DOI: doi:10.1016/j.spl.2007.01.001
Reference: STAPRO 4564

To appear in: *Statistics & Probability Letters*

Received date: 12 February 2004
Revised date: 19 June 2006
Accepted date: 17 January 2007

Cite this article as: Juha Vuolle-Apiala, Shiga-Watanabe's time inversion property for self-similar diffusion processes, *Statistics & Probability Letters* (2007), doi:10.1016/j.spl.2007.01.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



www.elsevier.com/locate/stapro

SHIGA-WATANABE'S TIME INVERSION PROPERTY FOR
SELF-SIMILAR DIFFUSION PROCESSES

Juha Vuolle-Apiala

Department of Mathematics and Statistics, University of Vaasa, P. O. Box
700, 65101 Vaasa, Finland, e-mail: jmva@uwasa.fi

Summary : Assume (X_t, P^0) is $\frac{1}{2}$ -self-similar, rotation invariant diffusion on \mathbb{R}^d , $d \geq 2$, starting at 0 and assume $\{0\}$ is a polar set. We will show, using the corresponding well-known result for the radial process, that Shiga-Watanabe's time inversion property holds for (X_t, P^0) . The generalization for an α -self-similar, rotation invariant diffusion, $\alpha > 0$, is also given.

Keywords : Time inversion, self-similar, Bessel process, diffusion, rotation invariant, skew product, radial process, spherical process

AMS Mathematics Subject Classification (2000): 60G18, 60J60, 60J25

0. Introduction. Theorem

The following time inversion property is well known for Brownian motion in \mathbb{R}^d , $d \geq 1$, and Bessel diffusions on $[0, \infty)$, starting at 0 (see Shiga, Watanabe (1973) and Watanabe (1975):

(X_t) has the same finite dimensional distributions as $(tX_{1/t})$ under P^0 . (1)

Graversen, Vuolle-Apiala (2000) show that (1) also holds for symmetrized Bessel processes on \mathbb{R} , starting at 0, in the case of the index $\nu \in (-1, 0)$. In this case (X_t) can both hit 0 and can be started at 0. Symmetrized Bessel processes form the class of one dimensional rotation invariant $\frac{1}{2}$ -self similar diffusions (see the definition below). If the index $\nu \leq -1$ then 0 is an exit boundary point. That is, (X_t) can hit 0 but cannot be started there. If $\nu \geq 0$ then 0 is an entrance boundary point, that is, (X_t) can be started there and it will never come back. Thus in this case (X_t) in fact lives either on $[0, \infty)$ or on $(-\infty, 0]$ and $\{0\}$ is a polar set. Obviously, (1) is valid. Result (1) has been generalized for α -self-similar diffusions on $[0, \infty)$ and for symmetric α -self-similar diffusions on \mathbb{R} ,

$\alpha > 0$, in Graversen, Vuolle-Apiala (2000). The corresponding generalization of (1) is then

$$(X_t) \text{ has the same finite dimensional distributions as } (t^{2\alpha} X_{1/t}) \text{ under } P^0. \quad (2)$$

In this note we will show that (2) holds for all rotation invariant, d -dimensional, α -self-similar diffusions (that is, strong Markov processes with continuous paths), $d \geq 2$, $\alpha > 0$, for which $\{0\}$ is a polar set. Our main tool is a skew product representation for rotation invariant diffusions starting at 0; see Ito, Mc Kean Jr. (1974), p. 274-276.

Let (X_t, P^x) be a rotation invariant (RI) α -self-similar (α -ss) diffusion on \mathbb{R}^d , $d \geq 2$, $\alpha > 0$, such that $\{0\}$ is a polar set. By α -self-similarity we mean that

$$(X_t) \text{ under } P^x \text{ has the same finite dimensional distributions as } (a^{-\alpha} X_{at}) \text{ under } P^{a^{-\alpha}x} \text{ for all } x \in \mathbb{R}^d, a > 0 \quad (3)$$

and by (RI) that

$$(X_t) \text{ under } P^x \text{ has the same finite dimensional distributions as } (T^{-1}(X_t)) \text{ under } P^{T(x)} \text{ for all } T \in O(d). \quad (4)$$

Self-similar diffusions on \mathbb{R} or on $[0, \infty)$ are defined similarly. Brownian motion fullfills both (3) and (4). See more about self-similar Markov processes in Graversen, Vuolle-Apiala (1986), Lamperti (1972) and Vuolle-Apiala, Graversen (1986). According to Graversen, Vuolle-Apiala (1986) and Vuolle-Apiala (2003), when $X_0 \neq 0$ the diffusion processes which fullfill (3) and (4) can be represented as skew products

$$(|X_t|, \theta_{A_t}), \quad (5)$$

where $A_t = \lambda \int_0^t |X_s|^{-\frac{1}{\alpha}} ds$ for some $\lambda > 0$, the radial part $(|X_t|)$ is an α -ss diffusion on $(0, \infty)$, and (θ_t) is a spherical Brownian motion on S^{d-1} independent of $(|X_t|)$.

REMARK: As showed in Graversen, Vuolle-Apiala (1986), (5) is valid for all strong Markov processes with cadlag paths fullfilling (3) and (4). However, as showed by a counterexample by J. Bertoin, W. Werner (1996), the independence between $(|X_t|)$ and (θ_t) is not necessarily true if the paths are only right continuous. There is an error in the proof of Proposition 2.4, p.19-20 in Graversen, Vuolle-Apiala (1986). It was shown in Vuolle-Apiala (2003), Lemma 2.1, that $(|X_t|)$ and (θ_t) are independent in the case of continuous paths.

We want to prove the following:

THEOREM: Let (X_t, P^0) be an (RI) α -ss diffusion on \mathbb{R}^d , $\alpha > 0$, $d \geq 2$, starting at 0, having $\{0\}$ as a polar set. Then the time inversion property (2) is valid.

1. The Proof of the Theorem

The proof will be based on

PROPOSITION: Let (r_t) be an α -ss diffusion on $[0, \infty)$, $\alpha > 0$, such that 0 is an entrance, non-exit boundary point. Then the skew product

$$[r_t, \theta_{\lambda \int_0^t r_s^{-\frac{1}{\alpha}} ds}], t > 0, r_0 > 0, \quad (6)$$

where (θ_t, Q^θ) is a spherical Brownian motion on S^{d-1} independent of (r_t) , such that $Q^\theta(\theta_0=0) = 1 \forall \theta \in S^{d-1}$,

can be completed to be an α -ss diffusion on \mathbb{R}^d by defining

$$[r_t, \nu_{\lambda \int_1^t r_s^{-\frac{1}{\alpha}} ds}], t > 0, \text{ when } r_0=0, \quad (7)$$

where (ν_t, Q) is an independent, spherical Brownian motion defined for $-\infty < t < +\infty$ and the law of ν_0 is the uniform spherical distribution $m(d\theta)$.

REMARK 1: Because of a uniqueness result of RI measures on S^{d-1} there is at most one way to complete (6) to be RI on the \mathbb{R}^d .

REMARK 2: It is obvious that (ν_t, Q) in fact is a stationary process and ν_t is uniformly distributed for all $t \in \mathbb{R}$ (see Kuznetsov, 1973).

We have

$$Q\{\nu(t_1) \in d\theta_1, \dots, \nu(t_n) \in d\theta_n\} = m(d\theta) Q^{\theta_1}(\theta(t_2-t_1) \in d\theta_2) \dots Q^{\theta_{n-1}}(\theta(t_n-t_{n-1}) \in d\theta_n) \quad (8)$$

for $-\infty < t_1 < \dots < t_n < +\infty$ (for typographical reasons we use in formula (8) notations $\nu(t_i)$ and $\theta(t_i)$ instead of ν_{t_i} and θ_{t_i})

In order to prove Proposition we need

LEMMA 1: $(\nu_{t_1}, \dots, \nu_{t_n})$ under Q has the same distribution as $(\nu_{-t_1}, \dots, \nu_{-t_n})$.

PROOF: Follows immediately from (8) and the fact that (θ_t, Q^θ) has a symmetric density with respect to the uniform measure $m(d\theta)$ on S^{d-1} (see Vuolle-Apiala, Graversen (1986), Lemma 3, p.329). \square

In the proof of Proposition we will use the result of Ito-McKean (1974), p.275, which says that the skew product (6) can be completed to be a diffusion

(which obviously is RI) on the whole \mathbb{R}^d having the skew product (7) when $r_0 = 0$ iff $A_{0+} = \infty$ a.s. P^0 . Here we need

LEMMA 2: Let (r_t) be an α -ss diffusion on $[0, \infty)$ such that 0 is an entrance, non-exit boundary point. Then

$$P^0\left\{\int_0^\epsilon r_s^{-\frac{1}{\alpha}} ds = \infty\right\} = 1 \quad \forall \epsilon > 0.$$

PROOF of LEMMA 2: The property (2) is valid for (r_t) which implies

$$P^0\left\{\int_0^\epsilon r_s^{-\frac{1}{\alpha}} ds = \infty\right\} = P^0\left\{\int_0^\epsilon (s^{2\alpha} r_{1/s})^{-\frac{1}{\alpha}} ds = \infty\right\} = P^0\left\{\int_0^\epsilon s^{-2} r_{1/s}^{-\frac{1}{\alpha}} ds = \infty\right\}.$$

Substituting $1/s = h$ gives this equal to

$$P^0\left\{\int_\infty^{1/\epsilon} (-r_h^{-\frac{1}{\alpha}}) dh = \infty\right\} = P^0\left\{\int_{1/\epsilon}^\infty r_h^{-\frac{1}{\alpha}} dh = \infty\right\}.$$

Using the Markov property we obtain

$$P^0\left\{\int_{1/\epsilon}^\infty r_h^{-\frac{1}{\alpha}} dh = \infty\right\} = E^0\{P^{r_{1/\epsilon}}\left\{\int_0^\infty r_h^{-\frac{1}{\alpha}} dh = \infty\right\}\}.$$

Because 0 is an entrance, non-exit boundary point, $r_{1/\epsilon} > 0$ a.s. (P^0). Now, according to Lamperti (1972),

$$P^r\left\{\int_0^\infty r_h^{-\frac{1}{\alpha}} dh = \infty\right\} = 1 \quad \text{for all } r > 0$$

and thus

$$E^0\{P^{r_{1/\epsilon}}\left\{\int_0^\infty r_h^{-\frac{1}{\alpha}} dh = \infty\right\}\} = 1$$

which implies

$$P^0\left\{\int_0^\epsilon r_h^{-\frac{1}{\alpha}} dh = \infty\right\} = 1. \quad \square$$

PROOF OF PROPOSITION: It only remains to prove that the skew product

$$\left[r_t, \nu_{\lambda \int_1^t r_s^{-\frac{1}{\alpha}} ds}\right], \quad r_0=0, \quad (7)$$

fulfills the α -self-similarity condition (3) under P^0 . Let I_1, \dots, I_n be Borel subsets of $[0, \infty)$ and J_1, \dots, J_n Borel subsets of S^{d-1} . We will show

$$P^0\{r_{t_1} \in I_1, \dots, r_{t_n} \in I_n, \nu_{\lambda \int_1^{t_1} r_s^{-\frac{1}{\alpha}} ds} \in J_1, \dots, \nu_{\lambda \int_1^{t_n} r_s^{-\frac{1}{\alpha}} ds} \in J_n\} = P^0\{a^{-\alpha} r_{at_1} \in I_1, \dots, a^{-\alpha} r_{at_n} \in I_n, \nu_{\lambda \int_1^{at_1} r_s^{-\frac{1}{\alpha}} ds} \in J_1, \dots, \nu_{\lambda \int_1^{at_n} r_s^{-\frac{1}{\alpha}} ds} \in J_n\} (*)$$

for all $t > 0$. For simplicity, assume $n = 2$, the general case is analogous.

Now the right hand side of (*) for $n = 2$ is equal to

$$\begin{aligned} & P^0\{a^{-\alpha}r_{at_1} \in I_1, a^{-\alpha}r_{at_2} \in I_2, \nu_{\lambda \int_{1/a}^{t_1} (a^{-\alpha} r_{as})^{-\frac{1}{\alpha}} ds} \in J_1, \nu_{\lambda \int_{1/a}^{t_2} (a^{-\alpha} r_{as})^{-\frac{1}{\alpha}} ds} \in J_2\} = \\ & P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \nu_{\lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds} \in J_1, \nu_{\lambda \int_{1/a}^{t_2} r_s^{-\frac{1}{\alpha}} ds} \in J_2\} \end{aligned}$$

because (r_t) fullfills (3) and because of independence between (r_t) and (ν_t) .

This is further equal to

$$\begin{aligned} & P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \nu_{\lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds + \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds} \in J_1, \nu_{\lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds + \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds} \in J_2\} = \\ & \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds \in du, \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dv, \\ & \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dw, \nu_{u+v} \in J_1, \nu_{u+w} \in J_2\} = \\ & \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds \in du, \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dv, \\ & \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dw\} Q(\nu_{u+v} \in J_1, \nu_{u+w} \in J_2) = \end{aligned}$$

because of independence between (r_t) and (ν_t) . Now (ν_t) is a stationary process and thus this is equal to

$$\begin{aligned} & \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds \in du, \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dv, \\ & \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dw\} Q(\nu_v \in J_1, \nu_w \in J_2) = \\ & \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds \in dv, \lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds \in dw\} Q(\nu_v \in J_1, \nu_w \in J_2) = \\ & P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \nu_{\lambda \int_{1/a}^{t_1} r_s^{-\frac{1}{\alpha}} ds} \in J_1, \nu_{\lambda \int_{t_1}^{t_2} r_s^{-\frac{1}{\alpha}} ds} \in J_2\}. \quad \square \end{aligned}$$

Now we can prove Theorem:

PROOF OF THEOREM: (X_t) has according to Graversen, Vuolle-Apiala (1986), Vuolle-Apiala (2003) and Proposition a skew product representation

$$[r_t, \theta_{\lambda \int_0^t r_s^{-\frac{1}{\alpha}} ds}] \text{ as } X_0 \neq 0 \quad (6)$$

and

$$[r_t, \nu_{\lambda \int_1^t r_s^{-\frac{1}{\alpha}} ds}] \text{ as } X_0 = 0, \quad (7)$$

where (r_t) is the radial process, (θ_t, Q^θ) is an independent spherical Brownian motion such that $Q^\theta(\theta_0 = \theta) = 1$ and (ν_h, Q) is an independent, stationary, spherical Brownian motion defined for $-\infty < h < +\infty$ and the law of (ν_h) is the uniform spherical distribution for all $h \in \mathbb{R}$. To show (2) let us consider the distribution of $\{t_1^{2\alpha} X_{1/t_1}, \dots, t_n^{2\alpha} X_{1/t_n}\}$ under P^0 . Assume for simplicity $n = 2$, $\alpha = \frac{1}{2}$, the general case is analogous.

Let I_i and J_i , $i = 1, 2$, be Borel subsets of $(0, \infty)$ and S^{d-1} , respectively. Now

$$\begin{aligned} P^0\{t_1 X_{1/t_1} \in (I_1, J_1), t_2 X_{1/t_2} \in (I_2, J_2)\} = \\ P^0\{t_1 r_{1/t_1} \in I_1, t_2 r_{1/t_2} \in I_2, \nu_{\lambda \int_1^{1/t_1} r_s^{-2} ds} \in J_1, \nu_{\lambda \int_1^{1/t_2} r_s^{-2} ds} \in J_2\} = \\ \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{t_1 r_{1/t_1} \in I_1, t_2 r_{1/t_2} \in I_2, \nu_u \in J_1, \nu_v \in J_2, \lambda \int_1^{1/t_1} r_s^{-2} ds \in du, \\ \lambda \int_1^{1/t_2} r_s^{-2} ds \in dv\} = \\ \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{t_1 r_{1/t_1} \in I_1, t_2 r_{1/t_2} \in I_2, \lambda \int_1^{1/t_1} r_s^{-2} ds \in du, \lambda \int_1^{1/t_2} r_s^{-2} ds \in dv\} Q(\nu_u \in J_1, \nu_v \in J_2) = \\ \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{t_1 r_{1/t_1} \in I_1, t_2 r_{1/t_2} \in I_2, -\lambda \int_1^{t_1} (sr_{1/s})^{-2} ds \in du, -\lambda \int_1^{t_2} (sr_{1/s})^{-2} ds \in dv\} Q(\nu_u \in J_1, \nu_v \in J_2). \end{aligned}$$

Because (1) is true for (r_t) this is equal to

$$\begin{aligned} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, -\lambda \int_1^{t_1} r_s^{-2} ds \in du, -\lambda \int_1^{t_2} r_s^{-2} ds \in dv\} Q(\nu_u \in J_1, \nu_v \in J_2) = \\ \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_1^{t_1} r_s^{-2} ds \in du, \lambda \int_1^{t_2} r_s^{-2} ds \in dv\} Q(\nu_{-u} \in J_1, \nu_{-v} \in J_2). \end{aligned}$$

Using Lemma 1 we get this equal to

$$\begin{aligned} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \lambda \int_1^{t_1} r_s^{-2} ds \in du, \lambda \int_1^{t_2} r_s^{-2} ds \in dv\} Q(\nu_u \in J_1, \nu_v \in J_2) = \\ P^0\{r_{t_1} \in I_1, r_{t_2} \in I_2, \nu_{\lambda \int_1^{t_1} r_s^{-2} ds} \in J_1, \nu_{\lambda \int_1^{t_2} r_s^{-2} ds} \in J_2\} = \\ P^0\{X_{t_1} \in (I_1, J_1), X_{t_2} \in (I_2, J_2)\}. \end{aligned}$$

□

REFERENCES

Bertoin, J., Werner, W. (1996), Stable windings, Ann. Probab. 24, n:o 3, 1269-1279

Graversen, S.E., Vuolle-Apiala, J. (1986), α -self-similar Markov processes, Probab. Th. Related Fields 71(1), 149-158

Graversen, S.E., Vuolle-Apiala, J. (2000), On Paul Levy's arc sine law and Shiga-Watanabe's time inversion result, *Probab. Math. Statist.* 20(1), 63-73

Ito, K., McKean Jr., H.P. (1974), *Diffusion Processes and Their Sample Paths* (Springer-Verlag Berlin Heidelberg New York)

Kuznetsov, S.E. (1973), Construction of Markov processes with random birth and death times, *Theor. Probability Appl.* 18, 571-575

Lamperti, J.W. (1972), Semi-stable Markov processes I, *Z. Wahrschein. verw. Gebiete* 22, 205-225

Shiga, T., Watanabe, S. (1973), Bessel diffusions as one parameter family of diffusion processes, *Z. Wahrschein. verw. Gebiete* 27, 37-46

Vuolle-Apiala, J., Graversen, S.E. (1986), Duality theory for self-similar processes, *Ann. Inst. H. Poincare Probab. Statist.* 22(3), 323-332

Vuolle-Apiala, J. (2003), On certain extensions of a rotation invariant Markov process, *J. Theoret. Probab.* 16(4), 957-969

Watanabe, S. (1975), On time inversion of one-dimensional diffusion process, *Z. Wahrschein. verw. Gebiete* 31, 115-124