



HAL
open science

Coxeter orbits and Brauer trees

Olivier Dudas

► **To cite this version:**

Olivier Dudas. Coxeter orbits and Brauer trees. *Advances in Mathematics*, 2012, 229 (6), pp.3398–3435. 10.1016/j.aim.2012.02.011 . hal-00538577v2

HAL Id: hal-00538577

<https://hal.science/hal-00538577v2>

Submitted on 13 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Coxeter orbits and Brauer trees

Olivier Dudas^{*†}

January 13, 2012

Abstract

We study the cohomology with modular coefficients of Deligne-Lusztig varieties associated to Coxeter elements. Under some torsion-free assumption on the cohomology we derive several results on the principal ℓ -block of a finite reductive group $\mathbf{G}(\mathbb{F}_q)$ when the order of q modulo ℓ is assumed to be the Coxeter number. These results include the determination of the planar embedded Brauer tree of the block (as conjectured by Hiss, Lübeck and Malle in [25]) and the derived equivalence predicted by the geometric version of Broué's conjecture [7].

Introduction

Let \mathbf{G} be a quasi-simple algebraic group defined over an algebraic closure of a finite field of characteristic p . Let F be the Frobenius endomorphism of \mathbf{G} associated to a rational \mathbb{F}_q -structure. The finite group $G = \mathbf{G}^F$ of fixed points under F is called a finite reductive group.

The ordinary representation theory of G is now widely understood: geometric methods have been developed by Deligne and Lusztig [15] and then extensively studied by Lusztig, leading to a complete classification of the irreducible characters of G [30]. One of the key step in Lusztig's fundamental work has been to determine explicitly the ℓ -adic cohomology of Deligne-Lusztig varieties associated to Coxeter elements [28]. This paper is an attempt to extend this work to the modular setting. To be more precise, we will be interested in the complex $R\Gamma_c(Y(\dot{c}), \Lambda)$ representing the cohomology of the variety $Y(\dot{c})$ with coefficients in a finite extension Λ of \mathbb{Z}_ℓ , and more specifically in the action of G and F on this complex. The representation theory of ΛG is highly dependent on the prime number ℓ . In this particular geometric situation – the Coxeter case – the corresponding primes ℓ are those which divide the cyclotomic polynomial $\Phi_h(q)$ where h is the Coxeter number. For such prime numbers, the principal

^{*}Mathematical Institute, Oxford.

[†]The author is supported by the EPSRC, Project No EP/H026568/1, by Magdalen College, Oxford and partly by the ANR, Project No JC07-192339.

part $bR\Gamma_c(Y(\dot{c}), \Lambda)$ of the cohomology complex should encode many aspects of the representation theory of the principal ℓ -block b of G , much of which remains conjectural.

In the Coxeter case, the principal ℓ -block of G has a cyclic defect group. From the work of Brauer we know that the category of modules over such a block has a combinatorial description, given by the *Brauer tree*. The shape of this tree is related to the decomposition matrix of the block whereas its planar embedding determines the module category up to Morita equivalence. By a case-by-case analysis, Hiss, Lübeck and Malle have underlined in [25] a deep connection between the Brauer tree of the principal ℓ -block and the ℓ -adic cohomology groups of the Deligne-Lusztig variety $X(c)$. They have conjectured that this connection remains valid even in the cases where the shape of the tree or its planar embedding is not explicitly known (see Conjectures (HLM) and (HLM+)). Furthermore, they have suggested that the cohomology with modular coefficients should give enough information to prove the conjecture in its full generality. This is the geometric approach that we will follow throughout this paper. It relies on the fact that the complex $bR\Gamma_c(Y(\dot{c}), \Lambda)$ is perfect and thus provides many projective modules.

In order to determine the shape of the Brauer tree, we must find every indecomposable projective ΛG -module lying in the principal block and compute their characters. Such projective modules admit no direct algebraic construction via Harish-Chandra induction, because they might have a cuspidal head. Drawing inspiration from Lusztig's work [28], we show that they can be obtained by taking suitable eigenspaces of F on $bR\Gamma_c(Y(\dot{c}), \Lambda)$. However, there is a price to pay since we have to make the following assumption:

- (W) For all minimal eigenvalues λ of F , the generalized (λ) -eigenspace of F on $bH_c^*(Y(\dot{c}), \Lambda)$ is torsion-free.

We call here "minimal" the eigenvalues of F on the cohomology group in middle degree. Under this precise assumption, in Section 3 we give a general proof of Conjecture (HLM) of Hiss-Lübeck-Malle.

There is strong evidence that the previous assumption should be valid for any eigenvalue of F [18] which means that the following should hold:

- (S) The Λ -modules $bH_c^i(Y(\dot{c}), \Lambda)$ are torsion-free.

This technical assumption will be discussed in a subsequent paper. Using generalized Gelfand-Graev representations we can actually prove that it holds in a majority of cases, which justifies our approach:

Theorem A ([17]). *Assume that \mathbf{G} has no factor of type E_7 or E_8 and that p is a good prime number. Then assumption (S) holds.*

Note that the torsion-free property is really specific to the case of Deligne-Lusztig varieties associated to Coxeter elements. The cohomology of varieties associated to other elements should have a non-trivial torsion component in general (see [18, Remark 5.37]).

In this paper, we shall instead focus on all the consequences we can draw from such an assumption. Our main result is the complete determination of the complex $b\mathrm{R}\Gamma_c(Y(\dot{c}), \Lambda)$ in terms of the projective modules lying in the block. Surprisingly, our representative turns out to be exactly the complex attached to a Brauer tree in [34]:

Theorem B. *Under the assumption (S), the complex $b\mathrm{R}\Gamma_c(Y(\dot{c}), \Lambda)$ is homotopy equivalent to the Rickard complex associated to the Brauer tree of the principal ℓ -block of G , shifted by the length of c .*

From that observation we deduce several important results:

- the cohomology complex induces the derived equivalence predicted by the geometric version of Broué’s conjecture [7] (see Theorem 4.13);
- this equivalence is perverse in the sense of [12] (see Theorem 4.20);
- the planar embedding of the Brauer tree can be read out from the eigenvalues of F (see Theorem 4.15).

Together with Theorem A, this gives new results for the geometric version of Broué’s conjecture. This extends significantly the previous work of Puig [33] (for $\ell \mid q - 1$), Rouquier [38] (for $\ell \mid \phi_h(q)$ and $r = 1$) and Bonnafé-Rouquier [3] (for $\ell \mid \phi_h(q)$ and (\mathbf{G}, F) of type A_n). We also obtain new planar embedded Brauer tree for groups of type 2G_2 and F_4 with $p \neq 2, 3$ (compare with [23] and [24]).

This paper is divided into four parts: in the first section, we introduce basic notation and recall standard constructions in the modular representation theory of finite reductive groups, formulated in the language of derived categories. In Section 2 we present what we call the *Coxeter case* and collect different results (both geometric and group-theoretic) that have been obtained so far for principal ℓ -blocks and their Brauer trees in this particular case. Section 3 is devoted to the proof of the conjecture (HLM) of Hiss, Lübeck and Malle. We show that under the assumption (W), the Brauer tree can be deduced from the ℓ -adic cohomology of $X(c)$. Finally, we use assumption (S) in Section 4 to determine an explicit representative of the cohomology complex. As a byproduct, we obtain a proof of the geometric version of Broué’s conjecture (as always in the Coxeter case) as well as and the planar embedding of the Brauer tree.

1 Preliminaries

1.1 Some homological algebra

We start by recalling some standard notions of homological algebra that we will use throughout this paper.

1.1.1. Module categories and usual functors. If \mathcal{A} is an abelian category, we will denote by $C(\mathcal{A})$ the category of cochain complexes, by $K(\mathcal{A})$ the corresponding homotopy category and by $D(\mathcal{A})$ the derived category. We shall use

the superscript notation $-$, $+$ and b to denote the full subcategories of bounded above, bounded below or bounded complexes. We will always consider the case where $\mathcal{A} = A\text{-Mod}$ is the module category over any ring A or the full subcategory $A\text{-mod}$ of finitely generated modules. This is actually not a strong restriction, since any small category can be embedded into some module category [31]. Since the categories $A\text{-Mod}$ and $A\text{-mod}$ have enough projective objects, one can define the usual derived bifunctors RHom_A^\bullet and ${}^L\otimes_A$.

Let H be a finite group and ℓ be a prime number. We fix an ℓ -modular system (K, Λ, k) consisting of a finite extension K of the field of ℓ -adic numbers \mathbb{Q}_ℓ , the integral closure Λ of the ring of ℓ -adic integers in K and the residue field k of the local ring Λ . We assume moreover that the field K is big enough for H , so that it contains the e -th roots of unity, where e is the exponent of H . In that case, the algebra KH is split semi-simple.

From now on, we shall focus on the case where $A = \mathcal{O}H$, with \mathcal{O} being any ring among (K, Λ, k) . By studying the modular representation theory of H we mean studying the module categories $\mathcal{O}H\text{-mod}$ for various \mathcal{O} , and also the different connections between them. In this paper, most of the representations will arise in the cohomology of some complexes and we need to know how to pass from one coefficient ring to another. The scalar extension and ℓ -reduction have a derived counterpart: if C is any bounded complex of ΛH -modules we can form $KC = C \otimes_\Lambda K$ and $\overline{C} = kC = C \otimes_\Lambda k$. Since K is a flat Λ -module, the cohomology of the complex KC is exactly the scalar extension of the cohomology of C . However this does not apply to ℓ -reduction, but the obstruction can be related to the torsion:

Theorem 1.1 (Universal coefficient theorem). *Let C be a bounded complex of ΛH -modules. Assume that the terms of C are free over Λ . Then for all $n \geq 1$ and $i \in \mathbb{Z}$, there exists a short exact sequence of ΛH -modules*

$$0 \longrightarrow \text{H}^i(C) \otimes_\Lambda \Lambda/\ell^n \Lambda \longrightarrow \text{H}^i(C \otimes_\Lambda \Lambda/\ell^n \Lambda) \longrightarrow \text{Tor}_1^\Lambda(\text{H}^{i+1}(C), \Lambda/\ell^n \Lambda) \longrightarrow 0.$$

In particular, whenever there is no torsion in both C and $\text{H}^*(C)$ then the cohomology of \overline{C} is exactly the ℓ -reduction of the cohomology of C .

1.1.2. Perfect complexes. In the derived category, any acyclic complex is isomorphic to zero. This can be generalized to the homotopy category if we restrict ourselves to a specific class of complexes. Recall that a complex C of $\mathcal{O}H$ -modules is said to be *perfect* if it is quasi-isomorphic to a bounded complex of finitely generated projective $\mathcal{O}H$ -modules. The value of a derived bifunctor on any perfect complex is obtained from the original functor: more precisely, if C is a perfect complex and C' is any complex in $C(\mathcal{O}H\text{-Mod})$ then there exists isomorphisms in $D(\mathbb{Z}\text{-Mod})$:

$$C \otimes_{\mathcal{O}H}^L C' \simeq C \otimes_{\mathcal{O}H} C' \quad \text{and} \quad \text{RHom}_{\mathcal{O}H}^\bullet(C, C') \simeq \text{Hom}_{\mathcal{O}H}^\bullet(C, C').$$

In particular the natural functor $K^b(\mathcal{O}H\text{-proj}) \rightarrow D(\mathcal{O}H\text{-Mod})$ induces an equivalence between the homotopy category of bounded complex of finitely generated projective modules and the full subcategory of $D(\mathcal{O}H\text{-Mod})$ of perfect complexes.

As in the derived case, one can obtain precise concentration properties for complexes in $K^b(\mathcal{O}H\text{-proj})$:

Lemma 1.2. *Let $C \in K^b(\mathcal{O}H\text{-proj})$. Assume that the cohomology of C vanishes outside the degrees m, \dots, M . Then C is homotopy equivalent to a complex*

$$\cdots \rightarrow 0 \rightarrow P_{m-1} \rightarrow P_m \rightarrow \cdots \rightarrow P_M \rightarrow 0 \rightarrow \cdots$$

whose terms are finitely generated projective modules, concentrated in degrees $m-1, m, \dots, M$. Moreover, P_{m-1} can be chosen to be zero in the following cases:

- (a) \mathcal{O} is a field;
- (b) $\mathcal{O} = \Lambda$ and the group $H^m(C)$ is torsion-free.

Proof. Let us write C as $\cdots \rightarrow P_r \xrightarrow{d_r} P_{r+1} \rightarrow 0$ with $r \geq M$. By assumption, the cohomology of C vanishes in degree $r+1$. The boundary map d_r is surjective and splits since P_{r+1} is a projective module. Therefore the complex $0 \rightarrow P_{r+1} \rightarrow P_{r+1} \rightarrow 0$ is a direct summand of C , and being homotopy equivalent to zero it can be removed.

If \mathcal{O} is one of the fields K or k , then every projective $\mathcal{O}H$ -module is injective, and we can again remove successively the terms P_i for $i < m$. The case where $\mathcal{O} = \Lambda$ requires more attention. The complex C can be written as $0 \rightarrow P_r \xrightarrow{d_r} P_{r+1} \rightarrow \cdots$ with $r < m$ and the map d_r being injective. We claim that there exists a retraction of d_r if $\text{Coker } d_r$ is torsion-free. Indeed, P_r is $(H, 1)$ -injective by [13, Theorem 19.12] so that the short exact sequence $0 \rightarrow P_r \rightarrow P_{r+1} \rightarrow \text{Coker } d_r \rightarrow 0$ splits over ΛH whenever it splits over Λ . Now, if $r < m-1$ then by assumption $\text{Im } d_r = \text{Ker } d_{r+1}$ so that $\text{Coker } d_r \simeq \text{Im } d_{r+1}$ is torsion-free (it is a submodule of a free module). If $m = r-1$, we can observe that the quotient $\text{Coker } d_{m-1}/H^m(C) \simeq \text{Im } d_m$ is torsion-free, and hence $\text{Coker } d_{m-1}$ is torsion-free whenever $H^m(C)$ is. \square

1.2 Cohomology of a quasi-projective variety

In the ordinary Deligne-Lusztig theory, representations of finite reductive groups arise from the cohomology of some varieties acted on by the group. In the modular setting, we shall be interested not only in the cohomology of these varieties but also in complexes representing the cohomology in the derived category. This gives much more information, some of which has already been collected by Broué [4], [7], and more recently by Bonnafé and Rouquier [2].

1.2.1. Rouquier's construction. Let X be a quasi-projective algebraic variety defined over $\bar{\mathbb{F}}_p$. We assume that X is acted on by a finite group H and by a monoid of endomorphisms Y normalizing H . We shall always assume that

the prime number p (the defining characteristic of all the varieties involved) is different from ℓ (associated to the modular system).

Let \mathcal{O} be any ring among (K, Λ, k) . Rouquier has constructed in [38] a bounded complex $\mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O})$ of $\mathcal{O}H \rtimes Y$ -modules representing the ℓ -adic cohomology with compact support of \mathbf{X} . In other words, we have $\mathrm{H}^*(\mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O})) \simeq \mathrm{H}_c^*(\mathbf{X}, \mathcal{O})$ as $\mathcal{O}H \rtimes Y$ -modules. This construction is particularly adapted to the modular setting, since the cohomology complexes behave well with respect to scalar extension and ℓ -reduction. We have indeed in $D^b(\mathcal{O}H \rtimes Y\text{-mod})$:

$$\mathrm{R}\Gamma_c(\mathbf{X}, \Lambda) \stackrel{\mathrm{L}}{\otimes}_{\Lambda} \mathcal{O} \simeq \mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O}).$$

In particular, the universal coefficient theorem (Theorem 1.1) will hold for ℓ -adic cohomology with compact support.

Let us forget about the action of Y for the moment. By construction, the terms of $\mathrm{R}\Gamma_c(\mathbf{X}, \Lambda)$ are far from being finitely generated. Nevertheless, we can, up to homotopy equivalence, find a representative with good finiteness properties [38, Section 2.5.1]:

Theorem 1.3 (Rouquier). *Let \mathbf{X} be a quasi-projective variety acted on by a finite group H . Denote by $\mathcal{S} = \{\mathrm{Stab}_H(x) \mid x \in \mathbf{X}\}$ the set of stabilizers of points of \mathbf{X} . Then $\mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O})$ is homotopy equivalent to a complex \mathcal{C} whose terms are direct summands of finite sums of permutation modules $\mathcal{O}[H/S]$ for various $S \in \mathcal{S}$.*

Remark 1.4. Note that \mathcal{C} is not a complex of $\mathcal{O}H \rtimes Y$ -modules for it can only inherit an action of Y up to homotopy.

Corollary 1.5. *Assume that the order of the stabilizer of any point is invertible in \mathcal{O} . Then $\mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O})$ is a perfect complex of $\mathcal{O}H$ -modules.*

Recall that the ℓ -adic cohomology with compact support of any irreducible affine variety of dimension d is concentrated in degrees $d, \dots, 2d$, and this for any coefficient ring among (K, Λ, k) . Using Lemma 1.2 and Theorem 1.1 we deduce the following:

Corollary 1.6. *Let \mathbf{X} be an irreducible affine variety of dimension d . Assume that the order of the stabilizer of any point is prime to ℓ . Then $\mathrm{R}\Gamma_c(\mathbf{X}, \Lambda)$ can be represented by a complex of finitely generated projective ΛH -modules concentrated in degrees $d, \dots, 2d$. In particular, $\mathrm{H}_c^d(\mathbf{X}, \Lambda)$ is torsion-free.*

Note that the result holds for any disjoint union of irreducible affine varieties with the same dimension, and therefore for any Deligne-Lusztig variety that has been proven to be affine.

1.2.2. Generalized eigenspaces of the Frobenius. We now study the case where $Y = \langle F \rangle_{\mathrm{mon}}$ is generated by the Frobenius endomorphism attached to some rational \mathbb{F}_q -structure on \mathbf{X} . We would like to factor out the complex $\mathrm{R}\Gamma_c(\mathbf{X}, \mathcal{O})$ with respect to the eigenvalues of F . For this purpose, we shall first review some basics about endomorphisms of finitely generated Λ -modules.

Let M be a $\Lambda[T]$ -module. Denote by f the endomorphism of M induced by T . Assume that M is finitely generated over Λ . Then there exists a monic polynomial $P \in \Lambda[T]$ such that $P(f) = 0$, and we are reduced to studying the action of the finite dimensional algebra $\Lambda[T]/(P)$ on M . We may assume without loss of generality that P splits over Λ . In that case, the factorization of $P = (T - \lambda_1)^{\alpha_1} \cdots (T - \lambda_n)^{\alpha_n}$ yields a decomposition of the module KM with respect to the generalized eigenspaces of f :

$$KM = \text{Ker}(f - \lambda_1)^{\alpha_1} \oplus \cdots \oplus \text{Ker}(f - \lambda_n)^{\alpha_n}.$$

In order to obtain an modular analog of this decomposition we have to group together the eigenvalues according to their ℓ -reduction (this becomes clear if we consider the module $\overline{M} = M \otimes_{\Lambda} k$). More precisely, if we define the polynomials

$$P_{\bar{\lambda}}(T) = \prod_{\bar{\lambda}_i = \bar{\lambda}} (T - \lambda_i)^{\alpha_i}$$

then the block decomposition of the algebra $\Lambda[T]/(P)$ is given by

$$\Lambda[T]/(P) \simeq \prod_{\bar{\lambda} \in k} \Lambda[T]/(P_{\bar{\lambda}}).$$

For $\lambda \in K$, we define the *generalized* (λ) -*eigenspace* of f in M to be $M_{(\lambda)} = e_{\lambda}M$ where e_{λ} is the idempotent associated to the term $\Lambda[T]/(P_{\bar{\lambda}})$. By construction, M decomposes into

$$M = \bigoplus_{\bar{\lambda} \in k} M_{(\lambda)}.$$

Remark 1.7. This definition does not depend on P since the module $e_{\lambda}M$ depends only on the image of e_{λ} in the algebra $\Lambda[T]/\text{Ann}(f)$.

Equivalently, one could have defined the module $M_{(\lambda)}$ to be the kernel of the endomorphism $P_{\bar{\lambda}}(f)$. One can easily deduce the following result using this description:

Lemma 1.8. *Let f and g be two endomorphisms of M . If $\bar{f} - \bar{g}$ is a nilpotent endomorphism of \overline{M} , then the generalized (λ) -eigenspaces of f and g on M coincide.*

The definition of (λ) -eigenpaces can be extended to the case where \mathcal{O} is one of the field K or k by setting $(M \otimes_{\Lambda} \mathcal{O})_{(\lambda)} = M_{(\lambda)} \otimes_{\Lambda} \mathcal{O}$. The following proposition describes the relation between these modules and the usual generalized eigenspaces:

Proposition 1.9. *Let $\lambda \in \Lambda$.*

- (i) *The $K[T]$ -module $(KM)_{(\lambda)} := M_{(\lambda)} \otimes_{\Lambda} K$ is isomorphic to the direct sum of all the generalized μ -eigenspaces where μ runs over the set of eigenvalues congruent to λ modulo ℓ .*
- (ii) *The $k[T]$ -module $\overline{M}_{(\lambda)} := M_{(\lambda)} \otimes_{\Lambda} k$ is isomorphic to the generalized $\bar{\lambda}$ -eigenspace corresponding to $\bar{\lambda} \in k$.*

More generally, if \mathcal{O} is any ring among (K, Λ, k) , one can define an endofunctor $C \mapsto C_{(\lambda)}$ of the category of bounded complexes of $\Lambda[T]$ -modules finitely generated over Λ . It is an exact functor, and as such it satisfies

$$H^\bullet(C_{(\lambda)}) \simeq H^\bullet(C)_{(\lambda)}.$$

Now, in order to apply this construction to the cohomology complex $R\Gamma_c(X, \mathcal{O})$ we need some finiteness conditions. These are given by Theorem 1.3: there exists a bounded complex \mathcal{C} of finitely generated $\mathcal{O}H$ -modules, together with H -equivariant morphisms $f : C \rightarrow \mathcal{C}$ and $g : \mathcal{C} \rightarrow C$ which are mutually inverse in the category $K^b(\mathcal{O}H\text{-Mod})$. Assume that the Frobenius F commutes with the action of H so that we can define a H -equivariant morphism on \mathcal{C} by setting $\mathcal{F} = f \circ F \circ g$. The definition of \mathcal{F} depends on the choice of the homotopy equivalence, but the images of \mathcal{F} and F coincide on the cohomology of X . In particular, there exists an isomorphism of $\mathcal{O}H$ -modules

$$H^\bullet(\mathcal{C}_{(\lambda)}) \simeq H_c^\bullet(X, \mathcal{O})_{(\lambda)}$$

where the eigenspace on the right side is taken with respect to F .

Moreover, if the terms of \mathcal{C} are projective modules (for example if the action of H is free) then the generalized (λ) -eigenspaces $\mathcal{C}_{(\lambda)}$ are in turn objects of the category $C^b(\mathcal{O}H\text{-proj})$ and have, besides, the advantage of being in general much smaller than \mathcal{C} itself.

1.3 Finite reductive groups

1.3.1. Algebraic groups. We keep the basic assumptions of the introduction, with some slight modification: \mathbf{G} is a connected reductive algebraic group, together with an isogeny F , some power of which is a Frobenius endomorphism. In other words, there exists a positive integer δ such that F^δ defines a split \mathbb{F}_{q^δ} -structure on \mathbf{G} for a certain power q^δ of the characteristic p (note that q might not be an integer). We will assume that δ minimal for this property. For all F -stable algebraic subgroup \mathbf{H} of \mathbf{G} , we will denote by H the finite group of fixed points \mathbf{H}^F .

We fix a Borel subgroup \mathbf{B} containing a maximal torus \mathbf{T} of \mathbf{G} such that both \mathbf{B} and \mathbf{T} are F -stable. They define a root system Φ with basis Δ , and a set of positive (resp. negative) roots Φ^+ (resp. Φ^-). Note that the corresponding Weyl group W is endowed with an action of F , compatible with the isomorphism $W \simeq N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$. Therefore, the image by F of a root is a positive multiple of some other root, which will be denoted by $\phi^{-1}(\alpha)$, defining thus a bijection $\phi : \Phi \rightarrow \Phi$. Since \mathbf{B} is F -stable, this map preserves Δ and Φ^+ . We will use the notation $[\Delta/\phi]$ for a set of representatives of the orbits of ϕ on Δ .

1.3.2. Deligne-Lusztig varieties. Following [2, Section 11.2], we fix a set of representatives $\{w\}$ of W in $N_{\mathbf{G}}(\mathbf{T})$ and we define, for $w \in W$, the Deligne-Lusztig varieties $X(w)$ and $Y(w)$ by:

$$\begin{array}{c} Y(w) = \{g\mathbf{U} \in \mathbf{G}/\mathbf{U} \mid g^{-1}F(g) \in \mathbf{U}w\mathbf{U}\} \\ \pi_w \downarrow / \mathbf{T}^{wF} \\ X(w) = \{g\mathbf{B} \in \mathbf{G}/\mathbf{B} \mid g^{-1}F(g) \in \mathbf{B}w\mathbf{B}\} \end{array}$$

where π_w denotes the restriction to $Y(w)$ of the canonical projection $\mathbf{G}/\mathbf{U} \rightarrow \mathbf{G}/\mathbf{B}$. They are both quasi-projective varieties endowed with a left action of G by left multiplication. Furthermore, \mathbf{T}^{wF} acts on the right of $Y(w)$ and π_w is isomorphic to the corresponding quotient map, so that it induces a G -equivariant isomorphism of varieties $Y(w)/\mathbf{T}^{wF} \simeq X(w)$.

The ℓ -adic cohomology of these varieties yields the so-called Deligne-Lusztig induction. More precisely, if θ is a character of \mathbf{T}^{wF} , one can look at the θ -isotypic component of the cohomology and define the following virtual character

$$R_w(\theta) = \sum_{i \in \mathbb{Z}} (-1)^i H_c^i(Y(w), K)_\theta.$$

Note that with our definition of the variety $Y(w)$ we have chosen to work with characters of \mathbf{T}^{wF} instead of characters of T_w for some torus \mathbf{T}_w of type w . Our aim is to understand a far-reaching generalization of this character in the case where w is a Coxeter element. It will be represented by a well-identified direct summand of the complex $R\Gamma_c(Y(w), \Lambda)$.

2 The principal ℓ -block in the Coxeter case

In this preliminary section we introduce the main object of our study: the principal ℓ -block b of G where the order of q modulo ℓ is assumed to be the Coxeter number h . We will refer to this case as the *Coxeter case*. This is in some sense the maximal interesting case in the modular representation theory of G , since h is also the largest integer d such that the cyclotomic polynomial $\Phi_d(q)$ divides the order of G .

The results in [8] express the irreducible characters of this block in terms of irreducible components of Deligne-Lusztig characters $R_c(\theta)$ induced from a Coxeter torus \mathbf{T}^{cF} . In this particular case, an explicit decomposition of these virtual characters is given by Lusztig's work on the cohomology of the Deligne-Lusztig variety $X(c)$ [28]. The characters of the block fall into two families:

- the characters $R_c(\theta)$ for θ a non-trivial ℓ -character of the torus. These are irreducible characters (up to a sign);
- the unipotent characters, coming from the cohomology of $X(c)$.

Here the defect group of the principal ℓ -block turns out to be a cyclic group and the distinction "non-unipotent/unipotent" translates into "exceptional/non-exceptional" in the theory of blocks with cyclic defect groups. The connection is actually much deeper: Hiss, Lübeck and Malle have observed in [25] that the cohomology of the Deligne-Lusztig variety $X(c)$ should not only give the characters of the block, but also its Brauer tree.

We shall first review the geometric objects and the fundamental results involved in their description before recalling their conjecture.

2.1 The Coxeter case

For the sake of simplicity, we first assume that \mathbf{G} has no twisted components of type ${}^2\mathbf{B}_2$, ${}^2\mathbf{F}_4$ or ${}^2\mathbf{G}_2$. The case of the Ree and Suzuki groups will be treated independently.

2.1.1. Coxeter elements. Let $V = X^\vee(\mathbf{T}) \otimes_{\mathbb{Z}} \mathbb{C}$ be the m -dimensional vector space generated by the cocharacters of \mathbf{T} . The Weyl group W can be seen as a subgroup of the linear automorphisms of this vector space; moreover, the linear map $\sigma = q^{-1}F$ has finite order δ and normalizes W (with the previous assumptions on (\mathbf{G}, F) , this is exactly the linear continuation of ϕ^\vee). By [40], there exist eigenvectors (f_1, \dots, f_m) of σ in $S(V)$ of degrees (d_1, \dots, d_m) with associated eigenvalues $(\varepsilon_1, \dots, \varepsilon_m)$ such that $S(V)^W$ is isomorphic to the polynomial algebra $\mathbb{C}[f_1, \dots, f_m]$. Up to permutation, the pairs (d_j, ε_j) are uniquely determined by σ . The order of G is then given by the following formula

$$|G| = q^N \prod_{j=1}^m (q^{d_j} - \varepsilon_j^{-1}) = q^N \prod_d \Phi_d(q)^{a(d)}$$

where $a(d)$ is the number of j such that $\varepsilon_j = \exp(2i\pi d_j/d)$ [6]. The largest integer such that $a(d)$ is non-zero will be denoted by h and referred as the *Coxeter number* of the pair (W, F) .

From now on, we assume that \mathbf{G} is semi-simple and W is irreducible. In this case, the \mathbb{C} -vector space $V = X^\vee(\mathbf{T}) \otimes_{\mathbb{Z}} \mathbb{C}$ can be identified with the reflection representation of W and the pairs (d_j, ε_j) have been explicitly computed in [10]. From these values one easily deduces the Coxeter numbers for each type

type	A_n	B_n	D_n	E_6	E_7	E_8	F_4	G_2	${}^2A_{2n}$	${}^2A_{2n+1}$	2D_n	3D_4	2E_6
h	$n+1$	$2n$	$2n-2$	12	18	30	12	6	$4n+2$	$4n+2$	$2n$	12	18

and one can check that $a(h)$ is always equal to 1.

The twisted counterpart of the usual notion of Coxeter elements for Weyl groups has been introduced in [39, Section 7]:

Definition 2.1. A Coxeter element of the pair (W, F) is a product $c = s_{\beta_1} \cdots s_{\beta_r}$ where $\{\beta_1, \dots, \beta_r\} = [\Delta/\phi]$ is any set of representatives of the orbits of the simple roots under the action of ϕ .

Such an element c has the same properties as usual Coxeter elements, provided that the conjugation under W is replaced by the F -conjugation. These properties become clear if we consider $c\sigma = q^{-1}cF \in \text{GL}(V)$ instead of c :

Theorem 2.2 (Springer). *Let c be a Coxeter element of (W, F) with W irreducible.*

- (i) $c\sigma$ is an h -regular element.

- (ii) Let c' be any element of W . If $c'\sigma$ has an eigenvalue of order h , it is h -regular and conjugated to $c\sigma$. In particular, the set of Coxeter elements is contained in a single F -conjugacy class.
- (iii) The eigenvalues of $c\sigma$ are $\varepsilon_j^{-1} \exp(2i\pi(d_j - 1)/h)$. Moreover, the eigenvalues of order h occur with multiplicity 1.
- (iv) The centralizer $C_W(c\sigma)$ is a cyclic group generated by $(c\sigma)^\delta = cF(c) \cdots F^{\delta-1}(c)$.

As a byproduct δ divides h . The quotient will be denoted by $h_0 = h/\delta$ in line with Lusztig's definition [28, Section 1.13]. For the sake of completeness, we give the different values of this number:

type	A_n	B_n	D_n	E_6	E_7	E_8	F_4	G_2	${}^2A_{2n}$	${}^2A_{2n+1}$	2D_n	3D_4	2E_6
h_0	$n+1$	$2n$	$2n-2$	12	18	30	12	6	$2n+1$	$2n+1$	n	4	9

Remark 2.3. One could have defined the Coxeter number h to be the maximal order of the eigenvalues of the elements of $W\sigma$. This is actually the original definition given by Springer [39], but it coincides with the previous one by the generic Sylow theorems [6].

2.1.2. Coxeter tori. Let c be a Coxeter element of (W, F) . We will be interested in rational tori \mathbf{T}_c of type c , which are usually called *Coxeter tori*. Recall that (\mathbf{T}_c, F) is isomorphic to (\mathbf{T}, cF) and that the order of the associated finite groups is given by

$$|T_c| = |\mathbf{T}^{cF}| = \det(qc\sigma - 1 | X^\vee(\mathbf{T}) \otimes_{\mathbb{Z}} \mathbb{C}).$$

Since $a(h) = 1$, the torus \mathbf{T}_c contains a unique Φ_h -Sylow subgroup \mathbf{S}_h of \mathbf{G} , as defined in [6]. The following proposition summarizes the different properties we will use later on. They are easily obtained by rephrasing Theorem 2.2 in the framework of [6] (see [18] for more details).

Proposition 2.4. *Let ℓ be a prime number different from p . We assume that ℓ divides $\Phi_h(q)$ but does not divide $|W^F|$. Then*

- (i) The set T_ℓ of ℓ -elements in \mathbf{S}_h is a cyclic ℓ -Sylow subgroup of G .
- (ii) $C_{\mathbf{G}}(T_\ell) = \mathbf{T}_c$ and $N_G(T_\ell)/C_G(T_\ell) \simeq (N_{\mathbf{G}}(\mathbf{T}_c)/\mathbf{T}_c)^F \simeq C_W(c\sigma)$.
- (iii) Any non-trivial ℓ -character θ of T_c (or \mathbf{T}^{cF}) is in general position. In other words, the centralizer $C_{C_W(c\sigma)}(\theta)$ is trivial.

2.1.3. The case of Ree and Suzuki groups. The previous proposition holds also when \mathbf{G} has type 2B_2 , 2F_4 or 2G_2 . The notion of Coxeter elements has indeed a natural generalization to these groups, taking into account that $\sigma = q^{-1}F$ does no longer stabilize $X^\vee(\mathbf{T})$ but only $X^\vee(\mathbf{T}) \otimes_{\mathbb{Z}} \mathbb{Z}[\sqrt{p}^{-1}]$ for $p = 2$ or 3 depending on the type of (\mathbf{G}, F) . The previous table can then be completed with the orders of $c\sigma$:

type	2B_2	2F_4	2G_2
h	8	24	12

Over the ring $\mathbb{Z}[\sqrt{p}]$, the cyclotomic polynomial Φ_h is no longer irreducible. For each type, the finite group T_c itself is a cyclic group, and its order is given by the evaluation at q of an irreducible factor of Φ_h . When q is positive, it is given by

type	${}^2\text{B}_2$	${}^2\text{F}_4$	${}^2\text{G}_2$
$ T_c $	$1 - q\sqrt{2} + q^2$	$1 - q\sqrt{2} + q^2 - q^3\sqrt{2} + q^4$	$1 - q\sqrt{3} + q^2$

Using a case-by-case analysis, one can also check that when ℓ divides one of these numbers without dividing the order of the corresponding Weyl group, the set of all ℓ -elements in T_c is again a Sylow ℓ -subgroup and it satisfies the assertions (ii) and (iii) in Proposition 2.4.

2.2 Characters in the principal block

In order to use the results stated in the previous section, we will, until further notice, assume \mathbf{G} to be a semi-simple group and W to be irreducible (we shall say that \mathbf{G} is *quasi-simple*). We fix a prime number ℓ not dividing the order of W^F and satisfying one of the two following assumptions, depending on the type of (\mathbf{G}, F) :

- "non-twisted" cases: ℓ divides $\Phi_h(q)$;
- "twisted" cases: ℓ divides the order of T_c for some Coxeter element c .

As in Section 1.1, the modular framework will be given by an ℓ -modular system (K, Λ, k) , which we require to be big enough for G . Note that the conditions on the prime number ℓ ensure that the class of q in k^\times is a primitive h -th root of unity. Indeed, q^h is congruent to 1 modulo ℓ and for any proper divisor m of h the class of q^m cannot be 1 otherwise $1 + q + \dots + q^{h-1}$ would be congruent to both 0 and $h/m(1 + q + \dots + q^{m-1})$. Taking m to be minimal would force ℓ to divide h/m . The principal block of ΛG for this particular class of primes will be at the center of our study.

We choose a Coxeter element c of (W, F) together with a maximal rational torus \mathbf{T}_c of type c (a Coxeter torus). This torus is the centralizer of a so-called Φ_h -torus \mathbf{S}_h (with $\mathbf{S}_h = \mathbf{T}_c$ for the Ree and Suzuki groups). As such, in the terminology of [8], it is a h -split Levi subgroup, and the h -cuspidal pair $(\mathbf{T}_c, 1)$ corresponds to the principal ℓ -block b of G . More precisely, it follows from [8, Theorem 5.24] that the characters in b are exactly the irreducible components of the Deligne-Lusztig characters $R_c(\theta)$ where θ runs over the set of ℓ -characters of \mathbf{T}^{cF} . Two families of characters occur in this description; using Lusztig's results on the variety $X(c)$, we now proceed with their parametrization.

2.2.1. The non-unipotent characters in the block. Proposition 2.4 says that any non-trivial ℓ -character of \mathbf{T}^{cF} is in general position. Consequently, the corresponding induced character is an actual irreducible character of G (up to a sign). It is worth pointing out that this result is a consequence of a deep property of the cohomology of the Deligne-Lusztig variety $Y(c)$ [15, Corollary 9.9]: for θ

a non-trivial ℓ -character of \mathbf{T}^{cF} , the θ -isotypic component $H_c^i(Y(\dot{c}), K)_\theta$ of the cohomology in degree i is non-zero for $i = \ell(c) = r$ only.

The Frobenius endomorphism F^δ acts on \mathbf{T}^{cF} . Moreover, since F^δ acts trivially on W , the representative \dot{c} of c can be chosen to be F^δ -stable. In that case, $\mathbf{T}^{cF} \rtimes \langle F^\delta \rangle_{\text{mon}}$ acts on the variety $Y(\dot{c})$, leading to an linear action on the cohomology which satisfies

$$F^\delta(H_c^r(Y(\dot{c}), K)_\theta) = H_c^r(Y(\dot{c}), K)_{F^\delta(\theta)} = H_c^r(Y(\dot{c}), K)_{v \cdot \theta}$$

where $v = cF(c) \cdots F^{\delta-1}(c)$. Note that v is a generator (of order $h_0 = h/\delta$) of the cyclic group $C_W(c\sigma)$. Since the actions of F^δ and G commute, the isotypic components associated to ℓ -characters lying in the same orbit under $C_W(c\sigma)$ are isomorphic (via some power of F^δ).

Notation 2.5. For θ a non-trivial ℓ -character of \mathbf{T}^{cF} , the θ -isotypic part of the bimodule $H_c^r(Y(\dot{c}), K)$ is a simple KG -module that will be denoted by Y_θ . The isomorphic class of this module depends only on the orbit of θ under $C_W(c\sigma)$; the corresponding character will be denoted by χ_θ .

Proposition 2.6. *When θ runs over $[\text{Irr}_\ell(\mathbf{T}^{cF})/C_W(c\sigma)]$ and is assumed to be non-trivial, the characters χ_θ are distinct irreducible cuspidal characters. Furthermore, they all have the same restriction to the set of ℓ -regular elements of G .*

Proof. The F -conjugacy class of a Coxeter element is cuspidal. In other words, c is not contained in any proper F -stable parabolic subgroup of W . An analogue property holds for the torus \mathbf{T}_c , and we deduce from [29, Corollary 2.19] that the characters χ_θ are cuspidal.

Let θ and θ' be two non-trivial ℓ -characters of \mathbf{T}^{cF} . The Mackey formula [16, Theorem 11.13], written for the torus (\mathbf{T}, cF) instead of (\mathbf{T}_c, F) yields

$$\langle \chi_\theta; \chi_{\theta'} \rangle_G = \sum_{w \in W^{cF}} \langle \theta; w \cdot \theta' \rangle_{\mathbf{T}^{cF}}.$$

Since both θ and θ' are in general position, we deduce that this sum is non-zero if and only if θ and θ' lie in the same orbit under $W^{cF} = C_W(c\sigma)$.

Finally, the value on ℓ -regular elements of any ℓ -character is trivial. By the character formula [16, Proposition 12.2], it follows that the restriction of χ_θ to the set of regular elements does not depend on θ . \square

2.2.2. The unipotent characters in the block. These are the irreducible components of the virtual character $R_c(1)$ attached to the Deligne-Lusztig variety $X(c)$. We review three main theorems in [28] giving the fundamental properties of the cohomology of this variety, with a view to establish a simple parametrization of the unipotent characters of the principal ℓ -block:

Theorem 2.7 (Lusztig). *The Frobenius F^δ acts semi-simply on $\bigoplus_i H_c^i(X(c), K)$ and its eigenspaces are mutually non-isomorphic simple KG -modules.*

Moreover, Lusztig has shown that any eigenvalue of F^δ on the cohomology can be written $\zeta q^{m\delta/2}$, where m is a non-negative integer and ζ is a root of unity. These eigenvalues are explicitly determined in [28, Table 7.3], and one can

check the following numerical property by a case-by-case analysis:

Fact 2.8. The ℓ -reduction $\Lambda \rightarrow k$ induces a bijection between the eigenvalues of F^δ and the h_0 -th roots of unity in k .

Besides, the assumption on the prime number ℓ forces the ℓ -reduction of q^δ to have order h_0 in k^\times , thus giving a canonical generator of the group of h_0 -th roots of unity. We now choose a particular square root of q^δ in K so that any eigenvalue of F^δ corresponds, via ℓ -reduction, to a unique power of q^δ . From this observation one can introduce the following notation:

Notation 2.9. For all $j = 0, \dots, h_0 - 1$, we denote by λ_j the unique eigenvalue of F^δ on $\bigoplus_i H_c^i(X(c), K)$ which is congruent to $q^{j\delta}$ modulo ℓ . Since a particular square root of q^δ has been chosen, there exists a unique root of unity ζ_j (in K) such that $\lambda_j = \zeta_j q^{m\delta/2}$ for some integer m . The eigenspace of F^δ associated to λ_j will be denoted by Y_j and its character by χ_j .

With this notation, the set $\{\chi_j \mid j = 0, \dots, h_0 - 1\}$ corresponds to the set of unipotent characters in the principal ℓ -block. Note that it is important to keep track of the root of unity $\zeta_j \in K$ occurring in the eigenvalue λ_j . It gives indeed the Harish-Chandra series in which the corresponding eigenspace Y_j falls:

Theorem 2.10 (Lusztig). *The simple KG -modules Y_i and Y_j lie in the same Harish-Chandra series if and only if $\zeta_i = \zeta_j$.*

In other words, the set $\{Y_j \mid \zeta_j = \zeta\}$ represents the (possibly empty) intersection of the principal ℓ -block with an Harish-Chandra series. The last result of this section tells us how these modules are precisely arranged in the cohomology of the Deligne-Lusztig variety $X(c)$:

Theorem 2.11. *Let $\zeta \in K$ be a root of unity. Assume that the set of integers j such that $\zeta_j = \zeta$ is non-empty. Then it is a set of consecutive integers $[[m_\zeta; M_\zeta]]$ and the corresponding eigenspaces of F^δ in the cohomology of $X(c)$ are arranged as follows:*

$$\begin{array}{c|c|c|c} H_c^r(X(c), K) & H_c^{r+1}(X(c), K) & \dots & H_c^{r+M_\zeta-m_\zeta}(X(c), K) \\ \hline Y_{m_\zeta} & Y_{m_\zeta+1} & \dots & Y_{M_\zeta} \end{array}$$

where $r = \ell(w)$. Moreover, Y_j is a cuspidal KG -module if and only if $j = m_{\zeta_j} = M_{\zeta_j}$.

More generally, the number $M_{\zeta_j} - m_{\zeta_j}$ measures the *depth* of Y_j as defined in [28], that is the obstruction of Y_j from being cuspidal.

2.3 Brauer tree of the principal block

The irreducible characters in the principal ℓ -block split into two distinct families: $\{\chi_\theta \mid \theta \in [\text{Irr}_\ell \mathbf{T}^{cF}/C_W(c\sigma)] \text{ and } \theta \neq 1\}$ and $\{\chi_i \mid i = 0, \dots, h_0 - 1\}$. The charac-

ters in the first set have the same restriction to $G_{\ell'}$ and as such, play the same role in the modular representation theory of G . If we define the *exceptional character* χ_{exc} to be the sum of these elements, then the structure of the block can be expressed in terms of the elements of the set $\mathcal{V} = \{\chi_0, \chi_1, \dots, \chi_{h_0-1}\} \cup \{\chi_{\text{exc}}\}$. More precisely, from the theory of blocks with cyclic defect groups one knows that the character of any indecomposable projective ΛG -module can be written as $[P] = \chi + \chi'$ where χ and χ' are two distinct elements of \mathcal{V} . Following Brauer, one can define a graph Γ encoding the structure of the block:

- the vertices of the graph are labeled by $\mathcal{V} = \{\chi_0, \chi_1, \dots, \chi_{h_0-1}\} \cup \{\chi_{\text{exc}}\}$. The vertex associated to χ_{exc} is called the *exceptional vertex* or the *exceptional node*. By extension, the other vertices are said to be *non-exceptional*;
- two vertices χ and χ' are connected by an edge if and only if $\chi + \chi'$ is the character of an indecomposable projective ΛG -module.

This graph Γ is actually a tree, which we refer as the *Brauer tree* of the block. The edges of the tree can be labeled either by the indecomposable projective ΛG -modules in the block or by the simple kG -modules in the block (for the indecomposable projective modules are exactly the projective covers of the simple modules).

Example 2.12. Let D be a cyclic ℓ -group and E an ℓ' -subgroup of $\text{Aut}(D)$. The Brauer tree of the unique ℓ -block of the group $H = D \rtimes E$ is a star. Indeed, the simple kH -modules are obtained by inflation of the simple kE -modules (with a trivial action of D). Since E is an ℓ' -group, such a module lifts uniquely to a ΛE -lattice \tilde{S} with projective cover $\text{Ind}_E^H \tilde{S}$. The character of the latter decomposes into

$$[\text{Ind}_E^H \tilde{S}] = [\tilde{S}] + \chi_{\text{exc}}$$

where χ_{exc} is the sum of the irreducible ordinary characters of H with trivial restriction to D (meaning that the restriction to D comes from a trivial representation of D). By construction, the Brauer tree has the following shape:




Figure 1: Brauer tree of $D \rtimes E$


Rickard [34], [35] and Linckelmann [26] have shown that one can "unfold" the Brauer tree of the principal ℓ -block of G in order to obtain this star, and that

this "unfolding" is the reflect of a splendid equivalence between the principal ℓ -blocks of G and $\mathbf{T}^{cF} \rtimes C_W(c\sigma)$. One of the main goals of this article is to show that this operation can be performed by means of the complex $R\Gamma_c(Y(\dot{c}), \Lambda)$ representing the cohomology of the Deligne-Lusztig variety $Y(\dot{c})$, thus giving a geometric explanation of Rickard-Linckelmann's result for finite reductive groups.

The conjecture of Hiss, Lübeck and Malle stated in [25] comes within this geometric framework. It predicts the shape of the Brauer tree of the principal ℓ -block of G in terms of the parametrization of the unipotent characters given previously:

Conjecture HLM (Hiss-Lübeck-Malle). *Let Γ^\bullet denote the graph obtained from the Brauer tree of the principal ℓ -block by removing the exceptional node and all edges incident to it. Then the following holds:*

- (i) *The connected components of Γ^\bullet are labeled by the Harish-Chandra series, hence by the roots of unity ζ_j 's.*
- (ii) *The connected component corresponding to a root ζ is:*



- (iii) *The vertices labeled by χ_{m_ζ} are the only nodes connected to the exceptional node.*

The validity of this conjecture has been checked in all cases where the Brauer tree was known, that is for all quasi-simple groups except the groups of type E_7 and E_8 . A general proof will be exposed in the next section, but under a precise assumption on the torsion in the cohomology of $X(c)$ (see the introduction or section 3.2 for more details).

As defined above, the Brauer tree of a block encodes only its decomposition matrix. However, as one can notice in the previous example, once the nodes have been labeled, there are different ways to draw the star. The planar embedding of the Brauer tree is actually a fundamental datum of the block: consider a vertex in the tree, together with all the edges incident to it, labeled by the simple kG -modules S_1, \dots, S_m . Then from the general theory there exist uniserial modules which have exactly the S_i 's as composition factors. The unique composition series of any of these determines an ordering of S_1, \dots, S_m which, up to cyclic permutation, does not depend on the module [21]. In the *planar embedded* Brauer tree, the edges incident to that vertex are labeled anti-clockwise according to this ordering. For a description of this ordering in terms of extentions, one can readily check that two edges labeled by the simple modules S and S' are (anti-clockwise) adjacent if and only if $\text{Ext}_{kG}^1(S', S) \neq 0$.

The Brauer tree, together with its planar embedding, fully encodes the representation theory of the block since it determines the block algebra up to Morita equivalence.

Example 2.13. We return to the previous example in the special case where E is a cyclic group, generated by an element $x \in \text{Aut}(D)$ of order m prime to ℓ . Recall that D is also assumed to be cyclic; consequently, there exists an integer n prime to $|D| = \ell^\alpha$, uniquely determined in $[[0; \ell^\alpha - 1]]$ such that x acts on any element $y \in D$ by raising y to the power of n . Since x has order m , the ℓ -reduction of n has order $d|m$. Besides, $\ell > m$ so that $v_\ell(n^d - 1) = v_\ell(n^m - 1) \geq \ell^\alpha$, which forces $d = m$.

By Hensel's lemma, there exists a (unique) primitive m -th root of unity $\zeta \in \mathbb{Z}_\ell^\times$ such that $\zeta \equiv n \pmod{\ell \mathbb{Z}_\ell}$. If we number the exceptional characters $\eta_j : H \rightarrow \mathbb{Z}_\ell^\times$ in such a way that $\eta_j(x) = \zeta^j$, then the planar embedded Brauer tree is given by




Figure 2: Planar embedded Brauer tree of $D \rtimes E$

In other words, if we denote by k_j the simple (one-dimensional) kH -module attached to the ordinary character η_j , then $\text{Ext}_{kH}^1(k_i, k_j)$ is non-trivial if and only if $i \equiv j + 1 \pmod{m}$.

The previous example has been thoughtfully chosen, since it gives also the planar embedded Brauer tree of the principal ℓ -block of $H = \mathbf{T}^{cF} \rtimes C_W(c\sigma)$. In this case, one should notice that the generator $x = (cF(c) \cdots F^{\delta-1}(c))^{-1}$ of $C_W(cF)$ acts on \mathbf{T}^{cF} via the Frobenius F^δ , thus raising any element to the power of q^δ . On the other side, the non-exceptional characters in the principal ℓ -block of G are labeled by the eigenvalues of F^δ on the cohomology of $X(c)$. Since the two actions of F^δ are compatible and give the same parametrization of the characters, it seems reasonable to complete Conjecture (HLM) with

Conjecture HLM+. *The ℓ -reduction of q^δ gives a canonical generator of the group of h_0 -th roots of unity and the corresponding order induces the cyclic ordering around the exceptional node of the Brauer tree of the principal ℓ -block of G .*

Such a result is of course only interesting for trees in which different planar embeddings are possible. This is the case for groups of type F_4 , 2F_4 , E_7 , E_8 and 2G_2 only. For the latter, according to Conjectures (HLM) and (HLM+) the Brauer tree should be given by Figure 3, where $i = \xi^3$ and ξ is the unique 12th

root of unity in Λ^\times congruent to q^5 modulo ℓ . The ordering becomes clear if we choose to label the non-exceptional vertices by the integers $q^{j\delta}$ congruent to the eigenvalues of F^δ .




Figure 3: Brauer tree for a group of type 2G_2

We will give in the last section a proof of this conjecture under the assumption (S) (see the introduction).

3 On the Hiss-Lübeck-Malle conjecture

The aim of this section is to present a general proof of Conjecture (HLM) under a precise assumption on the torsion in the cohomology. We follow here the geometric approach suggested in [25]. The geometry of the Deligne-Lusztig varieties $X(c)$ and $Y(\dot{c})$, and especially their cohomology with coefficients in Λ , should contain the information needed to understand the structure of the principal ℓ -block of G . The fundamental work of Lusztig on these varieties [28] will be the starting point for our proof.

The first part of the proof deals with the non-cuspidal kG -modules and their projective cover. We show that the contribution of these modules to the Brauer tree of the principal ℓ -block consists of the lines represented in the second assertion of the conjecture. The final part of the proof is obtained by showing that the edges labeled by the cuspidal modules are exactly the edges incident to the exceptional node. This is where the assumption (W) comes in, since it allows us, with the help of the tools introduced in Section 1.2, to single out projective modules with character $\chi_{\text{exc}} + \chi_{m_\zeta}$ in the cohomology of $Y(\dot{c})$, thus giving the missing edges.

3.1 Non-cuspidal kG -modules

Let $I \subset \Delta$ be a ϕ -stable subset of simple roots. The corresponding standard parabolic subgroup \mathbf{P}_I and Levi complement \mathbf{L}_I are both F -stable. Recall that

the Harish-Chandra induction and restriction functors are defined over any coefficient ring \mathcal{O} among (K, Λ, k) by

$$\begin{aligned} \mathbf{R}_{L_I}^G : \mathcal{O}_{L_I}\text{-mod} &\longrightarrow \mathcal{O}G\text{-mod} \\ N &\longmapsto \mathcal{O}[G/U_I] \otimes_{\mathcal{O}_{L_I}} N \end{aligned}$$

and

$$\begin{aligned} {}^*\mathbf{R}_{L_I}^G : \mathcal{O}G\text{-mod} &\longrightarrow \mathcal{O}_{L_I}\text{-mod} \\ M &\longmapsto M^{U_I}. \end{aligned}$$

We give the basic properties of these functors that we shall use in this section (see for example [19, Section 3.A]):

- $\mathbf{R}_{L_I}^G$ and ${}^*\mathbf{R}_{L_I}^G$ are exact functors. They stabilize the categories of finitely generated projective modules and finitely generated \mathcal{O} -free modules.
- $\mathbf{R}_{L_I}^G$ and ${}^*\mathbf{R}_{L_I}^G$ are compatible with scalar extension $-\otimes_{\Lambda} K$ and ℓ -reduction $-\otimes_{\Lambda} k$.

In particular, the induced morphisms between the Grothendieck groups are compatible with the decomposition maps. More precisely, if we denote by dec_G (resp. dec_{L_I}) the decomposition map between $K_0(KG\text{-mod})$ and $K_0(kG\text{-mod})$ (resp. between $K_0(KL_I\text{-mod})$ and $K_0(kL_I\text{-mod})$), then

$$\text{dec}_G \circ \mathbf{R}_{L_I}^G = \mathbf{R}_{L_I}^G \circ \text{dec}_{L_I} \quad \text{and} \quad \text{dec}_{L_I} \circ {}^*\mathbf{R}_{L_I}^G = {}^*\mathbf{R}_{L_I}^G \circ \text{dec}_G.$$

We shall first study the projective covers of the non-cuspidal simple kG -modules using their Harish-Chandra restriction. This method relies on the fact that the restriction of the cohomology of $X(c)$ can be expressed in terms of Coxeter varieties associated to "smaller" groups, for which the module categories over k are semi-simple.

3.1.1. Restriction of a kG -module. From now on, the Deligne-Lusztig variety associated to the Coxeter element c will be simply denoted by X . If I is any ϕ -stable subset of Δ , one obtains a Coxeter element c_I of (W_I, F) by removing from c the reflexions associated to the simple roots which are not in I . Written with the Borel subgroup $\mathbf{B}_I = \mathbf{B} \cap \mathbf{L}_I$ of \mathbf{L}_I , the Deligne-Lusztig variety X_I associated to c_I is by definition

$$X_I = X_{\mathbf{L}_I}(c_I) = \{g\mathbf{B}_I \in \mathbf{L}_I/\mathbf{B}_I \mid g^{-1}F(g) \in \mathbf{B}_I c_I \mathbf{B}_I\}.$$

When I is a maximal proper subset of Δ , the cohomology groups of X and X_I are closely related (see [28, Corollary 2.10]):

Proposition 3.1. *Assume that I is a maximal proper ϕ -stable subset of Δ . Then there exists an isomorphism of KL_I -modules, compatible with the action of F^δ*

$${}^*\mathbf{R}_{L_I}^G(\mathbf{H}_c^i(X, K)) \simeq \mathbf{H}_c^{i-1}(X_I, K) \oplus \mathbf{H}_c^{i-2}(X_I, K)(-1)$$

where the symbol (-1) indicates a Tate twist.

By successive applications of this proposition, one can extend the previous isomorphism to the case where I is not assumed to be maximal:

$${}^*R_{L_I}^G(\mathrm{H}_c^i(\mathrm{X}, K)) \simeq \bigoplus_{j \geq 0} \binom{r - r_I}{j} \mathrm{H}_c^{i - r + r_I - j}(\mathrm{X}_I, K)(-j)$$

with $r_I = |I/\phi|$ the number of ϕ -orbits in I . In particular, any eigenvalue of F^δ on the cohomology of X_I is also an eigenvalue of F^δ as an endomorphism of the cohomology of X . In order to rewrite the previous proposition in terms of eigenspaces, we introduce the following notation, valid for any ϕ -stable subset I of Δ :

Notation 3.2. Let $j \in \llbracket 0; h_0 - 1 \rrbracket$ and $\lambda_j = \zeta_j q^{m\delta/2}$ the corresponding eigenvalue of F^δ on $\mathrm{H}_c^\bullet(\mathrm{X}, K)$. We will denote by Y_j^I the λ_j -eigenspace of F^δ on $\mathrm{H}_c^\bullet(\mathrm{X}_I, K)$. It is a KL_I -module (which can be zero) and we will denote by χ_j^I its character.

Remark 3.3. As soon as the order of F on W_I is also equal to δ , Theorem 2.7 applies to the variety X_I , even if W_I is not irreducible or L_I is not semi-simple. The non-zero Y_j^I , which are exactly the eigenspaces of F^δ on $\mathrm{H}_c^\bullet(\mathrm{X}_I, K)$ are then mutually non-isomorphic simple KL_I -modules. The only non-trivial cases where the order of F on W_I is strictly lower than δ arise when W_I is split of type A_n . But in that special case, the eigenvalues of a Frobenius endomorphism are $1, q, \dots, q^n$ so that there cannot be any difference between the eigenspaces of F and F^δ . Consequently, the non-zero modules Y_j^I are eigenspaces of F and as such, are still simple and mutually non-isomorphic.

Using this notation, the previous proposition can be rephrased in terms of restriction of simple modules. Indeed, when I is assumed to be maximal, the restriction of a simple unipotent KG -module of the principal ℓ -block is given by

$${}^*R_{L_I}^G(Y_j) \simeq Y_j^I \oplus Y_{j-1}^I. \quad (3.4)$$

3.1.2. Finding the non-cuspidal simple kG -modules. Every simple kG -module M has a projective cover, which lifts uniquely (up to isomorphism) to an indecomposable projective ΛG -module that we will denote by P_M . The following lemma gives the character of this module in the case where M is non-cuspidal:

Lemma 3.5. *Let M be a simple kG -module in the principal ℓ -block. Assume that M is non-cuspidal. Then there exists a unique integer $j \in \llbracket 0; h_0 - 1 \rrbracket$ with $j > m_{\zeta_j}$ such that*

$$[P_M] = [Y_j] + [Y_{j-1}] = \chi_j + \chi_{j-1}.$$

*Moreover, if I is any maximal proper ϕ -stable subset of Δ , then the restriction ${}^*R_{L_I}^G(M)$ lifts uniquely (up to isomorphism) to a ΛL_I -lattice with character χ_j^I .*

Proof. Recall that ${}^*R_{L_I}^G$ is compatible with ℓ -reduction, so that the composition factors of any ℓ -reduction of a cuspidal KG -module are cuspidal. By Proposition 2.6, χ_{exc} is a cuspidal character and as such cannot be a component of the character of P_M . From the general theory of blocks with cyclic defect groups (see

Section 2.3) we deduce that there exist two distinct integers $i, j \in \llbracket 0; h_0 - 1 \rrbracket$ such that

$$[P_M] = [Y_j] + [Y_i].$$

In other words, M is a composition factors of the ℓ -reductions of both Y_j and Y_i .

Let I be a maximal proper ϕ -stable subset of Δ , such that the restriction ${}^*R_{L_I}^G(M)$ is non-zero (such a subset always exists since M is non-cuspidal). By the various properties of the restriction functors (listed at the beginning of the section), the composition factors of ${}^*R_{L_I}^G(M)$ occur as composition factors of the ℓ -reductions of both ${}^*R_{L_I}^G(Y_j)$ and ${}^*R_{L_I}^G(Y_i)$. Besides, by Formula 3.4, these modules decomposes into

$${}^*R_{L_I}^G(Y_j) \simeq Y_j^I \oplus Y_{j-1}^I$$

and

$${}^*R_{L_I}^G(Y_i) \simeq Y_i^I \oplus Y_{i-1}^I.$$

Since L_I is a proper Levi subgroup of G , the associated finite group has order prime to ℓ (otherwise L_I would contain a Coxeter torus, which is impossible by the results in Section 2.1). Therefore, the KL_I -modules remain simple after ℓ -reduction so that the modules ${}^*R_{L_I}^G(Y_i)$ and ${}^*R_{L_I}^G(Y_j)$ must have a common irreducible component. By Remark 3.3, this forces $|i - j| = 1$, $\zeta_i = \zeta_j$ and ${}^*R_{L_I}^G(M)$ to be a simple kL_I -module, isomorphic to any ℓ -reduction of $Y_{\min(i,j)}^I$. \square

Notation 3.6. Up to isomorphism, for any j there exists at most one module M satisfying the properties of the previous lemma. Such a module will be denoted by S_j . If it does not exist or if it is cuspidal, we will set $S_j = \{0\}$ by convention.

With this notation, Lemma 3.5 asserts that the non-cuspidal composition factors of any ℓ -reduction of Y_j are isomorphic to S_j or S_{j+1} . This gives the non-cuspidal part of the Brauer tree. It remains to determine whether S_j is always non-zero:

Lemma 3.7. *Let $j \in \llbracket 0; h_0 - 1 \rrbracket$ and assume that $j > m_{\zeta_j}$. Then S_j is non-zero and occurs with multiplicity one as a composition factor of the ℓ -reductions of both Y_j and Y_{j-1} .*

Proof. Our assumption on j implies that Y_j occurs as an eigenspace of F^δ on some cohomology group $H_c^i(X, K)$ of degree $i > r$. It is also non-cuspidal, and we can choose a maximal proper ϕ -stable subset I of simple roots such that the restriction ${}^*R_{L_I}^G(Y_j)$ is non-zero. By Formula 3.4, this forces one of the modules between Y_j^I and Y_{j-1}^I to be non-zero. The latter is actually always non-zero: indeed, by Proposition 3.1, the module Y_j^I is an eigenspace of F^δ on $H_c^{i-1}(X_I, K)$ with $i - 1 > r - 1$. But the Coxeter variety X_I has dimension $r - 1$, and Theorem 2.11 ensures that Y_{j-1}^I is non-zero as soon as Y_j^I is.

Denote by a (resp. b) the multiplicity of S_j (resp. S_{j+1}) in the composition series of the ℓ -reductions of Y_j (by convention, the multiplicity is set to zero if the

module S_m is zero). By the previous lemma, these modules are the only possible non-cuspidal composition factors. Since the restriction functor is compatible with the decomposition maps, we can write

$$\text{dec}_{L_I}([\ast R_{L_I}^G(Y_j)]) = a[\ast R_{L_I}^G(S_j)]_k + b[\ast R_{L_I}^G(S_{j+1})]_k$$

in $K_0(kG\text{-mod})$. Together with Formula 3.4, it becomes

$$\text{dec}_{L_I}([Y_j^I]) + \text{dec}_{L_I}([Y_{j-1}^I]) = a[\ast R_{L_I}^G(S_j)]_k + b[\ast R_{L_I}^G(S_{j+1})]_k$$


But by Lemma 3.5, we know that the character of the restriction of S_m is either zero or equal to $\text{dec}_{L_I}([Y_{m-1}^I])$. Since the latter is non-zero for $m = j$, the previous equality forces $\ast R_{L_I}^G(S_j) \neq \{0\}$ and $a = 1$. In particular, S_j is a simple kG -module which occurs with multiplicity one in any ℓ -reduction of Y_j . The same argument applies to the equation

$$\text{dec}_{L_I}([\ast R_{L_I}^G(Y_{j-1})]) = a'[\ast R_{L_I}^G(S_{j-1})]_k + b'[\ast R_{L_I}^G(S_j)]_k.$$

It shows that S_j occurs also as a composition factor with multiplicity $b' = 1$ in any ℓ -reduction of Y_{j-1} . \square

In particular, when j is not equal to m_{ζ_j} , the sum $\chi_j + \chi_{j-1}$ is always the character of an indecomposable projective ΛG -module. We will denote this module by P_j .

Consequences 3.8. From the two previous lemmas we deduce that the lines



are subtrees of the Brauer tree Γ of the principal ℓ -block of G . Moreover, the missing edges are labeled by the simple cuspidal kG -modules.

3.2 Cuspidal kG -modules

The most delicate step in the proof of the conjecture of Hiss-Lübeck-Malle consists in gluing the non-cuspidal branches to the exceptional node. If the conjecture holds, then by the previous work the missing edges are labeled by the simple cuspidal kG -modules, or equivalently by their projective cover. The character of the latter should therefore be given by $\chi_{\text{exc}} + \chi_{\zeta_m}$. We present here an explicit construction of these projective modules using the complex $\text{R}\Gamma_c(Y(\dot{c}), \Lambda)$ representing the cohomology of $Y(\dot{c})$. By the results in Section 1.2 we know that this complex is perfect and thus provides a bunch of projective modules. Unfortunately, these are in general too big to be computed explicitly, and we need to factor them out according to the action of F^δ . This can be achieved with the help of the tools introduced at the end of Section 1.2. However, to make this operation work perfectly, we will make the following assumption:

- (W) For all integer $j \in \llbracket 0; h_0 - 1 \rrbracket$ such that $j = m_{\zeta_j}$, the generalized (λ_j) -eigenspace of F^δ on $bH_c^\bullet(Y(\dot{c}), \Lambda)$ is torsion-free.

By Theorem 2.11, these eigenvalues correspond exactly to the eigenvalues of F^δ on the cohomology group in middle degree $H_c^r(X, K)$.

Under this assumption, the missing projective modules turn out to be homotopy equivalent to suitably chosen generalized eigenspaces of F^δ on the complex $C = bR\Gamma_c(Y(\dot{c}), \Lambda)$. We shall rather work with a particular representative of this complex, with good finiteness properties that are needed to apply the constructions detailed in Section 1.2.

By Corollary 1.5, there exists a bounded complex \mathcal{C} of finitely generated $(\Lambda G, \Lambda \mathbf{T}^{cF})$ -bimodules, projective as left and right modules, together with an equivariant map $f : C \rightarrow \mathcal{C}$ and its inverse $g : \mathcal{C} \rightarrow C$ in the category $K^b(\Lambda G \times (\mathbf{T}^{cF})^{\text{op}})$. The Frobenius F^δ induces a morphism on \mathcal{C} which we will denote by $\mathcal{F} = f \circ F^\delta \circ g$.

Proposition 3.9. *Let $j \in \llbracket 0; h_0 - 1 \rrbracket$ such that $j = m_{\zeta_j}$ and λ_j the associated eigenvalue of F^δ . Assume that λ_j satisfies the assumption (W). Then the complex $\mathcal{C}_{(\lambda_j)}$ is homotopy equivalent to an indecomposable projective module P concentrated in degree r , and its character is given by*

$$[P] = \chi_{\text{exc}} + \chi_{m_{\zeta_j}}.$$

Proof. For the sake of simplicity we will write $\lambda = \lambda_j$ and $\zeta = \zeta_j$. From Proposition 1.9 we deduce that the complex $\mathcal{C}_{(\lambda)}$ has the following properties:

- $\mathcal{C}_{(\lambda)}$ is a bounded complex of finitely generated projective ΛG -modules;
- the cohomology of the complex $\mathcal{C}_{(\lambda)} \otimes_\Lambda k$ is concentrated in degrees $r, \dots, 2r$ (it is already the case for $\mathcal{C} \otimes_\Lambda k$ since $Y(\dot{c})$ is an irreducible affine variety);
- the cohomology groups of the complex $\mathcal{C}_{(\lambda)} \otimes_\Lambda K$ vanish outside the degree r . This follows indeed from [15, Corollary 9.9], Theorem 2.11 and Fact 2.8:

$$H_c^i(\mathcal{C}_{(\lambda)} \otimes_\Lambda K) \stackrel{1.9}{=} bH_c^i(Y(\dot{c}), K)_{(\lambda)} \stackrel{[15]}{=} H_c^i(X, K)_{(\lambda)} \stackrel{2.8}{=} H_c^i(X, K)_\lambda \stackrel{2.11}{=} 0.$$

Since the assumption (W) ensures that the groups $H_c^i(\mathcal{C}_{(\lambda)})$ are torsion-free, they are in fact zero for $i \neq r$. Therefore, by Lemma 1.2, the complex $\mathcal{C}_{(\lambda)}$ can be represented up to homotopy by a projective module $P = H^r(\mathcal{C}_{(\lambda)})$ concentrated in degree r . The character of this module corresponds (up to a sign) to the total character of the complex and is given by

$$[P] = \sum_{i=r}^{2r} (-1)^{i-r} [\mathcal{C}_{(\lambda)}^i] = [bH_c^r(Y(\dot{c}), K)_{(\lambda)}].$$

Moreover, by Theorem 2.11, this character has only one non-exceptional (i.e. unipotent) irreducible component, namely $\chi_{m_{\zeta_j}}$. Consequently, the only possible choice for the character of P is $\chi_{\text{exc}} + \chi_{m_{\zeta_j}}$. \square

In view of the results in the previous section, we have constructed the projective covers of the simple cuspidal kG -modules in the block. Following the previous notation, these simple modules will be denoted by $S_{m_{\zeta_j}}$ and their cover

by P_{m_ζ} . They label the edges of the Brauer tree connecting the non-cuspidal branches (or equivalently, the connected components of Γ^\bullet) to the exceptional node. This gives exactly the first part of the conjecture of Hiss, Lübeck and Malle:

Corollary 3.10. *Under the assumption (W), Conjecture (HLM) holds.*

3.3 Some numerical applications

We conclude this section by recording two direct consequences of the previous study, always under the assumption that (W) holds.

Proposition 3.11. *The simple unipotent cuspidal KG -modules in the principal ℓ -block remain simple after ℓ -reduction.*

Proof. By Theorem 2.11, the simple unipotent cuspidal KG -modules in the block correspond to the modules Y_{m_ζ} such that $m_\zeta = M_\zeta$. According to Conjecture (HLM), the associated nodes in the Brauer tree are extremities of the branches, so that any ℓ -reduction of Y_{m_ζ} is isomorphic to the simple kG -module S_{m_ζ} . \square

In Lusztig's classification, the irreducible unipotent characters fall into families [30]. By a case-by-case analysis, it has been checked that Lusztig's α -function (defined as the valuation of the degree of the unipotent character as a polynomial in the variable q) is constant on each family \mathcal{F} . This value will be denoted by $a(\mathcal{F})$. Similarly, the degree A_χ of the polynomial degree of any unipotent character χ depends only on the family.

Proposition 3.12. *Let $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_m$ be the F -stable families of unipotent characters lying in the principal ℓ -block, ordered such that $a(\mathcal{F}_1) \leq \dots \leq a(\mathcal{F}_m)$. Then the irreducible Brauer characters in the block can be labeled such that the decomposition matrix has a lower unitriangular shape:*

$$D = \begin{pmatrix} I_{r_1} & 0 & \cdots & 0 \\ * & I_{r_2} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ * & \cdots & * & I_{r_m} \end{pmatrix}$$

where each diagonal block is the identity matrix of the appropriate size.

Proof. By [9, Lemma 5.11], the power of q^δ occurring in the eigenvalue of F^δ associated to χ is given by

$$n_\chi = 2r - \frac{\alpha_\chi + A_\chi}{h}.$$

Since these integers are constant on each family, we deduce from the shape of the tree that two distinct characters χ and η lying the same family belong to distinct branches. In particular, $\text{dec}_{kG}(\chi)$ and $\text{dec}_{kG}(\eta)$ have no common irreducible component, which explains the identity matrices in the diagonal of D .

On the other side, one can check by a case-by-case analysis that in the principal ℓ -block, $a_\chi \leq a_\eta$ if and only if $n_\eta \geq n_\chi$. Since n_χ increases along each branch of the tree (see assertion (ii) in Conjecture (HLM)), we deduce that the decomposition matrix has a lower triangular shape. \square

It is conjectured that these results (excluding the supercuspidality property) hold in general for any good prime number ℓ (see [20, Conjecture 3.4]). We will give in Section 4.4 a more conceptual explanation of the previous proposition.

4 Towards Broué's conjecture

This last section aims at finalizing the study of the principal ℓ -block of G . In all the results stated here, we assume that the following holds:

(S) The Λ -modules $bH_c^i(Y(\dot{c}), \Lambda)$ are torsion-free.

Such an assumption is known to be valid for groups with \mathbb{F}_q -rank 1 (since the corresponding Deligne-Lusztig variety is a irreducible affine curve) and for groups of type A_n [3]. Many other cases will be settled in a subsequent paper [17] (see also [18] for groups of type B_n , C_n and 2D_n). Under this assumption, the cohomology groups $bH_c^i(Y(\dot{c}), \Lambda)$ are integral versions of the groups $bH_c^i(Y(\dot{c}), K)$ and their ℓ -reduction correspond to the groups $bH_c^i(Y(\dot{c}), k)$. Therefore, most of the information contained in the modular cohomology of $Y(\dot{c})$ is again obtained from Lusztig's fundamental work.

We start by giving an explicit representative of the complex $bR\Gamma_c(Y(\dot{c}), \Lambda)$ in terms of the projective modules P_j defined in the previous section. More precisely, we give, for all eigenvalue λ of F^δ , a bounded complex of finitely generated projective ΛG -modules homotopy equivalent to the generalized (λ) -eigenspace of F^δ . Under the assumption (S), we know explicitly the shape of the Brauer tree of the principal ℓ -block. Surprisingly, this complex turns out to be exactly the Rickard complex associated to the node labeled by λ [34, Section 4]. From that observation we deduce that $bR\Gamma_c(Y(\dot{c}), \Lambda)$ induces a splendid equivalence between the principal ℓ -blocks of G and $\mathbf{T}^{cF} \rtimes C_W(c\sigma)$, as predicted in the geometric version of Broué's conjecture [7]. We will conclude by giving two main consequences of this equivalence, namely the planar embedding of the Brauer tree and the unitriangularity shape of the decomposition matrix. As we have shown in section 3.3, the latter property is already a consequence of Conjecture (HLM). Nevertheless, we shall give here a conceptual approach to this result using the recent work of Chuang and Rouquier on perverse equivalences [12].

4.1 Retrieving the complex

For the sake of notation, we shall simply denote by Y the Deligne-Lusztig variety associated to \dot{c} . We want to determine explicitly the contribution of each eigenspace of F^δ on the cohomology of Y . This is in some sense a generalization

of Proposition 3.9 which refers only to the "minimal" eigenvalues. We shall keep the same approach: using the torsion-free assumption one can find a small representative of the complex, in which only specific projective modules can show up. The determination of these modules is then achieved using the total character of the complex that we deduce from Lusztig's work.

As in Section 3.2, we shall work with a specific representative of the cohomology complex $C = b\mathrm{R}\Gamma_c(Y, \Lambda)$ with good finiteness properties: by Corollary 1.5, there exists a bounded complex \mathcal{C} of finitely generated $(\Lambda G, \Lambda \mathbf{T}^{cF})$ -modules, whose terms are projective as both ΛG and $\Lambda \mathbf{T}^{cF}$ -modules such that C is homotopy equivalent to \mathcal{C} . By transfer, the Frobenius F^δ induces an endomorphism \mathcal{F} of \mathcal{C} such that the images of F^δ and \mathcal{F} coincide under the isomorphism $\mathrm{End}_{K^b(\Lambda G)}(\mathcal{C}) \simeq \mathrm{End}_{K^b(\Lambda G)}(C)$.

Drawing inspiration from the case of $\mathrm{SL}_2(\mathbb{F}_q)$ detailed in [1], we express each relevant eigenspace of \mathcal{F} on \mathcal{C} in terms of the indecomposable projective ΛG -modules P_j :

Theorem 4.1. *Let $j \in \llbracket 0; h_0 - 1 \rrbracket$. The generalized (λ_j) -eigenspace of \mathcal{F} on \mathcal{C} is homotopy equivalent to the following complex, with non-zero terms in degrees $r, \dots, r + j - m_{\zeta_j}$ only:*

$$0 \longrightarrow P_{m_\zeta} \longrightarrow P_{m_\zeta+1} \longrightarrow \cdots \longrightarrow P_{j-1} \longrightarrow P_j \longrightarrow 0.$$

Moreover, the boundary maps $\bar{d}_i : \bar{P}_i \longrightarrow \bar{P}_{i+1}$ remain non-zero after ℓ -reduction.

Note that this complex is exactly the Rickard complex attached to the node labelled by χ_j in the Brauer tree [34]. This observation will be fundamental in the next sections.

Before going into the details of the proof, let us recall some notation and basic properties of the Brauer tree Γ of the principal ℓ -block of ΛG . The non-exceptional nodes are labelled by the unipotent characters χ_j lying in the block. There are as many simple kG -modules as unipotent characters, but we can distinguish

- the non-cuspidal modules S_j with $j > m_{\zeta_j}$. Their projective cover P_j has character $\chi_j + \chi_{j-1}$, and hence labels the edge connecting the nodes associated to χ_j and χ_{j-1} ;
- the cuspidal modules S_{m_ζ} . The character of the corresponding projective cover is given by $[P_{m_\zeta}] = \chi_{m_\zeta} + \chi_{\mathrm{exc}}$.

Moreover, from the particular shape of the tree (given by the conjecture of Hiss-Lübeck-Malle) one can deduce that for $j > m_{\zeta_j}$, the ℓ -reduction of the projective module P_j is given by

$$\bar{P}_j = \begin{array}{c} S_j \\ S_{j-1} \oplus S_{j+1} \\ S_j \end{array}$$

if $j < M_{\zeta_j}$, that is in the case where the node labelled by χ_j is not at an extremity of the tree. Otherwise it is given by

$$\overline{P}_{M_\zeta} = \begin{array}{c} S_{M_\zeta} \\ S_{M_{\zeta-1}} \\ S_{M_\zeta} \end{array}$$

On the other hand, if the planar embedding of Γ is not specified, one cannot know precisely what the modules \overline{P}_{m_ζ} will look like. We know, however, that they have simple head and simple socle, both isomorphic to S_{m_ζ} , and that their class in the Grothendieck group $K_0(kG\text{-mod})$ is given by

$$[\overline{P}_{m_\zeta}]_k = [S_{m_\zeta}]_k + [S_{m_{\zeta+1}}]_k + \frac{|\mathbf{T}^{cF}|_\ell - 1}{h_0} \sum_\xi [S_{m_\xi}]_k.$$

The number $(|\mathbf{T}^{cF}|_\ell - 1)/h_0$ corresponds actually to the multiplicity of the exceptional node, that is the number of irreducible components of χ_{exc} .

Definition 4.2. We define the height in the tree Γ of an indecomposable projective ΛG -module P lying in the principal block to be the minimal length of a path from the exceptional node to the edge labelled by P . It will be denoted by $h_\Gamma(P)$.

We shall adopt the convention $h_\Gamma(P_{m_\zeta}) = 0$. By extension, the height of any finitely generated projective module lying in the block will be the maximal height of its indecomposable factors. Also, the height of a simple kG -module in the block will be naturally defined as the height of its projective cover, so that $h_\Gamma(P_j) = h_\Gamma(S_j) = j - m_{\zeta_j}$.

Remark 4.3. If P is a projective module of height n , then the height of any of its composition factors is at most $n + 1$.

In order to prove Theorem 4.1, we start by scanning the complex from the left to the right by removing the highest indecomposable factors. We shall assume that j is different from m_{ζ_j} since this case has been treated in Proposition 3.9. We first obtain the following representative:

Lemma 4.4. Let $\zeta = \zeta_j$. The complex $\mathcal{C}_{(\lambda_j)}$ is homotopy equivalent to a bounded complex of finitely generated projective ΛG -modules

$$0 \longrightarrow R_{m_\zeta} \xrightarrow{\delta_{m_\zeta}} R_{m_{\zeta+1}} \longrightarrow \cdots \longrightarrow R_{j-1} \xrightarrow{\delta_{j-1}} R_j \longrightarrow 0$$

satisfying $h_\Gamma(R_i) \leq h_\Gamma(P_i)$ for all $i = m_{\zeta_j}, \dots, j$.

Proof. The assumption (S) together with the results [15, Corollary 9.9] and 2.11 ensure that the cohomology of the complex $\mathcal{C}_{(\lambda_j)}$ vanishes outside the degrees r and $r + j - m_\zeta$. By Lemma 1.2 we deduce that the latter can be represented by a bounded complex of finitely generated ΛG -modules

$$0 \longrightarrow Q_{m_\zeta} \xrightarrow{d_{m_\zeta}} Q_{m_{\zeta+1}} \longrightarrow \cdots \longrightarrow Q_{j-1} \xrightarrow{d_{j-1}} Q_j \longrightarrow 0$$

satisfying the following properties:

- $\text{Im } d_i = \text{Ker } d_{i+1}$ for all $i = m_\zeta, \dots, r + j - m_\zeta - 1$;

- $H^r(\mathcal{C}_{(\lambda_j)}) = \text{Ker } d_{m_\zeta}$ is a ΛG -lattice with character $n_j \chi_{\text{exc}}$ for some non-negative integer n_j ;
- $H^{r+j-m_\zeta}(\mathcal{C}_{(\lambda_j)}) = \mathcal{Q}_j / \text{Im } d_{j-1}$ is a ΛG -lattice with character χ_j .

The integer n_j is actually non-zero otherwise χ_j would be a linear combination of projective characters. Moreover, we know from Section 2.2 that the multiplicity of χ_{exc} in the cohomology of Y is $\sum_i n_i = |C_W(c\sigma)| = h_0$, which forces each integer n_i to be equal to 1.

Remark 4.5. This proves incidentally that the ℓ -reduction of any eigenvalue of F^δ on the cohomology of Y (and not only X) is an h_0 -th root of unity.

Let us prove by induction on n that $\mathcal{C}_{(\lambda_j)}$ is homotopy equivalent to a complex of the following form

$$0 \longrightarrow R_{m_\zeta} \xrightarrow{\delta_{m_\zeta}} R_{m_\zeta+1} \xrightarrow{\delta_{m_\zeta+1}} \cdots \xrightarrow{\delta_{n-2}} R_{n-1} \xrightarrow{d'_{n-1}} Q'_n \xrightarrow{d'_n} \cdots \xrightarrow{d'_{j-1}} Q'_j \longrightarrow 0$$

with the R_i 's satisfying $\text{h}_\Gamma(R_i) \leq \text{h}_\Gamma(P_i)$ for all $i < n$. The case $n = m_\zeta$ is obtained from the previous analysis.

Assume then that $\mathcal{C}_{(\lambda_j)}$ is homotopy equivalent to the previous complex for some integer $n \geq m_\zeta$ and let us try to simplify Q'_n . For the sake of notation, we shall write $d : A \longrightarrow B$ instead of $d'_n : Q'_n \longrightarrow Q'_{n+1}$. Let P_m be any indecomposable direct summand of A that is assumed to be strictly higher than P_n (and hence of non-zero height). If we decompose A into $A = P_m \oplus A'$ then one can check that the following properties hold:

- d restricts to an injective map from P_m to B : by construction the character of P_m is $\chi_m + \chi_{m-1}$ whereas the character of $\text{Ker } d$ equals

$$[\text{Ker } d] = [R_{n-1}] - [R_{n-2}] + \cdots + (-1)^{n-m_\zeta+1} [R_{m_\zeta}] + (-1)^{n-m_\zeta} \chi_{\text{exc}}.$$

Now, by assumption, neither χ_m nor χ_{m-1} can occur in this expression. Therefore, the module $P_m \cap \text{Ker } d$ has zero character; since it is torsion-free, it must be the zero module.

- the quotient module $B/d(P_m)$ is torsion-free: we can use the following exact sequence of kG -modules:

$$0 \longrightarrow \text{Tor}_1^\Lambda(B/d(P_m), k) \longrightarrow \bar{P}_m \xrightarrow{\bar{d}} \bar{B}.$$

Consequently, it is sufficient to show that $\text{Ker } \bar{d} \cap \bar{P}_m$ is zero. Let us consider the class of $\text{Ker } \bar{d}$ in $K_0(kG\text{-mod})$, which is given by

$$[\text{Ker } \bar{d}] = [\bar{R}_{n-1}] - [\bar{R}_{n-2}] + \cdots + (-1)^{n-m_\zeta+1} [\bar{R}_{m_\zeta}] + (-1)^{n-m_\zeta} [\text{Ker } \bar{\delta}_{m_\zeta}].$$

Using the assumption (S) and the universal coefficient formula, we can identify the kG -module $H^r(\mathcal{C}_{(\lambda_j)} \otimes_\Lambda k) = \text{Ker } \bar{\delta}_{m_\zeta}$ with the ℓ -reduction of $H^r(\mathcal{C}_{(\lambda_j)})$ and can thus write

$$\mathrm{Ker} \bar{\delta}_{m\zeta} \simeq \mathrm{H}^r(\mathcal{C}_{(\lambda_j)}) \otimes_{\Lambda} k = \overline{\mathrm{Ker} \delta_{m\zeta}}.$$

Since $\mathrm{H}^r(\mathcal{C}_{(\lambda_j)})$ is cuspidal (it is torsion-free and its character is χ_{exc}), the module $\mathrm{Ker} \bar{\delta}_{m\zeta}$ has only cuspidal composition factors. Therefore, the simple module S_m , which by definition has the same height as P_m , can occur as a composition factor

- neither in \bar{R}_i for $i < n$, for the height of any of its irreducible components is at most $h_{\Gamma}(P_n)$ by assumption (see Remark 4.3);
- nor in $\mathrm{Ker} \bar{\delta}_{m\zeta}$ whose irreducible components are cuspidal and have hence height zero.

From the expression of $[\mathrm{Ker} d]$ given previously we deduce that S_m is not a composition factor of $\mathrm{Ker} d$. Since it is the only simple module of the socle of \bar{P}_m , this forces $\mathrm{Ker} \bar{d} \cap \bar{P}_m$ to be zero.

Consequently, we can decompose B into $B = d(P_m) \oplus B'$ so that d induces an isomorphism between P_m and $d(P_m)$. If we define $d' : A' \rightarrow B'$ to be the composition of the restriction $d|_{A'}$ with the projection $B \rightarrow B'$, then one can construct the following complex

$$0 \longrightarrow R_{m\zeta} \xrightarrow{\delta_{m\zeta}} \cdots \longrightarrow R_{n-1} \longrightarrow A' \xrightarrow{d'} B' \longrightarrow \cdots \xrightarrow{d'_{j-1}} Q'_j \longrightarrow 0$$

which is clearly homotopy equivalent to $\mathcal{C}_{(\lambda_j)}$. By removing repeatedly all the indecomposable direct summands of A that are higher to P_n we obtain the projective module R_n . \square

In the previous lemma, we have modified the complex from the left to the right by removing the superfluous projective modules. We now use the same method in the other direction to finish the proof of Theorem 4.1.

Proof of the theorem. We argue once again by induction: we show that up to homotopy $\mathcal{C}_{(\lambda_j)}$ can be written as

$$0 \longrightarrow R_{m\zeta} \xrightarrow{\delta_{m\zeta}} R_{m\zeta+1} \longrightarrow \cdots \xrightarrow{\delta_{n-1}} R_n \xrightarrow{\delta} P_{n+1} \longrightarrow \cdots \longrightarrow P_j \longrightarrow 0$$

where the modules R_i 's satisfy the condition $h_{\Gamma}(R_i) \leq h_{\Gamma}(P_i)$. Note that the case $n = j$ has been treated in the previous lemma.

Assume that we are working with the previous complex for some integer $n \leq j$. In that case, the character of the ΛG -module $\mathrm{Coker} \delta_{n-1} = R_n / \mathrm{Im} \delta_{n-1}$ is given by

$$\begin{aligned} [\mathrm{Coker} \delta_{n-1}] &= [P_{n+1}] - [P_{n+2}] + \cdots + (-1)^{j-n+1} [P_j] + (-1)^{j-n} [\mathrm{H}^{r+j-m\zeta}(\mathcal{C}_{(\lambda_j)})] \\ &= (\chi_n + \chi_{n+1}) - (\chi_{n+1} + \chi_{n+2}) + \cdots + (-1)^{j-n} \chi_j \\ [\mathrm{Coker} \delta_{n-1}] &= \chi_n + (\chi_{n+1} - \chi_{n+1}) - (\chi_{n+2} - \chi_{n+2}) + \cdots = \chi_n. \end{aligned}$$

In addition, it is torsion-free: it is indeed isomorphic either to $\mathrm{Im} \delta$ when $n < j$ or to $\mathrm{H}^{r+j-m\zeta}(\mathcal{C}_{(\lambda_j)})$ when $n = j$. Therefore, the head of the ΛG -module $\mathrm{Coker} \delta_{n-1}$

consists of at most two simple modules, namely S_n and S_{n+1} . Let P be a projective cover of $\text{Coker } \delta_{n-1}$. The canonical projection $R_n \rightarrow \text{Coker } \delta_{n-1}$ factors through P so that $h_\Gamma(P) \leq h_\Gamma(R_n) \leq h_\Gamma(P_n)$. Consequently, S_{n+1} cannot be in the head of P which forces P to be exactly P_n . This allows us to decompose the module R_n into $R_n = P_n \oplus R'$ with $R' \subset \text{Im } \delta_{n-1}$.

For the sake of notation we shall now write $\partial : A \rightarrow B$ instead of $\delta_{n-1} : R_{n-1} \rightarrow R_n$. We can argue as in [1]: since R' is a projective module, the map $\partial^{-1}(R') \rightarrow R'$ splits. The image of the corresponding splitting map is a submodule R'' of A isomorphic to R' , such that the quotient A/R'' is torsion-free. Indeed, one can embed $(A/R'')/(\partial^{-1}(R')/R'')$ in B/R' via ∂ and both $B/R' \simeq P_n$ and $\partial^{-1}(R')/R'' \simeq \partial^{-1}(R') \cap \text{Ker } \partial \subset A$ are torsion-free. Since R'' is projective and A/R'' is torsion-free, then the map $R'' \hookrightarrow A$ must be a retraction (see the proof of Lemma 1.2 for more details), and A decomposes into $A = R'' \oplus A'$ as a ΛG -module. It follows that $\mathcal{C}_{(\lambda_j)}$ is homotopy equivalent to

$$0 \rightarrow R_{m_\zeta} \rightarrow \cdots \rightarrow A' \rightarrow P_n \rightarrow P_{n+1} \rightarrow \cdots \rightarrow P_j \rightarrow 0$$

where the heights of the modules $R_{m_\zeta}, \dots, R_{n-1}, A'$ satisfy the conditions given in Lemma 4.4. At the last step of the induction we have obtained a complex of the following form:

$$0 \rightarrow R \rightarrow P_{m_\zeta+1} \rightarrow \cdots \rightarrow P_j \rightarrow 0.$$

Finally, the character of R (and therefore R itself) can be deduced from the total character of the complex, which is here equal to $\chi_{\text{exc}} + (-1)^{j-m_\zeta} \chi_j$. \square

Remark 4.6. One can actually make the boundary maps $d_i : P_i \rightarrow P_{i+1}$ explicit. The characters of the kernel and the image of d_i can be deduced from the cohomology of the variety Y with coefficients in K . One gets respectively $[\text{Ker } d_i] = \chi_{i-1}$ and $[\text{Im } d_i] = \chi_i$. Besides, we have shown in the course of the proof that the cokernel of d_i is a torsion-free module with character $[\text{Coker } d_i] = \chi_{i+1}$. Consequently, the ℓ -reduction of d_i factors through

$$\begin{array}{ccc} S_i & & S_{i+1} \\ S_{i-1} \oplus S_{i+1} & \longrightarrow & S_i \quad \hookrightarrow \quad S_i \oplus S_{i+2} \\ S_i & & S_{i+1} \end{array}$$

and the complex $\overline{C} = b\text{R}\Gamma_c(Y, k)$ is, as expected, homotopy equivalent to the Rickard complex associated to the Brauer tree Γ [34].

As a byproduct, we obtain many properties of the cohomology of the Deligne-Lusztig variety Y . We shall use the followings:

Corollary 4.7. *The complex $C = b\text{R}\Gamma_c(Y, \Lambda)$ is a tilting complex for ΛGb . In other words, it is a perfect complex of ΛGb -modules satisfying the following properties:*

- $\text{Hom}_{K^b(\Lambda Gb)}(C, C[i]) = 0$ for $i \neq 0$;
- $\text{add } C$ generates $K^b(\Lambda Gb\text{-mod})$ as a triangulated category.

Moreover, the endomorphism algebra $\text{End}_{K^b(\Lambda Gb)}(C)$ is free over Λ and is homotopy equivalent to $\text{Hom}_{\Lambda Gb}^\bullet(C, C)$ as a complex of $(\Lambda \mathbf{T}^{cF}, \Lambda \mathbf{T}^{cF})$ -bimodules.

Proof. \overline{C} is homotopy equivalent to a Rickard complex and, as such, it is a tilting complex [34, Theorem 4.2]. The same holds for C since it is the unique tilting complex lifting \overline{C} (see [35, Proposition 3.1 and Theorem 3.3]). Consequently, the cohomology of both $E = \text{Hom}_{\Lambda G}^\bullet(C, C)$ and $\overline{E} = E \otimes_\Lambda k$ is zero outside the degree 0. Now C is a perfect complex, and therefore $\text{Hom}_{\Lambda G}^\bullet(C, C)$ is homotopy equivalent to a complex of finitely generated projective $(\Lambda \mathbf{T}^{cF}, \Lambda \mathbf{T}^{cF})$ -bimodules. By Lemma 1.2, we deduce that E is homotopy equivalent to $H^0(E) \simeq \text{End}_{K^b(\Lambda G)}(C)$. In particular, the latter module is free over Λ . \square

Corollary 4.8. *The natural homomorphism of Λ -algebra*

$$\text{End}_{K^b(\Lambda Gb)}(b\text{R}\Gamma_c(Y, \Lambda)) \longrightarrow \text{End}_{\Lambda Gb}^{\text{gr}}(\mathbf{H}_c^\bullet(Y, \Lambda))$$

is injective.

Proof. Since the category $KG\text{-mod}$ is semi-simple, we have the following commutative diagram, with $C = b\text{R}\Gamma_c(Y, \Lambda)$

$$\begin{array}{ccc} \text{End}_{D^b(\Lambda Gb)}(C) & \xrightarrow{\iota} & \text{End}_{D^b(KGb)}(C) \\ \downarrow & & \downarrow \\ \text{End}_{\Lambda Gb}^{\text{gr}}(\mathbf{H}^\bullet(C)) & \longrightarrow & \text{End}_{K^b(KGb)}^{\text{gr}}(\mathbf{H}^\bullet(C)) \end{array}$$

By the previous corollary, the Λ -module $\text{End}_{D^b(\Lambda Gb)}(C) \simeq \text{End}_{K^b(\Lambda Gb)}(C)$ is free over Λ . We deduce that the map ι , as well as the first vertical arrow, are into. \square

Corollary 4.9. *The image of $F^h - 1$ in $\text{End}_{K^b(kGb)}(\overline{b}\text{R}\Gamma_c(Y, k))$ is nilpotent.*

Proof. Let $A = \text{End}_{K^b(\Lambda Gb)}(b\text{R}\Gamma_c(Y, \Lambda))$. By Corollary 4.7, the ℓ -reduction of A is exactly $\text{End}_{K^b(kGb)}(\overline{b}\text{R}\Gamma_c(Y, k))$. Let us denote by χ the minimal polynomial of F^δ on $H_c^\bullet(Y, K)$. By the previous corollary and the assumption (S), the image of $\chi(F^\delta)$ in A is zero. But the eigenvalues of F^δ reduce to h_0 -th roots of unity modulo ℓ (see Remark 4.5) and hence the class of $F^h - 1$ in \overline{A} is a nilpotent element (recall that $h = h_0\delta$). \square

4.2 Broué's conjecture

The original version of *Broué's abelian defect group conjecture* [4] predicts that the module categories of a block and its Brauer correspondent are derived equivalent, provided that the defect of the block is an abelian group. More precisely, given a block with abelian defect group H , represented by an idempotent b , and $c = \text{Br}(b)$ the corresponding block of $N_G(H)$, there exists an equivalence

$$D^b(\Lambda Gb\text{-Mod}) \xrightarrow{\sim} D^b(\Lambda N_G(H)c\text{-Mod}).$$

Such an equivalence induces a *perfect isometry* between the Grothendieck groups carrying numerous arithmetical information. There are indeed many numerical consequences we can deduce from it, e.g. it preserves the number of irreducible characters (ordinary and modular) as well as the similarity invariants of the Cartan matrix [5]. Up to now, the version of this conjecture is known to hold in the following cases:

- with restrictions on the defect group: if H is a cyclic group [34], [26] and [37] or isomorphic to the Klein group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ [36] and [27];
- with restrictions on G : for ℓ -solvable groups [14], [32] and [22], symmetric groups and general linear groups [11] or for finite reductive groups when ℓ divides $q - 1$ [33].

Many other particular cases have been handled, and a lot of evidences for this conjecture to hold have been collected.

It is unclear whether there should exist a canonical way to construct this equivalence. However, when G is a finite reductive group, it is expected to be induced by the cohomology of certain Deligne-Lusztig varieties. This is known as the *geometric version of Broué's conjecture*, as stated in [7] and [9]. Note that varieties associated to Levi subgroups – and not only tori – can be involved in this description. However, if the order of q modulo ℓ is assumed to be a regular number d , then it is sufficient to consider Deligne-Lusztig varieties $Y(w)$ associated to elements that satisfy the following properties:

(B1) $w\sigma$ is a good d -regular element [9].

(B2) ℓ divides $|T_w|$ but does not divide $[G : T_w]$.

We will also assume that the prime number ℓ is large:

(B3) ℓ does not divide $|W^F|$

in order to ensure that $N_G(\mathbf{T}_w) = N_G(T_w)$.

In this geometric framework, the defect group H of the principal block (which is an Sylow ℓ -subgroup of G) can be chosen to be a subgroup of T_w , and the property of $w\sigma$ to be regular forces $C_G(H)$ to be the exactly the torus \mathbf{T}_w , leading to $N_G(\mathbf{T}_w) = N_G(H)$. Then the geometric version of Broué's conjecture predicts that the perfect complex $b\mathrm{R}\Gamma_c(Y(w), \Lambda)$ induces a splendid Rickard equivalence (and in particular a derived equivalence) between the principal ℓ -blocks of G and $N_G(T_w)$. More precisely,

Conjecture 4.10 (Broué). *Under the previous assumptions, there exists a bounded complex D of $(\Lambda Gb, \Lambda N_G(T_w))$ -bimodules such that*

- (i) *The restrictions of $b\mathrm{R}\Gamma_c(Y(w), \Lambda)$ and D to the category of bounded complex of $(\Lambda Gb, \Lambda T_w)$ -bimodules are homotopy equivalent.*
- (ii) *The complex D induces a splendid Rickard equivalence between the principal ℓ -blocks of G and $N_G(T_w)$.*

Remark 4.11. In the case where w is not assumed to be good, the disjunction property of the cohomology

$$\forall i \neq 0 \quad \mathrm{Hom}_{D^b(\Lambda Gb)}(b\mathrm{R}\Gamma_c(Y(\dot{w}), \Lambda), b\mathrm{R}\Gamma_c(Y(\dot{w}), \Lambda)[-i]) = 0 \quad (4.12)$$

does not always hold. When (\mathbf{G}, F) has no twisted components of type ${}^2\mathrm{B}_2$, ${}^2\mathrm{F}_4$ ou ${}^2\mathrm{G}_2$, the other assumptions on w tell us that $(\mathbf{T}_w, 1)$ is a d -cuspidal pair, and [8, Theorem 5.24] turns out to be nothing but the numerical reflect of this conjecture.

Among the elements w satisfying the previous assumptions, Coxeter elements of (W, F) play a particular role, for they have the following remarkable property:

(B4) $C_W(w\sigma)$ is a cyclic group generated by $wF(w)\cdots F^{\delta-1}(w)$.

In this case, the action of the Frobenius F^δ provides a natural way to extend the action of $T_w \simeq \mathbf{T}^{wF}$ on the complex $b\mathrm{R}\Gamma_c(Y(\dot{w}), \Lambda)$ to an action of the normalizer $N_G(T_w)$. Using this crucial observation, Rouquier has reduced Broué's conjecture to the disjunction property 4.12 for the cohomology with coefficients in k [38]. We know from the previous section that this property is satisfied whenever w is a Coxeter element, so that Conjecture 4.10 can now be directly deduced from Corollary 4.7 and [38, Theorem 4.5]:

Theorem 4.13. *Under the assumption (S), the geometric version of Broué's conjecture holds for Coxeter elements.*

Nevertheless, it is worth giving details of the proof of [38, Theorem 4.5] since it helps to understand how the actions of $v = cF(c)\cdots F^{\delta-1}(c)$ and F^δ on the cohomology of Y are related.

Let us consider the algebra $A = \mathrm{End}_{K^b(\Lambda Gb)}(C)$ associated to the complex $C = b\mathrm{R}\Gamma_c(Y, \Lambda)$. By the disjunction property, A is homotopy equivalent to $\mathrm{End}_{\Lambda Gb}^\bullet(C)$ as a complex of $(\Lambda \mathbf{T}^{cF}, \Lambda \mathbf{T}^{cF})$ -bimodules (see Corollary 4.7). Moreover, it is free over Λ and thus satisfies

$$\overline{A} = A \otimes_\Lambda k \simeq \mathrm{End}_{K^b(kG\overline{b})}(\overline{C}).$$

Since the action of F^δ on C commutes with the action of G , we get a canonical morphism $\phi : \Lambda \mathbf{T}^{cF} \rtimes \langle F^\delta \rangle_{\mathrm{mon}} \longrightarrow A$ which turns out to be surjective (for more details see the proof of [38, Theorem 4.5]). We will denote by τ the image of F^h and by $\langle \tau \rangle$ the subalgebra of A that it generates. By Corollary 4.9, $\overline{\tau} - 1$ is a nilpotent element of \overline{A} . Therefore, we can apply Hensel's lemma to the ideal $\mathfrak{m} = \mathrm{Nil}\langle \tau \rangle + \ell \langle \tau \rangle$ in order to obtain a element $\alpha \in \langle \tau \rangle$ such that $\alpha^{h_0} = \tau$ and $\overline{\alpha} - 1$ is nilpotent (recall that h is prime to ℓ). Then we can deform ϕ into a homomorphism of algebras

$$\psi : \Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma) \longrightarrow A$$

by setting $\psi(v^{-1}) = \alpha^{-1}\phi(F^\delta)$. This defines an action of v on C in the homotopy category such that the image of $v^{-1} - F^\delta$ in \overline{A} is a nilpotent element. To conclude,

we use [38, Lemma 4.9] to construct a complex D of $(\Lambda Gb, \Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma))$ -bimodules and a homotopy equivalence f between the restrictions of C and D to the category of complexes of $(\Lambda Gb, \Lambda \mathbf{T}^{cF})$ -bimodules such that the following diagram is commutative:

$$\begin{array}{ccc} \Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma) & \xrightarrow{\psi} & \text{End}_{K^b(\Lambda Gb)}(C) \\ & \searrow \text{can} & \uparrow f \\ & & \text{End}_{K^b(\Lambda Gb)}(D) \end{array}$$

We deduce that the functor $D^\vee \otimes_{\Lambda G} -$ induces the expected splendid Rickard equivalence between the principal ℓ -blocks of G and $\mathbf{T}^{cF} \rtimes C_W(c\sigma)$.

Remark 4.14. The projection $N_G(T_c) \twoheadrightarrow C_W(c\sigma)$ given by Proposition 2.4 does not split in general, and the groups $N_G(T_c)$ and $\mathbf{T}^{cF} \rtimes C_W(c\sigma)$ are not isomorphic. However, it is proven in [38] that the algebras $\Lambda N_G(T_c)$ and $\Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma)$ become isomorphic as soon as ℓ satisfies the assumptions given at the beginning of Section 2.2.

In the remaining sections, we shall investigate further properties of the functor $D^\vee \otimes_{\Lambda G} -$ using explicit representatives coming from Section 4.1.

4.3 Planar embedding of the Brauer tree

We start by constructing a representative \mathcal{D} of D with good finiteness properties, as we did for $C = b\text{R}\Gamma_c(Y, \Lambda)$. The restriction of D to the category of complexes of $(\Lambda G, \Lambda \mathbf{T}^{cF})$ -bimodules is homotopy equivalent to C , and hence to \mathcal{C} which is a bounded complex of projective modules.

It remains to define the action of $C_W(c\sigma)$ on this complex using the action of v on D . By transfer, there exists an endomorphism \tilde{v} of \mathcal{C} such that v and \tilde{v} coincide under the isomorphism $\text{End}_{K^b(\Lambda G)}(D) \simeq \text{End}_{K^b(\Lambda G)}(\mathcal{C})$. Note that there is no canonical choice for \tilde{v} , since it depends on the homotopy equivalence we choose between \mathcal{C} and D . The image of \tilde{v} under this isomorphism has order h_0 , which means that there exists a null-homotopic endomorphism n of \mathcal{C} such that $\tilde{v}^{h_0} = 1 + n$. We can then argue as in [1]: we consider the complex \mathcal{D} obtained from \mathcal{C} by removing any direct summand homotopy equivalent to zero. Since $\ell > h_0$, one can extract an h_0 -th root of $1 + n$ using the power series ${}^{h_0}\sqrt{1+X}$. In other words, there exists a null-homotopic endomorphism n' of \mathcal{D} such that $(1 + n')^{h_0} = 1 + n$. This allows us to define the action of $v \in C_W(c\sigma)$ on \mathcal{D} to be the endomorphism $\mathcal{V} = (1 + n')^{-1}\tilde{v}$.

In summary, we have constructed a bounded complex \mathcal{D} of finitely generated $(\Lambda G, \Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma))$ -bimodules such that:

- \mathcal{D} is a bounded complex whose terms are projective as both left and right modules (since the order of $C_W(c\sigma)$ is invertible in Λ);

- the restrictions of \mathcal{D} and $C = b\mathrm{R}\Gamma_c(Y, \Lambda)$ to the category of $(\Lambda G, \Lambda \mathbf{T}^{cF})$ -bimodules are homotopy equivalent;
- under the isomorphism $\mathrm{End}_{K^b(kG)}(\overline{\mathcal{D}}) \simeq \mathrm{End}_{K^b(kG)}(\overline{C})$, the images of v^{-1} and F^δ differ only by a nilpotent element.

In particular, we can identify the generalized (λ) -eigenspaces of v^{-1} and F^δ on the cohomology groups of Y . This means that the pair $(\mathcal{D}, \mathcal{V})$ has the same properties as the pair $(\mathcal{C}, \mathcal{F})$ that are required in the proof of Theorem 4.1. Consequently, for all $j = 0, \dots, h_0 - 1$, the generalized $(q^{-j\delta})$ -eigenspace \mathcal{D}_j of \mathcal{V} on \mathcal{D} is homotopy equivalent to the following complex of ΛG -modules:

$$0 \longrightarrow P_{m_\zeta} \longrightarrow P_{m_{\zeta+1}} \longrightarrow \cdots \longrightarrow P_{j-1} \longrightarrow P_j \longrightarrow 0$$

where ζ stands for ζ_j . Using this particular representative, we can now determine the planar embedding of the Brauer tree:

Theorem 4.15. *Under the assumption (S), Conjecture (HLM+) holds.*

Proof. It is sufficient to show that the condition $\mathrm{Ext}_{kG}^1(S_{m_\zeta}, S_{m_\xi}) \neq 0$ forces m_ξ to be congruent to $M_\zeta + 1$ modulo h_0 . Using the formalism of derived categories, we shall write this group as

$$\mathrm{Ext}_{kG}^1(S_{m_\zeta}, S_{m_\xi}) \simeq \mathrm{Hom}_{D^b(kG)}(S_{m_\zeta}, S_{m_\xi}[1])$$

which in turn is isomorphic to

$$\mathrm{Hom}_{D^b(k\mathbf{T}^{cF} \times C_W(c\sigma))}(\overline{\mathcal{D}}^\vee \otimes_{kG} S_{m_\zeta}, \overline{\mathcal{D}}^\vee \otimes_{kG} S_{m_\xi}[1])$$

via the fully-faithful functor $\overline{\mathcal{D}}^\vee \otimes_{kG} -$. Now, for $i \neq m_\zeta$, the module $\overline{P}_i^\vee \otimes_{kG} S_{m_\zeta} = \mathrm{Hom}_{kG}(\overline{P}_i, S_{m_\zeta})$ is zero. Consequently, we can use the action of v to obtain the following isomorphisms in the category $K^b(kC_W(c\sigma)\text{-mod})$:

$$\overline{\mathcal{D}}^\vee \otimes_{kG} S_{m_\zeta} \simeq \bigoplus_{j=0}^{h_0-1} \overline{\mathcal{D}}_j^\vee \otimes_{kG} S_{m_\zeta} \simeq \bigoplus_{j=m_\zeta}^{M_\zeta} \overline{\mathcal{D}}_j^\vee \otimes_{kG} S_{m_\zeta} \simeq \bigoplus_{j=m_\zeta}^{M_\zeta} k_j[-r]$$

where k_j is the one-dimensional simple $kC_W(c\sigma)$ module on which v acts by multiplication by $\overline{q}^{-j\delta}$. Note that v acts on \mathbf{T}^{cF} by raising any element to the power of $q^{-\delta}$ so that this notation is consistent to the one given in Example 2.13. From the previous isomorphism we deduce that the complex $\overline{\mathcal{D}}^\vee \otimes_{kG} S_{m_\zeta}$ is quasi-isomorphic to a $k\mathbf{T}^{cF} \times C_W(c\sigma)$ -module N_{m_ζ} concentrated in degree r satisfying

$$\mathrm{Res}_{kC_W(c\sigma)}^{k\mathbf{T}^{cF} \times C_W(c\sigma)}(N_{m_\zeta}) \simeq k_{m_\zeta} \oplus k_{m_\zeta+1} \oplus \cdots \oplus k_{M_\zeta}.$$

This shows that the composition factors of N_{m_ζ} are exactly $k_{m_\zeta}, \dots, k_{M_\zeta}$.

On the other side, the group $\mathrm{Ext}_{k\mathbf{T}^{cF} \times C_W(c\sigma)}^1(k_i, k_j)$ is non-zero if and only if $j \equiv i + 1$ modulo h_0 (see Example 2.13). Since

$$\mathrm{Ext}_{kG}^1(S_{m_\zeta}, S_{m_\xi}) \simeq \mathrm{Ext}_{k\mathbf{T}^{cF} \times C_W(c\sigma)}^1(N_{m_\zeta}, N_{m_\xi})$$

we deduce that this latter group is non-zero only if there exist integers $i \in \llbracket m_\zeta; M_\zeta \rrbracket$ and $j \in \llbracket m_\xi; M_\xi \rrbracket$ such that $j \equiv i + 1$ modulo h_0 . We conclude by observing that the only integers that can satisfy this condition are $i = M_\zeta$ and $j = m_\xi$. \square

Remark 4.16. One can be more precise about the module N_{m_ζ} . Being the image of an indecomposable, it is itself an indecomposable $k\mathbf{T}^{cF} \rtimes C_W(c\sigma)$ -module. Moreover, since its composition factors are $k_{m_\zeta}, \dots, k_{M_\zeta}$, it is necessarily uniserial and of the following shape:

$$N_{m_\zeta} = \begin{array}{c} k_{M_\zeta} \\ k_{M_\zeta-1} \\ \vdots \\ k_{m_\zeta} \end{array} .$$

Note however that there is still an ambiguity for groups of type A_1 .

Remark 4.17. For groups of \mathbb{F}_q -rank 1 such as the Ree group of type 2G_2 , the assumption (S) is automatically satisfied (the corresponding Deligne-Lusztig variety is an irreducible affine curve). As a consequence, the planar embedded Brauer tree of the principal ℓ -block when q has order 12 modulo ℓ is exactly the one given in Figure 3. This completes [23, Theorem 4.4].

Furthermore, we shall prove in a subsequent paper that the assumption (S) holds whenever conjecture (HLM) holds and p is assumed to be a good prime number [17]. In particular, from the knowledge of the shape of the Brauer tree we can deduce the planar embedding. We obtain therefore the planar embedding for groups of type F_4 when $p \neq 2, 3$, which completes [24, Theorem 2.1]. This would also apply to groups of type E_7 and E_8 if the shapes of the trees were known.

4.4 Perverse equivalence and decomposition matrix

To finish off with the principal ℓ -block, we observe that the equivalence induced by the cohomology of Y is perverse in the sense of [12]. This leads to a conceptual proof of the unitriangularity shape of the decomposition matrix of the block.

Let us fix some notation for the objects involved in the definition. Recall that a full subcategory \mathcal{B} of an abelian category \mathcal{A} is a *Serre subcategory* if for all short exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

in \mathcal{A} , B is an object of \mathcal{B} if and only if both A and C are objects of \mathcal{B} . In other words, the category \mathcal{B} is stable by quotients, submodules, and extensions by objects of \mathcal{B} .

Given such a category, we will denote by $D_{\mathcal{B}}^b(\mathcal{A})$ the full subcategory of $D^b(\mathcal{A})$ of complexes whose cohomology groups are objects of \mathcal{B} . We can then form the quotient categories \mathcal{A}/\mathcal{B} (abelian) and $D^b(\mathcal{A})/D_{\mathcal{B}}^b(\mathcal{A})$ (triangulated).

Definition 4.18. *Let \mathcal{A} and \mathcal{A}' be two abelian categories, \mathcal{S} and \mathcal{S}' the sets of isomorphism classes of simple objects of these categories. A derived equivalence $\Theta : D^b(\mathcal{A}) \xrightarrow{\sim} D^b(\mathcal{A}')$ is perverse if there exist*

- filtrations $\emptyset = \mathcal{S}_0 \subset \mathcal{S}_1 \subset \dots \subset \mathcal{S}_r = \mathcal{S}$ and $\emptyset = \mathcal{S}'_0 \subset \mathcal{S}'_1 \subset \dots \subset \mathcal{S}'_r = \mathcal{S}'$ of the sets \mathcal{S} and \mathcal{S}' ;
- a perversity function $p : \llbracket 0; r \rrbracket \rightarrow \mathbb{Z}$;

such that if \mathcal{A}_i (resp. \mathcal{A}'_i) denotes the Serre subcategory generated by \mathcal{S}_i (resp. \mathcal{S}'_i), then for all i

- (i) Θ restricts to an equivalence $D_{\mathcal{A}_i}^b(\mathcal{A}) \simeq D_{\mathcal{A}'_i}^b(\mathcal{A}')$;
- (ii) the functor $\mathcal{A}_i/\mathcal{A}_{i-1} \hookrightarrow D_{\mathcal{A}_i}^b(\mathcal{A})/D_{\mathcal{A}_{i-1}}^b(\mathcal{A}) \xrightarrow{\Theta} D_{\mathcal{A}'_i}^b(\mathcal{A}')/D_{\mathcal{A}'_{i-1}}^b(\mathcal{A}')$ factors through the following commutative diagram:

$$\begin{array}{ccc} D_{\mathcal{A}_i}^b(\mathcal{A})/D_{\mathcal{A}_{i-1}}^b(\mathcal{A}) & \xrightarrow{\Theta} & D_{\mathcal{A}'_i}^b(\mathcal{A}')/D_{\mathcal{A}'_{i-1}}^b(\mathcal{A}') \\ \text{can} \downarrow & & \downarrow [p(i)] \\ \mathcal{A}_i/\mathcal{A}_{i-1} & \overset{\sim}{\dashrightarrow} & \mathcal{A}'_i/\mathcal{A}'_{i-1} \end{array}$$

so that $\Theta[-p(i)]$ induces an equivalence $\mathcal{A}_i/\mathcal{A}_{i-1} \simeq \mathcal{A}'_i/\mathcal{A}'_{i-1}$.

From now on, we will assume that all the objects of the categories involved have finite composition series. This allows us to reformulate the assertions (i) and (ii) into less abstract terms:

- for any simple object L in \mathcal{S}_i , the composition factors of $H^n(\Theta(L))$ lie in \mathcal{S}'_i , and even in \mathcal{S}'_{i-1} whenever $n \neq -p(i)$;
- if $L \in \mathcal{S}_i \setminus \mathcal{S}_{i-1}$ then $H^{-p(i)}(\Theta(L))$ has a unique composition factor L' in $\mathcal{S}'_i \setminus \mathcal{S}'_{i-1}$ and the map $L \mapsto L'$ induces a bijection between the sets $\mathcal{S}_i \setminus \mathcal{S}_{i-1}$ and $\mathcal{S}'_i \setminus \mathcal{S}'_{i-1}$.

Roughly speaking, $\Theta(L)$ is quasi-isomorphic to $L'[p(i)]$ "modulo some composition factors in \mathcal{S}'_{i-1} ".

Let us go back to the derived equivalence we have been studying. Recall that we have constructed a representative \mathcal{D} of the cohomology of Y , as a bounded complex of finitely generated $(\Lambda G, \Lambda \mathbf{T}^{cF} \rtimes C_W(c\sigma))$ -bimodules, both projective as left and right modules. We can argue as in the proof of Theorem 4.15 and show that for all simple kG -module in the principal block, say S_j , we have

$$\overline{\mathcal{D}}^v \otimes_{kG} S_j \simeq N_j[m_\zeta - j - r] \quad (4.19)$$

where N_j is a $k\mathbf{T}^{cF} \rtimes C_W(c\sigma)$ -module satisfying

$$\mathrm{Res}_{kC_W(c\sigma)}^{k\mathbf{T}^{cF} \rtimes C_W(c\sigma)}(N_j) \simeq k_j \oplus k_{j+1} \oplus \cdots \oplus k_{M_{\zeta_j}}.$$

In particular, the composition factors of N_j are exactly the (inflation of) the simple modules $k_j, \dots, k_{M_{\zeta_j}}$.

From that observation we shall deduce that the functor $\Theta : \overline{\mathcal{D}} \otimes_{kG} -$ induces a perverse equivalence. We use the height function associated to the Brauer tree to define the filtrations

$$\mathcal{S}_i = \{S \in \mathrm{Irr} kG \mid h_\Gamma(S) \leq r - i\}$$

and

$$\mathcal{S}'_i = \{k_j \in \mathrm{Irr} k\mathbf{T}^{cF} \rtimes C_W(c\sigma) \mid S_j \in \mathcal{S}_i\}$$

on the set of isomorphism classes of simple modules (recall that $h_\Gamma(S_j) = j - m_{\zeta_j}$). Note that $\mathcal{S}_r = \mathrm{Irr} kG$ and $\mathcal{S}'_r = \mathrm{Irr} k\mathbf{T}^{cF} \rtimes C_W(c\sigma)$ are the last terms of these filtrations.

Finally, the image by Θ of a module $S_j \in \mathcal{S}_i \setminus \mathcal{S}_{i-1}$ has by 4.19 a unique composition factor in $\mathcal{S}'_i \setminus \mathcal{S}'_{i-1}$, namely k_j . Since it is concentrated in degree $r + j - m_{\zeta_j} = r + h_\Gamma(S_j) = 2r - i$, we may choose the perversity function to be $p(i) = i - 2r$ so that following holds:

Theorem 4.20. *Assume that the assumption (S) holds. Then the functor $\Theta = \overline{\mathcal{D}} \otimes_{kG} -$ together with the triple $(\mathcal{S}_\bullet, \mathcal{S}'_\bullet, p)$ induces a perverse equivalence between the principal blocks of kG and $k\mathbf{T}^{cF} \rtimes C_W(c\sigma)$. The corresponding bijection on the set of isomorphism classes of simple modules is $S_j \longleftrightarrow k_j$.*

In the case of the derived equivalences predicted by Broué's conjecture, one of the groups involved can be written as a semi-direct product $D \rtimes E$ with D an abelian ℓ -group and E an ℓ' -subgroup of $\mathrm{Aut}(D)$. For such an equivalence, the existence of a perverse datum provides information on the shape of the decomposition matrix of the block:

Corollary 4.21 (Chuang-Rouquier). *There exists an ordering of the simple KG -modules and the simple kG -modules such that the decomposition matrix of the principal block of ΛG has unitriangular shape.*

Proof. We claim that any ordering compatible with the perversity (here, the height in the tree) will be acceptable. Let $\tilde{\Theta} = \mathcal{D} \otimes_{\Lambda G} -$. The functor

$$K\tilde{\Theta} : D^b(KG\text{-mod}) \longrightarrow D^b(K\mathbf{T}^{cF} \rtimes C_W(c\sigma))$$

induces a bijection on the simples lying in the principal blocks. It is clear that this bijection maps χ_j to η_j (see Notation 2.9 and Example 2.13).

Let $\tilde{\Theta}^*$ be an inverse functor of $\tilde{\Theta}$. It induces again a perverse equivalence $\Theta^* = k\tilde{\Theta}^*$ with respect to the same filtrations but for an opposite perversity function. Since the filtration is given by the height in the tree, we obtain

$$\mathrm{dec}_G \chi_j = \mathrm{dec}_G \tilde{\Theta}^*(\eta_j) = \Theta^*([k_j]_k) = [S_j]_k + \sum_{h_\Gamma(S_i) \geq h_\Gamma(S_j)} a_{i,j} [S_i]_k$$

for some integers $a_{i,j}$, which proves that the decomposition matrix of ΛGb has a unitriangular shape. \square

Acknowledgements

I would like to thank the Oxford Mathematical Institute for providing me excellent research environment during the early stage of this work. I wish to thank Joe Chuang and Gerhard Hiss for our fruitful correspondence and Cédric Bonnafé, Jean-François Dat and Raphaël Rouquier for many valuable comments and suggestions.

References

- [1] C. Bonnafé. *Représentations du groupe $SL_2(q)$* .
- [2] C. Bonnafé and R. Rouquier. Catégories dérivées et variétés de Deligne-Lusztig. *Publ. Math. Inst. Hautes Études Sci.*, (97):1–59, 2003.
- [3] C. Bonnafé and R. Rouquier. Coxeter orbits and modular representations. *Nagoya Math. J.*, 183:1–34, 2006.
- [4] M. Broué. Isométries de caractères et équivalences de Morita ou dérivées. *Inst. Hautes Études Sci. Publ. Math.*, (71):45–63, 1990.
- [5] M. Broué. Isométries parfaites, types de blocs, catégories dérivées. *Astérisque*, (181-182):61–92, 1990.
- [6] M. Broué and G. Malle. Théorèmes de Sylow génériques pour les groupes réductifs sur les corps finis. *Math. Ann.*, 292(2):241–262, 1992.
- [7] M. Broué and G. Malle. Zyklotomische Heckealgebren. *Astérisque*, (212): 119–189, 1993. Représentations unipotentes génériques et blocs des groupes réductifs finis.
- [8] M. Broué, G. Malle, and J. Michel. Generic blocks of finite reductive groups. *Astérisque*, (212):7–92, 1993. Représentations unipotentes génériques et blocs des groupes réductifs finis.
- [9] M. Broué and J. Michel. Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées. In *Finite reductive groups (Luminy, 1994)*, volume 141 of *Progr. Math.*, pages 73–139.
- [10] R. W. Carter. *Simple groups of Lie type*. John Wiley & Sons, London-New York-Sydney, 1972. Pure and Applied Mathematics, Vol. 28.
- [11] J. Chuang and R. Rouquier. Calabi-yau algebras and perverse morita equivalences. *In preparation*.

- [12] J. Chuang and R. Rouquier. Derived equivalences for symmetric groups and \mathfrak{sl}_2 -categorification. *Ann. of Math. (2)*, 167(1):245–298, 2008.
- [13] C. W. Curtis and I. Reiner. *Methods of representation theory. Vol. I*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1990. With applications to finite groups and orders, Reprint of the 1981 original, A Wiley-Interscience Publication.
- [14] E. C. Dade. A correspondence of characters. In *The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979)*, volume 37 of *Proc. Sympos. Pure Math.*, pages 401–403.
- [15] P. Deligne and G. Lusztig. Representations of reductive groups over finite fields. *Ann. of Math. (2)*, 103(1):103–161, 1976.
- [16] F. Digne and J. Michel. *Representations of finite groups of Lie type*, volume 21 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1991.
- [17] O. Dudas. Coxeter Orbits and Brauer trees II. *preprint*, 2010.
- [18] O. Dudas. Géométrie des variétés de Deligne-Lusztig : décompositions, cohomologie modulo ℓ et représentations modulaire. *Thèse de doctorat*, 2010.
- [19] M. Geck. Modular Harish-Chandra series, Hecke algebras and (generalized) q -Schur algebras. In *Modular representation theory of finite groups (Charlottesville, VA, 1998)*, pages 1–66. de Gruyter, Berlin, 2001.
- [20] M. Geck and G. Hiss. Modular representations of finite groups of Lie type in non-defining characteristic. In *Finite reductive groups (Luminy, 1994)*, volume 141 of *Progr. Math.*, pages 195–249. Birkhäuser Boston, Boston, MA, 1997.
- [21] J. A. Green. Walking around the Brauer Tree. *J. Austral. Math. Soc.*, 17:197–213, 1974. Collection of articles dedicated to the memory of Hanna Neumann, VI.
- [22] M. E. Harris and M. Linckelmann. Splendid derived equivalences for blocks of finite p -solvable groups. *J. London Math. Soc. (2)*, 62(1):85–96, 2000.
- [23] G. Hiss. The Brauer trees of the Ree groups. *Comm. Algebra*, 19(3), 1991.
- [24] G. Hiss and F. Lübeck. The Brauer trees of the exceptional Chevalley groups of types F_4 and 2E_6 . *Arch. Math. (Basel)*, 70(1):16–21, 1998.
- [25] G. Hiss, F. Lübeck, and G. Malle. The Brauer trees of the exceptional Chevalley groups of type E_6 . *Manuscripta Math.*, 87(1):131–144, 1995.

-
- [26] M. Linckelmann. Derived equivalence for cyclic blocks over a p -adic ring. *Math. Z.*, 207(2):293–304, 1991.
- [27] M. Linckelmann. A derived equivalence for blocks with dihedral defect groups. *J. Algebra*, 164(1):244–255, 1994.
- [28] G. Lusztig. Coxeter orbits and eigenspaces of Frobenius. *Invent. Math.*, 38(2):101–159, 1976/77.
- [29] G. Lusztig. Representations of finite Chevalley groups. 39:v+48, 1978. Expository lectures from the CBMS Regional Conference held at Madison, Wis., August 8–12, 1977.
- [30] G. Lusztig. *Characters of reductive groups over a finite field*, volume 107 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1984.
- [31] B. Mitchell. The full imbedding theorem. *Amer. J. Math.*, 86:619–637, 1964.
- [32] L. Puig. Local block theory in p -solvable groups. In *The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979)*, volume 37 of *Proc. Sympos. Pure Math.*, pages 385–388.
- [33] L. Puig. Algèbres de source de certains blocs des groupes de Chevalley. *Astérisque*, (181-182):9, 221–236, 1990.
- [34] J. Rickard. Derived categories and stable equivalence. *J. Pure Appl. Algebra*, 61(3):303–317, 1989.
- [35] J. Rickard. Lifting theorems for tilting complexes. *J. Algebra*, 142(2):383–393, 1991.
- [36] J. Rickard. Splendid equivalences: derived categories and permutation modules. *Proc. London Math. Soc. (3)*, 72(2):331–358, 1996.
- [37] R. Rouquier. The derived category of blocks with cyclic defect groups. In *Derived equivalences for group rings*, volume 1685 of *Lecture Notes in Math.*, pages 199–220. Springer, Berlin, 1998.
- [38] R. Rouquier. Complexes de chaînes étales et courbes de Deligne-Lusztig. *J. Algebra*, 257(2):482–508, 2002.
- [39] T. A. Springer. Regular elements of finite reflection groups. *Invent. Math.*, 25:159–198, 1974.
- [40] R. Steinberg. *Lectures on Chevalley groups*. Yale University, New Haven, Conn., 1968. Notes prepared by John Faulkner and Robert Wilson.