



HAL
open science

New mutations in the RAB3GAP1 gene in patients with Warburg MICRO Syndrome and a possible founder effect in the Danish population

Deborah J Morris-Rosendahl, Reeval Segel, A. Peter Born, Christoph Conrad, Bart L. Loeys, Susan Sklower Brooks, Laura Müller, Christine Zeschnigk, Christina Botti, Ron Rabinowitz, et al.

► To cite this version:

Deborah J Morris-Rosendahl, Reeval Segel, A. Peter Born, Christoph Conrad, Bart L. Loeys, et al.. New mutations in the RAB3GAP1 gene in patients with Warburg MICRO Syndrome and a possible founder effect in the Danish population. *European Journal of Human Genetics*, 2010, 10.1038/ejhg.2010.79 . hal-00540039

HAL Id: hal-00540039

<https://hal.science/hal-00540039>

Submitted on 26 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Warburg MICRO Syndrome in patients from different ethnic backgrounds: new mutations in the *RAB3GAP1* gene and a possible founder effect in the Danish population

Deborah J. Morris-Rosendahl¹, Reeval Segel², A. Peter Born³, Christoph Conrad⁴, Bart Loeys⁵, Susan Skower Brooks⁶, Laura Müller¹, Christine Zeschnigk¹, Christina Botti⁶, Ron Rabinowitz⁷, Gökhan Uyanik⁸, Marc-Antoine Crocq⁹, Uwe Kraus¹⁰, Ingrid Degen⁴, Fran Fraes¹¹

¹Institute for Human Genetics, University Clinic Freiburg, Germany

²Medical Genetics Institute, Shaare Zedek Medical Center, Jerusalem, Israel

³Department of Paediatrics, Rigshospitalet, Copenhagen, Denmark

⁴Dept. of Child Neurology, Paediatric section, St. Elisabeth Hospital, Neuwied, Germany

⁵Center for Medical Genetics, Ghent University Hospital, Belgium

⁶Department of Pediatrics, Division Genetics, Robert Wood Johnson Medical School, New Brunswick, USA

⁷Ultrasound Unit, Shaare Zedek Medical Center, Jerusalem, Israel

⁸Institut for Human Genetics, University Medical Center, Hamburg-Eppendorff, Germany

⁹Centre Hospitalier, Rouffach, France

¹⁰Epilepsiezentrum Kehl-Kork, Kork, Germany

¹¹Paediatric Neurology, University of Gent, Belgium

Correspondence to: PD Dr. Deborah Morris-Rosendahl
Institute for Human Genetics
University Clinic Freiburg
Breisacherstr. 33
D-79106 Freiburg
Germany
Tel.: +49-761-270 7027

Fax: +49-761 270 7041

Email: deborah.morris-rosendahl@uniklinik-freiburg.de

Running Title: *RAB3GAP1* mutations in Warburg MICRO Syndrome

Total word count (excluding abstract, figures, tables and references): 3277

Abstract

Warburg MICRO Syndrome is a rare, autosomal recessive syndrome characterized by microcephaly, microphthalmia, microcornia, congenital cataracts, optic atrophy, cortical dysplasia, in particular corpus callosum hypoplasia, severe mental retardation, spastic diplegia, and hypogonadism. We have found five new mutations in the *RAB3GAP1* gene in seven patients with suspected MICRO Syndrome, from families with Turkish, Palestinian, Danish and Guatemalan backgrounds. A thorough clinical investigation of the patients has allowed the delineation of symptoms which are consistently present in the patients and may aid the differential diagnosis of MICRO Syndrome for patients in the future. All patients had postnatal microcephaly, microphthalmia, microcornia, bilateral congenital cataracts, short palpebral fissures, optic atrophy, severe mental retardation, and congenital hypotonia with subsequent spasticity. Only one patient had microcephaly at birth, highlighting the fact that congenital microcephaly is not a consistent feature of MICRO syndrome. Analysis of the brain MRIs revealed a consistent pattern of polymicrogyria in the frontal and parietal lobes, wide sylvian fissures, a thin hypoplastic corpus callosum and increased subdural spaces. All patients were homozygous for the mutations detected and all mutations were predicted to result in a truncated RAB3GAP1 protein. The analysis of nine polymorphic markers flanking the *RAB3GAP1* gene showed that the mutation c.1410C>A (p.Tyr470X), for which a Danish patient was homozygous, occurred on a haplotype that is shared by the unrelated heterozygous parents of the patient. This suggests a possible founder effect for this mutation in the Danish population.

238 words

Keywords: Warburg Micro Syndrome, *RAB3GAP1* mutation, founder effect

INTRODUCTION

Warburg MICRO Syndrome (WARBM, MIM 600118), also known as MICRO Syndrome, is characterized by symptoms of microcephaly, mental retardation, corpus callosum hypoplasia, diffuse cortical or subcortical atrophy, congenital cataracts, microcornea, microphthalmia, progressive joint contractures with growth failure, and hypothalamic hypogonadism^{1 2}. MICRO Syndrome was first described by Warburg *et al.*³ in a consanguineous Pakistani family with two affected sibs and an affected male cousin. Warburg *et al.*³ delineated it from the similar but distinct syndromes: Cerebrooculofacioskeletal Syndrome (COFS, MIM: 214150), Cockayne Syndrome (CS: CSA, MIM: 216400 and CSB MIM: 133540), Cataract, Microcephaly, Failure to thrive, Kyphoscoliosis syndrome (CAMAK and CAMFAK, MIM 212540), Martsolf Syndrome (MS, MIM 212720), lethal Rutledge syndrome (MIM: 268670) and lethal Neu-Laxova syndrome (256520), which all have microcephaly, severe mental retardation, childhood cataract, contractures and genital hypoplasia as symptoms.

COFS syndrome shares many clinical and cellular similarities with CS in that cultured cells from patients with both these syndromes demonstrate hypersensitivity to ultraviolet radiation due to impaired nucleotide excision repair (NER)^{4,5}. Various forms of COFS and CS type A (CSA) and CS type B (CSB) have been found to be due to mutations in excision-repair cross-complementing genes (*ERCC1, 2, 5* and *6*⁵⁻⁹). Impaired NER has not been demonstrated in patients with MICRO Syndrome². MICRO Syndrome differs from COFS and CS in that patients have less prenatal and postnatal growth deficiency, cortical dysplasia rather than progressive brain atrophy,

preservation of hearing, deficient visual perception despite early cataract removal, and a better prognosis for development and survival². Probably the most difficult differential diagnosis is between MICRO syndrome and MS, which are primarily differentiated by the higher life expectancy of patients with MS syndrome¹⁰.

The molecular genetic basis of MICRO Syndrome was unraveled by the finding of inactivating mutations in the Rab3 GTPase-activating protein, catalytic subunit gene (*RAB3GAP1*, chromosome 2q21.3, MIM *601536) in 12 of 14 families¹¹. Some of the Pakistani families described shared the same mutation¹¹. MICRO Syndrome is extremely rare, and the true incidence is not known. To date, a total of nine different *RAB3GAP1* mutations in patients from 14 different families have been described¹¹⁻¹³. Four of the mutations previously reported are recurrent and all the patients, with the exception of two from South Africa and Mexico, have consanguineous parents originating from the Middle East or Pakistan. Here, we provide a detailed clinical and molecular genetic analysis on seven patients from five families with Turkish, Palestinian, Guatemalan and Danish ethnic backgrounds and five new mutations in *RAB3GAP1*. The parents in the Turkish and Palestinian families are reported to be consanguineous, however the parents of the other two families are not closely related. The Danish patient is homozygous for a haplotype shared by the parents, suggesting a founder effect in the Danish population.

MATERIALS AND METHODS

Informed consent was obtained from the parents of all patients for the molecular genetic analysis of the patients and their parents, the publication of the clinical data, photos and brain MRIs.

DNA was extracted from EDTA blood using the Illustra BACC2 DNA extraction kit (GE Healthcare, Munich, Germany) and from buccal swabs from patient 6 using 5% Chelex 100 Resin (Biorad, Munich, Germany).

PCR and DNA Sequencing

The 24 coding exons of *RAB3GAP1* and at least 30 bp of intronic sequence flanking the exons were amplified by PCR using Taq Polymerase (Qiagen, Hilden, Germany), and cycle sequenced using the ABI Prism BigDye Terminator Cycle Sequencing Ready Reaction Kit Version 1.1 (Applied Biosystems, Darmstadt, Germany). Capillary electrophoresis was performed on an ABI 3100 sequencer (Applied Biosystems, Foster City, CA, USA). Sequence data were analysed using SeqPilot DNA sequence analysis software (JSI, Kippenheim, Germany). The database sequence NM_012233.1 for *RAB3GAP1*, was used as reference sequence and PCR primers were developed in our laboratory (Table 1, supplementary data).

cDNA analysis of the mutation c.648+5G>A in intron 7

RNA was extracted from lymphocytes that had been transformed with Epstein-Barr virus, from patient 2 and his parents, using the RNeasy kit (Qiagen, Hilden, Germany). Reverse transcription was performed using 1 µg RNA, an oligo DT Primer and Superscript III (Invitrogen, Darmstadt, Germany). cDNA was PCR-amplified using primers RT-E5F and RT-E9R2 in exons 5 and 9, respectively, and the PCR product was sequenced using the same primers.

Haplotype analysis of polymorphic markers around RAB3GAP1

DNA from patient 5 and her parents was genotyped for nine microsatellite markers surrounding *RAB3GAP1* on chromosome 2q21.3: D2S267, D2S368, AC110620GT24, AC016725TC23, AC020602CA17, AC012450TG23, AC012450AT37, D2S1334 and D2S2196. The primers used were according to Aligianis *et al.*¹¹. The forward primer of each primer pair was labeled with 6-Fam and fragment analysis was performed via capillary electrophoresis on the ABI 3100 automatic sequencer, followed by analysis using the Genescan program (Applied Biosystems, Foster City, USA).

Test for relatedness

The possible relatedness of the parents of patient 5, and those of the sibs 6 and 7, was tested using the PowerPlex16 Kit (Promega, Madison, USA) for individual identity. The kit contains 15 polymorphic short tandem repeat (STR) markers located on 14 different chromosomes and an additional marker which shows different fragments for the X and Y chromosomes. Two ng of sample DNA was amplified in a multiplex PCR reaction in which the primers were labelled with one of three different

fluorescent labels, and the resulting products were resolved on an ABI3100 automatic DNA sequencer. Fragment analysis and allele calling was performed using the Genescan Software (Applied Biosystems, Foster City, USA).

RESULTS

The clinical findings of the five patients are summarized in Table 2 (Supplementary data). The individual patients are described below and the brain magnetic resonance imaging (MRI) findings are described collectively thereafter.

Patient 1

Patient 1 (Fig. 1A) is the first child of consanguineous Turkish parents. She was born by caesarian section due to the breech position, after a normal pregnancy.

Ophthalmological evaluation showed bilateral microphthalmia and cataracts. Bilateral extirpation of the lenses was performed and glasses were prescribed. Further clinical evaluation showed axial hypotonia and peripheral hypertonia with rigidity of the lower limbs. Hypoplastic labia major were apparent in addition to some facial dysmorphic features including low-set ears, a small mouth, retrognathia, and a prominent forehead with bitemporal hollowing. Microcephaly was not present at birth, however was apparent by age 2 years (between 10th and 25th centile). Laboratory investigations including TORCH serology, and comparative genomic hybridization (array CGH) were all normal.

Mutation analysis in *RAB3GAP1* revealed the patient to be homozygous for the mutation c.1471C>T in exon 15 (Fig. 2E), which results in the nonsense mutation, p.Arg491X. Both parents are heterozygous carriers of the mutation.

Patient 2

This male patient (Fig. 1D, E, F) is the first child of Turkish parents who are first cousins. The pregnancy was uncomplicated and routine ultrasound investigations did not reveal any abnormalities. There was suspicion of a CMV infection during the pregnancy, however the child tested negative for CMV during the first week of life. The results of routine laboratory tests for newborn screening including TORCH serology, EBV serology, and the Griffith test, were all normal. Routine chromosome analysis revealed a normal male karyotype.

Mutation analysis in *RAB3GAP1* revealed homozygosity for the mutation c.648+5G>A in intron 7 (Fig. 2C). Both parents of the patient are heterozygous carriers of the mutation. We performed RT-PCR on RNA from the patient and his parents, followed by PCR and cDNA sequencing using primers in exons 5 and 9. The transcript from the patient showed the complete skipping of exon 7 (Fig. 2D), demonstrating the lack of recognition of the donor splice site, as predicted by two splice prediction programs, “Splice view” and “splice site prediction by neural network” from the Berkeley Drosophila genome project (<http://bioinfo.itb.cnr.it/oriel/splice-view.html> and www.fruitfly.org/seq_tools/splice.html). The transcripts from the parents were heterozygous for one normal transcript including exon 7 and one lacking exon 7. The loss of exon 7 eliminates amino acids 162 to 216 from the *RAB3GAP1* transcript and is predicted to result in a frameshift and stop codon after five altered codons in exon 8.

Patients 3 and 4

The Palestinian parents of both these patients are first degree cousins of Muslim origin. The paternal grandparents are also first cousins. Their oldest son is healthy and another son was born at 24 weeks of gestation and suffers from bronchopulmonary dysplasia and developmental delay, probably related to his premature birth.

In Patient 3 (Fig. 1G), bilateral cataracts were detected prenatally by ultrasound. At age 5 months he was operated on for cataract removal and glaucoma. At age 1 year microphthalmia, microcephaly, hypotonia, and a small phallus were described. Eye examination at 2 years demonstrated optic atrophy. At 7 years he had severe mental retardation, quadriplegia, mild hearing loss and reacted only to tactile stimulus. Chromosome analysis showed a normal male karyotype.

Patient 4 was found to have bilateral cataracts, a low nasal bridge and flat forehead on ultrasound examination at 22 weeks of gestation. His head measurements were normal at this gestational age. He was suspected of having the same syndrome as his brother, and the pregnancy was terminated at 28 weeks gestation. Mutation analysis in *RAB3GAP1* in patients 3 and 4 revealed that they were both homozygous for the mutation c.1786_1789delAAAAG in exon 17 (Fig. 2F). The mutation results in a frame-shift and stop codon: p.Lys596GlufsX613. The parents were not tested for the mutation.

Patient 5

Patient 5 (Fig. 1 H, I) is the first and only child of unrelated, Danish parents. The pregnancy and birth were unremarkable, and the female patient's eye abnormalities: microphthalmia, microcornia, absent lens and optic hypoplasia, were detected soon after birth. All routine postnatal screening including TORCH serology and chromosome analysis were normal. She failed to thrive, was hypotonic and feeding via a gastric tube was necessary from age 3 months; the tube has subsequently been removed. She developed seizures at age two months, was treated with oxcarbazepin and remained seizure-free after the cessation of therapy. She remains severely hypotonic, with slight spasticity and severely impaired vision. She is unable to sit without support.

Mutation analysis in *RAB3GAP1* revealed the patient to be homozygous for the mutation c.1410C>A in exon 15 (Fig. 2B). The mutation is predicted to result in a stop codon, p.Tyr470X and both parents were found to be heterozygous carriers of the mutation. Since the parents reported that they were unrelated at least to the generation of their great-grandparents, we decided to test whether the parents shared a common haplotype as the result of a possible founder effect in the Danish population. Haplotype analysis using nine microsatellite markers flanking *RAB3GAP1* showed that the affected child was homozygous for a haplotype shared by the heterozygous parents (Fig. 3). The analysis with 15 polymorphic STR markers (PowerPlex16 kit) located on chromosome arms other than chromosome 2q, showed that the parents shared only 6 of the 50 different alleles observed at the 15 loci (12%), thus confirming that the parents were not closely related.

Patients 6 and 7

Patients 6 (Fig. 1B) and 7 (Fig. 1C) are the first and second children of Spanish-speaking Guatemalan parents residing in the USA. The parents reported that they are unrelated, however the same analysis as described above for the parents of patient 5, using 15 polymorphic STR markers revealed that they shared 10 of the 42 observed alleles (23,8%), which may suggest that they are distantly related. Male Patient 6 was clinically diagnosed with MICRO Syndrome at five months. Pregnancy and delivery were unremarkable. Microcephaly, constricted pupils, excessive lacrimation and microphallus were evident at 2,5 months. Postnatal testing including TORCH serology, chromosomes and analysis of the 607 BAC array were normal. Cataract extraction was performed at six months. He showed failure to thrive with oral/pharyngeal dysphagia and required PEG feeding. He has cryptorchidism and a micropenis, is severely mentally retarded, he does not visually track, and at age 3 years 7 months is unable to sit unsupported.

MICRO Syndrome was suspected prenatally in patient 7 after the detection of bilateral cataracts. At age one year she shows profound developmental delay and cannot support her head. Mutation analysis in *RAB3GAP1* in both patients 6 and 7 showed that both were homozygous for the insertion/deletion mutation: c.264_270delAAAGGATinsTTATTA in exon 4 (Fig. 2A). The mutation results in a frameshift and stop codon, p.Lys89TyfsX90. Both parents of the patients are heterozygous for the mutation.

Brain MRI findings

Selected images from brain magnetic resonance from all patients except patient 4 (foetus) are shown in Figure 4. The brains of all patients showed a relatively uniform combination of cortical malformations including: a wider than normal sylvian fissure, as shown in patients 1, 2, 3 and 5 (Figs. 4A, C, E, I and O) and polymicrogyria in the frontal and parietal regions (Figs. 4C, G, K, O and S). In patient 3, the extensive polymicrogyria extending over the frontal, parietal and temporoal lobes was most clearly apparent (Fig. 4I), as was severe cerebellar hypoplasia (Fig. 4J). All patients had a thin corpus callosum (Figs. 4B, F, J, N, R) with the rostrum and splenium absent in all patients except patient 3. Mild cerebellar vermis hypoplasia was apparent in patients 2 (Fig. 4F), 5 (Fig. 4N) and 6 (Fig. 4R). All patients had increased subdural spaces, in particular the subarachnoid space on both temporal poles (Fig. 4P and T) and delayed myelinisation was apparent in patients 1, 2, and 5 (Fig. 4C, G, O).

DISCUSSION

One of the challenges for clinicians presented with patients with the combination of postnatal growth retardation, microcephaly, microphthalmia, cataracts and hypogonadism (when apparent), is the differential diagnosis and the direction for molecular genetic analysis. In order to define which symptoms appear to be consistently present in patients with MICRO Syndrome, we have extended the table of Graham *et al.*² with the clinical findings in previous cases of MICRO Syndrome and our patients (Table 3, supplementary data). Importantly, only one of our patients had microcephaly at birth. Microcephaly appeared to develop within the first year of life and was accompanied by general growth deficiency, so that postnatal growth deficiency is a distinguishing feature of Micro Syndrome. This is consistent with the findings of Ainsworth *et al.*¹⁴ and Graham *et al.*². Two of the patients described by Mégabarné *et al.*¹⁵ were microcephalic at birth, however congenital microcephaly is not a consistent feature of MICRO Syndrome. Microphthalmia, microcornia, bilateral congenital cataracts, short palpebral fissures, optic atrophy, severe mental retardation, and congenital hypotonia with subsequent spasticity, were present in all patients. The above eye pathology, together with the soft features of deep set eyes, a wide nasal bridge and relatively narrow mouth comprise common recognizable facial features. Hypogonadism was apparent in all male patients and one female patient; in the remaining two female patients it was uncertain. Skeletal malformations were variable: hallux valgus and over-extended and overlapping toes were reported in three of our patients and was shown in one of the patients reported by Graham *et al.*². Clinodactyly, camptodactyly and arachnodactyly have been reported in other patients^{2,12}.

Perhaps one of the most recognizable features of MICRO Syndrome, however, is the brain phenotype. All our patients showed a relatively consistent pattern of delayed myelinisation, polymicrogyria of the frontal and parietal lobes, wide sylvian fissures, corpus callosum hypogenesis and increased subdural spaces, particularly frontally and at the temporal poles. The brain has a particularly characteristic shape in axial slices, which is also apparent in the patients illustrated by Nassogne *et al.*¹⁶, Graham *et al.*² and patients 2 and 6 shown by Abdel-Salam *et al.*¹². Cerebellar vermis hypoplasia and cisterna magna were variably present.

Most mutations described to date have been found in Pakistani individuals and two recurrent mutations have been identified, one of which (c.2801delC) appeared to occur on a common haplotype¹¹. Two additional mutations have recurred in Turkish patients (c.748+1G>A), and Lebanese and Egyptian patients (c.2011C>T). The previously reported mutations have all occurred in patients with consanguineous parents from Middle Eastern ethnic backgrounds, with the exception of a patient from South Africa whose ethnic background was not provided, and one patient from Mexico¹¹. The patients in this report are from diverse ethnic backgrounds and the parents of four of the patients are consanguineous. The parents of patients 6 and 7 are reported not to be consanguineous, however they shared 10 of 42 (23,8%) of the different alleles observed between them for 15 different polymorphic STR markers on chromosome arms other than 2q. Although this is not an effective test for relatedness, the results indicate that the parents may be distantly related. The Danish parents of patient 5, however, shared only 6 of 50 (12%) of the different observed alleles between them and are certain that their great-grandparents were

not related. They nevertheless shared a common haplotype for markers surrounding the *RAB3GAP1* gene, for which their affected daughter is homozygous, thus suggesting a founder effect for the mutation c.1410C>A in the Danish population. Such a founder effect has previously been reported in the Danish population for a mutation in the *OPA1* gene in patients with dominant optic atrophy¹⁷, for *MLH1* and *MSH6* mutations in Lynch Syndrome¹⁸, and the Cys282Tyr *HFE* mutation in hereditary hemochromatosis¹⁹. A higher incidence of MICRO Syndrome in the Danish population has so far not been reported, however the targeted analysis of the c.1410C>A mutation in further Danish patients with suspected MICRO Syndrome should help to validate our finding.

The Rab family of small G proteins are regarded as principal classes of GTPases that play an important role in the regulation of intracellular vesicle transport^{20,21}. The Rab3 subfamily is associated with secretory granules or vesicles, and plays a crucial role in regulated exocytosis^{20,22,23}. Rab3A is the most abundant isoform in brain, where it is localized on synaptic vesicles, and evidence is accumulating that it plays a key role in neurotransmitter release and synaptic plasticity²⁴. RAB3GAP, which converts active Rab3-GTP to inactive Rab3-GDP, thereby regulating Rab3 activity, is a heterodimeric complex consisting of a 130 kDa catalytic subunit encoded by *RAB3GAP1*, and a 150 kDa non-catalytic subunit RAB3GAP2^{25,26}. The catalytic activity of RAB3GAP1 is localized in the C-terminal region (codons 601-981²⁵). The mutations in *RAB3GAP1* that have so far been detected in patients with MICRO Syndrome are all predicted to result in stop codons and are distributed throughout the coding region from codons 89 to 934¹¹ (and this study). Therefore, all the mutations are predicted to result in a truncated protein either before or within the

region important for catalytic activity. This is in line with the prediction that most *RAB3GAP1* mutations produce a mutant protein without catalytic activity¹¹. This may explain why there appears to be relatively little phenotypic variation between the patients so far described with different mutations in *RAB3GAP1*. A single splice mutation has been found in *RAB3GAP2* in three sibs of a consanguineous Pakistani family with Martsolf Syndrome¹⁰. These patients are able to walk and talk and show much milder ocular and neurodevelopmental defects than those with MICRO Syndrome and mutations in *RAB3GAP1*¹⁰. Currently it is still not clear whether the milder phenotype is due to the nature of the mutation or to possibly less important role of the non-catalytic *RAB3GAP2* subunit of the *RAB3GAP* protein.

There may be additional genes which when mutated cause MICRO Syndrome, since linkage to *RAB3GAP1* and *RAB3GAP2* has been excluded in families with patients with suspected MICRO Syndrome, and kindreds with or without mutations in *RAB3GAP1* were clinically indistinguishable^{11,12}. However, the addition of the seven patients with MICRO Syndrome and mutations in *RAB3GAP1* in this report to the previously described patients with a diagnosis of Micro Syndrome provides an even clearer picture of the symptoms essential to MICRO syndrome.

ACKNOWLEDGEMENTS

We are grateful to the parents of the patients in this study for their willing cooperation and Angela Steiert for technical assistance. We thank Dr. Beate Albrecht, Universitätsklinikum Essen, Germany, for stimulating our work on MICRO Syndrome.

Conflict of Interest

The authors declare no conflict of interest.

REFERENCES

1. Derbent M, Agras PI, Gedik S, Oto S, Alehan F, Saatci U: Congenital cataract, microphthalmia, hypoplasia of corpus callosum and hypogenitalism: report and review of Micro syndrome. *Am J Med Genet A* 2004; **128A**: 232-234.
2. Graham JM, Jr., Hennekam R, Dobyns WB, Roeder E, Busch D: MICRO syndrome: an entity distinct from COFS syndrome. *Am J Med Genet A* 2004; **128A**: 235-245.
3. Warburg M, Sjo O, Fledelius HC, Pedersen SA: Autosomal recessive microcephaly, microcornea, congenital cataract, mental retardation, optic atrophy, and hypogenitalism. Micro syndrome. *Am J Dis Child* 1993; **147**: 1309-1312.
4. Meira LB, Graham JM, Jr., Greenberg CR *et al*: Manitoba aboriginal kindred with original cerebro-oculo- facio-skeletal syndrome has a mutation in the Cockayne syndrome group B (CSB) gene. *Am J Hum Genet* 2000; **66**: 1221-1228.
5. Graham JM, Jr., Anyane-Yeboah K, Raams A *et al*: Cerebro-oculo-facio-skeletal syndrome with a nucleotide excision-repair defect and a mutated XPD

- gene, with prenatal diagnosis in a triplet pregnancy. *Am J Hum Genet* 2001; **69**: 291-300.
6. Jaspers NG, Raams A, Silengo MC *et al*: First reported patient with human ERCC1 deficiency has cerebro-oculo-facio-skeletal syndrome with a mild defect in nucleotide excision repair and severe developmental failure. *Am J Hum Genet* 2007; **80**: 457-466.
 7. Cleaver JE, Thompson LH, Richardson AS, States JC: A summary of mutations in the UV-sensitive disorders: xeroderma pigmentosum, Cockayne syndrome, and trichothiodystrophy. *Hum Mutat* 1999; **14**: 9-22.
 8. Nospikel T, Lalle P, Leadon SA, Cooper PK, Clarkson SG: A common mutational pattern in Cockayne syndrome patients from xeroderma pigmentosum group G: implications for a second XPG function. *Proc Natl Acad Sci U S A* 1997; **94**: 3116-3121.
 9. Mallery DL, Tanganelli B, Colella S *et al*: Molecular analysis of mutations in the CSB (ERCC6) gene in patients with Cockayne syndrome. *Am J Hum Genet* 1998; **62**: 77-85.
 10. Aligianis IA, Morgan NV, Mione M *et al*: Mutation in Rab3 GTPase-activating protein (RAB3GAP) noncatalytic subunit in a kindred with Martsolf syndrome. *Am J Hum Genet* 2006; **78**: 702-707.

11. Aligianis IA, Johnson CA, Gissen P *et al*: Mutations of the catalytic subunit of RAB3GAP cause Warburg Micro syndrome. *Nat Genet* 2005; **37**: 221-223.
12. Abdel-Salam GM, Hassan NA, Kayed HF, Aligianis IA: Phenotypic variability in Micro syndrome: report of new cases. *Genet Couns* 2007; **18**: 423-435.
13. Yuksel A, Yesil G, Aras C, Seven M: Warburg Micro syndrome in a Turkish boy. *Clin Dysmorphol* 2007; **16**: 89-93.
14. Ainsworth JR, Morton JE, Good P *et al*: Micro syndrome in Muslim Pakistan children. *Ophthalmology* 2001; **108**: 491-497.
15. Megarbane A, Choueiri R, Bleik J, Mezzina M, Caillaud C: Microcephaly, microphthalmia, congenital cataract, optic atrophy, short stature, hypotonia, severe psychomotor retardation, and cerebral malformations: a second family with micro syndrome or a new syndrome? *J Med Genet* 1999; **36**: 637-640.
16. Nassogne MC, Henrot B, Saint-Martin C, Kadhim H, Dobyns WB, Sebire G: Polymicrogyria and motor neuropathy in Micro syndrome. *Neuropediatrics* 2000; **31**: 218-221.
17. Thiselton DL, Alexander C, Morris A *et al*: A frameshift mutation in exon 28 of the OPA1 gene explains the high prevalence of dominant optic atrophy in the Danish population: evidence for a founder effect. *Hum Genet* 2001; **109**: 498-502.

18. Nilbert M, Wikman FP, Hansen TV *et al*: Major contribution from recurrent alterations and MSH6 mutations in the Danish Lynch syndrome population. *Fam Cancer* 2009; **8**: 75-83.
19. Milman N, Pedersen P: Evidence that the Cys282Tyr mutation of the HFE gene originated from a population in Southern Scandinavia and spread with the Vikings. *Clin Genet* 2003; **64**: 36-47.
20. Takai Y, Sasaki T, Matozaki T: Small GTP-binding proteins. *Physiol Rev* 2001; **81**: 153-208.
21. Zerial M, McBride H: Rab proteins as membrane organizers. *Nat Rev Mol Cell Biol* 2001; **2**: 107-117.
22. Geppert M, Sudhof TC: RAB3 and synaptotagmin: the yin and yang of synaptic membrane fusion. *Annu Rev Neurosci* 1998; **21**: 75-95.
23. Takai Y, Sasaki T, Shirataki H, Nakanishi H: Rab3A small GTP-binding protein in Ca(2+)-dependent exocytosis. *Genes Cells* 1996; **1**: 615-632.
24. Sakane A, Miyoshi J, Takai Y, Sasaki T: Analysis on the emerging role of Rab3 GTPase-activating protein in Warburg Micro and Martsof syndrome. *Methods Enzymol* 2008; **438**: 131-139.

25. Fukui K, Sasaki T, Imazumi K, Matsuura Y, Nakanishi H, Takai Y: Isolation and characterization of a GTPase activating protein specific for the Rab3 subfamily of small G proteins. *J Biol Chem* 1997; **272**: 4655-4658.
26. Nagano F, Sasaki T, Fukui K, Asakura T, Imazumi K, Takai Y: Molecular cloning and characterization of the noncatalytic subunit of the Rab3 subfamily-specific GTPase-activating protein. *J Biol Chem* 1998; **273**: 24781-24785.
27. Rodriguez Crido G, Rufo M, Gómez de Terreros I. A second family with Micro Syndrome. *Clin Dysmorphol* 1999;8:241-45.

Figure Legends

Figure 1. Photos of patients described in this study. **A:** Patient 1 at age 2 years 4 months, **B:** Patient 6 at age 5 years 6 months, **C:** Patient 7 (sister of Patient 6) at age 11 months, **D-F:** Patient 2 at age 1 year (D) and at 4 years 9 months (E). The patient's overlapping toes are shown in F, **G:** Profile of patient 3 at age 7 years, **H-I:** Patient 5, with her hyperextended big toes shown in I. Consistent facial features appear to be low set ears (apparent in A, D/E,) a wide nasal bridge, which may be due to the patients' small eyes (A, C, D/E,); anteverted nares of variable severity (C, G), a relatively narrow mouth with thin lips (A, B, C); hypertrichosis is apparent in patients 1 and (A and B).

Figure 2. DNA sequence of the new *RAB3GAP1* mutations found in this study. **A:** mutation c.264_270delAAGGATinsTTATTA in exon 4, found in patients 6 and 7; **B:** mutation c.1410C>A in exon 15 in patient 5; **C-D:** mutation c. 648+5G>A in intron 7, found in patient 2. The skipping of exon 7 in cDNA of patient 2 is shown in D; **E:** mutation c.1471C>T in exon 15 in patient 1; **F:** mutation c.1786_1789delAAAG in exon 17 found in patients 3 and 4.

Figure 3. Haplotype analysis of markers surrounding *RAB3GAP1* in patient 5 and her parents. The allele carrying the c.1410C>A mutation is shown in italics.

Figure 4. Brain magnetic resonance images (MRI) of patients 1 to 7, excluding patient 4 (foetus). Images **A-D** of patient 1 at age six months, images **E to H** of patient 2 at age nine months, images **I to L** of patient 3 at age 7 years 3 months, images **M to P** of patient 5 at age 15 months, images **Q to T** of patient 6 at age three months, images **U to X** of patient 7 (sister of patient 6) at postnatal day 1. A suitable parasagittal image was not available for patient 5. In images A, E, I Q and U the polymicrogyria around the frontal lobes, and in images A, E, I and Q, the wider-than-normal sylvian fissure, is apparent. The sagittal sections B, F, J, N,R and V show a consistently thin corpus callosum, shortened at the rostrum and genu (with the exception of patient 3, image J), cerebellar hypoplasia in patient 3 (J) and mild cerebellar vermis hypoplasia in patients 5, 6 and 7 (images N, R and V, respectively). The brainstem appears normal in all the patients. Increased subdural spaces are particularly apparent at the temporal poles, as shown in patients 5 and 7 (images M and X, respectively).

Figure 1






Figure 2





Fig. 3

D2S267	397	382
D2S368	245	231
AC110620GT24	444	436
AC016725TC23	169	173
AC020602CA17	213	207
RAB3GAP1 c.1410	C	A
AC012450TG23	217	220
AC012450AT37	265	292
D2S1334	298	290
D2S2196	241	243



378	382	D2S267
227	231	D2S368
436	436	AC110620GT24
171	173	AC016725TC23
205	207	AC020602CA17
C	A	RAB3GAP1 c.1410
217	220	AC012450TG23
292	292	AC012450AT37
290	290	D2S1334
241	243	D2S2196

D2S267	382	382
D2S368	231	231
AC110620GT24	436	436
AC016725TC23	173	173
AC020602CA17	207	207
RAB3GAP1 c.1410	A	A
AC012450TG23	220	220
AC012450AT37	292	292
D2S1334	290	290
D2S2196	243	243

Patient 5

Figure 4

