



Comparison of three template preparation methods for routine detection of beak and feather disease virus and avian polyomavirus with single and nested PCR in clinical specimens

Oldrich Tomasek, Oldrich Kubicek, Viktor Tukac

► To cite this version:

Oldrich Tomasek, Oldrich Kubicek, Viktor Tukac. Comparison of three template preparation methods for routine detection of beak and feather disease virus and avian polyomavirus with single and nested PCR in clinical specimens. *Avian Pathology*, 2008, 37 (02), pp.145-149. 10.1080/03079450801902047 . hal-00540110

HAL Id: hal-00540110

<https://hal.science/hal-00540110>

Submitted on 26 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Comparison of three template preparation methods for routine detection of beak and feather disease virus and avian polyomavirus with single and nested PCR in clinical specimens

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2007-0100.R2
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	23-Nov-2007
Complete List of Authors:	Tomasek, Oldrich; University of Veterinary and Pharmaceutical Sciences Brno, Avian and Exotic Animal Clinic Kubicek, Oldrich; National Institute for Nuclear, Chemical and Biological Protection Tukac, Viktor; University of Veterinary and Pharmaceutical Sciences Brno, Avian and Exotic Animal Clinic
Keywords:	APV, budgerigar fledgling disease, DNA isolation, PBFD
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
Backup of cavp-2007- 0100 style edited.wbk	

SCHOLARONE™
Manuscripts




Figure 1
Limits of single PCR (A) and nested PCR (B) detection of APV DNA tested with serial 10-fold dilutions of template DNA. Lanes 1 to 6, 10⁴ to 10⁻¹ copies/μL of template DNA. Lane 7, positive control; lane 8, negative control; M, weight marker.