



Comparison of three template preparation methods for routine detection of beak and feather disease virus and avian polyomavirus with single and nested PCR in clinical specimens

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2007-0100.R2
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	23-Nov-2007
Complete List of Authors:	Tomasek, Oldrich; University of Veterinary and Pharmaceutical Sciences Brno, Avian and Exotic Animal Clinic Kubicek, Oldrich; National Institute for Nuclear, Chemical and Biological Protection Tukac, Viktor; University of Veterinary and Pharmaceutical Sciences Brno, Avian and Exotic Animal Clinic
Keywords:	APV, budgerigar fledgling disease, DNA isolation, PBFD
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
Backup of cavp-2007- 0100 style edited.wbk	

SCHOLARONE™
Manuscripts




Figure 1
Limits of single PCR (A) and nested PCR (B) detection of APV DNA tested with serial 10-fold dilutions of template DNA. Lanes 1 to 6, 10⁴ to 10⁻¹ copies/μL of template DNA. Lane 7, positive control; lane 8, negative control; M, weight marker.