

Inhibition of 2A-mediated 'Cleavage' of Certain Artificial Polyproteins Bearing N-terminal Signal Sequences.

Martin Denis Ryan, Garry Alec Luke, Jeremy Brown, Pablo de Felipe

▶ To cite this version:

Martin Denis Ryan, Garry Alec Luke, Jeremy Brown, Pablo de Felipe. Inhibition of 2A-mediated 'Cleavage' of Certain Artificial Polyproteins Bearing N-terminal Signal Sequences.. Biotechnology Journal, 2009, 5 (2), pp.213. 10.1002/biot.200900134. hal-00540527

HAL Id: hal-00540527 https://hal.science/hal-00540527

Submitted on 27 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biotechnology Journal



Biotechnology Journal

Inhibition of 2A-mediated 'Cleavage' of Certain Artificial Polyproteins Bearing N-terminal Signal Sequences.

Journal:	Biotechnology Journal
Manuscript ID:	biot.200900134.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	01-Sep-2009
Complete List of Authors:	Ryan, Martin; Uni. St. Andrews, School of Biology Luke, Garry; University of St Andrews, Biology Brown, Jeremy; Uni. Newcastle, The Medical School de Felipe, Pablo; University of St Andrews, Biology
Keywords:	2A, polyprotein, targeting, slipstream, skipping



Research Article ((7047 words)) Inhibition of 2A-mediated 'Cleavage' of Certain Artificial Polyproteins Bearing N-terminal Signal Sequences.

AUTHORS: Pablo de Felipe^{1 §}, Garry A. Luke¹, Jeremy D. Brown² and Martin D. Ryan^{1*}

¹ Centre for Biomolecular Sciences, North Haugh, University of St. Andrews, KY16 9ST, St. Andrews, Scotland, U.K.

² RNA Biology Group and Institute for Cell and Molecular Biosciences, The Medical School, Newcastle University, NE2 4HH, Newcastle upon Tyne, U.K.

[§]Present address : Spanish Medicines Agency (AEMPS), Parque Empresarial "Las Mercedes", Campezo 1 - Edificio 8, 28022 Madrid, SPAIN.

Keywords : 2A, polyprotein, targeting, slipstream, skipping

*Correspondence:

Prof. Martin D. Ryan.

Centre for Biomolecular Sciences,

North Haugh,

University of St. Andrews,

KY16 9ST, St. Andrews,

Scotland, U.K.

Tel. 01334 463403

Facs. 01334 462595

e-mail: martin.ryan@st-and.ac.uk

ABBREVIATIONS

ACT, adoptive cell therapies; cDNA, complementary DNA; CMV, cytomegalovirus; DMEM, Dulbecco's Modified Eagle's Medium; DsRed, Discosoma sp red fluorescent protein; EYFP, enhanced yellow fluorescent protein; ECFP, enhanced cyan fluorescent protein; GFP, green fluorescent protein; ER, endoplasmic reticulum; eRF1, eukaryotic release factor 1, eRF3, eukaryotic release factor 3; GT, β-1,4 galactosyltransferase; HA, hemagglutinin; IDUA, iduronidase alpha-L; IL-12, interleukin-12; NA, neuraminidase; ORF, open reading frame; **PAC**, puromycin-resistance; **PAGE**, polyacrylamide gel electrophoresis; PCR, polymerase chain reaction; SDS, sodium dodecyl sulphate; tRNA, transfer RNA.

ABSTRACT

Where 2A oligopeptide sequences occur within open reading frames (ORFs), the formation of the glycyl-prolyl peptide bond at the C-terminus of (each) 2A does not occur. This property can be used to concatenate sequences encoding several proteins into a single ORF: each component of such an artificial polyprotein is generated as a discrete translation product. 2A and '2A-like' sequences have become widely utilised in biotechnology and biomedicine. Individual proteins may also be co- and post-translationally targeted to a variety of subcellular sites. In the case of polyproteins bearing N-terminal signal sequences we observed, however, that the protein downstream of 2A (no signal) was translocated into the endoplasmic reticulum (ER). We interpreted these data as a form of "slipstream" translocation: downstream proteins, without signals, were translocated through a translocon pore already formed by the signal sequence at the N-terminus of the polyprotein. Here we show this effect is, in fact, due to inhibition of the 2A reaction (formation of fusion protein) by the C-terminal region (immediately upstream of 2A) of some proteins when translocated into the ER. Solutions to this problem include the use of longer 2As (with a favourable upstream context), or, modifying the order of proteins comprising polyproteins.

1 INTRODUCTION

2A and '2A-like' oligopeptide (20-30aa) sequences ('2As') are widely used in biotechnology and biomedicine to co-express multiple different proteins within the same cell. Genes encoding proteins are linked *via* 2A sequences to form a single open reading frame (ORF). Translation of the single ORF results, however, in the biosynthesis of each protein as a discrete translation product [1-10]. The great utility here is that multiple proteins comprising heteromultimers, macromolecular assemblies, biochemical pathways *etc.*, plus (selectable) marker proteins may be assembled into, and co-expressed from, a single transgene. Since 2As function in all eukaryotic systems tested to date, they have found utility in animal, plant and microbial biotechnologies (http://www.st-andrews.ac.uk/ryanlab/Index.htm and reviewed in [11,12]). Notably, this system has recently been used in adoptive cell therapies (ACT) [13-18], genetic engineering of human stem cells [19-21] and the co-expression of transcription factors in the induction of pluripotent stem cells [22-27].

2As are neither proteolytic elements nor substrates for cellular proteinases, but mediate a newly discovered form of translational recoding event referred-to variously as ribosome 'skipping', 'stop-go' and 'stop-carry on' translation [5,8,9]. When a ribosome encounters 2A within an ORF it 'skips' the synthesis of the glycyl-prolyl peptide bond at the C-terminus of 2A. The nascent protein is released from the ribosome by eukaryotic release (termination) factors 1 and 3 (eRF1, eRF3), 2A forming its C-terminus [9]. The ribosome then resumes translation of the downstream sequences. As one test of our intra-ribosomal, co-translational, model for the mechanism of the 2A reaction, we assembled cDNA encoding a synthetic polyprotein comprising a protein targeted to the exocytic pathway (prepro-alpha factor; α F), 2A, and green fluorescent protein (GFP). Expression in yeast showed that the translation product upstream of 2A was, indeed, targeted to the ER whilst the downstream product

Biotechnology Journal

localised to the cytosol. The 2A reaction occurred even though the nascent protein was 'shielded' from cytosolic proteinases by the establishment of a ribosome:translocon complex [7].

To extend the utility of 2A in the co-expression of proteins targeted to different subcellular compartments, we constructed plasmids encoding a panel of polyproteins similar to those used in our yeast analyses. Our 'basic' polyprotein construct comprised enhanced yellow fluorescent protein (EYFP), 2A, enhanced cyan fluorescent protein (ECFP), a second 2A and puromycin resistance (PAC; plasmid pPDF20). All proteins were co-expressed and located to the cytoplasm [28]. This construct was modified by the insertion of a GalT type II signalanchor sequence (GT) onto the N-terminus, to encode a [GT-EYFP-2A-ECFP-2A-PAC] polyprotein (plasmid pPDF18). When analysed using translation systems *in vitro* the expected translation profile was observed: high-level 'cleavage' (>90%) at both of the 2As, producing the major translation products [GT-EYFP-2A], [ECFP-2A] and PAC. When this construct was used to transfect HeLa cells, however, whilst the fluorescence signal from EYFP (correctly) localised in the Golgi, the signal from ECFP localised to the ER [28]. The latter was unexpected since [ECFP-2A] did not bear a signal sequence. Furthermore, this result was at variance with our findings in yeast, where the protein downstream of 2A localised to the cytosol and was also at variance with reports in the literature where secreted heterodimers were co-expressed using 2A (discussed below).

Our conclusion from the data derived from the fluorescent proteins bearing N-terminal co-translational signal sequences (types I & II) was that (i) translation of the first protein (bearing a signal sequence) lead to the formation of the ribosome:translocon (Sec61) complex, (ii) the N-terminal protein was translocated into the ER and (iii) that the protein downstream of 2A (no signal sequence) simply "slipstreamed" through the pore of the translocon complex already formed [28].

Here we present data that shows this interpretation to be incorrect. We show that an interaction between the C-terminal region of *certain* nascent peptides and the translocon complex can affect the structure of the C-terminus of 2A within the peptidyl-transferase centre of the ribosome. This leads to inhibition of the 2A reaction, greatly increasing peptide bond formation and the production of 'uncleaved' fusion proteins. The fluorescence patterns we observed were primarily due, therefore, to the localisation and, to some extent, the fluorescent properties of these uncleaved forms.

these unctear.

2 MATERIALS AND METHODS

2.1 Cloning. Plasmids pPDF18, pPDF19, pPDF20 and pPDF67 were as described previously [28]. Plasmids encoding the fluorescent proteins EYFP and ECFP and the GT signal were purchased from Clontech. The intermediate plasmid pPDF15 [GT-EYFP-2A-PAC] was derived from pPDF9 [GFP-2A-PAC] [28], the GFP removed with NheI/XbaI and replaced with GT-ECFP amplified from pECFP-Golgi (Clontech) using the primers PDFK (5' -CAGATCCGCTAGCATGAGGCTTCGGGAG-3') and 3' egfp/XbaI (5'-GGCCTCTAGAC TTGTACAGCTCGTCCAT-3'). The ECFP was then removed by restriction with BamHI/XbaI and replaced with the PCR product EYFP (restricted with BglII and XbaI) from pEYFP-C1 (Clontech) using the primers PDFN (5'-GGCAGCAGATCTGGCTAG CATGAGGGTGAGCAAGGGCGAG-3') and 3' egfp/ XbaI. To obtain plasmid pPDF45 Δ B the 2A-GT-ECFP 'cassette' was first removed from pPDF19 by XbaI restriction then ligated into pTG394pac [29] doubly restricted with XbaI and SpeI. The 3' pac-2A-GT-ECFP portion of the insert was then removed by BglII restriction, blunt-ended (Klenow fill-in) and restricted with SacII. The restriction fragment was then ligated into pPDF15 restricted with XhoI, bluntended (Klenow fill-in), and SacII to produce plasmid pPDF41 (encodes [GT-EYFP-2A-PAC-2A-GT-ECFP]). Finally, restrictions with ApaI followed by BamHI removed the pac-2A and GT sequences, respectively.

Yeast signal sequences were amplified as α F-2A and $D_N \alpha$ F-2A fragments from plasmids pp α F-2A-GFP and $D_N \alpha$ F-2A-GFP [7] using the primers JN1 (5'-CCA<u>GCTAGCG</u> AATTCGTCGACCTCGAG-3') and JN2 (5'-AAAACCAT<u>GGGCCCAGGGTTGG-3')</u> [7]. The PCR products were restricted with *NheI/Apa*I and used to replace the GT-EYFP-2A of pPDF45, similarly restricted. The yeast signals were removed from these plasmids with *Nhel/Xba*I and used to replace the GT signal-anchor sequence excised from pPDF45 Δ B with *Nhel/Nhe*I, to give pPDF90 [D_N*a*F-EYFP-2A-ECFP] and pPDF91 [*a*F-EYFP-2A-ECFP], respectively. pPDF45 Δ B was restricted with *Nhe*I and *Xba*I to remove the EYFP and form plasmid pPDF87. To obtain pGL1, pPDF45 Δ B was doubly restricted with *Spe*I and *Xba*I to remove the 3' end of the CMV promoter, the T7 promoter, the GT signal and EYFP. This sequence was replaced with a similar fragment (with the shorter 5' end of EYFP), produced by PCR amplification of pPDF15 using primers PDFY (5'-GACAATTGCATGAAGAATC-3') and PDFX.30 (5'-CTCGCCTCTAGAGCTGAACTTGTGGCCGTTTAC-3'), similarly restricted. pPDF45 Δ B was restricted with *BspEI/Xba*I to remove EYFP, and replaced with a 3' deleted version of the gene generated by PCR using plasmid pPDF15 as the template and oligonucleotides T7 (5'-TAATACGACTCACTATAGGG-3') and PDFX.180 (5'-GTAGTG TCTAGAGAGCTGCACGCTGCCGTCTC-3') to form pAN1.9. Fortuitously, this reverse primer also hybridized 120 nucleotides downstream to give us an additional in-frame deleted version of the EYFP gene (pAN1.1)

Deletions in the EYFP gene in plasmid pPDF67 were introduced by replacing the *Bgl*II/*BsrG* I restriction fragment (containing CMV / T7 promoters, GT signal and EYFP) with equivalent fragments (having 3' deleted versions of EYFP) from pGL1 to obtain pPDF107 and from pPDF87 to obtain pGL3. Plasmid pGL11 was obtained from pPDF67 by deleting the full EYFP gene, plus 2A, with *BstXI/Bam*HI and the insertion of a small PCR fragment of EYFP, plus 2A, (amplified using primers PDFX.30b; 5'-TGGAGT<u>CCAAGGATCTG</u> <u>GCTCACAACGTCTATATCATGGC-3'</u> and SEQ4; 5'-ACCATGGTGGCGACCGGTGG AT-3'), similarly restricted, recreating the EYFP-2A junction present in pPDF116. Plasmid pPDF67 was restricted with *BstXI* and *SgrAI* to delete EYFP, which was replaced with the equivalent fragment from pAN1.9, or pAN1.1, to obtain pPDF115, or pPDF116, respectively.

Progressively smaller 3' deletions in EYFP were introduced in pPDF67 restricted with *BstX*I and *SgrA*I. The deleted EYFP sequences were generated from PCR of pPDF15 by using the primers T7 and PDFX.200 (5' –CAGGGC<u>CACCGGTG</u>CGCATGCTCTAGAGTTGTCGGG CAGCAGCACGGGGGCCG-3') producing pPDF117, or, using primers T7 and PDFX.220 (5'-GCGGC<u>CACCGGTG</u>CGCATGCTCTAGACATGTGATCGCGCTTCTCGTTGGG-3'), to produce pPDF118.

2.2 *In vitro* **translation.** Coupled transcription/translation assays were performed using rabbit reticulocyte lysates (Promega) as described [7].

2.3 Cell culture, transfection and imaging. HeLa cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% foetal calf serum. HeLa cells were seeded, transfected with Genejuice (Novagen), subsequently fixed (24hr), and imaged using DeltaVision microscope system (Applied Precision) as described [28].

2.4 Immunofluorescence. Transfected cells were fixed and then incubated with primary antibodies, either; (a) rabbit polyclonal anti-2A (kind gift of Dr D. Vignali) or (b) mouse monoclonal anti-V5 (kind gift of Prof. R. Randall). Texas red goat anti-rabbit and Texas red goat anti-mouse (Molecular Probes) were used as secondary antibodies.

2.5 Western blotting. HeLa cells were transfected and lysates collected 48hr later. Samples were run in 10% or 12.5% SDS-PAGE gels, transferred to IMMOBILON-P membranes (Millipore). Membranes were probed with primary antibodies, either (a) mouse monoclonal anti-GFP (Roche), (b) anti-V5, or, (c) anti-2A antibodies. Secondary antibodies used were ECL anti-mouse IgG-Peroxidase from sheep (GE Healthcare) and anti-rabbit IgG-Peroxidase

from goat (Sigma). Membranes were developed using ECL Plus Western Blotting Detection System (GE Healthcare).

3 RESULTS

3.1 Sub-cellular Localisation. To simplify certain plasmid constructions and to more closely mimic those constructs analysed in yeast [7], the [2A-PAC] portion of the pPDF18 polyprotein was deleted to create plasmid pPDF45 Δ B, encoding [GT-EYFP-2A-ECFP] (Fig.1A). Transfection of HeLa cells with pPDF45 Δ B produced the same pattern of fluorescence as that obtained with pPDF18: yellow fluorescence in the Golgi and cyan fluorescence in the ER (Fig. 1B). When the GT type II signal-anchor in pPDF45 Δ B was replaced by a type I signal sequence from the ER luminal protein calreticulin, both fluorescence signals were located in the ER (data not shown).

We suspected that the difference between the observations made in yeast and mammalian cells could lie either in; (i) some peculiarity of the signal sequences used in these analyses, or, (ii) the nature of the sequences upstream of 2A. To address the first possibility we replaced the GT Golgi targeting sequence with either of the two signal sequences used in the yeast experiments (α F and Dap2p - D_N α F), obtaining plasmids pPDF90 (encoding [D_N α F-EYFP-2A-ECFP]) and pPDF91 (encoding [α F-EYFP-2A-ECFP]). The yeast α F signal sequence has been shown to function within mammalian cells [30, 31]. Transfection of HeLa cells with these plasmids resulted in a similar pattern of fluorescence: both yellow and cyan signals in the ER. Again, it appeared that in both cases ECFP apparently 'slipstream' translocated into the ER (Fig.1B). This effect was not, therefore, due to the nature of the signal sequences.

Biotechnology Journal

To address the second possibility we analysed a panel of deletions within EYFP. In this, and in all subsequent cases below, the numerals given in plasmid designations indicates the residues present in the EYFP deletion forms. The 2A sequence used in these analyses was derived from plasmid pSTA1/34 [6]. Following the 2A reaction the upstream protein bears, therefore, a C-terminal (2A) extension of 23aa. Initially we deleted all EYFP sequences, creating plasmid pPDF87 (encodes [GT-2A-ECFP]; Fig.2A). We followed the fate of the [GT-2A] protein using an anti-2A monoclonal antibody (kind gift of Dr. Dario A. Vignali). Whilst this antibody recognises 2A C-terminal extension of proteins, it does not recognise 2A within "uncleaved" fusion proteins (Fig.1C). Transfection of HeLa cells with plasmid pPDF87 resulted in [GT-2A] localising to the Golgi and ER, with ECFP diffused throughout the cytoplasm but, inexplicably, the vast majority localising to the nucleus (Fig.2B). Whilst the protein upstream of 2A ([GT-2A]) had entered the exocytic pathway as expected, the protein downstream had not. We wished to confirm ECFP had not entered the exocytic pathway by intensifying the fluorescence signal from ECFP by targeting the protein to the nucleus. A modified construct, pGL3, encoding [GT-2A-ECFP-N-PAC] (Fig.3A) was expressed in HeLa cells. As before, [GT-2A] localised to the Golgi, whilst [ECFP-N-PAC] predominantly localised to the nucleus. In addition, a clear, but weak, signal could now be observed within the ER (Fig. 3B).

To further alter the sequence context immediately upstream of 2A, a range of deletions were made in the C-terminal region of EYFP. Plasmids pGL1 and pPDF107 both encode the same small fragment of the N-terminus of EYFP between the GT signal and the 2A. When either construct was transfected in HeLa cells, the ECFP (or, in the case of pPDF107 [ECFP-N-PAC]) apparently "slipstreamed" in the ER (as with pPDF45 Δ B), while the [GT- Δ EYFP₁. ₃₀-2A] localised to the Golgi (Figs.2A,B and 3A,B). In both cases a substantial amount of the full-length translation product was observed – the 2A reaction was inhibited (see below). Interestingly, when this N-terminal region of EYFP was also deleted (pPDF87, Fig2A; pGL3, Fig.3A), the 2A reaction was no longer inhibited (see below) and the vast majority of the protein downstream of 2A was not translocated into the ER (Figs 2B, 3B).

Plasmid pAN1.9 encodes EYFP with a much smaller deletion at the C-terminus ([GT- Δ EYFP₁₋₁₄₀-2A-ECFP]; Fig.2A). Here, the result obtained was similar to that using pPDF87: while the [GT- Δ EYFP₁₋₁₄₀-2A] product upstream of 2A localised to the Golgi, the downstream product (ECFP) localised to the cytoplasm and nucleus (Fig.2B). A similar plasmid, pPDF115, encodes [GT- Δ EYFP₁₋₁₄₀-2A-ECFP-N-PAC] (Figs.3&4A). Transfection with this plasmid showed the protein [GT- Δ EYFP₁₋₁₄₀-2A] upstream of 2A had, again, entered the exocytic pathway. Although bearing a GT signal, this protein was distributed in both the ER and Golgi. The protein downstream of 2A ([ECFP-N-PAC]) was localised in the nucleus, with a much weaker signal from the ER (Figs.3&4B).

If the context immediately upstream of 2A is crucial in producing this apparent "slipstream" translocation, then small deletions in the C-terminal region of EYFP might be sufficient to reduce or eliminate this effect. A series of constructs were made with progressive deletions within EYFP (plasmids pPDF118, pPDF117, pPDF116 and pPDF115; Fig.4A). In all cases, the protein upstream of 2A was translocated into the ER and the downstream [ECFP-N-PAC] localised to the nucleus (with a weak signal in the ER). The data from all of these plasmids contrasts with the ER localisation of [ECFP-N-PAC] in cases when the C-terminal region of EYFP was present immediately upstream of 2A (plasmid pPDF67; Figs.3&4B). Deletion of just the C-terminal 20aa of EYFP (pPFD118) relieved the inhibition of the 2A reaction.

3.2 Effect of the Upstream Context on the 2A Reaction *In vitro* **and** *In vivo*. We routinely monitor the 'cleavage' activity of 2A-containing artificial polyproteins using coupled Wilev-VCH

Biotechnology Journal

transcription/translation rabbit reticulocyte cell-free translation systems (Figs.1-4, Panel D). Using constructs encoding artificial polyproteins such as [EYFP-2A-ECFP] we had correlated the activities we had observed using *in vitro* translation systems with that observed within transfected cells (analysed by western blotting). In both cases cleavage was high – indeed higher (almost 100%) *in vivo*. We took it, therefore, that the *in vitro* analyses were a good indication of cleavage *in vivo*.

When we investigated the cleavage *in vivo* of the polyproteins we had created that gave rise to the apparent 'slipstream' sub-cellular localisation – those bearing N-terminal signal sequences – we found a very different picture. For example, in the case of pPDF45 ΔB and pPDF67, translation in vitro produced only a small amount of the full-length translation product (Figs. 2-4, Panel D), whilst expression in vivo showed this form to be the major product with the 'cleaved' forms being weak bands – sometimes visible only with prolonged exposure (Figs 2-4, Panel C). The *in vivo* analyses of cleavage showed a major difference in efficiency between those constructs which gave rise to the apparent "slipstream" effect and those which did not. The "slipstream" effect showed complete correlation with low efficiency of the 2A reaction – evidenced by the levels of the full-length translation products (Figs.1-4, panel D). Taken together these results strongly suggest that in the case of proteins targeted to the exocytic pathway the immediate upstream context of 2A may strongly influence the efficiency of the reaction. In the case of pGL1 and pPDF107 the same number of residues lie between the end of the transmembrane domain of the GT signal sequence and the N-terminal residue of 2A (Figs. 2A & 3A, respectively). We designed pGL11 ([GT-∆EYFP₁₅₀₋₁₈₀-2A-ECFP-N-PAC]) to encode the same *number* of residues between the GT signal and 2A as in pGL1/pPDF107, but in pGL11 the upstream context for 2A is identical to that in pPDF116 – a construct that showed a highly efficient 2A reaction (Fig.4C). Upon transfection of cells with plasmid pGL11, the translation product upstream of 2A ([GT- Δ EYFP₁₅₀₋₁₈₀-2A]) was detected

> Wiley-VCH

in the Golgi (immunofluorescence using anti-2A antibodies) whilst the translation product downstream of 2A ([ECFP-N-PAC]) was localised predominantly in the nucleus. The 2A reaction was highly efficient (using in vitro translation systems and in vivo; Fig.3C,D). These data are consistent with the notion that, in the case of proteins entering the exocytic pathway, the context immediately upstream of 2A may strongly inhibit the 2A reaction.

ι g in . . to that, in ι. . tream of 2A may str.

4 DISCUSSION

2A and 2A-like sequences are now widely used as a tool for co-expression in biomedicine and biotechnology due to (i) their shortness (20-30aa), (ii) highly efficient cleavage, (iii) the uniform stoichiometry of the cleavage products and (iv) their activity in all eukaryotic celltypes tested to date. Many laboratories have successfully used 2As to co-express proteins targeted to the exocytic pathway. For example, the p35 and p40 chains of interleukin-12 were cleaved highly efficiently. Both chains bore their native signal sequences and active IL-12 was secreted into the media [32]. Similarly, the heavy and light chains of antibodies cleaved highly efficiently, were assembled and secreted [33]. The α - and β -subunits of the T-cell receptor (in both orientations) cleaved highly efficiently, assembled and were localised to the plasma membrane [13]. Human iduronidase (IDUA) was targeted to the secretory pathway whilst a marker protein downstream of 2A (DsRed2) localised to the cytosol [34].

In our previous study with fluorescent polyproteins where the first protein was targeted to the secretory pathway and the protein downstream of 2A was cytosolic, we attributed the unexpected ER sub-cellular distribution of the downstream protein to "slipstream" translocation [28]. Here we show a complete correlation between those polyproteins displaying the apparent "slipstream" effect and those with low levels of the 2A reaction *in vivo* - monitored by western blotting. A large proportion of the translation products are uncleaved, leading to translocation of the fusion protein into the exocytic pathway.

Analyses of protein targeting using a control construct (pPDF93), encoding [GT-EYFP-ECFP], showed that (i) the fusion protein partitions between the Golgi and ER (mainly ER) and (ii) both proteins fluoresce in both compartments - producing complete co-localisation upon merging the images [28]. However, when 2A is present between the fluorescent proteins (pPDF45 Δ B), whilst the fusion protein partitions between the Golgi and

ER, the [GT-EYFP-2A] product formed by the 2A reaction (strongly fluorescent) localises to the Golgi, whilst we assume the [ECFP] product (much more weakly fluorescent than EYFP) is diffused throughout the cytoplasm and nucleus and not detectable.

In the case of constructs with N-terminal signal sequences, the source of the inhibitory effect of the EYFP C-terminal sequences (immediately upstream of 2A) must lie in the interaction with the translocon complex, since they cleave highly efficiently using *in vitro* translation systems. For example, the [GT-EYFP-2A-ECFP-N-PAC] polyprotein encoded by pPDF67 shows high level of cleavage *in vitro*, while *in vivo* most of the protein detected by anti-GFP antibodies was in the full-length, uncleaved, form [GT-EYFP-2A-ECFP-N-PAC], with very little detected as the cleavage products [GT-EYFP-2A] and [ECFP-N-PAC] (Figs.3,4, Panels C,D).

We have proposed that nascent 2A forms an α -helix with a tight-turn at its C-terminus. The helical portion is proposed to interact with the ribosome exit tunnel such that the C-terminal portion is sterically constrained within the peptidyl-transferase centre of the ribosome [4,5]. The ester linkage between 2A and tRNA^{gly} is precluded from nucleophilic attack by the prolyl-tRNA in the A site, effectively 'jamming' translation. We have recently shown that this block is relieved by the action of translation release factors 1 and 3 [9].

Residues that influence the 2A-mediated 'cleavage' of cytosolic proteins map to within 30aa of the cleavage site. This is consistent with many proteinase protection studies which have shown the length of nascent peptide within the ribosome exit tunnel to be 30-40aa (Fig.5; reviewed in [35]). The eukaryotic ribosome exit tunnel is ~100Å long and has an average diameter of ~20Å [36].

The C-terminal 20aa of EYFP upstream of 2A that inhibited the 2A reaction are 28-48aa distal from the peptidyl-transferase centre - since we know that the C-terminus of 2A (23aa long + 4aa linker) is in the P site when 'cleavage' occurs (Fig.5). Disulphide bond

Biotechnology Journal

formation, photochemical cross-linking, proteinase protection, glycosylation acceptor site and protein folding studies performed upon nascent proteins transiting into the lumen of the ER show that 65-70aa are present within the ribosome:translocon complex [37-42] (Fig.5). The consensus arising from structural studies on Sec61 or Sec61/ribosome complexes is that the protein conducting channel is formed at the interface of an oligomeric assembly - probably a tetramer - of the Sec61 α , γ , β complex. The channel is co-axial with the ribosome exit tunnel and is thought to be some 48-60Å long [43-48].

Theoretical work suggests that nascent proteins adopt an α -helical conformation [49] stabilised by the exit tunnel [50]: indeed, dynamic simulations suggested 2A adopted such a conformation [4]. Taken together, these data suggest that at the stage in elongation when 'cleavage' occurs (the C-terminal glycyl-tRNA^{gly} of 2A in the P site of the ribosome peptidyl-transferase centre), the C-terminal 20aa of the protein upstream (EYFP) most probably lie within a region defined by the interface between the ribosome and Sec61 complex and the translocon tunnel itself (Fig.5). Interactions between the nascent protein and the translocon tunnel may affect the conformation of 2A within the ribosome exit tunnel and, in consequence, the C-terminal tight-turn of 2A in the peptidyl-transferase centre itself – inhibiting the 2A reaction.

The problem we have identified in the co-expression of certain proteins targeted to the exocytic pathway has significance for the biotechnological utilities of 2A. The LOCATE subcellular localisation database indicates that of all human proteins currently analysed, ~40% are either secreted from the cell, located within the lumen /membranes of cytoplasmic vesicular structures (excluding mitochondria), or, are plasma membrane proteins (http://locate.imb.uq.edu.au/). Given that such a high proportion of cellular proteins are initially translocated into the ER, the ability to co-express multiple proteins targeted to such sites is essential.

Biotechnology Journal

In this study we identified two regions of EYFP which, *when placed immediately upstream of 2A*, inhibited the 2A reaction; (i) the C-terminal region (residues 220-240: pPDF118) and (ii) the N-terminal region (residues 1-29: pGL1/pPDF107). However, in constructs we assembled designed to co-express influenza HA or NA linked *via* 2A to fluorescent proteins ([HA-2A-CherryFP] or [NA-2A-CherryFP]), we observed the same effects we describe here: very high-level cleavage using translation systems *in vitro*, whilst expression *in vivo* showed the fluorescent proteins localised in the ER and not to the cytoplasm - again, the apparent 'slipstream' effect.

This inhibition of the 2A reaction may be overcome in two ways. Firstly, by the use of longer versions of 2A which incorporate a 39aa tract from the C-terminus of protein 1D, immediately upstream of 2A in the FMDV polyprotein [6]. This extension does not interact with the translocon pore to affect the activity of 2A in the ribosome - effectively 'insulating' the 2A sequence from the upstream protein. A number of studies suggest that cleavage efficiency may be improved by using a flexible Gly-Ser-Gly or Ser-Gly-Ser-Gly linker sequence separating the upstream protein from the 2A sequence [51-53]. Recent work by Yang and co-workers (2008) combined a furin cleavage site and a V5 oligopeptide spacer sequence immediately upstream of 2A [54]. Furin is a proteinase enriched in the Golgi, with a canonical recognition site of -R-X-(R/K)-R-. For proteins targeted to, or transiting through, the Golgi, incorporation of this site results in proteolysis. This strategy may provide an excellent solution since it removes the extended 2A linker [55]. We have used a linker comprising the furin proteinase cleavage site, 39aa of 1D, plus 2A (derived from plasmid pSTA1/31; [6]) to overcome this problem. In our influenza HA/NA co-expression studies the use of this linker resulted in the correct localisation of the cherry fluorescent protein to the cytoplasm, and not the exocytic pathway (S. Vater, pers.comm.).

The second solution lies in the basic design – the "gene" order – of the polyprotein. Previously described 2A-based multigene vectors have revealed the 2A region functions properly within different contexts, but the cleavage efficiency varies as flanking context changes. By swapping the order of

Biotechnology Journal

proteins in several artificial polyproteins the stoichiometry was affected by the gene upstream of 2A [56, 57]. If a problematic secreted/membrane protein is identified, then this could be incorporated as the C-terminal component of the polyprotein system. It should be noted in this context that in mammalian and yeast cells an N-terminal proline (such as that produced by 2A-mediated 'cleavage') confers a long half-life (>20hrs) on proteins (http://www.expasy.ch/tools/protparam-doc.html).

- [1] Ryan, M.D., King, A.M.Q., Thomas, G.P., Cleavage of foot-and-mouth disease virus polyprotein is mediated by residues located within a 19 amino acid sequence. *J. Gen. Virol.* 1991, 72, 2727-2732.
- [2] Ryan, M.D., Drew, J., Foot-and-mouth disease virus 2A oligopeptide mediated cleavage of an artificial polyprotein. *EMBO J.* 1994, *134*, 928-933.
- [3] Donnelly, M.L.L., Gani, D., Flint, M., Monoghan, S. *et al.*, The cleavage activity of aphtho- and cardiovirus 2A proteins. *J. Gen. Virol.* 1997, 78, 13-21.
- [4] Ryan, M.D., Donnelly, M.L.L., Lewis, A., Mehrotra, A.P. *et al.*, A model for nonstoichiometric, co-translational protein scission in eukaryotic ribosomes. *Bioorganic Chem.* 1999, 27, 55-79.
- [5] Donnelly, M.L.L., Luke, G., Mehrotra, A., Li, X. *et al.*, Analysis of the aphthovirus 2A/2B polyprotein 'cleavage' mechanism indicates not a proteolytic reaction, but a novel translational effect: a putative ribosomal 'skip'. *J. Gen. Virol.* 2001, 82, 1013-1025.
- [6] Donnelly, M.L.L., Hughes, L.E., Luke, G., Li, X. et al., The 'cleavage' activities of FMDV 2A site-directed mutants and naturally-occurring '2A-Like' sequences. J. Gen. Virol. 2001, 82, 1027-1041.
- [7] de Felipe, P. Hughes, L.E. Ryan, M.D., Brown, J.D., Co-translational, intraribosomal cleavage of polypeptides by the foot-and-mouth disease virus 2A peptide. *J. Biol. Chem.* 2003, 278, 11441-11448.
- [8] Atkins, J.F., Wills, N.M., Loughran, G., Wu, C-Y. *et al.*, A case for "StopGo": Reprogramming translation to augment codon meaning of GGN by promoting unconventional termination (Stop) after addition of glycine and then allowing continued translation (Go). *RNA* 2007, *13*, 1-8.

- [9] Doronina, V.A., Wu, C., de Felipe, P., Sachs, M.S. *et al.*, Site-specific release of nascent chains from ribosomes at a sense codon. *Mol. Cell. Biol.* 2008, *28*, 4227- 4239.
- [10] Luke, G.A., de Felipe, P., Lukashev, A., Kallioinen, S.E. *et al.*, The occurrence, function, and evolutionary origins of '2A-like' sequences in virus genomes. *J. Gen. Virol.* 2008, 89, 1036-1042.
- [11] de Felipe, P., Skipping the co-expression problem: the new 2A "CHYSEL" technology. *Genetic Vaccines Ther.* 2004, 2 13,
- [12] de Felipe, P., Luke, G.A., Hughes, L.E., Gani, D. et al., E unum pluribus: multiple proteins from a self-processing polyprotein. Trends Biotechnol. 2006, 5, 68-75.
- [13] Szymczak, A.L., Workman, C.J., Wang, Y., Vignali, K.M. *et al.*, Correction of multigene deficiency in vivo using a single 'self-cleaving' 2A peptide-based retroviral vector. *Nature Biotech.* 2004, 22, 589-94.
- [14] Huang, X., Wilber, A.C., Bao, L., Tuong, D. *et al.*, Stable gene transfer and expression in human primary T cells by the Sleeping Beauty transposon system. *Blood* 2006, *107*, 483-491.
- [15] Scholten, K.B., Kramer, D., Kueter, E.W., Graf, M. et al., Codon modification of T cell receptors allows enhanced functional expression in transgenic human T cells. *Clin. Immunol.* 2006, *119*, 135-145.
- [16] Hart, D.P., Xue, S-A., Thomas, S., Cesco-Gaspere, M. *et al.*, Retroviral transfer of a dominant TCR prevents surface expression of a large proportion of the endogenous TCR repertoire in human T cells. *Gene Therapy* 2008, *15*, 625-631.
- [17] Moisini, I., Nguyen, P., Fugger, L., Geiger. T.L., Redirecting therapeutic T cells against myelin-specific T lymphocytes using a humanized myelin basic protein-HLA-DR2-zeta chimeric receptor. *J. Immunol.* 2008, *180*, 3601-3611.

- [18] Chhabra, A., Yang, L., Wang, P., Comin-Anduix, B. *et al.*, CD4+ CD25- T Cells transduced to express MHC class I-restricted epitope-specific TCR synthesize Th1 cytokines and exhibit MHC class I-restricted cytolytic effector function in a human melanoma model. *J. Immunol.* 2008, *181*, 1063-1070.
- [19] Hasegawa, K., Cowan, A.B., Nakatsuji, N., Suemori, H., Efficient multicistronic expression of a transgene in human embryonic stem cells. *Stem Cells* 2007, 25, 1707-1712.
- [20] Unger, C., Kärner, E., Treschow, A., Stellan, B. *et al.*, Lentiviral-mediated HoxB4 expression in human embryonic stem cells initiates early hematopoiesis in a dose-dependent manner but does not promote myeloid differentiation. *Stem Cells* 2008, *26*, 2455-2466.
- [21] Karner, E., Unger, C., Cerny, R., Ahrlund-Richter, L. *et al.*, Differentiation of human embryonic stem cells into osteogenic or hematopoietic lineages: A dose-dependent effect of osterix over-expression. *J. Cellular Physiol.* 2009, 218, 323-333.
- [22] Okita, K., Nakagawa, M., Hyenjong, H., Ichisaka, T. *et al.*, Generation of mouse induced puripotent stem cells without viral vectors. *Science* 2008, *322*, 949-953.
- [23] Sommer, C.A., Stadtfeld, M., Murphy, G.J., Hochedlinger, K. *et al.*, iPS cell generation using a single lentiviral stem cell cassette. *Stem Cells* 2008, e-pub doi:10.1634/stemcells. 2008-1075
- [24] Carey, B.W., Markoulaki, S., Hanna, J., Saha, K., *et al.*, Reprogramming of murine and human somatic cells using a single polycistronic vector. *PNAS* 2009, *106*, 157-162.
- [25] Shao, L., Feng, W., Sun, Y., Bai, H., *et al.*, Generation of iPS cells using defined factors linked via the self-cleaving 2A sequences in a single open reading frame. *Cell Research* 2009, *19*, 296-306.

Biotechnology Journal

- [26] Woltjen, K., Michael, I.P., Mohseni, P., Desai, R., *et al.*, piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nature* 2009, 458, 715-716.
- [27] Kaji, K., Norrby, K., Paca, A., Mileikovsky, M., *et al.*, Virus-free induction of pluripotency and subsequent excision of reprogramming factors. *Nature* 2009, 458, 771-775.
- [28] de Felipe, P., Ryan, M.D., Targeting of proteins derived from self-processing polyproteins containing multiple signal sequences. *Traffic* 2004, *5*, 616-626.
- [29] de Felipe, P., Martin, V., Cortes, M.L., Ryan, M.D. *et al.*, Use of the 2A sequence from foot-and-mouth disease virus in the generation of retroviral vectors for gene therapy. *Gene Therapy*. 1999, 6, 198-208.
- [30] Garcia, P.D., Peter Walter, P., Full-length prepro-α-factor can be translocated across the mammalian microsomal membrane only if translation has not terminated. J. Cell Biol. 1988, 106, 1043-1048.
- [31] Su, K., Stoller, T., Rocco, J., Zemsky, J. *et al.*, Pre-Golgi degradation of yeast preproalpha-factor expressed in a mammalian cell. Influence of cell type-specific oligosaccharide processing on intracellular fate. *J. Biol. Chem.* 1993, 268, 14301-14309.
- [32] Chaplin, P.J., Camon, E.B., Villarreal-Ramos, B., Flint, M. et al., Production of interlukin-12 as a self-processing polypeptide. J. Interferon Cytokine Res. 1999, 19, 235-241.
- [33] Fang, J.M., Yi, S.L., Simmons, A., Tu, G.H. *et al.*, An antibody delivery system for regulated expression of therapeutic levels of monoclonal antibodies in vivo. *Mol. Therapy* 2007, , 1153-1159.
- [34] Osborn, M.J., Panoskaltsis-Mortari, A., McElmurry, R.T., Bell, S.K. *et al.*, A picornaviral '2A-like' sequence based tricistronic vector allowing for high level therapeutic gene expression coupled to a dual reporter system. *Mol. Ther.* 2005, *12*, 569-574.

- [35] Hardesty, B., Kramer, G., Folding of a nascent peptide on the ribosome. Prog. Nucleic Acid Res. Mol. Biol. 2001, 66, 41-66.
- [36] Morgan, D.G., Ménétret, J.F., Radermacher, M., Neuhof, A. *et al.*, A comparison of the yeast and rabbit 80 S ribosome reveals the topology of the nascent chain exit tunnel, inter-subunit bridges and mammalian rRNA expansion segments. *J. Mol. Biol.* 2000, *301*, 301-321.
- [37] Bergman L.W., Kuehl, W.M., Formation of intermolecular disulfide bonds on nascent immunoglobulin polypeptides. J. Biol. Chem. 1979, 254, 5690-5694.
- [38] Bergman L.W., Kuehl, W.M., Formation of an intrachain disulfide bond on nascent immunoglobulin light chains. J. Biol. Chem. 1979, 254, 8869-8876.
- [39] Mothes, W., Prehn, S., Rapoport, T.A., Systematic probing of the environment of a translocating secretory protein during translocation through the ER membrane. *EMBO J*. 1994, 13, 3973-3982.
- [40] Matlack, K.E., Walter, P., The 70 carboxyl-terminal amino acids of nascent secretory proteins are protected from proteolysis by the ribosome and the protein translocation apparatus of the endoplasmic reticulum membrane. *J. Biol. Chem.* 1995, 270, 6170-6180.
- [41] Whitley, P., Nilsson, I., von Heijne, G., A nascent secretory protein may traverse the ribosome/endoplasmic reticulum translocase complex as an extended chain. J. Biol. Chem. 1996, 271, 6241–6244.
- [42] Kowarik, M., Küng, S., Martoglio, B., Helenius, A., Protein folding during cotranslational translocation in the endoplasmic reticulum. *Mol Cell.* 2002, *10*, 769-78.
- [43] Hanein, D., Matlack, K.E., Jungnickel, B., Plath, K. *et al.*, Oligomeric rings of the Sec61p complex induced by ligands required for protein translocation. *Cell* 1996, 87, 593-595.

- [44] Beckmann, R., Bubeck, D., Grassucci, R., Penczek, P. *et al.*, Alignment of conduits for the nascent polypeptide chain in the ribosome-Sec61 complex. *Science* 1997, 278, 2123-2126.
- [45] Ménétret, J.F., Neuhof, A., Morgan, D.G., Plath, K. et al., The structure of ribosomechannel complexes engaged in protein translocation. Mol. Cell 2000, 6, 1219-1232.
- [46] Beckmann, R., Spahn, C.M.T., Eswar, N., Helmers, J. *et al.*, Architecture of the protein-conducting channel associated with the translating 80s ribosome. *Cell* 2001, *107*, 361–372.
- [47] Morgan, D.G., Ménétret, J.F., Neuhof, A., Rapoport, T.A. *et al.*, Structure of the mammalian ribosome-channel complex at 17Å resolution. *J. Mol. Biol.* 2002, *324*, 871-886.
- [48] Ménétret, J.F., Hegde, R.S., Heinrich, S.U., Chandramouli, P. *et al.*, Architecture of the ribosome-channel complex derived from native membranes. *J. Mol. Biol.* 2005, *348*, 445-557.
- [49] Lim, V.I., Spirin, A.S., Stereochemical analysis of ribosomal transpeptidation. Conformation of nascent peptide. J. Mol. Biol. 1986, 188, 565-574.
- [50] Ziv, G., Haran, G., Thirumalai, D., Ribosome exit tunnel can entropically stabilize αhelices. P.N.A.S. 2005, 102, 18956–18961.
- [51] Holst, J., Szymczak-Workman, A.L., Vignali, K.M., Burton, A.R. *et al.*, Generation of Tcell receptor retrogenic mice. *Nat. Protocols* 2006, 1, 406-417.
- [52] Provost, E., Rhee, J., Leach, S.D., Viral 2A peptides allow expression of multiple proteins from a single ORF in transgenic zebrafish embryos. *Genesis* 2007, 45, 625-629.
- [53] Holst, J., Wang, H., Eder, K.D., Workman, C.J. *et al.*, Scalable signalling mediated by T cell antigen receptor-CD3 ITAMs ensures effective negative selection and prevents autoimmunity. *Nat. Immunol.*, 9, 658-666.

- [54] Yang, S., Cohen, C.J., Peng, P.D., Zhao, Y. *et al.*, Development of optimal bicistronic lentiviral vectors facilitates high-level TCR gene expression and robust tumor cell recognition. *Gene Ther.*, 2008, 15, 1411-1423.
- [55] Fang, J., Qian, J-J., Yi, S., Harding, T.C. *et al.*, Stable antibody expression at therapeutic levels using the 2A peptide. *Nat. Biotech.* 2005, 23, 584-590.
- [56] Ma, C., Mitra, A., Expressing multiple genes in a single open reading frame with the 2A region of foot-and-mouth disease virus as a linker. *Mol. Breeding*, 2002, 9, 191-199.
- [57] Lengler, J., Holzmuller, H., Salmons, B., Gunzburg, W.H. *et al.*, FMDV-2A sequence and protein arrangement contribute to functionality of CYP2B1-reporter fusion protein. *Anal. Bioch.*, 2005, 343, 116-124.

Figure Legends

- Figure 1. Cleavage activities and sub-cellular localisation of products derived from 2Acontaining polyproteins. N-terminal signal sequences (stippled, boxed, areas) were derived from either β -1,4 galactosyltransferase (GT), yeast alpha factor (α F) or the yeast Dap2p signal sequence ($D_N \alpha F$). Single open reading frames were assembled to comprise; (i) an N-terminal signal sequence (GT - pPDF45 Δ B; D_N α F - pPDF90; α F pPDF91), (ii) a fluorescent protein marker (EYFP; open boxed area), (iii) 2A (shaded, boxed, area) and (iv) a second fluorescent protein marker (ECFP; open boxed area) (Panel A). In all cases protein expression was driven by a CMV promoter in the plasmid backbone. HeLa cells were transfected with plasmid constructs and images taken 24hr later (scale bar = $10\mu m$). The GT N-terminal signal-anchor directed EYFP to the Golgi and ECFP to the ER. This was also the case for proteins containing N-terminal yeast signal sequences (Panel B). Cell lysates were prepared 24hr post-transfection, proteins separated by SDS-PAGE and subsequently analysed by Western blotting using antibody probes as indicated (Panel C). Rabbit reticulocyte coupled transcription/translation in vitro systems were programmed with plasmid cDNA constructs, as indicated. The incorporation of ³⁵S-methionine into proteins synthesised *de novo* was determined by SDS-PAGE and phosphorimaging (Panel D).
- Figure 2. Effect of deletion of sequences upstream of 2A. EYPF sequences upstream of 2A were deleted (Panel A). The number of residues between the end of the transmembrane domain of the GT signal sequence and the N-terminal residue of 2A are shown above the appropriate images in panel B. HeLa cells were transfected with plasmid constructs and

images taken 24hr later (scale bar = 10µm). The localisation of EYFP and ECFP was determined by their native fluorescence, whilst the localisation of the deletion forms of EYFP was determined using anti-2A antibodies (Panel B). Cell lysates were prepared, separated by SDS-PAGE and subsequently Western blotted using anti-GFP antibodies (Panel C). *In vitro* translation systems programmed with constructs as indicated and the incorporation of ³⁵S-methionine determined by SDS-PAGE and phosphorimaging. The translation products are denoted as; 1 = [GT-EYFP-2A-ECFP], 2 = [GT- Δ EYFP-2A] and 4 = [ECFP] (Panel D). NOTE: the [GT- Δ EYFP-2A] products from pAN1.9 / pGL1 and the [GT-2A] product from pPDF87 were too small to be resolved in this gel system. Lanes in PAGE analyses of proteins (Panels C & D) were derived from constructs used to produce the cell images shown in panel B, above.

Figure 3. Effect of inclusion of a nuclear localisation signal. To visualise the partitioning of ECFP between the cytoplasm and the ER, we included a nuclear localisation signal (N; together with puromycin resistance – PAC) such that any cytoplasmic [ECFP-N-PAC] would be localised in the nucleus (Panel A). The number of residues between the end of the transmembrane domain of the GT signal sequence and the N-terminal residue of 2A are shown above the appropriate images in panel B. HeLa cells were transfected with plasmid constructs and images taken 24hr later (scale bar = 10μm). The localisation of EYFP and ECFP was determined by their native fluorescence, whilst the localisation of the deletion forms of EYFP was determined using anti-2A antibodies (Panel B). Cell lysates were prepared, separated by SDS-PAGE and subsequently Western blotted using anti-GFP antibodies (Panel C). Rabbit reticulocyte *in vitro* translation systems were programmed with constructs as indicated and the incorporation of ³⁵S-methionine determined by SDS-PAGE and phosphorimaging. Translation products are denoted as; 1

Wiley-VCH

Biotechnology Journal

= [GT-EYFP-2A-ECFP-N-PAC], 2 = [GT- Δ EYFP-2A-ECFP-N-PAC] and 3 = [ECFP-N-PAC]. NOTE: the gel system we used was designed to resolve the [GT- Δ EYFP-2A-ECFP-N-PAC] / [GT-EYFP-2A-ECFP-N-PAC] products from [ECFP-N-PAC], and the [GT- Δ EYFP-2A] products from pPDF115 / pPDF107 / pGL11 and the [GT-2A] product from pGL3 were not resolved in system. Lanes in PAGE analyses of proteins (Panels C & D) were derived from constructs used to produce the cell images shown in panel B, above.

Figure 4. Effects of deletion of sequences immediately upstream of 2A. A panel of constructs were assembled with a nested set of deletions upstream of 2A. As described above, the ECFP bore a nuclear localisation signal (Panel A). The number of residues between the end of the transmembrane domain of the GT signal sequence and the N-terminal residue of 2A are shown above the appropriate images in panel B. Cell lysates were prepared, separated by SDS-PAGE and subsequently Western blotted using anti-GFP antibodies. Rabbit reticulocyte in vitro translation systems were programmed with constructs as indicated and the incorporation of ³⁵S-methionine determined by SDS-PAGE and phosphorimaging. Translation products are denoted as: 1 = [GT-EYFP-2A-ECFP-N-PAC], $2 = [GT-\Delta EYFP-2A-ECFP-N-PAC]$, 3 = [ECFP-N-PAC], 4 = [GT-EYFP-2A] and 5 = [GT- Δ EYFP-2A]. NOTE: the higher molecular masses of the [GT- Δ EYFP-2A] products allowed complete resolution in this gel system. The 'extra' bands arise from internal initiation events in this batch of reticulocyte lysate. Lanes in PAGE analyses of proteins (Panels C & D) were derived from constructs used to produce the cell images shown in panel B, above.

Figure 5. Position of 2A and sequences immediately upstream in the ribosome:translocon complex. In the case of polyproteins bearing an N-terminal signal sequence, the 2A oligopeptide (white ovals) and the sequences immediately upstream (grey ovals) are predicted to be located within the ribosome exit tunnel and translocon pore, respectively. The lengths of the exit tunnel and translocon pore, plus the lengths of peptides (amino acids) which may be accommodated [44-49] are also shown.

ACKNOWLEDGEMENTS

We gratefully acknowledge D.A.A. Vignali (St. Jude Children's Research Hospital, Memphis, USA) for anti-2A antibodies and R.E. Randall (University of St. Andrews, St. Andrews, UK) for anti-V5 antibodies. We thank O.A. Vaughan and J. Nicholson for assistance with microscopy. This work was supported by the BBSRC (UK) and the Wellcome Trust. The authors have declared no conflict of interest.





Wiley-VCH

Figure 2





Figure 4



Figure 5

