



**HAL**  
open science

## Stochastic algorithms for computing means of probability measures

Marc Arnaudon, Clément Dombry, Anthony Phan, Le Yang

► **To cite this version:**

Marc Arnaudon, Clément Dombry, Anthony Phan, Le Yang. Stochastic algorithms for computing means of probability measures. 2010. hal-00540623v1

**HAL Id: hal-00540623**

**<https://hal.science/hal-00540623v1>**

Preprint submitted on 28 Nov 2010 (v1), last revised 24 Jun 2011 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# STOCHASTIC ALGORITHMS FOR COMPUTING MEANS OF PROBABILITY MEASURES

MARC ARNAUDON, CLÉMENT DOMBRY, ANTHONY PHAN, AND LE YANG

ABSTRACT. Consider a probability measure  $\mu$  supported by a regular geodesic ball in a manifold. For any  $p \geq 1$  we define a stochastic algorithm which converges almost surely to the  $p$ -mean  $e_p$  of  $\mu$ . Assuming furthermore that the functional to minimize is regular around  $e_p$ , we prove that a natural renormalization of the inhomogeneous Markov chain converges in law into an inhomogeneous diffusion process. We give an explicit expression of this process, as well as its local characteristic.

## CONTENTS

1. Introduction	1
2. Results	2
2.1. $p$ -means in regular geodesic balls	2
2.2. Fluctuations of the stochastic gradient algorithm	4
3. Proofs	5
3.1. Proof of Proposition 2.2	6
3.2. Proof of Theorem 2.3	7
3.3. Proof of Proposition 2.5	10
3.4. Proof of Theorem 2.6	11
4. Examples, simulations and pictures	17
4.1. Median of the uniform measure on an equilateral triangle in the plane	17
4.2. Uniform measure on the unit square in the plane	17
4.3. A non uniform measure on a square in the plane	17
4.4. A non uniform measure on the sphere $S^2$	18
4.5. Another non uniform measure on the unit square in the plane	19
References	19

## 1. INTRODUCTION

The geometric barycenter of a set of points is the point which minimizes the sum of the distances at the power 2 to these points. It is the most common estimator in statistics, however it is sensitive to outliers, and it is natural to replace power 2 by  $p$  for some  $p \in [1, 2)$ , which leads to the definition of  $p$ -mean. When  $p = 1$ , the minimizer is the median of the set of points, very often used in robust statistics. In many applications,  $p$ -means with some  $p \in (1, 2)$  give the best compromise.

The Fermat-Weber problem concerns finding the median  $e_1$  of a set of points in an Euclidean space. Numerous authors worked out algorithms for computing  $e_1$ . The first algorithm was proposed by Weiszfeld in [21]. It has been extended to

sufficiently small domains in Riemannian manifolds with nonnegative curvature by Fletcher and al in [7]. A complete generalization to manifolds with positive or negative curvature, including existence and uniqueness results (under some convexity conditions in positive curvature), has been given by one of the authors in [22].

The Riemannian barycenter or Karcher mean of a set of points in a manifold or more generally of a probability measure has been extensively studied, see e.g. [8], [10], [11], [5], [18], [2], where questions of existence, uniqueness, stability, relation with martingales in manifolds, behaviour when measures are pushed by stochastic flows have been considered. The Riemannian barycenter corresponds to  $p = 2$  in the above description. Computation of Riemannian barycenters by gradient descent has been performed by Le in [13].

In [1] Afsari proved existence and uniqueness of  $p$ -means,  $p \geq 1$  on geodesic balls with radius  $r < \frac{1}{2} \min \left\{ \text{inj}(M), \frac{\pi}{2\alpha} \right\}$  if  $p \in [1, 2)$ , and  $r < \frac{1}{2} \min \left\{ \text{inj}(M), \frac{\pi}{\alpha} \right\}$  if  $p \geq 2$ . Here  $\text{inj}(M)$  is the injectivity radius of  $M$  and  $\alpha > 0$  is such that the sectional curvatures in  $M$  are bounded above by  $\alpha^2$ . The point is that in the case  $p \geq 2$ , the functional to minimize is not convex any more, which makes the situation much more difficult to handle.

In this paper, under the assumptions of [1] we provide in Theorem 2.3 stochastic algorithms which converge almost surely to  $p$ -means in manifolds, which are easier to implement than gradient descent algorithm since computing the gradient of the function to minimize is not needed. The idea is at each step to go in the direction of a point of the support of  $\mu$ . The point is chosen at random according to  $\mu$  and the size of the step is a well chosen function of the distance to the point,  $p$  and the number of the step. For general convergence results on recursive stochastic algorithms, see [14] Theorem 1. However they do not cover the manifold case and nonlinearity of geodesics. Here we give a proof using martingale convergence theorem, and the main point consists in determining and estimating all the geometric quantities, checking that under our curvature conditions all the convergence assumptions are fulfilled, since our processes live in manifolds. See also [3] for convergence in probability of recursive algorithms.

The speed of convergence is studied, and in theorem 2.6 we prove that the renormalized inhomogeneous Markov chain of Theorem 2.3 converges in law to an inhomogeneous diffusion process. This is an invariance principle type result, see e.g. [9], [15], [4], [6] for related works. Here again the main point is to obtain the characteristics of the limiting process from the curvature conditions, the conditions on the support of the measure and estimates on Jacobi fields. Moreover we consider convergence in law for the Skorohod topology, and the limit depends in a crucial way on the decreasing steps of the algorithms.

Finally Section 4 is devoted to simulations.

## 2. RESULTS

**2.1.  $p$ -means in regular geodesic balls.** Let  $M$  be a Riemannian manifold with pinched sectional curvatures. Let  $\alpha, \beta > 0$  such that  $\alpha^2$  is a positive upper bound for sectional curvatures on  $M$ , and  $-\beta^2$  is a negative lower bound for sectional curvatures on  $M$ . Denote by  $\rho$  the Riemannian distance on  $M$ .

In  $M$  consider a geodesic ball  $B(a, r)$  with  $a \in M$ . Let  $\mu$  be a probability measure with support included in a compact convex subset  $K_\mu$  of  $B(a, r)$ . Fix  $p \in [1, \infty)$ . We will always make the following assumptions on  $(r, p, \mu)$ :

**Assumption 2.1.** The support of  $\mu$  is not reduced to one point. Either  $p > 1$  or the support of  $\mu$  is not contained in a line, and the radius  $r$  satisfies

$$(2.1) \quad r < r_{\alpha, p} \quad \text{with} \quad \begin{cases} r_{\alpha, p} &= \frac{1}{2} \min \left\{ \text{inj}(M), \frac{\pi}{2\alpha} \right\} & \text{if } p \in [1, 2) \\ r_{\alpha, p} &= \frac{1}{2} \min \left\{ \text{inj}(M), \frac{\pi}{\alpha} \right\} & \text{if } p \in [2, \infty) \end{cases}$$

Note that  $B(a, r)$  is convex if  $r < \frac{1}{2} \min \left\{ \text{inj}(M), \frac{\pi}{\alpha} \right\}$ .

Under assumption 2.1, it has been proved in [1] (Theorem 2.1) that the function

$$(2.2) \quad \begin{aligned} H_p : B(a, r) &\rightarrow \mathbb{R}_+ \\ x &\mapsto \int_M \rho^p(x, y) \mu(dy) \end{aligned}$$

has a unique minimizer  $e_p$  in  $M$ , the  $p$ -mean of  $\mu$ , and moreover  $e_p \in B(a, r)$ . If  $p = 1$ ,  $e_1$  is the median of  $\mu$ .

It is easily checked that if  $p \in [1, 2)$ , then  $H_p$  is strictly convex on  $B(a, r)$ . On the other hand, if  $p \geq 2$  then  $H_p$  is of class  $C^2$  on  $B(a, r)$ .

**Proposition 2.2.** *Let  $K$  be a convex subset of  $B(a, r)$  containing the support of  $\mu$ . Then there exists  $C_{p, \mu, K} > 0$  such that for all  $x \in K$ ,*

$$(2.3) \quad H_p(x) - H_p(e_p) \geq \frac{C_{p, \mu, K}}{2} \rho(x, e_p)^2.$$

Moreover if  $p \geq 2$  then we can choose  $C_{p, \mu, K}$  so that for all  $x \in K$ ,

$$(2.4) \quad \|\text{grad}_x H_p\|^2 \geq C_{p, \mu, K} (H_p(x) - H_p(e_p)).$$

In the sequel, we fix

$$(2.5) \quad K = \bar{B}(a, r - \varepsilon) \quad \text{with} \quad \varepsilon = \frac{\rho(K_\mu, B(a, r)^c)}{2}.$$

We now state our main result: we define a stochastic gradient algorithm  $(X_k)_{k \geq 0}$  to approximate the median  $e_p$  and prove its convergence.

**Theorem 2.3.** *Let  $(P_k)_{k \geq 1}$  be a sequence of independent  $B(a, r)$ -valued random variables, with law  $\mu$ . Let  $(t_k)_{k \geq 1}$  be a sequence of positive numbers satisfying*

$$(2.6) \quad \forall k \geq 1, \quad t_k \leq \min \left( \frac{1}{C_{p, \mu, K}}, \frac{\rho(K_\mu, B(a, r)^c)}{2p(2r)^{p-1}} \right),$$

$$(2.7) \quad \sum_{k=1}^{\infty} t_k = +\infty \quad \text{and} \quad \sum_{k=1}^{\infty} t_k^2 < \infty.$$

Letting  $x_0 \in K$ , define inductively the random walk  $(X_k)_{k \geq 0}$  by

$$(2.8) \quad X_0 = x_0 \quad \text{and for } k \geq 0 \quad X_{k+1} = \exp_{X_k} \left( -t_{k+1} \text{grad}_{X_k} F_p(\cdot, P_{k+1}) \right)$$

where  $F_p(x, y) = \rho^p(x, y)$ , with the convention  $\text{grad}_x F_p(\cdot, x) = 0$ .

The random walk  $(X_k)_{k \geq 1}$  converges in  $L^2$  and almost surely to  $e$ .

In the following example, we focus on the case  $M = \mathbb{R}^d$  and  $p = 2$  where drastic simplifications occur.

**Example 2.4.** In the case when  $M = \mathbb{R}^d$  and  $\mu$  is a compactly supported probability measure on  $\mathbb{R}^d$ , the stochastic gradient algorithm (2.8) simplifies into

$$X_0 = x_0 \quad \text{and for } k \geq 0 \quad X_{k+1} = X_k - t_{k+1} \operatorname{grad}_{X_k} F_p(\cdot, P_{k+1}).$$

If furthermore  $p = 2$ , clearly  $e_2 = \mathbb{E}[P_1]$  and  $\operatorname{grad}_x F_p(\cdot, y) = 2(x - y)$ , so that the linear relation

$$X_{k+1} = (1 - 2t_{k+1}X_k) + 2t_{k+1}P_{k+1}, \quad k \geq 0$$

holds true and an easy induction proves that

$$(2.9) \quad X_k = x_0 \prod_{j=0}^{k-1} (1 - 2t_{k-j}) + 2 \sum_{j=0}^{k-1} P_{k-j} t_{k-j} \prod_{\ell=0}^{j-1} (1 - 2t_{k-\ell}), \quad k \geq 1.$$

Now, taking  $t_k = \frac{1}{2k}$ , we have

$$\prod_{j=0}^{k-1} (1 - 2t_{k-j}) = 0 \quad \text{and} \quad \prod_{\ell=0}^{j-1} (1 - 2t_{k-\ell}) = \frac{k-j}{k}$$

so that

$$X_k = \sum_{j=0}^{k-1} P_{k-j} \frac{1}{k} = \frac{1}{k} \sum_{j=1}^k P_j.$$

The stochastic gradient algorithm estimating the mean  $e_2$  of  $\mu$  is given by the empirical mean of a growing sample of independent random variables with distribution  $\mu$ . In this simple case, the result of Theorem 2.3 is nothing but the strong law of large numbers. Moreover, fluctuations around the mean are given by the central limit theorem and Donsker's theorem.

**2.2. Fluctuations of the stochastic gradient algorithm.** The notations are the same as in the beginning of section 2.1. We still make assumption 2.1. Let us define  $K$  and  $\varepsilon$  as in (2.5) and let

$$(2.10) \quad \delta_1 = \min \left( \frac{1}{C_{p,\mu,K}}, \frac{\rho(K_\mu, B(a, r)^c)}{2p(2r)^{p-1}} \right).$$

We consider the time inhomogeneous  $M$ -valued Markov chain (2.8) in the particular case when

$$(2.11) \quad t_k = \min \left( \frac{\delta}{k}, \delta_1 \right), \quad k \geq 1$$

for some  $\delta > 0$ . The particular sequence  $(t_k)_{k \geq 1}$  defined by (2.11) satisfies (2.6) and (2.7), so Theorem 2.3 holds true and the stochastic gradient algorithm  $(X_k)_{k \geq 0}$  converges a.s. and in  $L^2$  to the  $p$ -mean  $e_p$ .

In order to study the fluctuations around the  $p$ -mean  $e_p$ , we define for  $n \geq 1$  the rescaled  $T_{e_p}M$ -valued Markov chain  $(Y_k^n)_{k \geq 0}$  by

$$(2.12) \quad Y_k^n = \frac{k}{\sqrt{n}} \exp_{e_p}^{-1} X_k.$$

We will prove convergence of the sequence of process  $(Y_{[nt]}^n)_{0 \leq t \leq 1}$  to a non-homogeneous diffusion process. The limit process is defined in the following proposition:

**Proposition 2.5.** *Assume that  $H_p$  is  $C^2$  in a neighborhood of  $e_p$ , and that  $\delta > C_{p,\mu,K}^{-1}$ . Define*

$$\Gamma = \mathbb{E} \left[ \text{grad}_{e_p} F_p(\cdot, P_1) \otimes \text{grad}_{e_p} F_p(\cdot, P_1) \right]$$

and  $G_\delta(t)$  the generator

$$(2.13) \quad G_\delta(t)f(y) := \langle d_y f, t^{-1}(y - \delta \nabla dH_p(y, \cdot)^\sharp) \rangle + \frac{1}{2} \text{Hess}_y f(\Gamma)$$

where  $\nabla dH_p(y, \cdot)^\sharp$  denotes the dual vector of the linear form  $\nabla dH_p(y, \cdot)$ .

There exists a unique inhomogeneous diffusion process  $(y_\delta(t))_{t>0}$  on  $T_{e_p}M$  with generator  $G_\delta(t)$  and converging in probability to 0 as  $t \rightarrow 0^+$ .

The process  $y_\delta$  is continuous, converges a.s. to 0 as  $t \rightarrow 0^+$  and has the following integral representation:

$$(2.14) \quad y_\delta(t) = \sum_{i=1}^d t^{1-\delta\lambda_i} \int_0^t s^{\delta\lambda_i-1} \langle \delta\sigma dB_s, e_i \rangle e_i, \quad t \geq 0,$$

where  $B_t$  is a standard Brownian motion on  $T_{e_p}M$ ,  $\sigma \in \text{End}(T_{e_p}M)$  satisfies  $\sigma\sigma^* = \Gamma$ ,  $(e_i)_{1 \leq i \leq d}$  is an orthonormal basis diagonalizing the symmetric bilinear form  $\nabla dH_p(e_p)$  and  $(\lambda_i)_{1 \leq i \leq d}$  are the associated eigenvalues.

Note that the integral representation (2.14) implies that  $y_\delta$  is the centered Gaussian process with covariance

$$(2.15) \quad \mathbb{E} \left[ y_\delta^i(t_1) y_\delta^j(t_2) \right] = \frac{\delta^2 \Gamma(e_i^* \otimes e_j^*)}{\delta(\lambda_i + \lambda_j) - 1} t_1^{1-\delta\lambda_i} t_2^{1-\delta\lambda_j} (t_1 \wedge t_2)^{\delta(\lambda_i + \lambda_j) - 1},$$

where  $y_\delta^i(t) = \langle y_\delta(t), e_i \rangle$ ,  $1 \leq i, j \leq d$  and  $t_1, t_2 \geq 0$ .

Our main result on the fluctuations of the stochastic gradient algorithm is the following:

**Theorem 2.6.** *Assume that  $H_p$  is  $C^2$  in a neighborhood of  $e_p$ , and that  $\delta > C_{p,\mu,K}^{-1}$ . The sequence of processes  $(Y_{[nt]}^n)_{t \geq 0}$  weakly converges in  $\mathbb{D}((0, \infty), T_{e_p}M)$  to  $y_\delta$ .*

**Remark 2.7.** From section 2.1 we know that, when  $p \in (1, 2]$ , the constant

$$C_{p,\mu,K} = p(2r)^{p-2} (\min(p-1, 2\alpha r \cot(2\alpha r)))$$

is explicit. The constraint  $\delta > C_{p,\mu,K}^{-1}$  can easily be checked in this case.

**Remark 2.8.** In the case  $M = \mathbb{R}^d$ ,  $Y_k^n = \frac{k}{\sqrt{n}}(X_k - e_p)$  and the tangent space  $T_{e_p}M$  is identified to  $\mathbb{R}^d$ . Theorem 2.6 holds and, in particular, when  $t = 1$ , we obtain a central limit Theorem:  $\sqrt{n}(X_n - e)$  converges as  $n \rightarrow \infty$  to a centered Gaussian  $d$ -variate distribution (with covariance structure given by (2.15) with  $t_1 = t_2 = 1$ ). This is a central limit theorem: the fluctuations of the stochastic gradient algorithm are of scale  $n^{-1/2}$  and asymptotically Gaussian.

### 3. PROOFS

For simplicity, let us write shortly  $e = e_p$  in the proofs.

### 3.1. Proof of Proposition 2.2.

For  $p = 1$  this is a direct consequence of [22] Theorem 3.7.

Next we consider the case  $p \in [1, 2)$ .

Let  $K \subset B(a, r)$  be a compact convex set containing the support of  $\mu$ . Let  $x \in K \setminus \{e\}$ ,  $t = \rho(e, x)$ ,  $u \in T_e M$  the unit vector such that  $\exp_e(\rho(e, x)u) = x$ , and  $\gamma_u$  the geodesic with initial speed  $u$  :  $\dot{\gamma}_u(0) = u$ . For  $y \in K$ , letting  $h_y(s) = \rho(\gamma_u(s), y)^p$ ,  $s \in [0, t]$ , we have since  $p > 1$

$$h_y(t) = h_y(0) + th'_y(0) + \int_0^t (t-s)h''_y(s) ds$$

with the convention  $h''_y(s) = 0$  when  $\gamma_u(s) = y$ . Indeed, if  $y \notin \gamma([0, t])$  then  $h_y$  is smooth, and if  $y \in \gamma([0, t])$ , say  $y = \gamma(s_0)$  then  $h_y(s) = |s - s_0|^p$  and the formula can easily be checked.

By standard calculation,

$$(3.1) \quad \begin{aligned} & h''_y(s) \\ & \geq p\rho(\gamma_u(s), y)^{p-2} \\ & \quad \times \left( (p-1)\|\dot{\gamma}_u(s)^{T(y)}\|^2 + \|\dot{\gamma}_u(s)^{N(y)}\|^2 \alpha\rho(\gamma_u(s), y) \cot(\alpha\rho(\gamma_u(s), y)) \right) \end{aligned}$$

with  $\dot{\gamma}_u(s)^{T(y)}$  (resp.  $\dot{\gamma}_u(s)^{N(y)}$ ) the tangential (resp. the normal) part of  $\dot{\gamma}_u(s)$  with respect to  $n(\gamma_u(s), y) = \frac{1}{\rho(\gamma_u(s), y)} \exp_{\gamma_u(s)}^{-1}(y)$ :

$$\dot{\gamma}_u(s)^{T(y)} = \langle \dot{\gamma}_u(s), n(\gamma_u(s), y) \rangle n(\gamma_u(s), y), \quad \dot{\gamma}_u(s)^{N(y)} = \dot{\gamma}_u(s) - \dot{\gamma}_u(s)^{T(y)}.$$

From this we get

$$(3.2) \quad h''_y(s) \geq p\rho(\gamma_u(s), y)^{p-2} (\min(p-1, 2\alpha r \cot(2\alpha r))).$$

Now

$$\begin{aligned} & H_p(\gamma_u(t')) \\ & = \int_{B(a, r)} h_y(\gamma_u(t')) \mu(dy) \\ & = \int_{B(a, r)} h_y(0) \mu(dy) + t' \int_{B(a, r)} h'_y(0) \mu(dy) + \int_0^{t'} (t' - s) \int_{B(a, r)} h_y(s)'' \mu(dy) \end{aligned}$$

and  $H_p(\gamma_u(t'))$  attains its minimum at  $t' = 0$ , so  $\int_{B(a, r)} h'_y(0) \mu(dy) = 0$  and we have

$$H_p(x) = H_p(\gamma_u(t)) = H_p(e) + \int_0^t (t-s) \int_{B(a, r)} h_y(s)'' \mu(dy).$$

Using Equation (3.2) we get

$$(3.3) \quad \begin{aligned} & H_p(x) \geq H_p(e) \\ & + \int_0^t \left( (t-s) \int_{B(a, r)} p\rho(\gamma_u(s), y)^{p-2} (\min(p-1, 2\alpha r \cot(2\alpha Kr))) \mu(dy) \right) ds. \end{aligned}$$

Since  $p \leq 2$  we have  $\rho(\gamma_u(s), y)^{p-2} \geq (2r)^{p-2}$  and

$$(3.4) \quad H_p(x) \geq H_p(e) + \frac{t^2}{2} p(2r)^{p-2} (\min(p-1, 2\alpha r \cot(2\alpha r))).$$

So letting

$$C_{p,\mu,K} = p(2r)^{p-2} (\min(p-1, 2\alpha r \cot(2\alpha r)))$$

we obtain

$$(3.5) \quad H_p(x) \geq H_p(e) + \frac{C_{p,\mu,K} \rho(e, x)^2}{2}.$$

To finish let us consider the case  $p \geq 2$ .

In the proof of [1] Theorem 2.1, it is shown that  $e$  is the only zero of the maps  $x \mapsto \text{grad}_x H_p$  and  $x \mapsto H_p(x) - H_p(e)$ , and that  $\nabla dH_p(e)$  is strictly positive. This implies that (2.3) and (2.4) hold on some neighbourhood  $B(e, \varepsilon)$  of  $e$ . By compactness and the fact that  $H_p - H_p(e)$  and  $\text{grad} H_p$  do not vanish on  $K \setminus B(e, \varepsilon)$  and  $H_p - H_p(e)$  is bounded, possibly modifying the constant  $C_{p,\mu,K}$ , (2.3) and (2.4) also holds on  $K \setminus B(e, \varepsilon)$ .  $\square$

### 3.2. Proof of Theorem 2.3.

Note that, for  $x \neq y$ ,

$$\text{grad}_x F(\cdot, y) = p\rho^{p-1}(x, y) \frac{-\exp_x^{-1}(y)}{\rho(x, y)} = -p\rho^{p-1}(x, y)n(x, y),$$

whith  $n(x, y) := \frac{\exp_x^{-1}(y)}{\rho(x, y)}$  a unit vector. So, with the condition (2.6) on  $t_k$ , the random walk  $(X_k)_{k \geq 0}$  cannot exit  $K$ : if  $X_k \in K$  then there are two possibilities for  $X_{k+1}$ :

- either  $X_{k+1}$  is in the geodesic between  $X_k$  and  $P_{k+1}$  and belongs to  $K$  by convexity of  $K$ ;
- or  $X_{k+1}$  is after  $P_{k+1}$ , but since

$$\begin{aligned} \|t_{k+1} \text{grad}_{X_k} F_p(\cdot, P_{k+1})\| &= t_{k+1} p \rho^{p-1}(X_k, P_{k+1}) \\ &\leq \frac{\rho(K_\mu, B(a, r)^c)}{2p(2r)^{p-1}} p \rho^{p-1}(X_k, P_{k+1}) \\ &\leq \frac{\rho(K_\mu, B(a, r)^c)}{2}, \end{aligned}$$

we have in this case

$$\rho(P_{k+1}, X_{k+1}) \leq \frac{\rho(K_\mu, B(a, r)^c)}{2}$$

which implies that  $X_{k+1} \in K$ .

First consider the case  $p \in [1, 2)$ .

For  $k \geq 0$  let

$$t \mapsto E(t) := \frac{1}{2} \rho^2(e, \gamma(t)),$$

$\gamma(t)_{t \in [0, t_{k+1}]}$  the geodesic satisfying  $\dot{\gamma}(0) = -\text{grad}_{X_k} F_p(\cdot, P_{k+1})$ . We have for all  $t \in [0, t_{k+1}]$

$$(3.6) \quad E''(t) \leq C(\beta, r, p) := p^2(2r)^{2p-1} \beta \coth(2\beta r)$$



(see e.g. [22]). By Taylor formula,

$$\begin{aligned} & \rho(X_{k+1}, e)^2 \\ &= 2E(t_{k+1}) \\ &= 2E(0) + 2t_{k+1}E'(0) + t_{k+1}^2 E''(t) \quad \text{for some } t \in [0, t_{k+1}] \\ &\leq \rho(X_k, e)^2 + 2t_{k+1} \langle \text{grad}_{X_k} F_p(\cdot, P_{k+1}), \exp_{X_k}^{-1}(e) \rangle + t_{k+1}^2 C(\beta, r, p). \end{aligned}$$

Now from the convexity of  $x \mapsto F_p(x, y)$  we have for all  $x, y \in B(a, r)$

$$(3.7) \quad F_p(e, y) - F_p(x, y) \geq \langle \text{grad}_x F_p(\cdot, y), \exp_x^{-1}(e) \rangle.$$

This applied with  $x = X_k, y = P_{k+1}$  yields

$$(3.8) \quad \rho(X_{k+1}, e)^2 \leq \rho(X_k, e)^2 - 2t_{k+1} (F_p(X_k, P_{k+1}) - F_p(e, P_{k+1})) + C(\beta, r, p)t_{k+1}^2.$$

Letting for  $k \geq 0$   $\mathcal{F}_k = \sigma(X_\ell, 0 \leq \ell \leq k)$ , we get

$$\begin{aligned} & \mathbb{E} [\rho(X_{k+1}, e)^2 | \mathcal{F}_k] \\ &\leq \rho(X_k, e)^2 - 2t_{k+1} \int_{B(a, r)} (F_p(X_k, y) - F_p(e, y)) \mu(dy) + C(\beta, r, p)t_{k+1}^2 \\ &= \rho(X_k, e)^2 - 2t_{k+1} (H_p(X_k) - H_p(e)) + C(\beta, r, p)t_{k+1}^2 \\ &\leq \rho(X_k, e)^2 + C(\beta, r, p)t_{k+1}^2 \end{aligned}$$

so that the process  $(Y_k)_{k \geq 0}$  defined by

$$(3.9) \quad Y_0 = \rho(X_0, e)^2 \quad \text{and for } k \geq 1 \quad Y_k = \rho(X_k, e)^2 - C(\beta, r, p) \sum_{j=1}^k t_j^2$$

is a bounded supermartingale. So it converges in  $L^1$  and almost surely. Consequently  $\rho(X_k, e)^2$  also converges in  $L^1$  and almost surely.

Let

$$(3.10) \quad a = \lim_{k \rightarrow \infty} \mathbb{E} [\rho(X_k, e)^2].$$

We want to prove that  $a = 0$ . We already proved that

$$(3.11) \quad \mathbb{E} [\rho(X_{k+1}, e)^2 | \mathcal{F}_k] \leq \rho(X_k, e)^2 - 2t_{k+1} (H_p(X_k) - H_p(e)) + C(\beta, r, p)t_{k+1}^2.$$

Taking the expectation and using Proposition 2.2, we obtain

$$(3.12) \quad \mathbb{E} [\rho(X_{k+1}, e)^2] \leq \mathbb{E} [\rho(X_k, e)^2] - t_{k+1} C_{p, \mu, K} \mathbb{E} [\rho(X_k, e)^2] + C(\beta, r, p)t_{k+1}^2.$$

An easy induction proves that for  $\ell \geq 1$ ,

$$(3.13) \quad \mathbb{E} [\rho(X_{k+\ell}, e)^2] \leq \prod_{j=1}^{\ell} (1 - C_{p, \mu, K} t_{k+j}) \mathbb{E} [\rho(X_k, e)^2] + C(\beta, r, p) \sum_{j=1}^{\ell} t_{k+j}^2.$$

Letting  $\ell \rightarrow \infty$  and using the fact that  $\sum_{j=1}^{\infty} t_{k+j} = \infty$  which implies

$$\prod_{j=1}^{\infty} (1 - C_{p, \mu, K} t_{k+j}) = 0,$$

we get

$$(3.14) \quad a \leq C(\beta, r, p) \sum_{j=1}^{\infty} t_{k+j}^2.$$

Finally using  $\sum_{j=1}^{\infty} t_j^2 < \infty$  we obtain that  $\lim_{k \rightarrow \infty} \sum_{j=1}^{\infty} t_{k+j}^2 = 0$ , so  $a = 0$ . This proves  $L^2$  and almost sure convergence.

Next assume that  $p \geq 2$ .

For  $k \geq 0$  let

$$t \mapsto E_p(t) := H_p(\gamma(t)),$$

$\gamma(t)_{t \in [0, t_{k+1}]}$  the geodesic satisfying  $\dot{\gamma}(0) = -\text{grad}_{X_k} F_p(\cdot, P_{k+1})$ . With a calculation similar to (3.6) we get for all  $t \in [0, t_{k+1}]$

$$(3.15) \quad E_p''(t) \leq 2C(\beta, r, p) := \frac{p^3}{2} (2r)^{3p-4} (2r\beta \coth(2\beta r) + 2p - 4).$$

(see e.g. [22]). By Taylor formula,

$$\begin{aligned} H_p(X_{k+1}) &= E_p(t_{k+1}) \\ &= E_p(0) + t_{k+1} E_p'(0) + \frac{t_{k+1}^2}{2} E_p''(t) \quad \text{for some } t \in [0, t_{k+1}] \\ &\leq H_p(X_k) + t_{k+1} \langle d_{X_k} H_p, \text{grad}_{X_k} F_p(\cdot, P_{k+1}) \rangle + t_{k+1}^2 C(\beta, r, p). \end{aligned}$$

We get

$$\begin{aligned} &\mathbb{E}[H_p(X_{k+1}) | \mathcal{F}_k] \\ &\leq H_p(X_k) - t_{k+1} \left\langle d_{X_k} H_p, \int_{B(a, r)} \text{grad}_{X_k} F_p(\cdot, y) \mu(dy) \right\rangle + C(\beta, r, p) t_{k+1}^2 \\ &= H_p(X_k) - t_{k+1} \langle d_{X_k} H_p, \text{grad}_{X_k} H_p(\cdot) \rangle + C(\beta, r, p) t_{k+1}^2 \\ &= H_p(X_k) - t_{k+1} \|\text{grad}_{X_k} H_p(\cdot)\|^2 + C(\beta, r, p) t_{k+1}^2 \\ &\leq H_p(X_k) - C_{p, \mu, K} t_{k+1} (H_p(X_k) - H_p(e)) + C(\beta, r, p) t_{k+1}^2 \end{aligned}$$

(by Proposition 2.2) so that the process  $(Y_k)_{k \geq 0}$  defined by

(3.16)

$$Y_0 = H_p(X_0) - H_p(e) \quad \text{and for } k \geq 1 \quad Y_k = H_p(X_k) - H_p(e) - C(\beta, r, p) \sum_{j=1}^k t_j^2$$

is a bounded supermartingale. So it converges in  $L^1$  and almost surely. Consequently  $H_p(X_k) - H_p(e)$  also converges in  $L^1$  and almost surely.

Let

$$(3.17) \quad a = \lim_{k \rightarrow \infty} \mathbb{E}[H_p(X_k) - H_p(e)].$$

We want to prove that  $a = 0$ . We already proved that

$$(3.18) \quad \begin{aligned} &\mathbb{E}[H_p(X_{k+1}) - H_p(e) | \mathcal{F}_k] \\ &\leq H_p(X_k) - H_p(e) - C_{p, \mu, K} t_{k+1} (H_p(X_k) - H_p(e)) + C(\beta, r, p) t_{k+1}^2. \end{aligned}$$

Taking the expectation we obtain

(3.19)

$$\mathbb{E}[H_p(X_{k+1}) - H_p(e)] \leq (1 - t_{k+1}C_{p,\mu,K})\mathbb{E}[H_p(X_k) - H_p(e)] + C(\beta, r, p)t_{k+1}^2$$

so that proving that  $a = 0$  is similar to the previous case.

Finally (2.3) proves that  $\rho(X_k, e)^2$  converges in  $L^1$  and almost surely to 0.  $\square$

**3.3. Proof of Proposition 2.5.** Fix  $\varepsilon > 0$ . Any diffusion process on  $[\varepsilon, \infty)$  with generator  $G_\delta(t)$  is solution of a sde of the type

$$(3.20) \quad dy_t = \frac{1}{t}L_\delta(y_t) dt + \delta\sigma dB_t$$

where  $L_\delta(y) = y - \delta\nabla dH_p(y, \cdot)^\sharp$  and  $B_t$  and  $\sigma$  are as in Proposition 2.5. This sde can be solved explicitly on  $[\varepsilon, \infty)$ . The symmetric endomorphism  $y \mapsto \nabla dH_p(y, \cdot)^\sharp$  is diagonalisable in the orthonormal basis  $(e_i)_{1 \leq i \leq d}$  with eigenvalues  $(\lambda_i)_{1 \leq i \leq d}$ . The endomorphism  $L_\delta = \text{id} - \delta\nabla dH_p(e)(\text{id}, \cdot)^\sharp$  is also diagonalisable in this basis

with eigenvalues  $(1 - \delta\lambda_i)_{1 \leq i \leq d}$ . The solution  $y_t = \sum_{i=1}^d y_t^i e_i$  of (3.20) started at

$y_\varepsilon = \sum_{i=1}^d y_\varepsilon^i e_i$  is given by

$$(3.21) \quad y_t = \sum_{i=1}^d \left( y_\varepsilon^i \varepsilon^{\delta\lambda_i - 1} + \int_\varepsilon^t s^{\delta\lambda_i - 1} \langle \delta\sigma dB_s, e_i \rangle \right) t^{1 - \delta\lambda_i} e_i, \quad t \geq \varepsilon.$$

Now by definition of  $C_{p,\mu,K}$  we clearly have

$$(3.22) \quad C_{p,\mu,K} \leq \min_{1 \leq i \leq d} \lambda_i.$$

So the condition  $\delta C_{p,\mu,K} > 1$  implies that for all  $i$ ,  $\delta\lambda_i - 1 > 0$ , and as  $\varepsilon \rightarrow 0$ ,

$$(3.23) \quad \int_\varepsilon^t s^{\delta\lambda_i - 1} \langle \delta\sigma dB_s, e_i \rangle \rightarrow \int_0^t s^{\delta\lambda_i - 1} \langle \delta\sigma dB_s, e_i \rangle \quad \text{in probability.}$$

Assume that a continuous solution  $y_t$  converging in probability to 0 as  $t \rightarrow 0^+$  exists. Since  $y_\varepsilon^i \varepsilon^{\delta\lambda_i - 1} \rightarrow 0$  in probability as  $\varepsilon \rightarrow 0$ , we necessarily have using (3.23)

$$(3.24) \quad y_t = \sum_{i=1}^d t^{1 - \delta\lambda_i} \int_0^t s^{\delta\lambda_i - 1} \langle \delta\sigma dB_s, e_i \rangle e_i, \quad t \geq 0.$$

Note  $y_\delta^i$  is Gaussian with variance  $\frac{t\delta^2\Gamma(e_i^* \otimes e_i^*)}{2\delta\lambda_i - 1}$ , so it converges in  $L^2$  to 0 as  $t \rightarrow 0$ .

Conversely, it is easy to check that equation (3.24) defines a solution to (3.20).

To prove the a.s. convergence to 0 we use the representation

$$\int_0^t s^{\delta\lambda_i - 1} \langle \delta\sigma dB_s, e_i \rangle = B_{\varphi_i(t)}^i$$

where  $B_s^i$  is a Brownian motion and  $\varphi_i(t) = \frac{\delta^2\Gamma(e_i^* \otimes e_i^*)}{2\delta\lambda_i - 1} t^{2\delta\lambda_i - 1}$ . Then by the law of iterated logarithm

$$\limsup_{t \downarrow 0} t^{1 - \delta\lambda_i} B_{\varphi_i(t)}^i \leq \limsup_{t \downarrow 0} t^{1 - \delta\lambda_i} \sqrt{2\varphi_i(t) \ln \ln(\varphi_i^{-1}(t))}$$

But for  $t$  small we have

$$\sqrt{2\varphi_i(t) \ln \ln (\varphi_i^{-1}(t))} \leq t^{\delta\lambda_i - 3/4}$$

so

$$\limsup_{t \downarrow 0} t^{1-\delta\lambda_i} B_{\varphi_i(t)}^i \leq \lim_{t \downarrow 0} t^{1/4} = 0.$$

This proves a.s. convergence to 0. Continuity is easily checked using the integral representation (3.24).  $\square$

**3.4. Proof of Theorem 2.6.** Consider the time homogeneous Markov chain  $(Z_k^n)_{k \geq 0}$  with state space  $[0, \infty) \times T_e M$  defined by

$$(3.25) \quad Z_k^n = \left( \frac{k}{n}, Y_k^n \right).$$

The first component has a deterministic evolution and will be denoted by  $t_k^n$ ; it satisfies

$$(3.26) \quad t_{k+1}^n = t_k^n + \frac{1}{n}, \quad k \geq 0.$$

Let  $k_0$  be such that

$$(3.27) \quad \frac{\delta}{k_0} < \delta_1.$$

Using equations (2.8), (2.12) and (2.11), we have for  $k \geq k_0$ ,

$$(3.28) \quad Y_{k+1}^n = \frac{nt_k^n + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt_k^n}} Y_k^n} \left( -\frac{\delta}{nt_k^n + 1} \text{grad}_{\frac{1}{\sqrt{nt_k^n}} Y_k^n} F_p(\cdot, P_{k+1}) \right) \right).$$

Consider the transition kernel  $P^n(z, dz')$  on  $(0, \infty) \times T_e M$  defined for  $z = (t, y)$  by

$$(3.29) \quad P^n(z, A) = \mathbb{P} \left[ \left( t + \frac{1}{n}, \frac{nt+1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt+1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, P_1) \right) \right) \right) \in A \right]$$

where  $A \in \mathcal{B}((0, \infty) \times T_e M)$ . Clearly this transition kernel drives the evolution of the Markov chain  $(Z_k^n)_{k \geq k_0}$ .

For the sake of clarity, we divide the proof of Theorem 2.6 into four lemmas.

**Lemma 3.1.** *Assume that for all  $x \in B(a, r)$  the application  $F_p(\cdot, x)$  is of class  $C^2$  in a neighbourhood of  $e$ . Fix  $\delta > 0$ . Let  $B$  be a bounded set in  $T_e M$  and let  $0 < \varepsilon < T$ . We have for all  $C^2$  function  $f$  on  $T_e M$*

$$(3.30) \quad \begin{aligned} & n \left( f \left( \frac{nt+1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt+1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, x) \right) \right) \right) - f(y) \right) \\ &= \left\langle d_y f, \frac{y}{t} \right\rangle - \sqrt{n} \langle d_y f, \delta \text{grad}_e F_p(\cdot, x) \rangle - \delta \nabla d F_p(\cdot, x) \left( \text{grad}_y f, \frac{y}{t} \right) \\ &+ \frac{\delta^2}{2} \text{Hess}_y f \left( \text{grad}_e F_p(\cdot, x) \otimes \text{grad}_e F_p(\cdot, x) \right) + O \left( \frac{1}{\sqrt{n}} \right) \end{aligned}$$

uniformly in  $y \in B$ ,  $x \in K_\mu$ ,  $t \in [\varepsilon, T]$ .

*Proof.* Let  $x \in K_\mu$ ,  $y \in T_e M$ ,  $u, v \in \mathbb{R}$  sufficiently close to 0, and  $q = \exp_e\left(\frac{uy}{t}\right)$ . For  $s \in [0, 1]$  denote by  $a \mapsto c(a, s, u, v)$  the geodesic with endpoints  $c(0, s, u, v) = e$  and

$$c(1, s, u, v) = \exp_{\exp_e\left(\frac{uy}{t}\right)}\left(-us \operatorname{grad}_{\exp_e\left(\frac{uy}{t}\right)} F_p(\cdot, x)\right) :$$

$$c(a, s, u, v) = \exp_e \left\{ a \exp_e^{-1} \left[ \exp_{\exp_e\left(\frac{uy}{t}\right)}\left(-sv \operatorname{grad}_{\exp_e\left(\frac{uy}{t}\right)} F_p(\cdot, x)\right) \right] \right\}.$$

This is a  $C^2$  function of  $(a, s, u, v) \in [0, 1]^2 \times (-\eta, \eta)^2$ ,  $\eta$  sufficiently small. It also depends in a  $C^2$  way of  $x$  and  $y$ . Letting  $c(a, s) = c\left(a, s, \frac{1}{\sqrt{n}}, \frac{\delta}{nt+1}\right)$ , we have

$$\exp_e^{-1} \left( \exp_{\exp_e\left(\frac{1}{\sqrt{nt}}y\right)} \left( -\frac{\delta}{nt+1} \operatorname{grad}_{\exp_e\left(\frac{1}{\sqrt{nt}}y\right)} F_p(\cdot, x) \right) \right) = \partial_a c(0, 1).$$

So we need a Taylor expansion up to order  $n^{-1}$  of  $\frac{nt+1}{\sqrt{n}} \partial_a c(0, 1)$ .

We have  $c(a, s, 0, 1) = \exp_e(-as \operatorname{grad}_e F_p(\cdot, x))$  and this implies

$$\partial_s^2 \partial_a c(0, s, 0, 1) = 0, \quad \text{so} \quad \partial_s^2 \partial_a c(0, s, u, 1) = O(u).$$

On the other hand the identities  $c(a, s, u, v) = c(a, sv, u, 1)$  yields  $\partial_s^2 \partial_a c(a, s, u, v) = v^2 \partial_s^2 \partial_a c(a, s, u, 1)$ , so we obtain

$$\partial_s^2 \partial_a c(0, s, u, v) = O(uv^2)$$

and this yields

$$\partial_s^2 \partial_a c(0, s) = O(n^{-5/2}),$$

uniformly in  $s, x, y, t$ . But since

$$\|\partial_a c(0, 1) - \partial_a c(0, 0) - \partial_s \partial_a c(0, 0)\| \leq \frac{1}{2} \sup_{s \in [0, 1]} \|\partial_s^2 \partial_a c(0, s)\|$$

we only need to estimate  $\partial_a c(0, 0)$  and  $\partial_s \partial_a c(0, 0)$ .

Denoting by  $J(a)$  the Jacobi field  $\partial_s c(a, 0)$  we have

$$\frac{nt+1}{\sqrt{n}} \partial_a c(0, 1) = \frac{nt+1}{\sqrt{n}} \partial_a c(0, 0) + \frac{nt+1}{\sqrt{n}} \dot{J}(0) + O\left(\frac{1}{n^2}\right).$$

On the other hand

$$\frac{nt+1}{\sqrt{n}} \partial_a c(0, 0) = \frac{nt+1}{\sqrt{n}} \frac{y}{\sqrt{nt}} = y + \frac{y}{nt}$$

so it remains to estimate  $\dot{J}(0)$ .

The Jacobi field  $a \mapsto J(a, u, v)$  with endpoints  $J(0, u, v) = 0_e$  and

$$J(1, u, v) = -v \operatorname{grad}_{\exp_e\left(\frac{uy}{t}\right)} F_p(\cdot, x)$$

satisfies

$$\nabla_a^2 J(a, u, v) = -R(J(a, u, v), \partial_a c(a, 0, u, v)) \partial_a c(a, 0, u, v) = O(u^2 v).$$

This implies that

$$\nabla_a^2 J(a) = O(n^{-2}).$$

Consequently, denoting by  $P_{x_1, x_2} : T_{x_1} M \rightarrow T_{x_2} M$  the parallel transport along the minimal geodesic from  $x_1$  to  $x_2$  (whenever it is unique) we have

$$(3.31) \quad P_{c(1,0), e} J(1) = J(0) + \dot{J}(0) + O(n^{-2}) = \dot{J}(0) + O(n^{-2}).$$

But we also have

$$\begin{aligned} P_{c(1,0,u,v),e}J(1,u,v) &= P_{c(1,0,u,v),e} \left( -v \operatorname{grad}_{c(1,0,u,v)} F_p(\cdot, x) \right) \\ &= -v \operatorname{grad}_e F_p(\cdot, x) - v \nabla_{\partial_a c(0,0,u,v)} \operatorname{grad}_e F_p(\cdot, x) + O(vu^2) \\ &= -v \operatorname{grad}_e F_p(\cdot, x) - v \nabla dF_p(\cdot, x) \left( \frac{uy}{t}, \cdot \right)^\sharp + O(vu^2) \end{aligned}$$

where we used  $\partial_a c(0,0,u,v) = \frac{uy}{t}$  and for vector fields  $A, B$  on  $TM$  and a  $C^2$  function  $f_1$  on  $M$

$$\begin{aligned} \langle \nabla_{A_e} \operatorname{grad} f_1, B_e \rangle &= A_e \langle \operatorname{grad} f_1, B_e \rangle - \langle \operatorname{grad} f_1, \nabla_{A_e} B \rangle \\ &= A_e \langle df_1, B_e \rangle - \langle df_1, \nabla_{A_e} B \rangle \\ &= \nabla df_1(A_e, B_e) \end{aligned}$$

which implies

$$\nabla_{A_e} \operatorname{grad} f_1 = \nabla df_1(A_e, \cdot)^\sharp.$$

We obtain

$$P_{c(1,0),e}J(1) = -\frac{\delta}{nt+1} \operatorname{grad}_e F_p(\cdot, x) - \frac{\delta}{\sqrt{n}(nt+1)} \nabla dF_p(\cdot, x) \left( \frac{y}{t}, \cdot \right)^\sharp + O(n^{-2}).$$

Combining with (3.31) this gives

$$J(0) = -\frac{\delta}{nt+1} \operatorname{grad}_e F_p(\cdot, x) - \frac{\delta}{nt+1} \nabla dF_p(\cdot, x) \left( \frac{y}{\sqrt{nt}}, \cdot \right)^\sharp + O\left(\frac{1}{n^2}\right).$$

So finally

$$(3.32) \quad \frac{nt+1}{\sqrt{n}} \partial_a c(0,1) = y + \frac{y}{nt} - \frac{\delta}{\sqrt{n}} \operatorname{grad}_e F_p(\cdot, x) - \delta \nabla dF_p(\cdot, x) \left( \frac{y}{nt}, \cdot \right)^\sharp + O\left(n^{-3/2}\right).$$

To get the final result we are left to make a Taylor expansion of  $f$  up to order 2.  $\square$

Define the following quantities:

$$(3.33) \quad b_n(z) = n \int_{\{|z'-z|\leq 1\}} (z' - z) P^n(z, dz')$$

and

$$(3.34) \quad a_n(z) = n \int_{\{|z'-z|\leq 1\}} (z' - z) \otimes (z' - z) P^n(z, dz').$$

The following property holds:

**Lemma 3.2.** *Assume that there exists a neighbourhood  $V_e$  of  $e$  such that for all  $x \in B(a, r)$  the application  $F_p(\cdot, x)$  is of class  $C^2$  in  $V_e$ .*

- (1) *For all  $R > 0$  and  $\varepsilon > 0$ , there exists  $n_0$  such that for all  $n \geq n_0$  and  $z \in [\varepsilon, T] \times B(0_e, R)$ , where  $B(0_e, R)$  is the open ball in  $T_e M$  centered at the origin with radius  $R$ ,*

$$(3.35) \quad \int 1_{\{|z'-z|>1\}} P^n(z, dz') = 0.$$

(2) For all  $R > 0$  and  $\varepsilon > 0$ ,

$$(3.36) \quad \lim_{n \rightarrow \infty} \sup_{z \in [\varepsilon, T] \times B(0_\varepsilon, R)} |b_n(z) - b(z)| = 0$$

with

$$(3.37) \quad b(z) = \left(1, \frac{1}{t} L_\delta(y)\right) \quad \text{and} \quad L_\delta(y) = y - \delta \nabla dH(y, \cdot)^\sharp.$$

(3) For all  $R > 0$  and  $\varepsilon > 0$ ,

$$(3.38) \quad \lim_{n \rightarrow \infty} \sup_{z \in [\varepsilon, T] \times B(0_\varepsilon, R)} |a_n(z) - a(z)| = 0$$

with

$$(3.39) \quad a(z) = \delta^2 \text{diag}(0, \Gamma) \quad \text{and} \quad \Gamma = \mathbb{E}[\text{grad}_e F_p(\cdot, P_1) \otimes \text{grad}_e F_p(\cdot, P_1)].$$

*Proof.* (1) We use the notation  $z = (t, y)$  and  $z' = (t', y')$ . We have

$$\begin{aligned} & \int 1_{\{|z' - z| > 1\}} P^n(z, dz') \\ &= \int 1_{\{\max(|t' - t|, |y' - y|) > 1\}} P^n(z, dz') \\ &= \int 1_{\{\max(\frac{1}{n}, |y' - y|) > 1\}} P^n(z, dz') \\ &= \mathbb{P} \left[ \left| \frac{nt + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt + 1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, P_1) \right) \right) - y \right| > 1 \right]. \end{aligned}$$

On the other hand, since  $F_p(\cdot, x)$  is of class  $C^2$  in a neighbourhood of  $e$ , we have by (3.32)

$$(3.40) \quad \left| \frac{nt + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt + 1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, P_1) \right) \right) - y \right| \leq \frac{C\delta}{\sqrt{n\varepsilon}}$$

for some constant  $C > 0$ .

(2) Equation (3.35) implies that for  $n \geq n_0$

$$\begin{aligned} & b_n(z) \\ &= n \int (z' - z) P^n(z, dz') \\ &= n \left( \frac{1}{n}, \mathbb{E} \left[ \frac{nt + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{y}{\sqrt{nt}}} \left( -\frac{\delta}{nt + 1} \text{grad}_{\exp_e \frac{y}{\sqrt{nt}}} F_p(\cdot, P_1) \right) \right) \right] - y \right). \end{aligned}$$

We have by lemma 3.1

$$\begin{aligned} & n \left( \frac{nt + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt + 1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, P_1) \right) \right) - y \right) \\ &= \frac{1}{t} y - \delta \sqrt{n} \text{grad}_e F_p(\cdot, P_1) - \delta \nabla dF_p(\cdot, P_1) \left( \frac{1}{t} y, \cdot \right)^\sharp + O \left( \frac{1}{n^{1/2}} \right) \end{aligned}$$

a.s. uniformly in  $n$ , and since

$$\mathbb{E} [\delta \sqrt{n} \text{grad}_e F_p(\cdot, P_1)] = 0,$$

this implies that

$$n \left( \mathbb{E} \left[ \frac{nt + 1}{\sqrt{n}} \exp_e^{-1} \left( \exp_{\exp_e \frac{1}{\sqrt{nt}} y} \left( -\frac{\delta}{nt + 1} \text{grad}_{\exp_e \frac{1}{\sqrt{nt}} y} F_p(\cdot, P_1) \right) \right) \right] - y \right)$$

converges to

$$(3.41) \quad \frac{1}{t}y - \mathbb{E} \left[ \delta \nabla dF_p(\cdot, P_1) \left( \frac{1}{t}y, \cdot \right) \right]^\# = \frac{1}{t}y - \delta \nabla dH_p \left( \frac{1}{t}y, \cdot \right)^\#.$$

Moreover the convergence is uniform in  $z \in [\varepsilon, T] \times B(0_e, R)$ , so this yields (3.36).

(3) In the same way, using lemma 3.1,

$$\begin{aligned} & n \int (y' - y) \otimes (y' - y) P^n(z, dz') \\ &= \frac{1}{n} \mathbb{E} [(-\sqrt{n}\delta \operatorname{grad}_e F_p(\cdot, P_1)) \otimes (-\sqrt{n}\delta \operatorname{grad}_e F_p(\cdot, P_1))] + o(1) \\ &= \delta^2 \mathbb{E} [\operatorname{grad}_e F_p(\cdot, P_1) \otimes \operatorname{grad}_e F_p(\cdot, P_1)] + o(1) \end{aligned}$$

uniformly in  $z \in [\varepsilon, T] \times B(0_e, R)$ , so this yields (3.38).  $\square$

**Lemma 3.3.** *Suppose that  $t_n = \frac{\delta}{n}$  for some  $\delta > 0$ . For all  $\delta > C_{p,\mu,K}^{-1}$ ,*

$$(3.42) \quad \sup_{n \geq 1} n \mathbb{E} [\rho^2(e, X_n)] < \infty.$$

*Proof.* First consider the case  $p \in [1, 2)$ .

We know by (3.12) that there exists some constant  $C(\beta, r, p)$  such that

$$(3.43) \quad \mathbb{E} [\rho^2(e, X_{k+1})] \leq \mathbb{E} [\rho^2(e, X_k)] \exp(-C_{p,\mu,K} t_{k+1}) + C(\beta, r, p) t_{k+1}^2.$$

From this (3.42) is a consequence of Lemma 0.0.1 (case  $\alpha > 1$ ) in [16]. We give the proof for completeness. We deduce easily by induction that for all  $k \geq k_0$ ,

$$(3.44) \quad \begin{aligned} & \mathbb{E} [\rho^2(e, X_k)] \\ & \leq \mathbb{E} [\rho^2(e, X_{k_0})] \exp \left( -C_{p,\mu,K} \sum_{j=k_0+1}^k t_j \right) + C(\beta, r, p) \sum_{i=k_0+1}^k t_i^2 \exp \left( -C_{p,\mu,K} \sum_{j=i+1}^k t_j \right), \end{aligned}$$

where the convention  $\sum_{j=k+1}^k t_j = 0$  is used. With  $t_n = \frac{\delta}{n}$ , the following inequality holds for all  $i \geq k_0$  and  $k \geq i$ :

$$(3.45) \quad \sum_{j=i+1}^k t_j = \delta \sum_{j=i+1}^k \frac{1}{j} \geq \delta \int_{i+1}^{k+1} \frac{dt}{t} \geq \delta \ln \frac{k+1}{i+1}.$$

Hence,

$$(3.46) \quad \begin{aligned} & \mathbb{E} [\rho^2(e, X_k)] \\ & \leq \mathbb{E} [\rho^2(e, X_{k_0})] \left( \frac{k_0+1}{k+1} \right)^{\delta C_{p,\mu,K}} + \frac{\delta^2 C(\beta, r, p)}{(k+1)^{\delta C_{p,\mu,K}}} \sum_{i=k_0+1}^k \frac{(i+1)^{\delta C_{p,\mu,K}}}{i^2}. \end{aligned}$$

For  $\delta C_{p,\mu,K} > 1$  we have as  $k \rightarrow \infty$

$$(3.47) \quad \frac{\delta^2 C(\beta, r, p)}{(k+1)^{\delta C_{p,\mu,K}}} \sum_{i=k_0+1}^k \frac{(i+1)^{\delta C_{p,\mu,K}}}{i^2} \sim \frac{\delta^2 C(\beta, r, p)}{(k+1)^{\delta C_{p,\mu,K}}} \frac{k^{\delta C_{p,\mu,K}-1}}{\delta C_{p,\mu,K}-1} \sim \frac{\delta^2 C(\beta, r, p)}{\delta C_{p,\mu,K}-1} k^{-1}$$



and

$$\mathbb{E} [\rho^2(e, X_{k_0})] \left( \frac{k_0 + 1}{k + 1} \right)^{\delta C_{p,\mu,K}} = o(k^{-1}).$$

This implies that the sequence  $k\mathbb{E} [\rho^2(e, X_k)]$  is bounded.

Next consider the case  $p \geq 2$ .

Now we have by (3.19) that

$$(3.48) \quad \mathbb{E} [H_p(X_{k+1}) - H_p(e)] \leq \mathbb{E} [H_p(X_k) - H_p(e)] \exp(-C_{p,\mu,K} t_{k+1}) + C(\beta, r, p) t_{k+1}^2.$$

From this, arguing similarly, we obtain that the sequence  $k\mathbb{E} [H_p(X_k) - H_p(e)]$  is bounded. We conclude with (2.3).  $\square$

**Lemma 3.4.** *Assume  $\delta > C_{p,\mu,K}^{-1}$  and that  $H_p$  is  $C^2$  in a neighbourhood of  $e$ . For all  $0 < \varepsilon < T$ , the sequence of processes  $\left( Y_{[nt]}^n \right)_{\varepsilon \leq t \leq T}$  is tight in  $\mathbb{D}([\varepsilon, T], \mathbb{R}^d)$ .*

*Proof.* Denote by  $\left( \tilde{Y}_\varepsilon^n = \left( Y_{[nt]}^n \right)_{\varepsilon \leq t \leq T} \right)_{n \geq 1}$ , the sequence of processes. We prove that from any subsequence  $\left( \tilde{Y}_\varepsilon^{\phi(n)} \right)_{n \geq 1}$ , we can extract a further subsequence  $\left( \tilde{Y}_\varepsilon^{\psi(n)} \right)_{n \geq 1}$  that weakly converges in  $\mathbb{D}([\varepsilon, 1], \mathbb{R}^d)$ .

Let us first prove that  $\left( \tilde{Y}_\varepsilon^{\phi(n)}(\varepsilon) \right)_{n \geq 1}$  is bounded in  $L^2$ .

$$\left\| \tilde{Y}_\varepsilon^{\phi(n)}(\varepsilon) \right\|_2^2 = \frac{[\phi(n)\varepsilon]^2}{\phi(n)} \mathbb{E} [\rho^2(e, X_{[\phi(n)\varepsilon]})] \leq \varepsilon \sup_{n \geq 1} (n\mathbb{E} [\rho^2(e, X_n)])$$

and the last term is bounded by lemma 3.3.

Consequently  $\left( \tilde{Y}_\varepsilon^{\phi(n)}(\varepsilon) \right)_{n \geq 1}$  is tight. So there is a subsequence  $\left( \tilde{Y}_\varepsilon^{\psi(n)}(\varepsilon) \right)_{n \geq 1}$  that weakly converges in  $T_e M$  to the distribution  $\nu_\varepsilon$ . Thanks to Skorohod theorem which allows to realize it as an a.s. convergence and to lemma 3.2 we can apply Theorem 11.2.3 of [20], and we obtain that the sequence of processes  $\left( \tilde{Y}_\varepsilon^{\psi(n)} \right)_{n \geq 1}$  weakly converges to a diffusion  $(y_t)_{\varepsilon \leq t \leq T}$  with generator  $G_\delta(t)$  given by (2.13) and such that  $y_\varepsilon$  has law  $\nu_\varepsilon$ . This achieves the proof of lemma 3.4.  $\square$


*Proof of Theorem 2.6.* Let  $\tilde{Y}^n = \left( Y_{[nt]}^n \right)_{0 \leq t \leq T}$ . It is sufficient to prove that any subsequence of  $\left( \tilde{Y}^n \right)_{n \geq 1}$  has a further subsequence which converges in law to  $(y_\delta(t))_{0 \leq t \leq T}$ . So let  $\left( \tilde{Y}^{\phi(n)} \right)_{n \geq 1}$  a subsequence. By lemma 3.4 with  $\varepsilon = 1/m$  there exists a subsequence which converges in law on  $[1/m, T]$ . Then we extract a sequence indexed by  $m$  of subsequence and take the diagonal subsequence  $\tilde{Y}^{\eta(n)}$ . This subsequence converges in  $\mathbb{D}((0, T], \mathbb{R}^d)$  to  $(y'(t))_{t \in (0, T]}$ . On the other hand, as in the proof of lemma 3.4, we have

$$\|\tilde{Y}^{\eta(n)}(t)\|_2^2 \leq Ct$$

for some  $C > 0$ . So  $\|\tilde{Y}^{\eta(n)}(t)\|_2^2 \rightarrow 0$  as  $t \rightarrow 0$ , which in turn implies  $\|y'(t)\|_2^2 \rightarrow 0$  as  $t \rightarrow 0$ . The unicity statement in Proposition 2.5 implies that  $(y'(t))_{t \in (0, T]}$  and  $(y_\delta(t))_{t \in (0, T]}$  are equal in law. This achieves the proof.  $\square$


## 4. EXAMPLES, SIMULATIONS AND PICTURES

## 4.1. Median of the uniform measure on an equilateral triangle in the plane.




Here  $M$  is the Euclidean plane  $\mathbb{R}^2$ ,  $\mu$  is the uniform measure on an equilateral triangle. The red path represents one trajectory of the inhomogeneous Markov chain  $(X_k)_{k \geq 0}$  corresponding to  $p = 1$ , with linear interpolation between the different steps. The red point in the middle is  $e_1$ .

## 4.2. Uniform measure on the unit square in the plane.




Again  $M$  is the Euclidean plane  $\mathbb{R}^2$ , but here  $\mu$  is the uniform measure on the square  $[0, 1] \times [0, 1]$ . The red path represents one trajectory of the inhomogeneous Markov chain  $(X_k)_{k \geq 0}$  corresponding to  $p = 1$ , with linear interpolation between the different steps. The red point in the middle is  $e_1$ .

## 4.3. A non uniform measure on a square in the plane.



$M$  is still the Euclidean plane  $\mathbb{R}^2$ , now here  $\mu$  is the renormalized restriction to the square  $[0, 4] \times [0, 4]$  of an exponential law on  $[0, \infty) \times [0, \infty)$ . The red path represents one trajectory of the inhomogeneous Markov chain  $(X_k)_{k \geq 0}$  corresponding to  $p = 1$ , with linear interpolation between the different steps. The red point is  $e_1$ . Black circles represent the values of  $(P_k)_{k \geq 1}$


#### 4.4. A non uniform measure on the sphere $S^2$ .



(4.4)


Here  $M$  is the embedded sphere  $S^2$  and  $\mu$  is a non uniform law. Again the red path represents one trajectory of the inhomogeneous Markov chain  $(X_k)_{k \geq 0}$  corresponding to  $p = 1$ , with linear interpolation between the different steps. The red point is  $e_1$ . Black circles represent the values of  $(P_k)_{k \geq 1}$ . One can observe that even if the convexity assumptions are not fulfilled, convergence still holds.

4.5. Another non uniform measure on the unit square in the plane.



(4.5)

This is an example of density on the square  $[0, 1] \times [0, 1]$ .



(4.6)

Here  $\mu$  is given by the density of figure (4.5). The red path represents one trajectory of the inhomogeneous Markov chain  $(X_k)_{k \geq 0}$  corresponding to  $p = 1$ , with linear interpolation between the different steps. The red point is  $e_1$ . Black circles represent the values of  $(P_k)_{k \geq 1}$ .

#### REFERENCES

- [1] B. Afsari, *Riemannian  $L^p$  center of mass : existence, uniqueness, and convexity*, Proceedings of the American Mathematical Society, S 0002-9939(2010)10541-5, Article electronically published on August 27, 2010.

- [2] M. Arnaudon and X.M. Li, *Barycenters of measures transported by stochastic flows*, The Annals of Probability, 33 (2005), no. 4, 1509–1543
- [3] A. Benveniste, M. Goursat and G. Ruget, *Analysis of stochastic approximation schemes with discontinuous and dependent forcing terms with applications to data communication algorithm*, IEEE Transactions on Automatic Control, Vol. AC-25, no. 6, December 1980.
- [4] E. Berger, *An almost sure invariance principle for stochastic approximation procedures in linear filtering theory*, the Annals of applied probability, vol. 7, no. 2 (May 1997), pp. 444–459.
- [5] M. Emery and G. Mokobodzki, *Sur le barycentre d'une probabilité dans une variété*, Séminaire de Probabilités XXV, Lecture Notes in Mathematics 1485 (Springer, Berlin, 1991), pp. 220–233
- [6] S. Gouëzel, *Almost sure invariance principle for dynamical systems by spectral methods* The annals of Probability, Vol 38, no. 4 (2010), pp. 1639–1671
- [7] P.T. Fletcher, S. Venkatasubramanian, S. Joshi, *The geometric median on Riemannian manifolds with application to robust atlas estimation*, NeuroImage, 45 (2009), pp. S143–S152
- [8] H. Karcher, *Riemannian center of mass and mollifier smoothing*, Communications on Pure and Applied Mathematics, vol XXX (1977), 509–541
- [9] R.Z. Khas'Minskii, *On stochastic processes defined by differential equations with a small parameter*, Theory Prob. Appl., vol. XI-2 (1968), pp. 211–228
- [10] W.S. Kendall, *Probability, convexity and harmonic maps with small image I: uniqueness and fine existence*, Proc. London Math. Soc. (3) 61 no. 2 (1990) pp. 371–406
- [11] W.S. Kendall, *Convexity and the hemisphere*, J. London Math. Soc., (2) 43 no.3 (1991), pp. 567–576
- [12] H.W. Kuhn, *A note on Fermat's problem*, Mathematical Programming, 4 (1973), pp. 98–107
- [13] H. Le, *Estimation of Riemannian barycentres*, LMS J. Comput. Math. 7 (2004), pp. 193–200
- [14] L. Ljung, *Analysis of recursive stochastic algorithms*, IEEE Transactions on Automatic Control, Vol. AC-22, no. 4, August 1977.
- [15] D. L. MCLeish, *Dependent central limit theorems and invariance principles*, the Annals of Probability, vol. 2, no. 4 (1974), pp. 620–628
- [16] A. Nedic and D.P. Bertsekas, *Convergence rate of incremental subgradient algorithms*, Stochastic optimization : algorithms and applications (S. Uryasev and P.M. Pardalos, Editors), Kluwer Academic Publishers (2000), pp. 263–304
- [17] L.M.JR. Ostresh, *On the convergence of a class of iterative methods for solving Weber location problem*, Operation. Research, 26 (1978), no. 4
- [18] J. Picard, *Barycentres et martingales dans les variétés*, Ann. Inst. H. Poincaré Probab. Statist. 30 (1994), pp. 647–702
- [19] A. Sahib, *Espérance d'une variable aléatoire á valeurs dans un espace métrique*, Thèse de l'université de Rouen (1998)
- [20] D.W. Stroock and S.R.S. Varadhan, *Multidimensional diffusion processes*, Grundlehren der mathematischen Wissenschaften 233, Springer, 1979.
- [21] E. Weiszfeld, *Sur le point pour lequel la somme des distances de n points donnés est minimum*, Tohoku Math. J. 43 (1937), pp. 355–386
- [22] L.Yang, *Riemannian median and its estimation*, to appear in LMS Journal of Computation and Mathematics

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS  
CNRS: UMR 6086  
UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179  
F-86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE  
*E-mail address:* [marc.arnaudon@math.univ-poitiers.fr](mailto:marc.arnaudon@math.univ-poitiers.fr)

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS  
CNRS: UMR 6086  
UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179  
F-86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE  
*E-mail address:* [clement.dombry@math.univ-poitiers.fr](mailto:clement.dombry@math.univ-poitiers.fr)

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS  
CNRS: UMR 6086  
UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179  
F-86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE  
*E-mail address:* [anthony.phan@math.univ-poitiers.fr](mailto:anthony.phan@math.univ-poitiers.fr)

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS  
CNRS: UMR 6086  
UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179  
F-86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE  
*E-mail address:* [le.yang@math.univ-poitiers.fr](mailto:le.yang@math.univ-poitiers.fr)