



**HAL**  
open science

## Role of the HBx oncoprotein in carbonic anhydrase 9 induction

Tereza Holotnakova, Lucia Tylkova, Martina Takacova, Juraj Kopacek, Juraj Petrik, Silvia Pastorekova, Jaromir Pastorek

► **To cite this version:**

Tereza Holotnakova, Lucia Tylkova, Martina Takacova, Juraj Kopacek, Juraj Petrik, et al.. Role of the HBx oncoprotein in carbonic anhydrase 9 induction. *Journal of Medical Virology*, 2009, 82 (1), pp.32. 10.1002/jmv.21671 . hal-00541152

**HAL Id: hal-00541152**

**<https://hal.science/hal-00541152>**

Submitted on 30 Nov 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Role of the HBx oncoprotein in carbonic anhydrase 9 induction**

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-09-1390.R1
Wiley - Manuscript type:	Research Article
Keywords:	hypoxia, hepatitis B virus, hypoxia-inducible factor 1, hepatocarcinogenesis, hepatocellular carcinoma



Review

HBx and CA9 induction

**Role of the HBx oncoprotein in carbonic anhydrase 9 induction****Shortened title: HBx and CA9 induction**

Tereza Holotnakova<sup>1</sup>, Lucia Tylkova<sup>2</sup>, Martina Takacova<sup>1</sup>, Juraj Kopacek<sup>1</sup>, Juraj Petrik<sup>3</sup>,  
Silvia Pastorekova<sup>1</sup> and Jaromir Pastorek<sup>1</sup>

<sup>1</sup>Institute of Virology and <sup>2</sup>Institute of Molecular Physiology and Genetics, Slovak Academy of  
Sciences, Bratislava, Slovak Republic, <sup>3</sup>Scottish National Blood Transfusion Service, NHS  
National Services Scotland, Edinburgh, UK

Correspondence to:

Jaromir Pastorek, Institute of Virology, Slovak Academy of Sciences, Dubravská cesta 9, 845 05

Bratislava, E-mail: [virupast@savba.sk](mailto:virupast@savba.sk)

## HBx and CA9 induction

**Abstract**

Carbonic anhydrase 9 (CA9), as one of the most hypoxia-responsive genes, has been associated almost exclusively with hypoxic tumours. Its principal role is in pH regulation which helps tumour cells overcome intracellular acidosis and survive extended periods of time with low oxygen. Hypoxia-inducible factor 1 (HIF-1) is the main transcriptional activator of CA9. Hepatitis B virus X protein (HBx) has been shown to increase the transcriptional activity of HIF-1. HBx is often expressed from the gene integrated in the hepatocytes infected persistently and contributes significantly to alterations in host gene expression that can lead to the development of hepatocellular carcinoma (HCC) associated with HBV. The aim of this study was to determine the effect of HBx on expression of CA9. Transient transfection of HBx led to an increase in the expression of CA9 as assessed by RT-PCR and Western blotting. HBx was able to increase CA9 promoter activity significantly in several cell lines. The effect was mediated via HIF-1 and a functional HRE element located  $-10/-3$  bp upstream of the CA9 transcription initiation site. These data suggest that CA9 may be involved in the development of HCC by contributing to the survival of hepatocytes infected with HBV in liver tissue with fibrosis.

HBx and CA9 induction

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Key Words**

Hypoxia; hepatitis B virus; hypoxia-inducible factor 1; hepatocarcinogenesis; hepatocellular carcinoma

For Peer Review

## HBx and CA9 induction

**Introduction**

Hepatitis B virus (HBV) infection is one of the most common viral infections. It is self-limiting in 90-95% of adults infected acutely, while the majority of neonatal infections become persistent. Similarly, horizontal transmission to children or immunocompromised adults will result in a significant risk of developing persistent infection. Perinatal transmission accounts for most infections in highly endemic areas. Over 350 million people worldwide are infected chronically with HBV, with one million deaths per annum resulting from the complications of chronic infection [Ganem and Prince, 2004; Guidotti and Chisari, 2006].

HBV is an enveloped hepatotropic virus. It is a noncytopathic virus and most of the liver damage occurs as a result of immune responses, especially the cytotoxic T lymphocyte (CTL) response, with contributions from antigen-nonspecific inflammatory cells and possibly platelets facilitating the intrahepatic accumulation of CTLs [Iannacone et al., 2007]. The 3.2-kb-long, partially double-stranded circular DNA genome of HBV encodes the surface, core, polymerase, and X (HBx) genes. HBx is a portion of HBV DNA integrated most frequently into hepatocyte chromosomes during the development of hepatocellular carcinoma (HCC), which is a highly vascularized and malignant solid tumour. Such a tumour phenotype is often associated with an insufficient supply of oxygen, which creates selective pressure in favour of tumour cells that can adapt to this microenvironmental stress. Hypoxia-inducible factor 1 (HIF-1) is the main mediator of hypoxic regulation facilitating the transcription of a battery of genes implicated in angiogenesis, erythropoiesis, anaerobic glycolysis, pH control, and more [Semenza et al., 1991; Semenza et al., 1994; Ke and Costa, 2006; Brahimi-Horn and Pouyssegur, 2007].

HIF-1 is a heterodimer that consists of an oxygen-sensitive  $\alpha$  subunit and a constitutively expressed  $\beta$  subunit [Wang et al., 1995]. Under normoxic conditions, HIF-1 $\alpha$  is hydroxylated by prolyl hydroxylases (PHDs) on specific proline residues localized within an oxygen-dependent degradation domain (ODDD) and by Factor inhibiting HIF-1 (FIH-1) on an asparagine residue in the C-terminal transactivation domain (C-TAD) [Masson et al., 2001; Epstein et al., 2001; Lando et al., 2002]. Von Hippel Lindau protein (pVHL) recognizes the prolyl hydroxylated form of HIF-1 $\alpha$  and recruits ubiquitin ligase for its ubiquitination [Hon et al., 2002]. HIF-1 $\alpha$  is then destroyed in

## HBx and CA9 induction

1  
2  
3 the proteasome. Hydroxylated asparagine residue in the C-TAD domain on the other hand  
4 cannot recruit the p300 transcription co-activator. In hypoxia, the HIF-1 $\alpha$  subunit is not  
5 hydroxylated efficiently by PHDs and FIH-1 because of shortage of oxygen, which serves as the  
6 hydroxylation substrate, and thus HIF-1 $\alpha$  is able to escape degradation mediated by pVHL in the  
7 proteasome. It is then accumulated in the cytoplasm and translocates to the nucleus where it  
8 dimerizes with the constitutive  $\beta$  subunit to form the HIF-1 transcription factor that binds to  
9 hypoxia-response elements (HREs) in the promoters of hypoxia-inducible genes and induces  
10 their transcription [Wang et al., 1995].  
11  
12

13  
14  
15  
16  
17  
18  
19 Although HIF-1 is regulated mainly by oxygen tension, other factors also modulate HIF-1  
20 expression and consequent function, such as nitric oxide, cytokines, growth factors, etc. [Stroka  
21 et al., 2001]. HBx stimulates HIF-1 $\alpha$  stability by blocking the proteasomal degradation mediated  
22 by pVHL and also increases its transcriptional activity [Moon et al., 2004]. The C-terminus of HBx  
23 binds the bHLH/PAS, ODDD and C-TAD domains of HIF-1 $\alpha$  [Yoo et al., 2004]. By binding to  
24 bHLH/PAS, HBx potentiates HIF-1 interaction with its target promoters. Through the C-TAD  
25 domain, HBx can block the association of FIH-1 with HIF-1 $\alpha$ , and inhibit Asn<sup>803</sup> hydroxylation,  
26 thus facilitating binding of the p300/CBP co-activator.  
27  
28  
29  
30  
31  
32  
33  
34  
35

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Several hypoxic tumour markers are direct targets of HIF-1, such as vascular endothelial  
growth factor (VEGF), glucose transporter 1 (GLUT1) or carbonic anhydrase IX (CA IX).  
However, there is variability between expression patterns of these markers in tumours, and out of  
these three genes, CA IX is the most reliable marker of hypoxia [Rafajova et al., 2004]. CA IX is  
one of the most hypoxia-responsive genes and was proposed to serve as an intrinsic marker of  
hypoxia [Potter and Harris, 2003]. It catalyzes the reversible conversion of CO<sub>2</sub> to bicarbonate  
ion and proton like other active carbonic anhydrases, and participates in ion transport and pH  
control [Svastova et al, 2004]. Additionally, CA IX also reduces E-cadherin-mediated adhesion by  
binding to  $\beta$ -catenin [Svastova et al, 2003]. CA IX is a transmembrane isoform associated mainly  
with tumours, in contrast to other CAs that are present mostly in differentiated cells of normal  
tissues [Pastorek et al., 1994]. CA IX in the normal human liver is expressed on the basolateral  
surfaces of all ductal epithelial cells in the portal tracts and is absent from hepatocytes, while

## HBx and CA9 induction

1  
2  
3 some hepatocellular carcinomas exhibit positive immunostaining for CA IX [Saarnio et al, 2001].  
4  
5 Expression pattern of CA IX is determined principally by the strong activation of the CA9 gene  
6  
7 transcription via HIF-1 that binds to the HRE localized in the CA9 promoter proximal to the  
8  
9 transcription start site at the -10/-3 position [Wykoff et al., 2000]. CA IX expression contributes  
10  
11 actively to adaptation to a hypoxic microenvironment, and is crucial during carcinogenesis.  
12

13 Since HIF-1 is the main transcriptional activator of CA9, and HBx possesses the ability to  
14  
15 stimulate HIF-1 $\alpha$ , the reports cited above were the rationale behind the interest to investigate the  
16  
17 influence of HBx on CA9 expression.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Materials and methods

### Cell culture

Rat2 thymidine kinase-deficient immortalized fibroblasts, BP6 fibrosarcoma cells (provided kindly by Dr. J. Zavada, Institute of Molecular Genetics, Prague, Czech Republic), H9c2 rat embryonic cardiomyocytes (provided kindly by Prof. I. Morano, Charité University Medicine, Berlin, Germany), rat liver epithelial stem-like WB-F344 cells (provided kindly by Dr. M. Machala, Veterinary Research Institute, Brno, Czech Republic), HIF-1 $\beta$ -deficient mouse Hepa 1c4 hepatoma cells derived from the wild-type Hepa 1c1c7 cell line (provided kindly by Prof. L. Poellinger, Karolinska Institute, Stockholm, Sweden), immortalized HIF-1 $\alpha$  knock-out mouse embryonic MEF HIF-1 $\alpha$ <sup>-/-</sup> fibroblasts (also provided kindly by Prof. L. Poellinger), and human hepatocellular carcinoma cell lines HepG2 (provided kindly by Prof. L. Poellinger) and PLC/PRF/5 (HPA Culture Collections, Porton Down, UK, Cat No 85061113) were cultivated in Dulbecco's modified Eagle's medium supplemented with 10% FCS (BioWhittaker, Verviers, Belgium) and 40  $\mu$ g/ml gentamicin (Lek, Ljubljana, Slovenia) in a humidified 5% CO<sub>2</sub> incubator at 37°C. Hypoxic treatments were performed in a hypoxic workstation (Ruskin Technology, Bridgend, UK) in 2% O<sub>2</sub>, 5% CO<sub>2</sub>, 2% H<sub>2</sub> and 91% N<sub>2</sub> at 37°C.

### Constructs

As a source of the HBx nucleic acid sequence, we used the pREP4D/HBV991A plasmid containing the complete HBV genome (provided kindly by Dr. G. Alexander, Cambridge University Hospitals, NHS Foundation Trust, Cambridge, UK). HBx was amplified by Phusion polymerase (Finnzymes, Espoo, Finland) with primers flanking the HBx ORF (F – caaaactcatcgggactgacaa; R – acagcttgaggcttgaacag). The resultant 565-bp PCR product was ligated into the pBKS- vector (Stratagene, La Jolla, CA) linearized by EcoRV (Invitrogen, Carlsbad, CA). The insert was subcloned into pcDNA3.1 (Invitrogen) eukaryotic expression plasmid using EcoRI and XhoI (both from Invitrogen). The pcDNA3.1 HBx plasmid was sequenced to ensure that the HBx open reading frame was unaltered.

## HBx and CA9 induction

1  
2  
3 The pGL3 PR5 and pGL3 PR1 plasmids containing the human CA9 core and minimal  
4 promoters were generated by insertions of -173/+37 (PR5) and -50/+37 (PR1) CA9 genomic  
5 regions amplified by PCR upstream of the firefly luciferase gene in pGL3-Basic luciferase reporter  
6 vector (Promega, Madison, WI) [Kopacek et al., 2005]. The pGL3 PR5 muthRE plasmid coding  
7 the human core CA9 promoter with a mutated HRE upstream of the firefly luciferase gene was  
8 created from the original pBMN5HREmut [Kaluz et al., 2002]. Renilla pRL-TK vector (Promega)  
9 served as a transfection efficiency control. Human HIF-1 $\alpha$  cDNA cloned in pcDNA1 expression  
10 plasmid was provided kindly by Prof. P. Maxwell, Imperial College of Science, Technology and  
11 Medicine, London, UK [Wood et al., 1998]. HIF-1 $\alpha$  dominant negative mutant (mut HIF-1 $\alpha$ ) was  
12 generated from pcDNA1-HIF-1 $\alpha$  by a deletion of a large part of ODDD and N-TAD (aa 402-564)  
13 [Holotnakova et al, 2008]. Empty pcDNA3.1 plasmid (Invitrogen) was used as a negative control  
14 for mock transfection. Plasmid containing full-length HIF-1 $\beta$  (pARNT/GEM7) was provided kindly  
15 by Prof. L. Poellinger, Karolinska Institute, Stockholm, Sweden [Lindebro et al., 1995]. Empty  
16 pCMV (Genlantis, San Diego, CA) vector was used as a negative control.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

**Reverse transcription PCR**

33  
34  
35 Rat2 and HepG2 cells were transfected transiently with 1  $\mu$ g of pcDNA3.1 HBx or an  
36 empty pcDNA3.1 vector using GenePorterII reagent (Genlantis) according to the manufacturer's  
37 recommendations, and split the following day. The parallel transfectants were incubated in  
38 normoxia and hypoxia for 24 or 48 h followed by RNA isolation using InstaPure (Eurogentec,  
39 Seraing, Belgium) according to the manufacturer's instructions. Reverse transcription was  
40 performed with M-MuLV reverse transcriptase (Finnzymes) and random heptameric primers (400  
41 ng/ $\mu$ l). The mixture of 3  $\mu$ g total RNA and random primers was heated for 10 min at 70°C, cooled  
42 on ice and supplemented with dNTPs (each at 0.5 mM concentration, Finnzymes), M-MuLV  
43 reverse transcriptase buffer (Finnzymes), and with 200 U of reverse transcriptase M-MuLV,  
44 incubated for 1 h at 42°C, heated to 70°C for 15 min and stored at -20°C until further use.  
45 Reverse transcription PCR (RT-PCR) was performed with Taq polymerase (Promega) in an  
46 automatic DNA thermal cycler (Eppendorf AG, Hamburg, Germany) using gene-specific primers  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## HBx and CA9 induction

1  
2  
3 and primers for  $\beta$ -actin that served as internal standards (Table 1). The PCR protocol consisted  
4  
5 of 94 °C for 3 min followed by 30-35 cycles of: denaturation at 94 °C for 30 s, annealing for 40 s  
6  
7 ( $T_m$  in Table 1) and extension at 72 °C for 40 s, followed by final extension at 72 °C for 5 min. The  
8  
9 intensity of bands corresponding to individual PCR products was evaluated with ImageJ 1.34s  
10  
11 software (Rasband, W.S., ImageJ, U.S. National Institutes of Health, Bethesda, Maryland, USA,  
12  
13 <http://rsb.info.nih.gov/ij/>, 1997-2007). Amount of gene-specific PCR products was expressed  
14  
15 semiquantitatively as the ratio of the intensity of each band to the intensity of the related  $\beta$ -actin  
16  
17 internal standard.  
18  
19

**Immunoblotting**

20  
21  
22 Rat2 and HepG2 cells were transfected transiently with 1  $\mu$ g of pcDNA3.1 HBx or an  
23  
24 empty pcDNA3.1 vector with GenePorterII reagent and split the following day. The parallel  
25  
26 transfectants were incubated in normoxia and hypoxia for 24 or 48 h followed by protein  
27  
28 extraction with RIPA buffer (1% Triton X-100 and 0.1% sodium deoxycholate in PBS) containing  
29  
30 inhibitors of proteases Complete mini (Roche Applied Science, Mannheim, Germany) for 15 min  
31  
32 on ice. The extracts were then centrifuged for 15 min at 13000 rpm in an Eppendorf  
33  
34 microcentrifuge at 4 °C, and total protein concentration was determined by bicinchoninic acid  
35  
36 assay (Pierce, Rockford, IL, USA) according to the manufacturer's instructions. Total protein  
37  
38 extracts (20  $\mu$ g/lane – Rat2, 50  $\mu$ g/lane – HepG2) were separated in 10% sodium dodecyl sulfate-  
39  
40 polyacrylamide gel electrophoresis. After separation, the proteins were blotted onto  
41  
42 polyvinylidene fluoride membrane (Immobilon™, Millipore, Billerica, MA, USA). For CA IX  
43  
44 detection, the membrane was incubated with the M75 primary antibody and secondary anti-  
45  
46 mouse peroxidase-conjugated antibody (Sevapharma, Prague, Czech Republic). For loading  
47  
48 control, the membrane was probed with anti-actin antibody (Santa Cruz Biotechnology, Santa  
49  
50 Cruz, CA, USA) and polyclonal rabbit anti-goat IgG-HRP (horseradish peroxidase; Dako,  
51  
52 Glostrup, Denmark). After repeated washing, the membranes were developed with the ECL  
53  
54 detection system. The intensity of bands was evaluated with ImageJ 1.34s software and  
55  
56  
57  
58  
59  
60

## HBx and CA9 induction

1  
2  
3 expressed as the ratio of the intensity of each CA IX band to the intensity of the related actin  
4  
5 internal standard.  
6  
7

**Luciferase assay**

8  
9  
10  
11 Cells were plated into 35-mm Petri dishes to reach ~70% density on the following day.  
12  
13 Transient transfection was performed with 1  $\mu$ g of luciferase construct containing promoter and  
14  
15 50 ng of pRL-TK plasmid DNA using GenePorterII reagent. To analyze the effect of HBx either  
16  
17 with or without transcription factors (HIF-1 $\beta$ , HIF-1 $\alpha$  mut HIF-1 $\alpha$ ), the plasmids containing  
18  
19 corresponding cDNAs were co-transfected. Cells transfected with control empty vectors served  
20  
21 as negative controls. The day after transfection, cells were trypsinized and plated in triplicates  
22  
23 into 24-well plates. Cells were allowed to attach for 24 h, and then they were either transferred to  
24  
25 hypoxia or maintained in normoxia for additional 24 h. Reporter gene expression was assessed  
26  
27 72 h after transfection using Dual-Luciferase Reporter Assay System (Promega), and luciferase  
28  
29 activity was normalized against renilla activity. Results were analyzed by two-tailed unpaired  
30  
31 *t* test (Student's test) with  $P < 0.05$  considered significant.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Results

### HBx increases CA9 expression

It was demonstrated that the expression of VEGF and the formation of new blood vessels were increased in hepatoma cells transfected with HBx [Lee et al., 2000]. VEGF and CA9 are both hypoxia-regulated genes, and their main transcriptional activator HIF-1 has been reported to be influenced by HBx. The HBx ORF was cloned into pcDNA3.1 under the control of a constitutive CMV promoter, and Rat2 immortalized fibroblasts were transfected transiently with HBx or an empty pcDNA3.1 vector. After splitting the cells the following day, the parallel transfectants were incubated in normoxia or hypoxia for 48 h. Afterwards, the expression of hypoxia-inducible genes was analyzed by RT-PCR and immunoblotting (Fig.1). Increased normoxic transcription can be seen only with VEGF and slightly with GLUT4 in cells transfected with HBx (Fig.1A). In hypoxia, HBx was able to increase the transcription of *Car9* (more than 1.5-fold), GLUT4, PDK1, and slightly of VEGF. HBx had no effect on expression of iNOS or GLUT1 in Rat2 cells. As expected, the levels of HIF-1 $\alpha$  transcript were roughly the same in all the samples. At the protein level, HBx was also able to increase CA IX expression significantly in Rat2 cells (Fig.1B). Each of the examined hypoxia-regulated genes showed different magnitude of induction, most probably as a consequence of different molecular mechanisms involved in their regulation. For example, the regulation of VEGF, GLUT1 and CA9 has several common features but there are also clear differences. The CA9 promoter contains HRE at a position close to the transcription initiation site, serving as its main transcriptional regulator, whereas VEGF possesses HRE almost 1 kb upstream of the transcription start and utilizes it more as an enhancer element [Levy et al, 1995], as well as the GLUT1 promoter, which contains HRE in the enhancer 1 region, more than 2 kb upstream of the transcription site [Ebert et al, 1995]. On the other hand, VEGF mRNA is regulated at the level of posttranscriptional stability, and GLUT1 expression is additionally influenced by intracellular glucose levels.

### HBx induces CA9 promoter activity

## HBx and CA9 induction

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

After realization that HBx was able to increase CA9 expression in Rat2 cells, the effect of HBx on CA9 promoter activity was studied by employing dual luciferase reporter assay (Promega) (Fig.2). Rat cells of different origin were transfected transiently with the pGL3 PR5 construct containing the core human CA9 promoter linked to luciferase reporter. They were also co-transfected with renilla plasmid used as a transfection efficiency control and either with HBx or empty pcDNA3.1. Two days later, sparse (Fig.2A) or dense (Fig.2B) cell cultures were incubated in normoxia and hypoxia for 24 h. HBx elevated CA9 promoter activity significantly in Rat2, BP6 and WB-F344 cell lines under hypoxia (Fig.2A). The promoter activity increased slightly also in hypoxic H9c2 cardiomyocytes. Under normoxic conditions, HBx was able to induce CA9 promoter activity only in WB-F344 cells. In dense Rat2 cell culture, HBx increased the CA9 promoter activity also under normoxia (Fig.2B) probably due to pericellular hypoxia. In this experiment, the effects of HBx on the core (pGL3 PR5) and minimal (pGL3 PR1) CA9 promoters were tested. The effect was more prominent on the minimal promoter most likely because the core promoter contains a silencer element [Kaluz et al., 1999]. Since HBx also increased the activity of the minimal promoter, it can be assumed that the effect is mediated via the functional HRE contained within this region. Indeed, the effect was abolished on the core promoter with mutated HRE (pGL3 PR5 mutHRE). Empty pGL3 vector served as negative control.

**HBx requires both HIF-1 subunits for CA9 promoter activity induction**

The mechanism of HBx action in HIF-1 transactivation is understood incompletely. Its C-terminus binds the bHLH/PAS and C-TAD domains of HIF-1 $\alpha$  and thus potentiates HIF-1 interaction with its target promoters [Yoo et al., 2004]. The aim of this series of experiments was to confirm that HBx mediates CA9 promoter induction through HIF-1 and to find out whether HBx, by binding to the bHLH/PAS domain of HIF-1 $\alpha$ , could block HIF-1 subunit dimerization and whether HIF-1 $\alpha/\beta$  and HBx alone could possibly induce activity of the CA9 promoter.

Mouse Hepa 1c4 hepatoma cells that lack natural expression of HIF-1 $\beta$  were co-transfected transiently with the pGL3 PR1 minimal CA9 promoter luciferase construct, renilla plasmid, and HBx (Fig.3A). They were also co-transfected with an empty pCMV vector or the

## HBx and CA9 induction

1  
2  
3 same amount of HIF-1 $\beta$ . As can be seen in Fig.3A, HIF-1 $\beta$  is indispensable to the induction of  
4  
5 the CA9 promoter mediated by HBx.  
6

7 Just like HIF-1 $\alpha$ , HIF-1 $\beta$  also possesses the bHLH/PAS domain. A similar experiment on  
8  
9 mouse embryonic fibroblasts knocked-out for HIF-1 $\alpha$  (MEF HIF-1 $\alpha$ <sup>-/-</sup>) was performed (Fig.3B).  
10  
11 MEF HIF-1 $\alpha$ <sup>-/-</sup> cells were co-transfected transiently with the pGL3 PR1 construct, renilla plasmid,  
12  
13 HBx, and an empty pCMV vector or the same amount of HIF-1 $\alpha$ . From this experiment, it is  
14  
15 evident that HIF-1 $\alpha$  is crucial to HBx-mediated CA9 promoter induction (Fig.3B). It can thus be  
16  
17 concluded that HBx does not block HIF-1 subunit dimerization and influences the CA9 promoter  
18  
19 through a functional HIF-1 dimer.  
20  
21  
22  
23

**HBx regulation of CA IX expression in human hepatocellular carcinoma cells**

24  
25 All the previous experiments were done on rat and mouse cellular models. The possible  
26  
27 regulation of CA IX expression by HBx in human hepatocellular carcinoma cells was tested first  
28  
29 on a human HepG2 HCC cell line, which was transfected transiently with HBx, or an empty  
30  
31 pcDNA3.1 vector. After splitting the cells the following day, the parallel transfectants were  
32  
33 incubated in normoxia or hypoxia for 24 h. Afterwards, the expression of hypoxia-inducible genes  
34  
35 was analyzed by RT-PCR and immunoblotting (Fig.4). In contrast to Rat2 cells, CA IX expression  
36  
37 was increased significantly in HepG2 cells even under normoxic conditions, at both mRNA and  
38  
39 protein levels. The increase was significant also in hypoxia. The same pattern was seen with  
40  
41 other hypoxia-regulated genes (Fig.4A). HBx stimulated the transcription of GLUT1 and PDK1 in  
42  
43 normoxia, as well as hypoxia, and slightly of VEGF and MCT4 in hypoxia.  
44

45 The hypoxic induction of the CA9 promoter activity was increased significantly by HBx in  
46  
47 HepG2 cells as assessed by dual luciferase reporter assay (Fig.5A). The induction was even  
48  
49 more prominent in the PLC/PRF/5 HCC cell line (Fig.5B). Apart from being transfected  
50  
51 transiently with the pGL3 PR1 construct, renilla plasmid, HBx or empty pcDNA3.1, the cells were  
52  
53 also co-transfected with the dominant-negative HIF-1 $\alpha$  mutant (mut HIF-1 $\alpha$ ) or control pcDNA3.1.  
54  
55 Mut HIF-1 $\alpha$  can dimerize with HIF-1 $\beta$  and bind to DNA but it is not capable of transactivating HIF-  
56  
57 target genes as it lacks the N-TAD domain [Holotnakova et al, 2008]. As can be seen in Fig.5,  
58  
59  
60

## HBx and CA9 induction

1  
2  
3 mut HIF-1 $\alpha$  decreased the CA9 promoter activity stimulated by HBx significantly in both HCC cell  
4  
5 lines. In PLC/PRF/5 cells, the effect of HBx was completely abolished in cells transfected with  
6  
7 mut HIF-1 $\alpha$  (Fig.5B). This experiment provides further proof that HBx influences the CA9  
8  
9 promoter through HIF-1 $\alpha$ .  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review



## Discussion

Apart from cancer and ischaemic diseases, the HIF system plays an important role in embryogenesis, wound healing, exercise, inflammatory diseases, anaemia, and many other situations. There are also viral proteins capable of stabilizing or transactivating HIF and promoting changes in cells reminiscent of hypoxic conditions. One such viral protein is HBx. It has a role in hepatocarcinogenesis because HCC incidence has been reported in animals infected with mammalian hepadnaviruses that have a conserved X-open reading frame in their genomes but not in birds infected with avian hepadnaviruses where the X-ORF is absent [Tang et al., 2006]. Knock-down of HBx reduces the tumourigenicity of HCC cells [Chan and Ng, 2006]. HBx has been regarded as a multifunctional viral regulator that modulates a wide variety of host functions in a rather promiscuous way. It is a trans-activating protein that stimulates HBV gene expression and replication, and also alters host gene expression important to the pathogenesis of chronic liver disease and the development of HCC associated with HBV [Feitelson, 1999].

This study showed for the first time that HBx stimulated the CA9 promoter activity significantly in several different cell types, and that this stimulation was mediated via the hypoxia-response element located in the CA9 promoter proximal to the transcription start site. It also demonstrated that HBx induces the CA9 promoter activity through a functional HIF-1 dimer. The previously proposed mechanisms include HBx-mediated HIF-1 $\alpha$  stabilization in the cytoplasm [Moon et al., 2004] and increased transcriptional activity of HIF-1 $\alpha$  in the nucleus [Yoo et al., 2004]. The C-terminus of HBx interacts physically with the bHLH/PAS, C-terminal ODDD, and C-TAD domains of HIF-1 $\alpha$  [Moon et al., 2004; Yoo et al., 2004]. The interactions with ODDD and C-TAD constrain the access to proline and asparagine residues for the PHD and FIH-1 dioxygenases. The association with bHLH/PAS enhances HIF-1 DNA binding. In the experimental settings described here, the effect of HBx on the CA9 promoter was seen only in conditions of hypoxia and pericellular hypoxia (dense cultures); in normoxia, HBx increased the CA9 promoter activity only in the stem cell-like hepatic WB-F344 cells and hepatocellular carcinoma cells. This could mean that the reported HIF-1 $\alpha$  stabilization by HBx in normoxic conditions is possible only in cells of hepatic origin, which might be connected to the

## HBx and CA9 induction

1  
2  
3 hepatotropism of HBV. It is not unreasonable to assume that a factor, which could help stabilize  
4 the interactions between HBx and HIF-1 $\alpha$  in normoxia, is present in cells of hepatic origin. This  
5 possibility opens a new window of opportunity to explore this area for an important component in  
6 hepatocarcinogenesis induced by HBV.  
7  
8  
9

10  
11 The effect of HBx that was manifested in hypoxia in all the cell lines could be explained  
12 by its ability to stimulate activated transcription [Lin et al., 1998]. Since HBx cannot bind double  
13 stranded DNA directly, protein-protein interactions are crucial for HBx transactivation. Lin et al.  
14 (1998) proposed that HBx could act as a co-activator in transcription through modulation of the  
15 transcriptional machinery and distal binding activators. They suggested that HBx might bridge  
16 the activators (in our case HIF-1) and the transcription machinery or release co-repressors from  
17 the machinery. HBx has also been reported to enhance transcriptional activity of HIF-1 $\alpha$  through  
18 activation of the MAPK pathway [Yoo et al., 2003], which has been implicated previously in CA9  
19 expression induced by density and hypoxia [Kopacek et al., 2005].  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Clearly, HIF target genes must be important in hepatocarcinogenesis induced by HBV  
30 since HBx has an additive effect on HIF-transactivation. One proof is the hypervascularity of  
31 HCC correlating with expression of VEGF [Lee et al., 2000]. Angiogenesis has also been  
32 observed in chronic hepatitis and in liver and biliary cirrhosis [Medina et al., 2004], all preceding  
33 HCC. Most chronic liver diseases are characterized by fibrosis and inflammation. Cirrhosis is the  
34 end stage of liver fibrosis and is related highly to the development of HCC. HBx expression in  
35 hepatocytes induces paracrine activation of surrounding cells and contributes actively to the  
36 development of fibrosis in patients infected by HBV chronically [Martin-Vilchez et al., 2008].  
37 Fibrotic tissue offers resistance to blood flow and to the delivery of oxygen, thus becoming  
38 ischaemic. The activation of acid buffering systems in response to intracellular acidosis in hypoxia  
39 causes an irreversible accumulation of Na<sup>+</sup> that precipitates osmotic hepatocyte lysis by altering  
40 cell volume regulation [Alchera et al., 2008]. Expression of CA IX induced by HIF-1 thus  
41 increases hepatocyte tolerance to hypoxia by preventing the alteration of intracellular pH and Na<sup>+</sup>  
42 homeostasis that lead to ischaemic cell death [Alchera et al., 2008]. Silencing of anion  
43 exchanger 2 (AE2), a member of the chloride-bicarbonate transporter family which has an  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## HBx and CA9 induction

1  
2  
3 important function in pH regulation and is overexpressed in HCC tissues, indicated its role in the  
4 survival of poorly differentiated HCC cells [Hwang et al., 2009]. CA IX is also capable of  
5 interacting directly with AEs and improving their bicarbonate transport activity [Morgan et al.,  
6 2007]. CA IX in the normal human liver is expressed on the basolateral surfaces of all ductal  
7 epithelial cells in the portal tracts and is absent from hepatocytes [Saarnio et al., 2001]. In the  
8 study of Saarnio et al. (2001), 33% of HCCs exhibited positive immunostaining for CA IX. The  
9 role of CA IX in hepatocarcinogenesis is yet unknown, and in vivo data about CA IX expression in  
10 chronic hepatitis and liver fibrosis is missing completely.

11  
12  
13 Jin et al. (2001) found that cells positive for HBx were localized preferentially in the  
14 periportal region of liver from patients with chronic hepatitis or on the periphery of cirrhotic  
15 nodules where high necroinflammatory activity was observed. Only 4% of HCC showed HBx  
16 positivity, whereas 30% of surrounding non-tumour tissue was still HBx-positive. They suggested  
17 that HBx might play a major role at the promotion stage of carcinogenesis because it was related  
18 to the progression of chronic hepatitis and accumulated preferentially in the active pathological  
19 stages. Therefore, it would be interesting to see the correlation between HBx and CA IX  
20 expression, which might be overlapping in the periportal regions during chronic hepatitis.  
21 Upregulated HIF-1 $\alpha$  in HCC is correlated well with portal vein invasion, lymph node metastasis,  
22 and is an adverse prognostic factor for HCC patients [Xie et al., 2008].

23  
24  
25 The significant co-activating effect of HBx on CA9 expression was also prominent in cell  
26 lines of non-hepatic origin. That means that its effects are more general and not solely tissue-  
27 specific. Many patients with HBV infection experience a spectrum of extrahepatic disorders (e.g.  
28 dermatological disease, arthritis, glomerulonephritis, aplastic anaemia, neuropathy, vasculitis,  
29 myocarditis), and the virus has extensive reservoirs of extra-hepatic replication (e.g. lymph  
30 nodes, spleen, bone marrow, kidney, colon, stomach, skin, pancreas, brain, heart, lung) [Rong et  
31 al., 2007]. It is therefore conceivable that chronic HBV infection could upregulate the expression  
32 of hypoxia-inducible genes even in tissues other than the liver. The extrahepatic disorders  
33 associated with HBV have all been linked to hypoxia before.

## HBx and CA9 induction

1  
2  
3 This study demonstrated that HBx increased the CA9 promoter activity efficiently in  
4 several cell lines. This could have implications in chronic hepatitis and during the development of  
5 HCC, where CA IX could help increase hepatocyte tolerance to ischaemia caused by fibrosis.  
6  
7 Targeting CA IX, as well as other hypoxia-regulated genes that favour tumour cells able to adapt  
8 to this microenvironmental stress, might help decrease the risk of developing HCC. Early  
9 detection still remains the most important factor in prevention.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

HBx and CA9 induction

**Acknowledgements**

We thank Dr. Graeme Alexander from the Department of Medicine, Cambridge University Hospitals, NHS Foundation Trust and Dr. Christopher Prowse from the Scottish National Blood Transfusion Service, NHS National Services Scotland for the critical reading of the manuscript.

This work was supported by grants from EU (7<sup>th</sup> Framework Program of the European Commission – METOXIA FP7-HEALTH-2007-222741) and from Research and Development Support Agency (Contract APVV-51-024805).

For Peer Review

## HBx and CA9 induction

## References

- Alchera E, Tacchini L, Imarisio C, Dal Ponte C, De Ponti C, Gammella E, Cairo G, Albano E, Carini R. 2008. Adenosine-dependent activation of hypoxia-inducible factor-1 induces late preconditioning in liver cells. *Hepatology* 48:230-239.
- Brahimi-Horn MC, Pouyssegur J. 2007. Oxygen, a source of life and stress. *FEBS Lett* 581:3582-3591.
- Chan DW, Ng IO. 2006. Knock-down of hepatitis B virus X protein reduces the tumorigenicity of hepatocellular carcinoma cells. *J Pathol* 208:372-380.
- Ebert BL, Firth JD, Ratcliffe PJ. 1995. Hypoxia and mitochondrial inhibitors regulate expression of glucose transporter-1 via distinct Cis-acting sequences. *J Biol Chem* 250:29083-29089.
- Epstein AC, Gleadle JM, McNeil LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Metzen E, Wilson MI, Dhanda A, Tian YM, Masson N, Hamilton DL, Jaakkola P, Barstead R, Hodgkin J, Maxwell PH, Pugh CW, Schofield CJ, Ratcliffe PJ. 2001. *C.elegans* EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell* 107:43-54.
- Feitelson M. 1999. Hepatitis B virus in hepatocarcinogenesis. *J Cell Physiol* 181:188-202.
- Ganem D, Prince AM. 2004. Hepatitis B virus infection—natural history and clinical consequences. *N Engl J Med* 350:1118-1129.
- Guidotti LG, Chisari FV. 2006. Immunobiology and pathogenesis of viral hepatitis. *Annu Rev Pathol* 1:23-61.
- Holotnakova T, Ziegelhoffer A, Ohradanova A, Hulikova A, Novakova M, Kopacek J, Pastorek J, Pastorekova S. 2008. Induction of carbonic anhydrase IX by hypoxia and chemical disruption of oxygen sensing in rat fibroblasts and cardiomyocytes. *Pflugers Arch* 456:323-337.
- Hon WC, Wilson MI, Harlos K, Claridge TD, Schofield CJ, Pugh CW, Maxwell PH, Ratcliffe PJ, Stuart DI, Jones EY. 2002. Structural basis for the recognition of hydroxyproline in HIF-1 by pVHL. *Nature* 417:975-978.

## HBx and CA9 induction

- 1  
2  
3 Hwang JM, Kao SH, Hsieh YH, Li KL, Wang PH, Hsu LS, Liu JY. 2009. Reduction of anion  
4 exchanger 2 expression induces apoptosis of human hepatocellular carcinoma cells. *Mol*  
5 *Cell Biochem* [Epub ahead of print].  
6  
7  
8  
9 Iannacone M, Sitia G, Ruggeri ZM, Guidotti LG. 2007. HBV pathogenesis in animal models:  
10 recent advances on the role of platelets. *J Hepatol* 46:719-726.  
11  
12 Jin YM, Yun C, Park C, Wang HJ, Cho H . 2001. Expression of hepatitis B virus X protein is  
13 closely correlated with the high periportal inflammatory activity of liver diseases. *J Viral*  
14 *Hepatits* 8:322-330.  
15  
16  
17  
18 Kaluz S, Kaluzova M, Chrastina A, Olive PL, Pastorekova S, Pastorek J, Lerman MI, Stanbridge  
19 EJ. 2002. Lowered oxygen tension induces expression of the hypoxia marker MN/Carbonic  
20 anhydrase IX in the absence of hypoxia-inducible factor 1 $\alpha$  stabilization: a role for  
21 phosphatidylinositol 3'-kinase. *Cancer Res* 62:4469-4477.  
22  
23  
24  
25  
26  
27 Kaluz S, Kaluzova M, Opavsky R, Pastorekova S, Gibadulinova A, Dequiedt F, Kettmann R,  
28 Pastorek J. 1999. Transcriptional regulation of the MN/CA 9 gene coding for the tumor-  
29 associated carbonic anhydrase IX. Identification and characterization of a proximal silencer  
30 element. *J Biol Chem* 274:32588-32595.  
31  
32  
33  
34  
35 Ke Q, Costa M, 2006. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol* 70:1469-1480.  
36  
37 Kietzmann T, Cornesse Y, Brechtel K, Modaressi S, Jungermann K. 2001. Perivenous expression  
38 of the mRNA of the three hypoxia-inducible factor  $\alpha$  subunits, HIF1 $\alpha$ , HIF2 $\alpha$  and HIF3 $\alpha$ , in  
39 rat liver. *Biochem J* 354:531-537.  
40  
41  
42  
43 Kopacek J, Barathova M, Dequiedt F, Sepelakova J, Kettmann R, Pastorek J, Pastorekova S.  
44 2005. MAPK pathway contributes to density- and hypoxia-induced expression of the tumor-  
45 associated carbonic anhydrase IX. *Biochim Biophys Acta* 1729:41-49.  
46  
47  
48  
49 Lando D, Peet DJ, Whelan DA, Gorman JJ, Whitelaw ML. 2002. Asparagine hydroxylation of the  
50 HIF transactivation domain a hypoxic switch. *Science* 295:858-861.  
51  
52  
53 Lee SW, Lee YM, Bae SK, Murakami S, Yun Y, Kim KW. 2000. Human hepatitis B virus X protein  
54 is a possible mediator of hypoxia-induced angiogenesis in hepatocarcinogenesis. *Biochemi*  
55 *and Biophys Res Commun* 268:456-461.  
56  
57  
58  
59  
60

## HBx and CA9 induction

- 1  
2  
3 Levy AP, Levy NS, Wegner S, Goldberg MA. 1995. Transcriptional regulation of the rat vascular  
4 endothelial growth factor gene by hypoxia. *J Biol Chem* 270:13333-13340.  
5  
6  
7 Li SK, Ho SF, Tsui KW, Fung KP, Waye MYM. 2008. Identification of functionally important amino  
8 acid residues in the mitochondria targeting sequence of Hepatitis B virus X protein.  
9 *Virology* 381:81-88.  
10  
11  
12 Lin Y, Tang H, Nomura T, Dorjsuren D, Hayashi N, Wei W, Ohta T, Roeder R, Murakami S. 1998.  
13 The hepatitis B virus X protein is a co-activator of activated transcription that modulates the  
14 transcription machinery and distal binding activators. *JBC* 273:27097-27103.  
15  
16  
17 Lindebro MC, Poellinger L, Whitelaw ML. 1995. Protein-protein interaction via PAS domains: role  
18 of the PAS domain in positive and negative regulation of the bHLH/PAS dioxin receptor-  
19 Arnt transcription factor complex. *EMBO J* 14:3528-3539.  
20  
21  
22 Martin-Vilchez S, Sanz-Cameno P, Rodriguez-Munoz Y, Majano PL, Molina-Jimenez F, Lopez-  
23 Cabrera M, Moreno-Otero R, Lara-Pezzi E. 2008. The hepatitis B virus X protein induces  
24 paracrine activation of human hepatic stellate cells. *Hepatology* 47:1872-1883.  
25  
26  
27 Masson N, Willam C, Maxwell PH, Pugh CW, Ratcliffe PJ. 2001. Independent function of two  
28 destruction domains in hypoxia-inducible factor- $\alpha$  chains activated by prolyl  
29 hydroxylation. *EMBO J* 20:5197-5206.  
30  
31  
32 Medina J, Arroyo AG, Sanchez-Madrid F, Moreno-Otero R. 2004. Angiogenesis in chronic  
33 inflammatory liver disease. *Hepatology* 39:1185-1195.  
34  
35  
36 Moon EJ, Jeong CH, Jeong JW, Kim KR, Yu DY, Murakami S, Kim CW, Kim KW. 2004. Hepatitis  
37 B virus X protein induces angiogenesis by stabilizing hypoxia-inducible factor-1 $\alpha$ . *FASEB J*  
38 18:382-384.  
39  
40  
41 Morgan PE, Pastorekova S, Stuart-Tilley AK, Alper SL, Casey JR. 2007. Interactions of  
42 transmembrane carbonic anhydrase, CAIX, with bicarbonate transporters. *Am J Physiol*  
43 *Cell Physiol* 293:C738-748.  
44  
45  
46 Pastorek J, Pastorekova S, Callebaut I, Mornon JP, Zelnik V, Opavsky R, Zatovicova M, Liao S,  
47 Portetelle D, Stanbridge EJ, et al. 1994. Cloning and characterization of MN, a human  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## HBx and CA9 induction

1  
2  
3 tumor-associated protein with a domain homologous to carbonic anhydrase and a putative  
4  
5 helix-loop-helix DNA binding segment. *Oncogene* 9:2877-2888.

6  
7 Potter CP, Harris AL. 2003. Diagnostic, prognostic and therapeutic implications of carbonic  
8  
9 anhydrases in cancer. *Br J Cancer* 89:2-7.

10  
11 Rafajova M, Zatovicova M, Kettmann R, Pastorek J, Pastorekova S. 2004. Induction by hypoxia  
12  
13 combined with low glucose or low bicarbonate and high posttranslational stability upon  
14  
15 reoxygenation contribute to carbonic anhydrase IX expression in cancer cells. *Int J Oncol*  
16  
17 24:995-1004.

18  
19 Rong Q, Huang J, Su E, Li J, Li J, Zhang L, Cao K. 2007. Infection of hepatitis B virus in  
20  
21 extrahepatic endothelial tissues mediated by endothelial progenitor cells. *Virology* 4:1-14.

22  
23 Saarnio J, Parkkila S, Parkkila AK, Pastorekova S, Haukipuro K, Pastorek J, Juvonen T,  
24  
25 Karttunen TJ. 2001. Transmembrane carbonic anhydrase, MN/CA IX, is a potential  
26  
27 biomarker for biliary tumours. *J Hepatol* 35:643-649.

28  
29 Semenza GL, Neufeldt MK, Chi SM, Antonarakis SE. 1991. Hypoxia-inducible nuclear factors bind  
30  
31 to an enhancer element located 3' to the human erythropoietin gene. *Proc Natl Acad Sci*  
32  
33 88:5680-5684.

34  
35 Semenza GL, Roth PH, Fang HM, Wang GL. 1994. Transcriptional regulation of genes encoding  
36  
37 glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem* 269:23757-23763.

38  
39 Stroka DM, Burkhardt T, Desbaillets I, Wenger RH, Neil DA, Bauer C, Gassmann M, Candinas D.  
40  
41 2001. HIF-1 is expressed in normoxic tissue and displays an organ-specific regulation  
42  
43 under systemic hypoxia. *FASEB J* 15:2445-2453.

44  
45 Svastova E, Hulikova A, Rafajova M, Zatovicova M, Gibadulinova A, Casini A, Cecchi A,  
46  
47 Scozzafava A, Supuran CT, Pastorek J, Pastorekova S. 2004. Hypoxia activates the  
48  
49 capacity of tumor-associated carbonic anhydrase IX to acidify extracellular pH. *FEBS Lett*  
50  
51 577:439-445.

52  
53 Svastova E, Zilka N, Zatovicova M, Gibadulinova A, Ciampor F, Pastorek J, Pastorekova S. 2003.  
54  
55 Carbonic anhydrase IX reduces E-cadherin-mediated adhesion of MDCK cells via  
56  
57 interaction with beta-catenin. *Exp Cell Res* 290:332-345.

## HBx and CA9 induction

- 1  
2  
3 Tang H, Oishi N, Kaneko S, Murakami S. 2006. Molecular functions and biological roles of  
4 hepatitis B virus x protein. *Cancer Sci* 97:977-983.  
5  
6  
7 Wang GL, Jiang BH, Rue EA, Semenza GL. 1995. Hypoxia-inducible factor 1 is a basic-helix-  
8 loop-helix-PAS heterodimer regulated by cellular O<sub>2</sub> tension. *PNAS* 92:5510-5514.  
9  
10  
11 Wood SM, Wiesener MS, Yeates KM, Okada N, Pugh CW, Maxwell PH, Ratcliffe PJ. 1998.  
12 Selection and analysis of a mutant cell line defective in the hypoxia-inducible factor-1  
13 alpha-subunit (HIF-1alpha). Characterization of hif-1alpha-dependent and -independent  
14 hypoxia-inducible gene expression. *J Biol Chem* 273:8360-8368.  
15  
16  
17  
18  
19 Wykoff C, Beasley N, Watson P, Turner L, Pastorek J, Sibtain A, Wilson G, Turley H, Talks K,  
20 Maxwell PH, Pugh C, Ratcliffe P, Harris AL. 2000. Hypoxia-inducible regulation of tumor-  
21 associated carbonic anhydrases. *Cancer Res* 60:7057-7083.  
22  
23  
24  
25 Xie H, Song J, Liu K, Ji H, Shen H, Hu S, Yang G, Du Y, Zou X, Jin H, Yan L, Liu J, Fan D. 2008.  
26 The expression of hypoxia-inducible factor-1 $\alpha$  in hepatitis B virus-related hepatocellular  
27 carcinoma: correlation with patients' prognosis and hepatitis B virus X protein. *Dig Dis Sci*  
28 53:3225-3233.  
29  
30  
31  
32  
33 Yoo YG, Cho S, Park S, Lee MO. 2004. The carboxy-terminus of the hepatitis B virus X protein is  
34 necessary and sufficient for the activation of hypoxia-inducible factor-1 $\alpha$ . *FEBS Lett*  
35 577:121-126.  
36  
37  
38  
39 Yoo YG, Oh SH, Park ES, Cho H, Lee N, Park H, Kim DK, Yu DY, Seong JK, Lee MO. 2003.  
40 Hepatitis B virus X protein enhances transcriptional activity of hypoxia-inducible factor-1 $\alpha$   
41 through activation of mitogen-activated protein kinase pathway. *J Biol Chem* 278:39076-  
42 39084.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## HBx and CA9 induction

## Tables

Table 1

## List of primers used for RT PCR analysis

Rat mRNA	GenBank® accession no.	Forward primer	Reverse primer	T <sub>m</sub> (°C)	Product size (bp)
β-actin (r)	NM_031144	atcgtgggccgccctaggcac	gtacatggctggggtttgaagg	61	300
β-actin (h)	NM_001101	tcctccctggagaagagcta	acatctgctggaaggaggac	60	367
Car9 (r)	XM_233380	atcaccaggctcagaacacac	gctggggcagcaaagagaa	64	587
CA9 (h)	NM_001216	cggagcgcagcagcctttg	ggctccagctcggctacc	62	253
HIF-1α (r,h)	NM_024359 (r) NM_001530 (h)	gcttgggtgctgatttgaacc	gcatcctgtactgtcctgtggtg	64	267
VEGF (r)	NM_031836	ggagaagttcatggacgtctacc	ctctccgacgtgggcacgcac	61	180
VEGF (h)	NM_001025366	cagcacggtccctcttgaa	cctcctctccctgtcagga	61	313
GLUT1 (r)	NM_138827	cagaaggtaattgaggagtctaca	acaaaggccaacagggtcatcatc	61	206
GLUT1 (h)	NM_006516	ctccttctccagccagcaatg	ccagcagaacgggtggccatag	64	355
GLUT4 (r)	NM_012751	ttccttctatttgccgtcctctg	ttccttcatcctggaggtaaggg	66	280
iNOS (r)	NM_012611	cgcaccaccctcctgttcaac	acaaggcctccaacctctgcct	67	340
MCT4 (h)	NM_004207	ctcaccatcctgggcttcat	agaagaagttgccagcagca	62	427
PDK1 (r,h)	NM_053826 (r) NM_002610 (h)	attggaagcataaatccaaactg	cggtcactcatcttcacagtc	58.5	311

Car9, rat carbonic anhydrase 9; CA9, human carbonic anhydrase 9; GLUT, glucose transporter; (h), human gene; HIF, hypoxia-inducible factor; iNOS, inducible nitric oxide synthase; MCT4, monocarboxylate transporter 4; PDK1, pyruvate dehydrogenase kinase 1; (r), rat gene; VEGF, vascular endothelial growth factor




Figure 1

Effect of HBx on expression of hypoxia-inducible genes. Rat2 cells were transfected transiently with HBx or an empty pcDNA3.1 vector and incubated for 48 h in normoxia or hypoxia. A Expression of hypoxia-inducible genes was analyzed by RT-PCR. Numbers given below the figures represent relative levels of PCR products normalized against the internal standard  $\beta$ -actin. B Expression of CA IX protein was analyzed by Western blotting using the specific M75 monoclonal antibody. Anti-actin antibody was used for loading controls. Relative levels of CA IX protein, calculated relative to actin, are shown below the blot.




Figure 2

Effects of HBx on CA9 promoter activity. A Sparse Rat2, BP6, H9c2, and WB-F344 cells were co-transfected transiently with pGL3 PR5, renilla plasmid and either a plasmid containing HBx or an empty vector (pcDNA3.1). The cells were incubated in normoxia and hypoxia for 24 h. Luciferase activity was normalized by renilla activity and expressed as fold induction. B Dense Rat2 cells were co-transfected transiently with HBx or empty pcDNA3.1 and pGL3 PR5, pGL3 PR1, empty pGL3 vector or pGL3 PR5 mutHRE and renilla plasmid. The cells were incubated in normoxia and hypoxia for 24 h. Luciferase activity was normalized by renilla activity. \*P < 0.05 (HBx vs. pcDNA3.1)




Figure 3

Effect of HIF-1 $\beta$  and HIF-1 $\alpha$  on CA9 promoter activity induced by HBx. A Sparse mouse Hepa 1c4 hepatoma cells that lack natural HIF-1 $\beta$  expression and B mouse embryonic fibroblasts knocked-out for HIF-1 $\alpha$  (MEF HIF-1 $\alpha$ <sup>-/-</sup>) were co-transfected transiently with pGL3 PR1, renilla plasmid and either a plasmid containing HIF-1 $\beta$  (A) or HIF-1 $\alpha$  (B) or an empty vector (pCMV) (A, B). They were also co-transfected with HBx or an empty pcDNA3.1 vector. The cells were incubated in normoxia and hypoxia for 24 h. Luciferase activity was normalized by renilla activity. \*P < 0.05 (HBx vs. pcDNA3.1)




Figure 4

Effect of HBx on expression of hypoxia-inducible genes in HepG2 hepatocellular cell line. HepG2 cells were transfected transiently with HBx or an empty pcDNA3.1 vector and incubated for 24 h in normoxia or hypoxia. A Expression of hypoxia-inducible genes was analyzed by RT-PCR. Numbers given below the figures represent relative levels of PCR products normalized against the internal standard  $\beta$ -actin. B Expression of CA IX protein was analyzed by Western blotting using the specific M75 monoclonal antibody. Anti-actin antibody was used for loading controls. Relative levels of CA IX protein, calculated relative to actin, are shown below the blot.




Figure 5

Effect of HBx on hypoxic induction of CA9 promoter in HCC cell lines. A Human HepG2 and B PLC/PRF/5 hepatocellular carcinoma cells were co-transfected transiently with pGL3 PR1, renilla plasmid and either a plasmid containing the dominant-negative mut HIF-1 $\alpha$  or an empty vector (pcDNA3.1). They were also co-transfected with HBx or an empty pcDNA3.1 vector. The cells were incubated in normoxia and hypoxia for 24 h. Luciferase activity was normalized by renilla activity and expressed as fold induction. Mut HIF-1 $\alpha$  vs. pcDNA3.1 values were all significant ( $P < 0.05$ ). \* $P < 0.05$  (HBx vs. pcDNA3.1)