



HAL
open science

The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in young chickens from 7-28 days of age

Tom Acamovic, Deborah Cross

► To cite this version:

Tom Acamovic, Deborah Cross. The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in young chickens from 7-28 days of age. *British Poultry Science*, 2007, 48 (04), pp.496-506. 10.1080/00071660701463221 . hal-00545318


HAL Id: hal-00545318

<https://hal.science/hal-00545318>

Submitted on 10 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in young chickens from 7-28 days of age

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2006-228.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	01-Mar-2007
Complete List of Authors:	Acamovic, Tom; SAC, ASRC Cross, Deborah; SAC, ASRC
Keywords:	Broilers, Nutrition, Feedstuffs, Anti-nutrients, Microbiology



CBPS-2006-228

ed. MacLeod, May 2007

The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 days to 28 days of age

D.E. CROSS^{1,2}, R.M. McDEVITT¹, K. HILLMAN³ AND T. ACAMOVIC^{1,4}

¹*Avian Sciences Research Centre, SAC, West Mains Road, Edinburgh, EH9 3JG and*

³*Hillman Consultancy, Fairway Avenue, Inverurie AB51 3WY*

RUNNING TITLE: PLANT EXTRACTS IN BROILER NUTRITION

²Current address: Nutrition and Food Science Group, Department of Biological and Molecular Sciences, Oxford Brookes University, Gypsy Lane, Oxford, OX3 0BP

⁴Correspondence address: Avian Sciences Research Centre, SAC, Auchincruive Estate, Ayr, UK, KA6 5HW

Telephone: +44 1292 525100

Fax: +44 1292 525098

Email: Thomas.acamovic@sac.ac.uk

Accepted for publication 25th April 2007

1
2
3
4 28 **Abstract** 1. The effect of the dietary inclusion of 5 culinary herbs or their essential
5
6 29 oils on the growth, digestibility and intestinal microflora status in female broiler
7
8 30 chicks was assessed. From 7-28 d of age, either a basal control diet without
9
10 31 supplement was given or one of 10 others, consisting of the basal diet with either
11
12 32 10g/kg herb (thyme, oregano, marjoram, rosemary or yarrow) or 1 g/kg of essential
13
14 33 oil.
15
16
17 34 2. Body mass (BM) and feed consumption (AFC) were measured on a weekly basis
18
19 35 and used to calculate chick performance. Total viable counts of lactic acid bacteria,
20
21 36 coliforms, anaerobes and *Clostridium perfringens* were determined at 25 d.
22
23 37 Apparent nutrient digestibilities were calculated from the measured values for gross
24
25 38 energy, nitrogen (N), dry matter (DM) and organic matter, and sialic acid
26
27 39 concentration was also measured.
28
29 40 3. Generally, dietary thyme oil or yarrow herb inclusion had the most positive effects
30
31 41 on chick performance, while oregano herb and yarrow oil were the poorest
32
33 42 supplements. Only thyme and yarrow in these diets had a different effect when used
34
35 43 as an herb or oil on weight gain and BM.
36
37 44 4. Dietary treatment had no effect on the intestinal microflora populations, apparent
38
39 45 metabolisable energy (AME) or the calculated coefficients of digestibility. Sialic
40
41 46 acid concentration was greatest in the birds given dietary thyme oil, compared with
42
43 47 all other treatments except those birds receiving marjoram oil, rosemary herb and the
44
45 48 controls. However, less sialic acid was excreted in those birds given diets with
46
47 49 oregano or rosemary oils, or oregano herb, than in the controls.
48
49 50 5. Plant extracts in diets may therefore affect chick performance, gut health and
50
51 51 endogenous secretions, although the chemical composition of the extract appears to
52
53 52 be important in obtaining the optimal effects.
54
55
56
57
58
59
60

1
2
3 53

INTRODUCTION

4
5 54 Plants contain an extensive variety of phytochemical compounds with antimicrobial
6
7
8 55 activity (Cowan, 1999), which may have either beneficial or detrimental effects in
9
10 56 animals (Panter *et al.*, 2004; Wink, 2004; Acamovic and Brooker, 2004), depending
11
12 57 on both the compound used and its concentration. The removal of antimicrobial
13
14 58 growth promoters (AGP) from poultry diets has triggered a search for suitable
15
16 59 natural alternatives, to combat the increased potential for bacterial disease
17
18 60 development in growing flocks, especially under conditions of average management
19
20 61 quality. The use of various plant materials as dietary supplements, including
21
22 62 culinary herbs, may positively affect poultry health and productivity. Essential oils
23
24 63 and herbs are used in human aromatherapy for their holistic mode of action. The
25
26 64 large number of active components in these supplements may therefore present a
27
28 65 more acceptable defence against bacterial attack than synthetic antimicrobials. The
29
30 66 main constituents of the herbs and essential oils are terpenes, which are responsible
31
32 67 for the bulk of the antimicrobial activity (Charai *et al.*, 1996). The oils of *Thymus*
33
34 68 *vulgaris* L. (thyme), *Rosmarinus officinalis* L. (rosemary), *Origanum majorana* L.
35
36 69 (marjoram) (Hammer *et al.*, 1999), *O. vulgare* subsp. *hirtum* (oregano) (Hammer *et*
37
38 70 *al.*, 1999; Dorman and Deans, 2000) and *Achillea millefolium* L. (yarrow) (Candan *et*
39
40 71 *al.*, 2003) have *in vitro* antimicrobial properties. Yarrow has also been reported to
41
42 72 reduce the effects of viruses and helminths (Cowan, 1999). However, there is only
43
44 73 limited evidence for a bioactive effect of herbal compounds in live poultry. It is
45
46 74 unknown whether the inclusion of terpenes in different forms, *i.e.* as solid herb
47
48 75 material or as the extracted oil, would have the same antimicrobial or growth
49
50 76 promoting effects *in vivo*.

1
2
3 77 The present experiment aimed to assess the suitability of a number of
4
5 78 culinary herbs; thyme, marjoram, rosemary, oregano and yarrow, and their extracted
6
7 79 oils as dietary supplements for poultry. The effects of these herbs and oils on the
8
9 80 growth, nutrient utilisation and intestinal microflora in growing broiler chickens were
10
11 81 determined.
12

13 MATERIALS AND METHODS

14 **15 Provision of plant supplements and extraction of essential oils**

16
17 83
18 84 The oregano (*O. vulgare* subsp. *hirtum*), marjoram (*O. majorana* L.), rosemary (*R.*
19
20 85 *officinalis* L.) and thyme (*T. vulgaris* L.) were purchased commercially as dried herb
21
22 86 supplements (Green City, Glasgow, UK) in early 2001. Yarrow (*A. millefolium* L.
23
24 87 var. *Alba*) seeds were selected in Slovakia and imported to the UK, where the yarrow
25
26 88 plants were grown in experimental plots at SAC Ayr (Dr K. Svoboda). The
27
28 89 flowering heads of the yarrow plants were collected in the summer of 2000 and dried
29
30 90 on the day of harvest in an oven at 40°C. Essential oil was distilled from the ground
31
32 91 plant material using Clevenger distillation apparatus in accordance with BS 4585
33
34 92 (1985), following British Pharmacopoeia methods. The oil samples were stored in
35
36 93 sealed glass vials below 4°C until required.
37
38

39 **40 Birds and housing environment**

41
42 94
43 95 The experiment was subjected to an assessment for its ethical acceptability and
44
45 96 approved by the Animal Experimental Committee (AEC) within SAC. The birds
46
47 97 were housed in cages in an environmentally controlled room. Heating was provided
48
49 98 by a single gas brooder, where the initial temperature (study d 0) was set at 32°C and
50
51 99 decreased linearly by 0.5°C per d to a final temperature of 21°C at 21 d (d 21).
52
53 100 Supplementary heating was provided as required by mobile butane gas heaters. The
54
55 101 birds were provided with 1 h of darkness following a period of 23 h light, from the
56
57
58
59
60

1
2
3 102 start to the finish of the study. Each cage was provided with a single feed trough and
4
5 103 nipple line, with 3 nipples per cage, to provide *ad libitum* access to feed and water.
6
7

8 104 **Experimental design**

9
10 105 The chickens (165 Ross 308 female birds) were placed at d-old in a single colony in
11
12 106 a floor pen containing fresh wood shavings to a depth of 10 cm, and all were fed on a
13
14 107 starter ration. At 7 d old, 5 replicate cages containing 3 chicks selected at random
15
16 108 were assigned to each of 11 dietary treatments using a randomised block design,
17
18 109 giving a total of 55 experimental cages. The study was completed at 28 d of age.
19
20 110 Body mass (BM) was measured on an individual basis when the birds were 7, 14, 21
21
22 111 and 28 d of age. The average feed consumption (AFC) was determined on a weekly
23
24 112 basis from 7-28 d. The feed conversion efficiency (FCE) per cage was derived from
25
26 113 the average weight gain (AWG) and AFC per bird on a weekly basis. The treatment
27
28 114 diets contained titanium dioxide (TiO₂) as an indigestible marker and excreta
29
30 115 samples from each cage were collected on d 25. These samples, along with samples
31
32 116 of each diet, were stored at -20°C until required for laboratory analysis. Excreta
33
34 117 samples were then freeze-dried prior to analysis. At the end of the study, a bird from
35
36 118 each cage was randomly selected and killed using 1 ml/kg BM of sodium
37
38 119 pentobarbitone (euthatal), to allow caecal contents collection.
39
40
41
42
43
44

45 120 **Diet formulation**

46
47 121 The basal starter ration was formulated to be marginal in energy and nutrients and
48
49 122 also low in tocopherol (<50 mg/kg), in order to provide the birds with slightly sub-
50
51 123 optimal conditions for growth. This diet contained mainly wheat and soybean meal,
52
53 124 and was manufactured at SAC as a mash (Table 1). The dietary treatments therefore
54
55 125 comprised the basal diet only (control), or, alternatively, the basal diet supplemented
56
57 126 with one of the 5 herbs (10 g/kg) or one of the distilled essential oils from these herbs
58
59
60

1
2
3 127 (1 g/kg), resulting in 11 dietary treatments in total. Since the herbs were assumed to
4
5 128 contain up to about 100 g oil/kg it was decided to include the oils in the diets at 1
6
7
8 129 g/kg and the herbs at 10g/kg thus ensuring the presence of adequate amounts of the
9
10 130 terpenes in the diets. All the birds were given the control diet from d 0-6 of the
11
12 131 experiment, and then either the control diet on its own or one of the 10 supplemented
13
14 132 diets from d 7-29. During mixing, each plant herb was incorporated directly on a
15
16 133 weight for weight basis into the dry mix of the diet. The essential oils were first
17
18 134 mixed into the vegetable oil component of the ration, and then the oil mixture was
19
20 135 added to the basal ration. No in-feed antimicrobials, anti-coccidial drugs or feed
21
22 136 enzymes were included.

Table 1 near here

137 **Analytical methodology**

138 Enumeration of microbial populations in the caecal and faecal contents of the
139 chickens was performed using various dehydrated media (Oxoid Ltd., UK), which
140 were prepared and sterilised according to the manufacturer's instructions before
141 being poured into sterile Petri dishes. The media used included MRS (DeMan,
142 Rogosa, Sharpe), Wilkins Chalgren anaerobe agar (with 10% defibrinated horse
143 blood), MacConkey agar No. 3 and Perfringens OPSP (Oleandomycin-polymixin-
144 sulphadiazine-perfringens). Caecal and faecal samples were serially diluted in
145 Maximum Recovery Diluent, and all plates were inoculated according to the method
146 of Miles and Misra, (1938) except those containing Perfringens OPSP agar. The
147 Perfringens OPSP agar plates were inoculated using 1ml bacterial suspensions by
148 means of the pour plate method. The MRS and WCA plates (48 h) and perfringens
149 OPSP plates (24 h) were incubated anaerobically at 37°C, while MacConkey plates
150 were incubated for 24 h aerobically at 37°C.

1
2
3 151 The dry matter (DM) and organic matter (OM) contents of each diet and of
4
5 152 the excreta samples collected from each cage were determined according to BS 5766,
6
7
8 153 Parts 1 and 8 (1983). Gross energy was determined by adiabatic bomb calorimetry
9
10 154 (Autobomb A. Gallenkamp and Co Ltd., England) in accordance with BS 1016, Part
11
12 155 105 (1992), using a benzoic acid standard (VWR Ltd., UK), with a known heat
13
14 156 capacity. Nitrogen contents were determined using the Dumas combustion method
15
16 157 (Leco FP-428/328, St. Louis, USA). The concentration of titanium dioxide in diet
17
18 158 and excreta samples were determined according to the method of Short *et al.*, (1996),
19
20 159 and sialic acid concentrations in excreta samples according to Jourdian *et al.*, (1971).
21
22 160 The condensed tannin (CT) contents in the herb samples were measured by the
23
24 161 butanol/HCl method, following Makkar, (2003).
25
26
27
28

29 162 **Statistical analysis**

30
31 163 This experiment was designed as a 5 X 2 factorial experiment, incorporating a non-
32
33 164 supplemented control, to give a total of 11 treatments. The statistical analysis was
34
35 165 carried out using a balanced randomised block ANOVA in General Linear Model
36
37 166 (GLM), using Fishers Least Significant difference (LSD) test to separate means
38
39 167 (Genstat, Release 7). A factorial analysis of the experiment was enabled when the
40
41 168 term Cont/(herb_oil*type) was entered into Genstat, allowing the separation of
42
43 169 effects due to plant species and due to the supplementation of herbs or essential oils.
44
45
46
47

48 170 **RESULTS**

49 171 **Diet compositional analysis**

50
51 172 The proximate analysis carried out on treatment diets showed that there was no
52
53 173 difference in the nutritional composition of the dietary treatments. The dietary DM
54
55 174 concentration ranged from 882-890 g/kg over all treatments and N content was
56
57 175 between 198-214 g/kg. Ash content ranged from 64-75 g/kg DM in all treatment
58
59
60

1
2
3 176 diets, and the GE of the treatment diets was within the range 18.06-18.63 MJ/kg DM.
4
5 177 The OM content in each treatment diet varied from 925-936 g/kg DM. The content
6
7 178 of extractable CT in the oregano herb supplement was 16 g/kg, while each of the
8
9 179 other herb supplements contained <1 g/kg CT. The terpene compositions in the
10
11 180 essential oils were determined by gas chromatography: marjoram oil contained (g/kg
12
13 181 of oil) carvacrol (560), terpinene-4-ol (70) and *trans*-cinnamaldehyde (60), oregano
14
15 182 oil contained γ -terpinene (170), 1,8-cineole (60) and an unidentified terpene, thyme
16
17 183 oil contained α -terpineol (297), thymol (100) and carvacrol (70), rosemary oil
18
19 184 contained primarily 1,8-cineole (470), camphor (180) and α -pinene (110), and the
20
21 185 yarrow oil contained β -pinene (180), linalol (170), sabinene (100) and α -terpineol
22
23 186 (60).
24
25
26
27
28

29 187 **Effect of herbs and essential oils on broiler growth performance**

30
31 188 In this experiment, 2 chicks died in the first week from different dietary treatments.
32
33 189 Generally, those birds given diets including thyme oil and yarrow herb had the
34
35 190 greatest overall body mass (BM), average gain, average feed consumption and FCE
36
37 191 in the study, compared to the other dietary treatments (Table 2). The average weight
38
39 192 of chicks at 7 d of age was 79.2 g (standard deviation, SD=0.982), which was low,
40
41 193 but a large proportion of these birds weighed less than 30g on arrival at d of hatch
42
43 194 and the diets were in mash form, which may have subsequently reduced overall
44
45 195 weights achieved in the trial. No oil-herb interactions were observed between
46
47 196 supplements at any stage of this experiment. In each case, the differences described
48
49 197 are for individual plant supplements in comparison with each other or the control.
50
51 198 Only the birds given diets with thyme oil had a greater BM than those fed on the
52
53 199 control diet (d 21, $P<0.01$; d 28, $P=0.001$). There was also a differential effect in
54
55 200 feeding essential oil and herb supplements from some plant species on BM. The
56
57
58
59
60

1
2
3 201 birds given yarrow herb in their diet had a greater BM than those receiving yarrow
4
5 202 oil (d 21, $P<0.01$; d 28, $P=0.001$), whereas those birds given thyme oil in their diet
6
7 203 had a greater BM than those with thyme herb supplements (d 21, $P<0.01$; d 28,
8
9 204 $P=0.001$). The birds fed on diets with thyme oil and yarrow herb also had the largest
10
11 205 overall weight gain (AWG) over the study period, although only thyme oil was
12
13 206 significantly greater than the control treatment (15-21 d, $P<0.01$; 7-28 d, $P\leq 0.001$).
14
15 207 Differences were again observed between oil and herb supplements from the same
16
17 208 plant, with a higher AWG in birds receiving dietary thyme oil rather than thyme herb
18
19 209 supplements (15-21 d, $P<0.01$; 22-28 d and 7-28 d, both $P\leq 0.001$) and in those birds
20
21 210 receiving yarrow herb rather than yarrow oil supplements (15-21 d, $P<0.01$; 22-28 d
22
23 211 and 7-28 d, both $P\leq 0.001$). Average feed consumption (AFC) was reduced for the
24
25 212 birds given dietary oregano herb, when compared with those receiving the control
26
27 213 treatment diet over the trial period ($P<0.05$). The birds fed on diets with thyme oil
28
29 214 consumed more feed than the control birds, but this was not significant. Inclusion of
30
31 215 rosemary oil in the diets reduced the AFC, when compared with those birds given the
32
33 216 control treatment (22-28 d, $P<0.01$; 7-28 d, $P<0.05$). The AFC was lower in the
34
35 217 birds given diets containing thyme herb when compared with those containing thyme
36
37 218 oil (15-21 d, $P\leq 0.05$; 22-28 d, $P=0.006$, not shown). However, the birds given
38
39 219 supplementary rosemary herb in the diet had higher AFC than those fed on diets with
40
41 220 rosemary oil (22-28 d, $P<0.01$; 7-28 d, $P<0.05$), and those fed on diets with yarrow
42
43 221 herb had a higher AFC than those given diets containing yarrow oil (22-28 d,
44
45 222 $P<0.01$) in the study. During the study period overall (7-28 d), rosemary oil had a
46
47 223 positive effect on chick FCE compared with rosemary herb, whilst yarrow herb also
48
49 224 had a positive effect compared with yarrow oil (7-28 d, $P\leq 0.05$).
50
51
52
53
54
55
56
57
58
59
60 225

Table 2 near here

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

226 **Effect of dietary inclusion of herbs on intestinal microbial status**

227 The bacteriological data required transformation before statistical analysis. There
228 were no effects of treatment on the log₁₀ population counts of the main intestinal
229 microflora species, when the birds were sampled at 28 d of age (Table 3). The
230 calculated lactic acid: coliform ratios of the original bacterial counts were
231 transformed prior to statistical analyses and were not affected by dietary herbal
232 supplementation; nor were the proportions of total lactic acid bacteria, total coliforms
233 or *Clostridium perfringens* in relation to total anaerobes (Table 4). Tables 3,4 near here

234 **Effect of herbs and essential oil supplementation on nutrient digestibility**

235 The apparent digestibility coefficients of DM (ADMD), OM (DOMD) and the
236 apparent metabolisability of nitrogen (AMN) and the apparent metabolisable energy
237 (AME) and also AME corrected to zero nitrogen retention (AME_N), contents were
238 determined. The digestible OM content and also metabolisable nitrogen content in
239 the treatment diets were calculated. The metabolisability of the dietary energy
240 (AME:GE) was also determined. Concentrations of sialic acid in broiler excreta
241 were used to calculate sialic acid excreted/kg food intake for the birds as a measure
242 of endogenous losses.

243 No treatment effects were observed with any plant supplement within the
244 diets in relation to the nutrient digestibility (Table 5). As a whole, the calculated
245 nutrient digestibility content and coefficients were rather low in these diets when
246 compared with commercial values. However, the low values for dry matter and
247 organic matter digestibility are in accordance with the low values for AME and
248 AME_N. One treatment replicate of the birds fed the diet with yarrow essential oil
249 was removed as a statistical outlier (>3.5 standard deviations from mean) in the
250 analysis of the coefficient of AMN and also the metabolisable N content in the diet.

1
2
3 251 This bird was measured with an abnormally low concentration of TiO₂ in the excreta
4
5 252 and an error was assumed. In the adjusted dataset, there was a tendency ($P=0.07$) for
6
7
8 253 the chicks to utilise nitrogen less effectively from the diet with rosemary herb when
9
10 254 compared to the others, which resulted in a lower dietary metabolisable N content.

11
12 255 Sialic acid was measured to assess the effects of treatment on endogenous
13
14 256 losses. Dietary treatment did not influence the excretion of sialic acid, whether this
15
16 257 was expressed as concentration in the excreta, per kg of diet intake or related to
17
18
19 258 metabolic liveweight.

Table 5 near here

22 259 DISCUSSION

23
24 260 These plant supplements are common dietary additives for humans, and were chosen
25
26 261 for their non-toxic chemical composition, relatively low cost and easy availability.
27
28 262 Phytochemical supplements are known to have considerable variability in their
29
30 263 chemical composition. With the exception of oregano, the composition of each
31
32 264 essential oil used in this experiment was consistent with those described in the
33
34 265 literature. Oregano and thyme oils are usually composed of the monoterpenes
35
36 266 thymol and carvacrol in varying proportions (Daferera *et al.*, 2000). Neither thymol
37
38 267 nor carvacrol were present in the oregano oil used in this study, which they may have
39
40 268 originated from senescent plant tissue since it had a similar composition (Russo *et*
41
42 269 *al.*, 1998). Terpene compositions in peppermint plants are known to change over
43
44 270 time (Burbott and Loomis, 1969), which may be important when selecting the best
45
46 271 plant supplements for use in poultry feeding trials. Although the mechanisms behind
47
48 272 terpene interactions are unknown, synergistic (Cox *et al.*, 2001) or antagonistic
49
50 273 interactions in a plant extract may affect its antimicrobial potential. The diet with
51
52 274 oregano herb also contained 0.16g of extractable CT/kg, which is very low, but the
53
54 275 reduction in AWG by 10-13% in birds fed this diet over the experimental period

1
2
3 276 compared with the control birds, suggests that there may have been bound tannin
4
5 277 which may have had some influence as well as other components of the herb
6
7
8 278 (Oduguwa *et al.*, 2007). This reduction in AWG is similar to that reported when
9
10 279 tannic acid and sorghum were fed in chick diets (Dale *et al.*, 1980). The affinities of
11
12 280 CT compounds for proteins and carbohydrates have been well characterised, and
13
14 281 those in this oregano herb may have bound to other dietary substrates, thereby
15
16 282 reducing nutrient availability (Bento *et al.*, 2005; Oduguwa *et al.*, 2007). The
17
18 283 measurement of chemical composition in a plant supplement prior to its use may
19
20 284 prevent reduced performance in broiler chicks.
21
22
23

24 285 **Effects on bird performance**

26 286 The different herbs and essential oils had variable effects on chick performance.
27
28 287 Both the oregano herb and oil treatments were among the poorest dietary treatments,
29
30 288 and were associated with reduced values for AFC and AWG in these chicks. Orego-
31
32 289 stim is a commercial supplement based on oregano, and has been shown to improve
33
34 290 AWG and FCR in pigs (Gill, 1999). However, no effect of Orego-Stim or oregano
35
36 291 has been reported on performance in growing broilers (Botsoglou *et al.*, 2002; Lewis
37
38 292 *et al.*, 2003), or turkeys between 12 and 16 weeks of age (Papageorgiou *et al.*, 2003).
39
40 293 Orego-Stim has reduced the incidence of coccidiosis in chickens (Gill, 1999), which
41
42 294 may indicate a protective effect of this supplement. In this study, the inclusion level
43
44 295 of essential oil was higher than in previous experiments (Botsoglou *et al.*, 2002;
45
46 296 Papageorgiou *et al.*, 2003), and was around 10 times that used in the diets fed to the
47
48 297 pigs. Although oregano oil is a potent antimicrobial, these studies indicate that
49
50 298 oregano in the diets of broiler chicks may not improve growth, and may even restrict
51
52 299 it if the oil quality is less than optimal. Broiler AFC and BM has been increased in
53
54 300 birds fed diets with Orego-stim without an effect on FCR, when fed in combination
55
56
57
58
59
60

1
2
3 301 with a vaccination programme against coccidiosis (Waldenstedt, 2004), but the
4
5 302 vaccination programme may have produced the bulk of the effect. The increased
6
7 303 rates of intestinal transit in chickens may not allow for the same performance
8
9 304 response as that observed in pigs, especially if microbial challenge is low.

10
11
12 305 In this experiment, the birds given diets with thyme oil performed best,
13
14 306 although thyme oil in the diet did not affect performance in male Wistar rats
15
16 307 (Youdim and Deans, 1999). Blends of essential oils have improved broiler
17
18 308 performance when given as dietary supplements (Suk *et al.*, 2003; Hernandez *et al.*,
19
20 309 2004). In other studies, only very limited information is available concerning the
21
22 310 composition of the blend. Blended supplements containing carvacrol, capsaicin and
23
24 311 cinnamaldehyde have improved both BM and FCR in broilers at 21 d of age, to a
25
26 312 greater extent than avilamycin (Jamroz *et al.*, 2003). Yarrow appears to be more
27
28 313 effective when fed in these study diets as a herb, which may be due to the precursors
29
30 314 of the sesquiterpene chamazulene, which is present in the herb but not the oil, or
31
32 315 another phytochemical component. Lewis *et al.* (2003) observed an improved
33
34 316 growth rate and AFC in broilers with yarrow herb inclusion in diets between 18 and
35
36 317 36 d of age. The dietary inclusion of yarrow at 30 g/kg improved AWG in broiler
37
38 318 chicks (Fritz *et al.*, 1993). In diets diluted by 10% in energy provision, yarrow herb
39
40 319 inclusion increased AFC and tended to increase AWG in broilers (Lewis *et al.*,
41
42 320 2004). Lewis *et al.* (2004) observed no positive effects of yarrow inclusion when the
43
44 321 diets were not diluted, and concluded that the benefits of yarrow inclusion may be
45
46 322 dependent on dietary quality. These experimental diets had a reduced compositional
47
48 323 quality, which may have optimised the effects of yarrow herb, but these results
49
50 324 suggest that both thyme oil and yarrow herb have potential as dietary supplements in
51
52 325 growing poultry.

1
2
3 326 The form of herbal supplementation may also be influential in determining
4
5 327 bioactivity. The present paper indicates that the terpene composition within essential
6
7
8 328 oil and herb extracts from the same plant, or the presence of additional secondary
9
10 329 plant compounds, may result in differential effects for some supplements on broiler
11
12 330 performance. Only marjoram supplements produced an equivalent effect on AWG
13
14
15 331 over the study, irrespective of whether the herb or the oil was included in the diet,
16
17 332 which suggests that the chemical composition of both in this marjoram may be
18
19 333 similar. The marjoram used in this study contained a high proportion of carvacrol.
20
21 334 Carvacrol has reduced both AFC and AWG when included in broiler diets at 200
22
23 335 mg/kg (Lee *et al.*, 2003a). Thymol and carvacrol have been reported to differ in their
24
25 336 effects on bird performance and lipid metabolism (Lee *et al.*, 2003a), which is
26
27 337 surprising, as the two terpenes are isomers with identical chemical formulae.
28
29 338 Complex interactions appear to exist between the terpenes in essential oils and herbs,
30
31 339 and may also occur between terpenes and the remainder of the dietary ration. As the
32
33 340 herbs used in this experiment were not composed of pure terpenes, the yarrow herb
34
35 341 should contain compounds that are either absent in the oil or are damaged during
36
37 342 extraction. It is possible that the reduced quality of this oil may have resulted in an
38
39 343 overly negative perception of yarrow oil as a supplement. These compounds were
40
41 344 purchased commercially, so any effects relating to the drying process are unknown.

42
43 345 Active terpenes in plants may be trapped within secretory gland structures, as
44
45 346 suggested by Dorman and Deans (2000), which may favour the antimicrobial activity
46
47 347 of essential oils rather than herbs. However, sage herb has shown a greater
48
49 348 inhibitory effect against bacteria than sage oil in a meat broth medium (Shelef *et al.*,
50
51 349 1984). In future, milling the herb fraction may release the active terpenes more
52
53 350 effectively into the gut where digestion and absorption are taking place. In this
54
55
56
57
58
59
60

1
2
3 351 study, all herbs were fed as chopped leaf material, apart from yarrow, which came as
4
5 352 complete flower heads and was then milled. The herbs were not milled as it was
6
7
8 353 considered that any volatile components would be lost through vapourisation. In an *in*
9
10 354 *vitro* study, extracted terpinene-4-ol has been reported to have a higher bioactivity
11
12 355 separately than as part of an essential oil (Cox *et al.*, 2001), suggesting that separate
13
14
15 356 terpenes may be the most bioactive when suitable compounds are identified.

17 357 **Effect of essential oils on digestibility**

18
19 358 In order to allow the effects of the herbs and oils to be more clearly observed, these
20
21 359 experimental diets were formulated to be marginal in their energy and protein
22
23 360 provision, which should at least partly explain the slow growth and poor digestibility
24
25 361 values in these birds. The wheat used in this experiment was of a lower quality than
26
27 362 expected, as the determined AME value in the excreta was about 3 MJ lower than the
28
29 363 formulated dietary ME content. A reduced protein digestibility when diets with low
30
31 364 AME were fed (Choct *et al.*, 1995), and an increased starch digestibility when AME
32
33 365 is increased has been reported in poultry (Svihus and Hetland, 2001). Arabinoxylan
34
35 366 and lectins can also be associated with poor quality wheat samples. The birds were
36
37 367 also fed on mash instead of pelleted diets; mash is less dense in energy and have a
38
39 368 greater bulk. This was done to allow easy incorporation of the plant supplements.
40
41 369 No dietary enzymes were included, as some may have an independent antimicrobial
42
43 370 effect (Fuglsang *et al.*, 1995), and the reduced vitamin E concentration will further
44
45 371 have reduced dietary feeding value. Thymol and carvacrol have a known antioxidant
46
47 372 activity *in vitro* (Deighton *et al.*, 1993). Previously, antioxidant effects have been
48
49 373 demonstrated in poultry given dietary oregano, but this has not been associated with
50
51 374 any effect on bird performance (Botsoglou *et al.*, 2002; Papageorgiou *et al.*, 2003).
52
53
54
55
56
57
58
59
60

1
2
3 375 Exogenous xylanases in wheat diets have improved broiler performance by
4
5
6 376 reducing arabinoxylan contents, improving AME and nutrient digestibility, reducing
7
8 377 variation between birds and increasing the rate of dietary passage by an associated
9
10 378 reduction in the digesta viscosity (Almirall and Esteve-Garcia, 1994; Choct *et al.*,
11
12 379 1995). A reduced intestinal viscosity is associated with a reduction in the amount
13
14
15 380 and water holding capacity of NSP (Choct *et al.*, 1995), and enzymes are most
16
17 381 efficacious in younger chicks (Almirall and Esteve-Garcia, 1994). A carbohydrase in
18
19 382 future experimental diets may be beneficial to ameliorate any anti-nutrient effects.

20
21
22 383 Herbs and spices have traditionally been used to stimulate the production of
23
24 384 endogenous secretions in the small intestinal mucosa, pancreas and liver, and thus
25
26 385 aid digestion. Plant spices, when fed together with organic acids, have stimulated
27
28 386 proteolysis, by lowering the pH to the optimal level for pepsin activity (Kamel,
29
30 387 2001). In rats, the dietary inclusion of spices has increased the endogenous secretion
31
32 388 of enzymes, bile acids and pancreatic juice (Platel and Srinivasan, 2000, 2001 and
33
34 389 2003), and reduced the transit time for digesta passage (Platel and Srinivasan, 2001).
35
36 390 Some spices were observed to reduce endogenous secretions, without an effect on
37
38 391 food consumption (Platel and Srinivasan, 2000). An increased amylase activity in
39
40 392 intestinal chyme was observed with dietary inclusion of Crina® Poultry, an essential
41
42 393 oil blend, at 21 d of age in broilers, but the effect decreased with increasing age (Lee
43
44 394 *et al.*, 2003b). In the present experiment, the herb and oil supplement had no effect
45
46 395 on sialic acid excretion suggesting that endogenous losses were not a major factor
47
48 396 associated with the dietary supplements. It is possible that terpene and phenolic
49
50 397 compounds are excreted as a conjugate with other components, which may increase
51
52 398 the endogenous losses from the bird and reduce the utilisation of nutrients (Cowieson
53
54 399 *et al.*, 2006; Mansoori and Acamovic, 2007; Oduguwa *et al.*, 2007).

1
2
3 400 **Effects of herbal supplementation on the intestinal microflora**
4
5

6 401 The results in the present study were disappointing, with no clear effects observed on
7
8 402 the intestinal microflora populations, which may have been due to an insufficient
9
10 403 degree of replication since only 3 of the 5 treatments could be sampled within the
11
12 404 time frame of 24 h. Thyme, marjoram and rosemary inclusion in these diets reduced
13
14 405 the numbers of caecal *Cl. perfringens* by >1 log, a trend which may potentially be
15
16 406 significant under a different experimental methodology. It is possible that the birds
17
18 407 should be fed on the test diets from the day of hatch, to influence the intestinal
19
20 408 bacterial composition more completely. A decreased exposure time to the air may
21
22 409 have influenced the bacterial population but unfortunately, the samples had to be
23
24 410 transported before analysis.
25
26
27
28

29 411 Several studies have reported effects on intestinal microflora when herbs and
30
31 412 essential oils have been included in broiler diets. The dietary supplementation of
32
33 413 XTRACT, an encapsulated product containing capsaicin, carvacrol and
34
35 414 cinnamaldehyde, reduced the numbers of *E. coli* and *Cl. perfringens* in broiler rectal
36
37 415 contents to the same extent as avilamycin (Jamroz *et al.*, 2003). *Cl. perfringens* has
38
39 416 been reduced in number when the blended essential oil supplement CRINA@poultry,
40
41 417 was fed in poultry diets (Losa and Kohler, 2001; Mitsch *et al.*, 2002). Selective
42
43 418 inhibitions in the growth of the hyper-ammonia-producers *Cl. sticklandii* and
44
45 419 *Peptostreptococcus anaerobius* were observed when a blend of dietary terpenes
46
47 420 containing thymol, eugenol, vanillin and limonene were fed to ruminants (McIntosh
48
49 421 *et al.*, 2003). The growth of *E. coli* and *C. perfringens* was reduced in broilers, when
50
51 422 blends of essential oils were fed in industry trials (Losa and Kohler, 2001; Tucker,
52
53 423 2002), while numbers of *Lactobacillus* spp. increased (Tucker, 2002). Thus,
54
55
56
57
58
59
60

1
2
3 424 essential oils may act differently compared with synthetic antimicrobials, which tend
4
5 425 to depress bacterial numbers across species.
6
7

8 426 In this study, the caecal coliform populations in the birds given any of the
9
10 427 herbal treatments compared to those fed the control diet were higher, which may
11
12 428 suggest that these dietary herbs may select more against Gram-positive rather than
13
14 429 Gram-negative bacteria. Conventional antimicrobials used in broiler production
15
16 430 have preferentially targeted Gram-positive bacteria, especially *C. perfringens*
17
18 431 (Engberg and Petersen, 2001). If this is true, conditions suitable for the presence of
19
20 432 *Campylobacter* spp. or *Salmonella* spp. may exist in birds fed on diets supplemented
21
22 433 with herbs. If terpenes bind to dietary components rather than disable bacteria, and
23
24 434 are then released as dietary components are absorbed, the pH in the distal tract may
25
26 435 be decreased, thus preventing the growth of pathogenic bacteria. Each herb contains
27
28 436 an extensive variety of phytochemicals, all with variable bioactivity, so the use of
29
30 437 herbal combinations in poultry diets may be most effective when suitable terpenes
31
32 438 have been identified. In broiler chicken production, the cage environment has been
33
34 439 shown to be best in reducing populations of both helminth (Permin *et al.*, 1999) and
35
36 440 red mite (Höglund *et al.*, 1995), whereas floor litter provides a substrate for
37
38 441 pathogenic bacterial growth (Pope and Cherry, 2000). It is concluded that responses
39
40 442 to phytochemicals may be greater in a more challenging environment, in agreement
41
42 443 with Lee *et al.*, (2003b).
43
44
45
46
47
48
49

50 444 This study has demonstrated that various herbs and oils have different effects
51
52 445 on broiler performance, which may be primarily related to differences in their
53
54 446 terpene composition. The inclusion of a single herb or its extracted essential oil in
55
56 447 diets may not always have a similar effect on broiler performance, therefore it is
57
58 448 necessary to measure the chemical quality in a plant extract to identify optimal
59
60

1
2
3 449 compositions of secondary plant compounds for future reference. Dietary quality
4
5 450 and environmental conditions are likely to be important in the assessment of
6
7 451 responses to terpene inclusion. In conclusion, herbs and essential oils may influence
8
9
10 452 both the performance and the production of endogenous secretions in broilers.
11

12 ACKNOWLEDGEMENTS

13
14
15 454 This study was funded by the Scottish Executive for Environment, Rural Affairs and
16
17 455 Development Department (SEERAD; NCR 643701). Special thanks are due to Ian
18
19 456 Nevison from Biostatistics and Information Services Scotland (BioSS), for his
20
21 457 assistance in statistical analysis.
22
23

24 REFERENCES

- 25
26
27 459 ACAMOVIC, T and BROOKER, J.D. (2005) Biochemistry of plant secondary
28
29 460 metabolites and their effects in animals. *Proceedings of the Nutrition Society*,
30
31 461 **64**; 403-412.
32
33 462 ALMIRALL, M. and ESTEVE-GARCIA, E. (1994) Rate of passage of barley diets
34
35 463 with chromium oxide: Influence of age and poultry strain and effect of β -
36
37 464 glucanase supplementation. *Poultry Science*, **73**:1433-1440.
38
39 465 BALOG, J.M. and MILLAR, R.I. (1989) Influence of the sense of taste on broiler
40
41 466 chick feed consumption. *Poultry Science*, **68**:1519-1526.
42
43 467 BENTOSGLOU, N.A., FLOROU-PANERI, P., CHRISTAKI, E., FLETOURIS, D.J.
44
45 468 tannin, pectin and polyethylene glycol on attachment of N-labelled
46
47 469 microorganisms to cellulose. *Animal Feed Science and Technology*, **122**: 29-
48
49 470 40.
50
51 471 BOTSOGLOU, N.A., FLOROU-PANERI, P., CHRISTAKI, E., FLETOURIS, D.J.
52
53 472 and SPAIS, A.B. (2002) Effect of dietary oregano essential oil on performance
54
55 473 of chickens and on iron-induced lipid oxidation of breast, thigh and abdominal
56
57 474 fat tissues. *British Poultry Science*, **43**:223-230.
58
59 475 BURBOTT, A.J. and, LOOMIS, W.D. (1969) Evidence of metabolic turnover of
60
476 monoterpenes in peppermint. *Plant Physiology*, **44**:173-179.
477
478 CANDAN, F., UNLU, M., TEPE, B., DAFERERA, D., POLISSIOU, M.,
SÖKMEN, A. and AKPULAT, A. (2003) Antioxidant and antimicrobial

- 1
2
3 479 activity of the essential oil and methanol extracts of *Achillea millefolium*
4 subsp. *millefolium* Afan. (Asteraceae). *Journal of Ethnopharmacology*, **87**:215-
5 480
6 220.
7
8 482 CHARAI, M., MOSADDAK, M. and FAID, M. (1996) Chemical composition and
9 antimicrobial activities of two aromatic plants: *Origanum majorana* L. and *O.*
10 483
11 *compactum* Benth. *Journal of Essential Oil Research*, **8**:657-664.
12 484
13 485 CHOCT, M., HUGHES, R.J., TRIMBLE, R.P., ANGKANAPORN, K. and
14 486
15 ANNISON, G. (1995) Non-starch polysaccharide-degrading enzymes increase
16 487
17 the performance of broiler chickens fed wheat of low apparent metabolizable
18 488
19 energy. *Journal of Nutrition*, **125**:485-492.
20 489
21 COWAN, M.M. (1999) Plant products as antimicrobial agents. *Clinical*
22 490
23 *Microbiology Reviews*, **12**:564-582.
24 491
25 COWIESON, A.J., ACAMOVIC, T and BEDFORD, M.R. (2006) Phytic acid and
26 492
27 phytase: implications for protein utilisation by poultry. *Poultry Science* **85**:
28 493
29 878-885.
30 494 COX, S.D., MANN, C.M. and MARKHAM, J.L. (2001) Interactions between
31 495
32 components of the essential oil of *Melaleuca alternifolia*. *Journal of Applied*
33 496
34 *Microbiology*, **91**:492-497.
35 497
36 DAFERERA, D.J., ZIOGAS, B.N. and POLISSIOU, M.G. (2000) GC-MS analysis
37 498
38 of essential oils from some Greek aromatic plants and their fungitoxicity on
39 499
40 *Penicillium digitatum*. *Journal of Agriculture and Food Chemistry*, **48**:2576-
41 500
42 2581.
43 501 DALE, N.M., WYATT, R.D. and FULLER, H.L. (1980) Additive toxicity of
44 502
45 aflatoxin and dietary tannins in broiler chicks. *Poultry Science*, **59**:2417-2420.
46 503
47 DEIGHTON, N., GLIDEWELL, S.M., DEANS, S.G. and GOODMAN, B.A. (1993)
48 504
49 Identification by EPR spectroscopy of carvacrol and thymol as the major
50 505
51 sources of free radicals in the oxidation of plant essential oils. *Journal of the*
52 506
53 *Science of Food and Agriculture*, **63**:221-225.
54 507
55 DORMAN, H.J.D. and DEANS, S.G. (2000) Antimicrobial agents from plants:
56 508
57 antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*,
58 509
59 **88**:308-316.
60 510
511 ENGBERG, R.M. and PETERSEN, J.S. (2001) Poultry production with and without
512 512
questionable feed additives and ingredients. *Proceedings of the 13th*
Symposium on Poultry Nutrition, Blankenberge, Belgium, pp. 118-124.

1
2
3 513

4
5 514 FRITZ, Z., SCHLEICHER, A. and KINAL, S. (1993) Effect of substituting milfoil,
6
7 515 St. John's wort and lovage for antibiotics on chicken performance and meat
8
9 516 quality. *Journal of Animal and Feed Sciences*, **2**:189-195.

10 517 FUGLSANG, C.C., JOHANSEN, C., CHRISTGAU, S. and ADLER-NISSEN, J.
11
12 518 (1995) Antimicrobial enzymes: Applications and future potential in the food
13
14 519 industry. *Trends Food Science and Technology*, **6**:390-395.

15 520 GILL, C. (1999) Herbs and plant extracts as growth enhancers. *Feed International*,
16
17 521 (April), pp. 20-23.

18
19 522

20
21 523 GUO, F.C., KWAKKEL, R.P., SOEDE, J., WILLIAMS, B.A. and VERSTEGEN,
22
23 524 M.W.A. (2004) Effect of a Chinese herb medicine formulation, as an
24
25 525 alternative for antibiotics on growth performance in broilers. *British Poultry*
26
27 526 *Science*, **45**:793-797.

28
29 527

30 528 HAMMER, K.A., CARSON, C.F. and RILEY, T.V. (1999) Antimicrobial activity of
31
32 529 essential oils and other plant extracts. *Journal of Applied Microbiology*,
33
34 530 **86**:985-990.

35 531 HÖGLUND, J., NORDENFORS, H. and UGGLA, A. (1995) Prevalence of the
36
37 532 poultry red mite, *Dermanyssus gallinae*, in different types of production
38
39 533 systems for egg layers in Sweden. *Poultry Science*, **74**:1793-1798.

40 534 HERNANDEZ, F., MADRID, J., GARCIA, V., ORENGO, J. and MEGIAS, M.D.
41
42 535 (2004) Influence of two plant extracts on broiler performance, digestibility and
43
44 536 digestive organ size. *Poultry Science*, **83**:169-174.

45
46 537

47
48 538 JAMROZ, D., ORDA, J., KAMEL, C., WILICZKIEWICZ, A., WERTELECKI, T.
49
50 539 and SKORUPINSKA, J. (2003) The influence of phytogetic extracts on
51
52 540 performance, nutrient digestibility, carcass characteristics, and gut microbial
53
54 541 status in broiler chickens. *Journal of Animal and Feed Sciences*, **12**:583-596.

54 542 JOURDIAN, G.W., DEAN, L. and ROSEMAN, S. (1971) The Sialic Acids XI. A
55
56 543 periodate-resorcinol method for the quantitative estimation of free sialic acids
57
58 544 and their glycosides. *Journal of Biological Chemistry*, **246**:430-435.

- 1
2
3 545 KAMEL, C. (2001) Tracing modes of action and the roles of plant extracts in non-
4
5 546 ruminants, in: Garnsworthy, P.C. and Wiseman, J. (Eds.) *Recent advances in*
6
7 547 *animal nutrition*, pp. 135-150. (Nottingham University Press, Nottingham).
- 8
9 548 LEE, K.W., EVERTS, H., KAPPERT, H.J., KEOM, K.-H. and BEYNEN, A.C.
10
11 549 (2003a) Dietary carvacrol lowers body weight gain but improves feed
12
13 550 conversion in female broiler chickens. *Journal of Applied Poultry Research*,
14
15 551 **12**:394-399.
- 16
17 552 LEE, K.W., EVERTS, H., KAPPERT, H.J., FREHNER, M., LOSA, R. and
18
19 553 BEYNEN, A.C. (2003b). Effects of dietary essential oil components on growth
20
21 554 performance, digestive enzymes and lipid metabolism in female broiler
22
23 555 chickens. *British Poultry Science*, **44**:450-457.
- 24
25 556 LEWIS, M.R., ROSE, S.P., MACKENZIE, A.M., SMITH, J. and ESKANAZI, S.
26
27 557 (2004) Dietary yarrow (*Achillea millefolium*) and the growth performance and
28
29 558 nutrient digestibility in broiler chickens. *Proceedings of the 22nd World*
30
31 559 *Poultry Congress, Istanbul*, pp487 (CD ROM).
- 32
33 560
- 34
35 561 LEWIS, M.R., ROSE, S.P., MACKENZIE, A.M. and TUCKER, L.A. (2003) Effects
36
37 562 of dietary inclusion of plant extracts on the growth performance of male broiler
38
39 563 chickens. *British Poultry Science*, **44**:S43-S44.
- 40
41 564 LOSA, R. and KÖHLER, B. (2001) Prevention of colonisation of *Clostridium*
42
43 565 *perfringens* in broiler intestine by essential oils. *Proceedings of the 13th*
44
45 566 *European Symposium of Poultry Nutrition, Blankenberge, Belgium*, pp. 133-
46
47 567 134.
- 48
49 568
- 50
51 569 MAKKAR, H.P.S. (2003). *Quantification of tannins in tree and shrub foliage*.
52
53 570 Kluwer Academic Publishers, Dordrecht, The Netherlands.
- 54
55 571
- 56
57 572 MANSOORI, B. and ACAMOVIC, T. (2007) Tannic acid increases the excretion of
58
59 573 endogenous methionine, histidine and lysine from broilers. *Animal Feed*
60
574 *Science and Technology* (in press)
- 575
576
577 576 MANSOORI, B. and ACAMOVIC, T. (1998) The influence of tannic acid on amino
acid digestibility in broilers. In: Garland, T. and Barr, A.C. (Eds.) *Toxic Plants*

- 1
2
3 578 *and other Natural Toxicants*, CAB International, Wallingford, UK., pp106-
4 110.
5 579
6 580
7
8 581 MCINTOSH, F.M., WILLIAMS, P., LOSA, R., WALLACE, R.J., BEEVER, D.A.
9
10 582 and NEWBOLD, C.J. (2003) Effects of essential oils on ruminal micro-
11 583 organisms and their protein metabolism. *Applied Environmental Microbiology*,
12 584 **69**:5011-5014.
13
14 585 MILES, A.A. and MISRA, S.S. (1938) Estimation of the bactericidal power of blood.
15 586 *Journal of Hygiene, Cambridge*, **38**:732-749.
16
17 587 MITSCH, P., KÖHLER, B., GABLER, C., LOSA, R. and ZITTERL-EGLESEER, K.
18 588 (2002) CRINA Poultry reduces the colonisation and proliferation of
19 589 *Clostridium perfringens* in the intestine and faeces of broiler chickens.
20 590 *Proceedings of the 11th European Poultry Conference, Bremen.*, pp. 113.
21 591 ODUGUWA, O.O., PIRGOZLIEV, V. and ACAMOVIC, T. (2007) Energy
22 592 metabolisability and digestibility of amino acids by broilers fed malted
23 593 sorghum sprouts supplemented with polyethylene glycol, charcoal, phytase and
24 594 xylanase. *British Poultry Science* **48**:55-63
25
26 595 PANTER, K.E., JAMES, L.F., WANG, S., GARDNER, D.R., GAFFIELD, W.,
27 596 MOLYNEUX, R.J., STEGELMEIER, B.L. and BUNCH, T.D. (2004)
28 597 Screening poisonous plant toxins for cytotoxicity using bovine embryos
29 598 produced by *in vitro* fertilization techniques, in: ACAMOVIC, T., STEWART,
30 599 C.S. and PENNYCOTT, T.W. (Eds). *Poisonous Plants and Related Toxins*, pp.
31 600 347-353 (CABI International, Wallingford, Oxon, UK).
32
33 601 PAPAGEORGIOU, G., BOTSOGLOU, N., GOVARIS, A., GIANNENAS, I.,
34 602 ILIADIS, S. and BOTSOGLOU, E. (2003) Effect of dietary oregano oil and
35 603 α -tocopherol acetate supplementation on iron-induced lipid oxidation of
36 604 turkey breast, thigh, liver and heart tissues. *Journal of Animal Physiology and*
37 605 *Animal Nutrition*, **87**:324-335.
38
39 606 PERMIN, A., BISGAARD, M., FRANDBSEN, F., PEARMAN, M., KOLD, J. and
40 607 NANSEN, P. (1999) Prevalence of gastrointestinal helminths in different
41 608 poultry production systems. *British Poultry Science*, **40**:439-443.
42
43 609 PLATEL, K. and SRINIVASAN, K. (2000) Influence of dietary spices and their
44 610 active principles on pancreatic digestive enzymes in albino rats. *Nahrung*,
45 611 **44**:42-46.

- 1
2
3 612 PLATEL, K. and SRINIVASAN, K. (2001) Studies on the influence of dietary
4 spices on food transit time in experimental rats. *Nutrition Research*, **21**:1309-
5 1314.
6
7
8 615 PLATEL, K. and SRINIVASAN, K. (2003) Stimulatory influence of select spices on
9 bile secretion in rats. *Nutrition Research*, **20**:1493-1503.
10
11 617 POPE, M.J. and CHERRY, T.E. (2000) Evaluation of the presence of pathogens on
12 broilers raised on poultry litter treatment®-treated litter. *Poultry Science*,
13 **79**:1351-1355.
14
15
16 620 RUSSO, M., GALLETTI, G.C., BOCCHINI, P. and CARNACINI, A. (1998)
17 Essential oil chemical composition of wild populations of Italian oregano spice
18 (*Origanum vulgare* ssp. *hirtum* (Link) Ietswaart): A preliminary evaluation of
19 their use in chemotaxonomy by cluster analysis. 1. Inflorescences. *Journal of*
20 *Agriculture and Food Chemistry*, **46**:3741-3746.
21
22
23 625 SHELEF, L.A., JYOTHI, E.K. and BULGARELLI, M.A. (1984) Growth of
24 enteropathogenic and spoilage bacteria in sage-containing broths and foods.
25 *Journal of Food Science*, **49**:737-740.
26
27
28 628 SHORT, F.J., GORTON, P., WISEMAN, J. and BOORMAN, K.N. (1996)
29 Determination of titanium dioxide added as an inert marker in chicken
30 digestibility studies. *Animal Feed Science and Technology*, **59**:215-221.
31
32
33 631 SUK, J.C., LIM, H.S. and PAIK, I.K. (2003) Effects of blended essential oil
34 (CRINA®) supplementation on the performance, nutrient digestibility, small
35 intestinal microflora and fatty acid composition of meat in broiler chickens.
36 *Journal of Animal Science and Technology (Korea)*, **45(5)**:777-786.
37
38
39 635 SVIHUS, B. and HETLAND, H. (2001) Ileal starch digestibility in growing broiler
40 chickens fed on a wheat-based diet is improved by mash feeding, dilution with
41 cellulose or whole wheat inclusion. *British Poultry Science*, **42**:633-637.
42
43
44 638 TUCKER, L.A. (2002) Maintaining poultry performance in antibiotic-free diets by
45 supplementation with commercial botanical feed ingredients. *Proceedings of*
46 *the 7th WPSA Asian Pacific Federation Conference, Gold Coast, Australia*, pp.
47 227-230
48
49
50
51 642
52
53 643 WALDENSTEDT, L. (2004) Effect of vaccination against coccidiosis in
54 combination with an antibacterial oregano (*Origanum vulgare*) compound in
55
56
57
58
59
60

- 1
2
3 645 organic broiler production. *Acta Agriculturae Scandinavica Section A – Animal*
4 *Science*, **53**:101-109.
5 646
6
7 647 WINK, M. (2004) Evolution of toxins and anti-nutritional factors in plants with
8 special emphasis on Leguminosae, in: ACAMOVIC, T., STEWART, C.S. and
9 648 PENNYCOTT, T.W. (Eds). *Poisonous Plants and Related Toxins*, pp. 1-25. (CABI
10 649 International, Wallingford, Oxon, UK).
11 650
12
13 651 YOUDIM, K.A. and DEANS, S.G. (1999) Dietary supplementation of thyme
14 652 (*Thymus vulgaris* L.) essential oil during the lifetime of the rat: its effects on
15 the antioxidant status in liver, kidney and heart tissues. *Mechanisms of Ageing*
16 653 *and Development*, **109**:163-175.
17 654
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

655 **Table 1.** *Diet formulation and calculated chemical composition of the basal ration*656 *(as fed)*

657

<i>Feed Ingredient</i>	<i>Amount in diet (g/kg)</i>	<i>Calculated chemical composition</i>	<i>(g/kg)</i>
Wheat	671.0	ME (MJ/kg)	13.4
Soya bean meal	251.0	Crude Protein	201.6
Soya oil	35.0	Ether Extract/Fat	52.8
Monocalcium phosphate	15.0	Crude Fibre	27.8
Limestone	15.0	Calcium	9.6
Sodium chloride	3.0	Phosphorus	7.5
Lysine	2.0	Lysine	12.7
Methionine	3.0	Methionine + Cysteine	8.9
Vit/ Min premix ¹	5.0		

658 Diets were also mixed with TiO₂ (5 g/kg) as a dietary marker and with the plant supplements.659 ¹Supplied per kg diet: vitamin A 4128 µg (retinyl acetate), vitamin D3 125 µg cholecalciferol,

660 vitamin E (as α-tocopherol) 50 mg, vitamin K 3 mg, folic acid 1mg, nicotinic acid 50 mg, vitamin B1

661 (Thiamine) 2 mg, vitamin B2 (Riboflavin) 7 mg, vitamin B6 (Pyridoxine) 5 mg, vitamin B12 15 µg,

662 biotin 200 µg, calcium pantothenate 15 mg, iodine 1mg, molybdenum 0.5 mg, selenium 200 µg,

663 cobalt 0.5 mg, copper 10 mg, iron 80 mg, manganese 100 mg, zinc 80 mg, limestone 4.18 g.

664

665

666

667

668

669

670

671

672

673

674

675

676

677 **Table 2.** *Effect of herbs or their essential oils on growth performance, when*
 678 *included in broiler chick diets from 7-28 d of age*

679

<i>Chick growth performance data from 7-28 d</i>				
<i>Treatment</i>	<i>BM at 28d</i>	<i>Average gain (g/bird/d)</i>	<i>Average feed cons (g/bird/d)</i>	<i>FCE (gain/cons)</i>
Control	727 ^{bcde}	30.9 ^{bcdef}	53.1 ^{ab}	0.588 ^{abc}
Marjoram herb	754 ^{abcd}	32.1 ^{abcd}	53.5 ^{ab}	0.602 ^{ab}
Oregano herb	645 ^e	26.9 ^{ef}	45.1 ^c	0.604 ^{ab}
Rosemary herb	734 ^{abcde}	31.2 ^{abcdef}	53.9 ^{ab}	0.580 ^{bc}
Yarrow herb	789 ^{ab}	33.8 ^{ab}	55.5 ^{ab}	0.610 ^{ab}
Thyme herb	691 ^{cde}	29.2 ^{cdef}	52.0 ^{abc}	0.564 ^{bc}
Marjoram oil	764 ^{abc}	32.6 ^{abc}	55.2 ^{ab}	0.594 ^{ab}
Oregano oil	680 ^{cde}	28.6 ^{cdef}	47.9 ^{bc}	0.594 ^{abc}
Rosemary oil	664 ^{de}	27.9 ^{def}	44.6 ^c	0.634 ^a
Yarrow oil	641 ^e	26.8 ^f	49.0 ^{bc}	0.544 ^c
Thyme oil	821 ^a	35.3 ^a	58.7 ^a	0.600 ^{ab}
sed	45.14	2.2	3.9	0.0277
Plant species (herb+oil)	P=0.021	P=0.021	P=0.015	NS
Herb v Oil	P=0.001	P=0.001	P<0.05	P<0.05

680 Means were tested at $P<0.05$, but the largest treatment differences were significant to the level shown.

681 Means within a column without a common superscript differ at $P<0.05$ (NS: $P>0.05$).

682 Values are means for 5 replicates of each treatment.

683

684

685

686

687

688

689

690

691

692

693

694

695 **Table 3.** *Bacterial population counts in samples of faecal and caecal material taken*
 696 *from broilers aged 28 d, when birds were fed diets with herbs from 7-28 d*
 697

<i>Microfloral populations in samples from the intestinal contents of broilers (log₁₀ c.f.u./g)</i>								
<i>Treatment</i>	<i>Coliforms</i>		<i>Lactic Acid Bacteria</i>		<i>Total Anaerobes</i>		<i>C. perfringens</i>	
	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>
Control	8.91	6.94	8.91	8.33	9.42	11.24	3.54	3.94
Marjoram	8.79	7.25	8.67	8.68	9.06	10.98	4.19	3.61
Oregano	8.83	7.50	8.21	8.27	9.13	11.36	3.77	3.57
Rosemary	8.62	7.24	8.51	8.15	9.21	11.12	2.41	2.80
Yarrow	8.60	8.40	8.76	8.63	8.62	10.99	2.57	2.00
Thyme	8.32	7.82	8.74	8.05	9.21	11.15	3.68	2.27
sed	0.70	0.78	0.49	0.69	0.35	0.33	0.76	0.75
	NS	NS	NS	NS	NS	NS	NS	NS

698 Data comparisons are done within each column and were statistically compared after transformation.

699 NS ($P>0.05$).

700 Values are means in colony forming units per g sample of the logarithms of 3 treatment replicates.

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717 **Table 4.** *Proportions of bacterial species in relation to each other in broiler chicks*718 *aged 28 d, when the birds were fed diets with plant herbs from 7-28 d of age*

719

<i>Proportions of intestinal microflora species in broilers</i>								
<i>Treatment</i>	<i>LAB: Coliforms</i>		<i>Coliforms: Anaerobes</i>		<i>LAB: Anaerobes</i>		<i>C. perfringens: Anaerobes</i>	
	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>	<i>Faecal</i>	<i>Caecal</i>
Control	1.000	1.217	0.947	0.618	0.947	0.742	0.373	0.350
Marjoram	0.987	1.206	0.970	0.661	0.957	0.794	0.462	0.332
Oregano	0.933	1.117	0.967	0.661	0.901	0.727	0.412	0.314
Rosemary	0.992	1.124	0.936	0.651	0.925	0.734	0.263	0.252
Thyme	1.077	1.061	0.901	0.700	0.949	0.722	0.403	0.203
Yarrow	1.020	1.028	1.001	0.765	1.020	0.785	0.296	0.185
s.e.d.	0.095	0.150	0.059	0.068	0.044	0.069	0.081	0.068
	NS	NS	NS	NS	NS	NS	NS	NS

720 Significance comparisons are done within each column and statistically analysed after transformation.

721 NS ($P>0.05$).

722 LAB = Lactic acid bacteria.

723 Values are proportions of 3 replicate sample means for each count measured in \log_{10} c.f.u./g.

724

725 **Table 5.** *Effect of dietary herb or essential oil supplementation from 7-28 d of age on nutrient digestibility in broiler chicks aged 25 d, measured as*
 726 *apparent metabolisable energy (AME) and the coefficient of energy metabolisability (AME:GE), with correction to zero nitrogen retention (AMEn*
 727 *and AMEn:GE), and also the coefficients of the digestibility of dry and organic matter (ADMD and DOMD) and nitrogen metabolisability (AMN) in*
 728 *the ration, presented with the digestible OM and metabolisable N content in each treatment diet*

<i>Energy and its metabolisability, dry and organic matter digestibility and the metabolisability of nitrogen in the treatment rations</i>									
<i>Treatment</i>	<i>AME</i>	<i>AMEn</i>	<i>AME:GE</i>	<i>AMEn:GE</i>	<i>ADMD</i>	<i>DOMD</i>	<i>Digestible OM content</i>	<i>AMN</i>	<i>Metabolisable N content</i>
	<i>(MJ/kg DM)</i>		<i>(MJ/MJ)</i>	<i>(MJ/MJ)</i>	<i>(g/g)</i>	<i>(g/g)</i>	<i>(g/kg DM)</i>	<i>(g/g)</i>	<i>(g/kg DM)</i>
Control	10.46	9.96	0.567	0.579	0.534	0.574	531	0.405	13.61
Marjoram herb	10.39	9.89	0.568	0.583	0.539	0.578	536	0.399	13.64
Oregano herb	10.59	10.14	0.572	0.592	0.538	0.580	540	0.371	12.45
Rosemary herb	10.11	9.76	0.550	0.574	0.515	0.562	526	0.307	9.72
Yarrow herb	10.78	10.26	0.582	0.608	0.544	0.589	547	0.421	14.29
Thyme herb	10.34	9.94	0.558	0.588	0.516	0.558	520	0.331	10.89
Marjoram oil	10.05	9.56	0.556	0.570	0.526	0.568	527	0.382	13.25
Oregano oil	10.65	10.21	0.576	0.607	0.536	0.573	534	0.359	11.96
Rosemary oil	10.59	10.12	0.576	0.607	0.539	0.576	533	0.377	12.89
Yarrow oil	10.31	9.96	0.553	0.569	0.523	0.561	525	0.365	11.49
Thyme oil	10.63	10.15	0.572	0.593	0.539	0.579	540	0.386	13.12
sed 1	0.553	0.498	0.030	0.036	0.029	0.029	27.09	0.045	1.556
sed 2								0.048	1.651
Plant spp. (herb+oil)	NS	NS	NS	NS	NS	NS	NS	NS	NS
Herb v Oil	NS	NS	NS	NS	NS	NS	NS	NS	P=0.07

729 Values are the means of 5 replicates for each dietary treatment.

730 Data comparisons are done within each column. NS ($P>0.05$)

731 For most treatment comparisons, sed 1 is used, but sed 2 should be used for nitrogen metabolisability, when comparing yarrow oil with all other treatments.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

732

For Peer Review Only