



**HAL**  
open science

# EFFECT OF FEEDING SILAGES OR CARROTS AS SUPPLEMENTS TO LAYING HENS ON PRODUCTION PERFORMANCE, NUTRIENT DIGESTIBILITY, GUT STRUCTURE, GUT MICROFLORA AND FEATHER PECKING BEHAVIOUR

Sanna Steinfeldt, Joergen Kjaer, Ricarda Margarete Engberg

► **To cite this version:**

Sanna Steinfeldt, Joergen Kjaer, Ricarda Margarete Engberg. EFFECT OF FEEDING SILAGES OR CARROTS AS SUPPLEMENTS TO LAYING HENS ON PRODUCTION PERFORMANCE, NUTRIENT DIGESTIBILITY, GUT STRUCTURE, GUT MICROFLORA AND FEATHER PECKING BEHAVIOUR. *British Poultry Science*, 2007, 48 (04), pp.454-468. 10.1080/00071660701473857. hal-00545319


**HAL Id: hal-00545319**

**<https://hal.science/hal-00545319>**

Submitted on 10 Dec 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**EFFECT OF FEEDING SILAGES OR CARROTS AS SUPPLEMENTS TO LAYING HENS ON PRODUCTION PERFORMANCE, NUTRIENT DIGESTIBILITY, GUT STRUCTURE, GUT MICROFLORA AND FEATHER PECKING BEHAVIOUR**

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2006-164.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	25-Jan-2007
Complete List of Authors:	Steenfeldt, Sanna; University of Aarhus, Faculty of Agricultural Sciences, Animal Health, Welfare and Nutrition Kjaer, Joergen; Federal Agricultural Research Centre, Institute for Animal Welfare and Animal Husbandry Engberg, Ricarda; University of Aarhus, Faculty of Agricultural Sciences, Animal Health, Welfare and Nutrition
Keywords:	Feather pecking, Microbiology, Laying hens, Nutrition



1  
2  
3 1 **Effect of feeding silages or carrots as supplements to laying hens on production**  
4  
5 2 **performance, nutrient digestibility, gut structure, gut microflora and feather pecking**  
6  
7 3 **behaviour**  
8  
9

10  
11 4  
12  
13 5 S. STEENFELDT, J. B. KJAER<sup>1</sup> AND R. M. ENGBERG  
14

15 6  
16  
17 7 *University of Aarhus, Dept. of Animal Health, Welfare and Nutrition, Research Centre*  
18 *Foulum, Tjele, Denmark and <sup>1</sup>Federal Agricultural Research Centre, Institute for Animal*  
19 *Welfare and Animal Husbandry, Celle, Germany*  
20  
21

22  
23  
24  
25 10  
26  
27 11 Short title: FORAGING MATERIAL FOR LAYING HENS  
28

29  
30  
31  
32 13 Correspondence to Sanna Steinfeldt, University of Aarhus, Faculty of Agricultural Sciences,  
33  
34 14 Dept. of Animal Health, Welfare and Nutrition, Research Centre Foulum, P.O. Box 50,  
35  
36 15 DK-8830 Tjele, Denmark.  
37

38  
39 16 Tel: +45 8999 1147  
40

41 17 Fax.: +45 8999 1378  
42

43 18 [Sanna.Steenfeldt@agrsci.dk](mailto:Sanna.Steenfeldt@agrsci.dk)  
44  
45

46 19  
47

48 20  
49

50  
51 21

52  
53 22 **Abstract.** 1. An experiment was carried out to examine the suitability of using maize  
54  
55 23 silage, barley-peas silage and carrots as foraging materials for egg-laying hens. Production  
56  
57  
58  
59  
60

1  
2  
3 24 performance, nutrient digestibility, gastrointestinal characteristics, including the  
4  
5 25 composition of the intestinal microflora as well as feather pecking behaviour were the  
6  
7  
8 26 outcome variables.  
9  
10 27 2. The protein content of the foraging material (g/kg DM) was on average 69 g in carrots  
11  
12 28 94 g in maize silage and 125 g in barley-pea silage. The starch content was highest in the  
13  
14 29 maize silage (312 g/kg DM), and the content of NSP varied from 196-390 g/kg, being  
15  
16 30 lowest in carrots. Sugars were just traceable in the silages, whereas carrots contained on  
17  
18 31 average 496 g/kg DM.  
19  
20 32 3. Egg production was highest in hens fed either carrots or maize-silage, whereas hens fed  
21  
22 33 barley-pea silage produced less (219 vs. 208). Although the consumption of foraging  
23  
24 34 material was high (33%, 35% and 48% of the total feed intake on "as fed" basis for maize  
25  
26 35 silage, barley-pea silage and carrots, respectively) only a minor effect on AME<sub>n</sub> and  
27  
28 36 apparent digestibility was seen. At 53 weeks of age, hens fed maize silage had AME<sub>n</sub> and  
29  
30 37 apparent digestibility values close to the control group (12.61 and 12.82, respectively),  
31  
32 38 whereas access to barley-pea silage and carrots resulted in slightly lower values (12.36 and  
33  
34 39 12.42, respectively). Mortality was reduced dramatically in the three groups given  
35  
36 40 supplements (0.5-2.5%) compared to the control group (15.2%).  
37  
38 41 4. Hens receiving silage had greater relative gizzard weights than the control or carrot-fed  
39  
40 42 groups. At 53 weeks of age, the gizzard-content pH of hens receiving silage was about 0.7-  
41  
42 43 0.9 units lower than that of the control or carrot-fed hens. Hens fed both types of silage had  
43  
44 44 higher concentrations of lactic acid (15.6 vs. 3.2 µmol/g ) and acetic acid (3.6 vs. 6.1  
45  
46 45 µmol/g ) in the gizzard contents than the other two groups. The dietary supplements had a  
47  
48 46 minor effect on the composition of the intestinal microflora of the hens.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 47 5. Access to all three types of supplements decreased damaging pecking in general (to  
5  
6 48 feathers as well as skin/cloaca), reduced severe feather pecking behaviour and improved  
7  
8 49 the quality of the plumage at 54 weeks of age.

9  
10 50 6. In conclusion, access to different types of foraging material such as silages and carrots  
11  
12 51 improved animal welfare.

## 13 14 15 INTRODUCTION

16  
17 53 Non cage housing systems for egg production (single or multiple tier systems with or  
18  
19 54 without access to range area) have expanded markedly in Denmark within the last 15  
20  
21 55 years, due mainly to changes in consumer preference. In Denmark, the market share of  
22  
23 56 organic eggs was 14.1% of all eggs in retail in 2005 (Danish Poultry Council, 2006), a  
24  
25 57 trend which reflects an increased interest and concern related to our environment, animal  
26  
27 58 welfare and product quality. Unfortunately, it has been demonstrated in many cases that  
28  
29 59 egg production in alternative housing systems is facing welfare problems in terms of  
30  
31 60 feather pecking and cannibalism. Outbreaks of feather pecking and cannibalism can be  
32  
33 61 caused by a number of factors such as genetic background of the high productive strains  
34  
35 62 used (Kjaer *et al.*, 2001), dietary deficiencies (Ambrosen and Petersen, 1997) or  
36  
37 63 environmental conditions (Savory, 1995; Kjaer and Sørensen, 2002). The Danish law  
38  
39 64 requires that hens in organic egg production have access to some high-fibre feed  
40  
41 65 ingredients such as silages, straw, grass, sugar beets or carrots. In general, feeding with  
42  
43 66 forage seems to have a positive effect on behaviour, and available foraging material in the  
44  
45 67 housing system motivates the hens to spend more time foraging and less feather pecking  
46  
47 68 (Blokhuys, 1986; Nørgaard-Nielsen *et al.*, 1993; Aerni *et al.*, 2000).

48  
49 69 Feeding poultry with whole cereals, oat hulls or wood shavings having a  
50  
51 70 large particle size and more structure than pelleted feed stimulates the upper part of the  
52  
53 71 intestinal system and increases gizzard weight and activity (Gabriel *et al.*, 2003; Hetland

1  
2  
3  
4 72 and Svihus, 2001; Hetland *et al.*, 2003, Engberg *et al.*, 2004). Furthermore, an increased  
5  
6 73 content of structural components, such as whole grains, is claimed to have a beneficial  
7  
8 74 effect on gastrointestinal health, probably due to increased hydrochloric acid secretion and  
9  
10 75 a fall in gizzard pH, which inhibits acid-sensitive potentially-pathogenic bacteria like  
11  
12 76 *Salmonella* from entering the lower intestinal tract (Engberg *et al.*, 2004; Bjerrum *et al.*,  
13  
14 77 2005). Other coarsely-structured feeds may have a comparable effect on the  
15  
16 78 gastrointestinal tract, as shown for maize silage by Idi *et al.* (2005).

17  
18  
19  
20 79 The nutritive value of silages to poultry may be minimal, because it has a  
21  
22 80 high content of dietary fibre (non-starch polysaccharides ~ NSP + lignin) which, due to the  
23  
24 81 lack of endogenous fibre-degrading enzymes in the intestine (Scott *et al.*, 1982), is poorly  
25  
26 82 digested by poultry (Longstaff and McNab, 1989; Carré *et al.*, 1990; Steinfeldt *et al.*,  
27  
28 83 1998; Lazaro *et al.*, 2003). A high intake of soluble NSP can depress performance and the  
29  
30 84 digestibility of other nutrients in broilers (Choct and Annison, 1992a; Steinfeldt, 2001), in  
31  
32 85 part because of increased microbial activity (Choct *et al.*, 1996; Smits *et al.*, 1998;  
33  
34 86 Langhout *et al.*, 1999). In contrast to soluble fibre, the insoluble fibre fraction is not much  
35  
36 87 fermented by the microflora (Choct *et al.*, 1996).

37  
38  
39  
40  
41 88 Diets with moderate amounts of insoluble fibre can improve starch  
42  
43 89 digestibility in broilers or laying hens (Svihus and Hetland, 2001; Hetland *et al.*, 2002,  
44  
45 90 2003) and reduce pecking at other birds and mortality from cannibalism in layers (Hartini  
46  
47 91 *et al.*, 2003). If the use of foraging material is to be recommended in practice in alternative  
48  
49 92 production systems for laying hens, it is important to know how it may influence different  
50  
51 93 aspects of production. The purpose of the present study was to examine the effect of using  
52  
53 94 different foraging materials as feed supplements to laying hens on egg production, nutrient  
54  
55 95 digestibility, gastrointestinal characteristics including the composition of the intestinal  
56  
57  
58  
59  
60

1  
2  
3 96 microflora, as well as aspects of feather pecking behaviour and plumage condition. In the  
4  
5 97 present investigation maize silage, barley-pea silage and carrots were used as forages.  
6  
7  
8 98  
9

10 99 MATERIAL AND METHODS

11  
12  
13 100 **Diet, foraging material and treatments**

14  
15 101 The basal layer diet was formulated to contain the nutrients recommended (NRC, 1994) for  
16  
17 102 commercial layer diets but was based on Danish-grown raw ingredients, with the exception  
18  
19 103 of soyabean meal (Table 1.)  
20  
21

22 104 **Table 1**

23  
24  
25 105 The control group received the layer diet without any supplement, whereas  
26  
27 106 hens in the other three treatments were fed supplemental foraging material, provided as  
28  
29 107 maize-silage, or barley-pea-silage, or carrots. The supplements were given unprocessed  
30  
31 108 (carrots as cleaned, uncooked roots), and the daily amounts were slightly increased during  
32  
33 109 the experimental period in order to achieve an approximate *ad libitum* intake during the  
34  
35 110 light hours of the day. The layer diet was provided as 3 mm pellets and fed *ad libitum* over  
36  
37 111 the entire experimental period. The treatments will be referred to as LD (layer diet without  
38  
39 112 supplements), LD+MS (layer diet + maize silage), LD+BS (layer diet + barley-pea silage)  
40  
41 113 and LD+C (layer diet + carrots). The experiment was performed over a period of 238 d,  
42  
43 114 from the age of 20 to 54 weeks.  
44  
45  
46  
47

48 115 **Birds, housing and experimental design**

49  
50 116 A total of 800 ISA Brown pullets were obtained from a commercial company at 16 weeks  
51  
52 117 of age and placed in 16 identical indoor floor pens with 50 birds/pen, each with a floor area  
53  
54 118 of 8.9 m<sup>2</sup>. The hens were not beak trimmed because this process would probably hamper  
55  
56 119 the hens ability to fully exploit the foraging material. In addition, beak trimming is not  
57  
58 120 allowed in organic egg production in Denmark (Danish Ministry of Justice, 1998:  
59  
60

1  
2  
3 121 Bekendtgørelse nr. 210 af 6/4-1998). The lighting programme was 12L:12D at 16 weeks.  
4  
5 122 Day length was gradually increased by 1 h per week to 16L:8D at 19 weeks. The 16L:8D  
6  
7 123 programme was planned to continue until termination of the experiment at 54 weeks of  
8  
9 124 age. However, a 23L:1D programme was introduced in error when the hens were 27 weeks  
10  
11 125 of age. The mistake was discovered when the hens were 39 weeks; thereafter the light  
12  
13 126 programme was corrected gradually by reducing day length by 1 h per week until a  
14  
15 127 16L:8D programme was reached at 46 weeks and continued for the rest of the experiment.  
16  
17 128 The number of hens was reduced to 43 hens per pen after 5 weeks, when the hens were 23  
18  
19 129 weeks old. The 4 experimental treatments were allocated at random to the 16 floor pens, to  
20  
21 130 provide 4 replicates. The bedding material was wood shavings. The basal diet was given *ad*  
22  
23 131 *libitum* in one round feeder (34 cm diameter) and water was supplied by nipple drinkers.  
24  
25 132 Each pen was equipped with 9 single nests in a battery, each 28 cm wide. Adjacent pens  
26  
27 133 were separated by thin wooden walls 160 cm high and with wire mesh above (total 200  
28  
29 134 cm), which allowed auditory contact but prevented visual contact between hens from  
30  
31 135 separate pens and treatments. Supplementation with maize silage or barley-pea silage or  
32  
33 136 carrots started at 20 weeks of age, with fresh material fed each morning in a wooden box  
34  
35 137 (height: 40 cm, length: 50 cm, width: 40 cm). Eggs were collected and recorded from nests  
36  
37 138 every day, and the collection of one day per week was weighed. Feed consumption was  
38  
39 139 recorded every second week. Egg production and feed consumption were calculated on a  
40  
41 140 hen-day basis (taking into account the actual number of hens housed per floor pen at the  
42  
43 141 time of recording). Mortality was noted daily, and body weight was recorded at 25, 39  
44  
45 142 and 54 weeks of age. The experiment complied with the guidelines of the Danish Ministry  
46  
47 143 of Justice with respect to animal experimentation and care of birds under study.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

#### 144 **Digestibility experiments and calculations**



1  
2  
3 145 At the age of 23 and 53 weeks, respectively, digestibility experiments were performed in  
4  
5 146 battery cages with raised floors, each of the three batteries contained 12 cages (50x50x50  
6  
7 147 cm), each with a feeding trough outside and two water cups inside. Two hens were taken at  
8  
9 148 random from each pen and placed in one cage, representing one replicate. The 4 treatments  
10  
11 149 were assigned at random, each treatment being represented in each battery and on each tier.  
12  
13 150 Supplements were given fresh in small portions, once in the morning and once in the  
14  
15 151 afternoon and after the adaptation period of 7 d, the excreta were collected quantitatively  
16  
17 152 on three consecutive days. Pelleted feed and supplements were weighed separately.  
18  
19 153 Excreta were collected three times per day and stored in closed containers at 20°C,  
20  
21 154 immediately after collection to prevent microbial degradation. In addition, two hens per  
22  
23 155 replicate were fed with the silages or carrots alone without access to layer diet to give an  
24  
25 156 indication of the DC of organic matter and AME of the supplements.  
26  
27  
28  
29  
30  
31

32 157 The content of polysaccharide residues was calculated as anhydrosugars, and apparent  
33  
34 158 digestibility coefficients were calculated according to the analysed content of nutrients (g/kg  
35  
36 159 DM) in feed (layer diet plus supplements, analysed separately) and in excreta, taking into  
37  
38 160 account the amount of feed eaten (g) and excreta voided (g) on a dry matter basis. The AME<sub>n</sub>  
39  
40 161 content was calculated as the difference between gross energy content in feed eaten (layer diet  
41  
42 162 and supplement, separately) and the gross energy in excreta, and corrected to zero N-retention  
43  
44 163 using a value of 34 kJ/g N retained (Hill and Anderson, 1958). N-retention was determined as  
45  
46 164 (N in feed – N in excreta)/g feed intake.  
47  
48  
49

#### 50 165 **Chemical analyses**

51  
52 166 The dry matter (DM) content of the layer diet, foraging material and excreta was determined  
53  
54 167 by drying at 105°C for 8 h. Protein (N x 6.25) was determined by the Kjeldahl method 984.13  
55  
56 168 (AOAC, 1990a) using a Kjell-Foss 16200 autoanalyser and energy by a LECO AC 300  
57  
58 169 automated calorimeter system 789-500 (LECO, St. Joseph, Michigan, USA). Ash was  
59  
60

1  
2  
3 170 analysed according to method 923.03 (AOAC, 1990b), and fat (hydrochloric acid-fat) was  
4  
5 171 extracted with diethyl ether after acid-hydrolysis (Stoldt, 1952). Amino acids were analysed  
6  
7 172 as described by Mason *et al.* (1980). The sugars (glucose, fructose and sucrose), and the  
8  
9 173 oligosaccharides (raffinose, stachyose and versbascose) were extracted with 50% (v/v)  
10  
11 174 ethanol at 60°C and quantified by gas-liquid chromatography (GLC) using the method of  
12  
13 175 Bach Knudsen and Li (1991). Starch was analysed by the enzymatic-colorimetric method of  
14  
15 176 Bach Knudsen (1997). Total NSP and their constituent sugars were determined as alditol  
16  
17 177 acetates by gas-liquid chromatography for neutral sugars and by the colorimetric method for  
18  
19 178 uronic acids using a modification of the Uppsala procedure (Theander *et al.*, 1994) as  
20  
21 179 described by Back Knudsen (1997). Cellulose was determined as the difference in glucose  
22  
23 180 content of NSP when the swelling step with 12 M sulphuric acid was included and omitted,  
24  
25 181 respectively, and the content of cellulose, non-cellulosic (NCP) and soluble NSP was  
26  
27 182 calculated as described by Bach Knudsen (1997). Klason lignin was measured gravimetrically  
28  
29 183 as the residue obtained of the treatment with 12 M sulphuric acid (Theander *et al.*, 1994). All  
30  
31 184 analyses were performed in duplicate.

### 185 **Gastrointestinal characteristics and intestinal microflora**

186 At 23 and 53 weeks of age, 5 hens from each pen were killed by cervical dislocation and  
187 the contents of the gizzard, jejunum, ileum, caeca and rectum were quantitatively collected  
188 and pooled by segment before further analyses. Jejunum and ileum were defined as the  
189 intestinal segments cranial and caudal to the Meckels diverticulum. The weight of the  
190 contents in gizzard, jejunum, ileum and caeca as well the weight of the empty  
191 gastrointestinal segments were measured. The pH-value of the contents in all segments was  
192 measured with a combined glass/reference electrode (GK 2401C, Radiometer, Denmark),  
193 and the dry matter of intestinal contents was analysed after freeze drying of the samples. In  
194 gastrointestinal contents, coliform bacteria, lactose-negative enterobacteria, lactic acid

1  
2  
3 195 bacteria and enterococci were enumerated as described by Engberg *et al.* (2004).  
4

5 196 Concentrations of short chain fatty acids and lactic acid were analysed (Jensen *et al.*,  
6  
7 197 1995).  
8  
9

### 10 11 198 **Behavioural and clinical recordings** 12

13  
14 199 Recordings of allo-pecking (pecking of one bird by another) were made at 24, 38 and 53  
15  
16 200 weeks of age. Every pen was observed for 20 min at each age. All pecks at other chickens  
17  
18 201 were recorded. Each peck was counted, and recorded in groups (events) called bouts. A  
19  
20 202 bout was defined as pecks in a continuous series directed to the same chicken and body  
21  
22 203 part (plumage, beak or feet) and of the same type (see below). The inter-bout interval was  
23  
24 204 10 s. Pecks were classified: 1) severe feather pecking, 2) gentle feather pecking, 3)  
25  
26 205 aggressive pecking, 4) beak pecking, 5) object pecking and 6) pecking at dust particles on  
27  
28 206 dustbathing birds. The two last categories were rarely seen and discarded from further  
29  
30 207 analyses. The distinction between non-aggressive and aggressive pecks was in most cases  
31  
32 208 quite clear, aggressive pecks being vigorous, directed towards the head region and forcing  
33  
34 209 the receiver to react (escape or fight) (Hoffmeyer, 1969; Wennrich, 1975). At 25, 39 and  
35  
36 210 54 weeks of age a random sample of 15 birds per pen were individually scored for the  
37  
38 211 condition of plumage and skin, as well as for health status of feet and comb. Body weight  
39  
40 212 was also recorded. Plumage and skin condition were scored individually (Tauson *et al.*,  
41  
42 213 2005), assigning scores of 1, 2, 3 and 4 points for each character. Higher scores means  
43  
44 214 better health. Plumage condition was scored for each of 5 body parts (neck, breast, back,  
45  
46 215 wings and tail). The sum of these 5 scores ('total score', 5 to 20 points) was also used for  
47  
48 216 analyses. Less than 1% of birds had wounds (skin score of body part less than 4) and  
49  
50 217 therefore these data were discarded from further analysis.  
51  
52  
53  
54  
55  
56  
57

### 58 218 **Statistical analyses** 59 60

1  
2  
3 219 A randomised complete block design was used, a single pen representing the experimental  
4  
5 220 unit (replicate). The results are presented as means and standard error of means (SEM)  
6  
7 221 calculated by standard procedures. Analysis of variance by the general linear model  
8  
9 222 (GLM) procedure (SAS Institute Inc., 1990) was used to determine the significance of  
10  
11 223 treatment and block effects on the dependent variables (production, apparent digestibility,  
12  
13 224 AME<sub>n</sub>, intestinal values, feather pecking behaviour and plumage score at 54 weeks of  
14  
15 225 age). Normality of data was tested by a univariate procedure, and the means of the  
16  
17 226 variables were tested for variance homogeneity by Bartlett's test. An outlier test based on  
18  
19 227 externally Studentised residuals was performed. No significant outliers were found in the  
20  
21 228 data, and no effect of block. The LS-Means were calculated and differences regarded as  
22  
23 229 significant at  $P < 0.05$ . With respect to production, apparent digestibility and AME<sub>n</sub>,  
24  
25 230 differences were classified by the Ryan-Einot-Gabriel-Welsch (REGW) multiple range test  
26  
27 231 (SAS Institute Inc., 1990). The results obtained from measurements in the intestinal  
28  
29 232 content of gizzard, duodenum and jejunum, ileum, caeca and rectum (pH, dry matter and  
30  
31 233 bacterial counts) were analysed separately for each segment. In cases in which the overall  
32  
33 234 effect was significant ( $P < 0.05$ ) means were compared pairwise (pdiff). Data on plumage  
34  
35 235 condition at 25 and 39 weeks of age were tested using the Wilcoxon signed rank test (non-  
36  
37 236 normally distributed data).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 238 RESULTS

### 239 **Chemical analysis of basal diet and foraging material**

240 The silages and carrots were chemically analysed at the beginning of the experiment at 23  
241 weeks of age (I) and at the end at 54 weeks of age (II), which represented the periods  
242 where the digestibility experiments were performed (Table 2).

#### 243 **Table 2**

1  
2  
3 244 The analyses were performed at these times to examine any possible changes in chemical  
4  
5 245 composition of the foraging material over time. The silages and carrots used were from the  
6  
7  
8 246 same harvest year (2000). The DM content of the silages was on average 323 g/kg DM  
9  
10 247 (maize I and II) and 221 g/kg DM (barley-pea I and II), respectively, whereas carrots had a  
11  
12 248 DM content as low as 84 and 125 g/kg DM (I and II). Protein content varied from 65 to  
13  
14  
15 249 127 g/kg DM, the lowest values measured in carrots. Fat content was generally low,  
16  
17 250 ranging between 12-32 g/kg DM. Starch content averaged 312 g/kg DM in maize silage  
18  
19 251 and 126 g/kg DM in barley-pea silage, but was not detectable in carrots. Overall, the  
20  
21 252 results of the nutrient analyses in samples from the two periods agreed closely.  
22  
23

24 253 Only traces of sugars were detected in the silages. In carrots, high  
25  
26 254 concentrations of glucose, fructose and sucrose were measured, constituting 458 g/kg DM  
27  
28 255 in sample I and 533 g/kg DM in sample II. The highest content of total NSP was found in  
29  
30 256 the silages, containing average values of 333 g/kg DM in maize-silage and 376 g/kg DM in  
31  
32 257 barley-pea silage. Cellulose and arabinoxylans (AX), the sum of arabinose and xylose  
33  
34 258 residues, were the dominant NSP residues, constituting approximately 88% and 77% of the  
35  
36 259 total NSP in maize- and barley-pea silage respectively. In barley-pea silage the content of  
37  
38 260 uronic acid averaged 57 g/kg DM. Soluble NSP as a percentage of total NSP averaged 6%  
39  
40 261 in the maize-silage and 12% in the barley-pea silage. The lignin content was high in the  
41  
42 262 silages, with dietary fibre ranging between 403-483g/kg DM. In carrots the dominant NSP  
43  
44 263 residues were uronic acids and glucose monosaccharides, cellulose content averaged 63  
45  
46 264 g/kg DM, total NSP content was 196 and 203 g/kg DM, and in contrast to the silages,  
47  
48 265 soluble NSP was a very high proportion of total NSP, constituting more than 50%. The  
49  
50 266 total dietary fibre content averaged 232g/kg DM. The two silages were similar with respect  
51  
52 267 to the dietary fibre fraction, but differed with regard to protein and starch. The carrots  
53  
54 268 contained less protein than the silages.  
55  
56  
57  
58  
59  
60

1  
2  
3 269 The protein content of the layer diet was close to that calculated (Table 1).  
4  
5 270 The starch content was 365 g/kg DM and fat was 76 g/kg DM. The ME of the layer diet  
6  
7  
8 271 was 11.79 MJ/kg DM, calculated on the basis of the chemical analyses and close to the ME  
9  
10 272 given in Table 1.

### 11 273 **Production**

12  
13  
14 274 Average egg weight was not influenced by the use of silages or carrots (Table 3). The  
15  
16  
17 275 number of eggs and total egg mass (kg) per hen was highest in the group given carrots,  
18  
19  
20 276 whereas hens with access to barley-pea silage (LD+BS) produced fewer eggs per hen than  
21  
22 277 the other groups given foraging material ( $P<0.03$ ).

### 23 278 **Table 3**

24  
25  
26  
27 279 Rate of lay (%) was highest for hens fed carrots or maize-silage when compared to the  
28  
29 280 control group, whereas hens fed barley-pea-silage produced less ( $P<0.04$ ). The  
30  
31 281 unintended change in the light programme at 27 weeks of age did not influence  
32  
33 282 production during the remaining part of the experimental period. The consumption  
34  
35 283 of the layer feed was reduced ( $P<0.004$ ) by supplementation with carrots and barley-pea  
36  
37 284 silage as compared to the LD group. The intake of maize silage, barley-pea silage and  
38  
39 285 carrots was 33%, 35% and 48 %, respectively, as a percentage of total feed intake on "as  
40  
41 286 fed" basis. On a dry matter (DM) basis, the figures were 15%, 11% and 8%. Due to the  
42  
43 287 lower DM content of the foraging material the total feed consumption in kg per kg egg  
44  
45 288 produced was higher in the groups given the silages and carrots compared to the LD group  
46  
47 289 on "as fed" basis ( $P<0.0001$ ). Litter samples were taken at the end of the experiment and  
48  
49 290 the DM content was 78% (LD), 76% (LD+MS), 71% (LD+BS) and 68% (LD+C), where  
50  
51 291 the DM content for the last two groups was lower than for the control ( $P<0.05$ ).  
52  
53 292 Supplemental foraging material reduced mortality dramatically ( $P<0.02$ ), (Table 3).  
54  
55  
56 293 Approximately 50% of the mortality recorded in the LD group was cannibalism directed  
57  
58  
59  
60

1  
2  
3 294 against the cloacal region. Body weight was higher in the LD group and in hens fed LD+C  
4  
5 295 at 25 and 39 weeks and at 54 weeks the hens fed LD+C were heavier than all other hens.  
6  
7

8 296 **Digestibility and AME<sub>n</sub>**  
9

10 297 The coefficients of total tract apparent digestibility (DC) of different dietary constituents  
11  
12 298 and the apparent metabolisable energy, nitrogen corrected (AME<sub>n</sub>) are given in Table 4.  
13  
14

15 299 **Table 4**  
16

17 300 Three weeks after the hens had been introduced to the supplements (23 weeks of age),  
18  
19 301 there was no significant difference between treatments with respect to the DC of organic  
20  
21 302 matter (average 0.749) and fat (average 0.853). The hens digested starch very efficiently in  
22  
23 303 all groups with DC averaging 0.997, however, the DC of starch was lower for hens fed  
24  
25 304 LD+BS compared to the other groups ( $P<0.01$ ). The DC of total NSP was low, ranging  
26  
27 305 from 0.131 (LD+C) to 0.309 (LD), whereas hens fed with carrots had a significantly lower  
28  
29 306 NSP digestibility as compared to hens fed LD and LD+MS. The N-retention varied from  
30  
31 307 0.412 (LD+C) to 0.521 (LD+MS), ( $P<0.01$ ). The highest AME<sub>n</sub> was seen in hens fed the  
32  
33 308 LD diet and LD+MS, both groups being better than the LD+C ( $P<0.01$ ), reflecting the  
34  
35 309 generally lower DC obtained in hens fed carrots as supplement at this stage of the  
36  
37 310 experiment.  
38  
39  
40  
41  
42

43 311 At 53 weeks of age, the DC of organic matter was lowest for the group  
44  
45 312 receiving LD+BS. The DC of fat was significantly lower when hens were fed LD+C as  
46  
47 313 compared to LD + MS (0.804 vs 0.861). No significant difference was observed between  
48  
49 314 groups with respect to DC of starch and total NSP. The overall digestibility of total NSP  
50  
51 315 was low, however, the different residues of NSP were digested to a varying degree (data  
52  
53 316 not shown). The major NSP residues arabinose, xylose and galactose appeared to be  
54  
55 317 digested to some extent in all groups (average DC arabinose: 0.320, xylose: 0.202 and  
56  
57 318 galactose: 0.248). Uronic acids, being an important NSP-residue in carrots were poorly  
58  
59  
60



1  
2  
3 319 digested in hens fed the LD+C diet with a DC close to zero. In general, all groups digested  
4  
5 320 the soluble part of NSP to a higher extent compared with the insoluble part, being on  
6  
7 321 average 0.334 (S-NSP) and 0.219 (I-NSP) at 23 weeks of age, and 0.392 (S-NSP) and  
8  
9 322 0.208 (I-NSP) at 53 weeks of age (Table 4). The N-retention was found to be in the range  
10  
11 323 of 0.363 (LD+C) to 0.419 (both group LD and LD+BS) with no significant difference. The  
12  
13 324 AME<sub>n</sub> varied from 12.1 to 12.8 MJ/kg DM and no significant difference between groups  
14  
15 325 was observed. During the 30 weeks between the two digestibility experiments, the daily  
16  
17 326 intake of the three supplements increased from 31 (MS), 29 (BS) and 45 (C) g/day to 74,  
18  
19 327 50 and 120 g/day, respectively. The results indicate that high supplemental intake of  
20  
21 328 silages or carrots had only minor effects on the apparent digestibility of the dietary  
22  
23 329 constituents and AME<sub>n</sub>. The DM content in excreta (Table 4) was lowest in hens fed  
24  
25 330 LD+C. With respect to the digestibility of the supplements fed without the layer diet (data  
26  
27 331 not shown), on average 55 % of organic matter in the foraging material was digested. The  
28  
29 332 AME<sub>n</sub> obtained reached an average value of 11 MJ/kg DM. The results indicate that the  
30  
31 333 supplements to some extent contribute with nutrient and energy to the hens.  
32  
33  
34  
35  
36  
37  
38  
39

#### 334 **Gastrointestinal characteristics and intestinal microflora**

40  
41 335 Birds receiving supplements of maize silage or barley pea silage had higher relative  
42  
43 336 gizzard weights as compared to the control and birds fed carrots (Table 5).  
44

#### 337 **Table 5**

45  
46  
47  
48 338 This difference was evident after three weeks on the experimental diets ( $P < 0.05$ ) and was  
49  
50 339 even more pronounced at 53 weeks of the experiment ( $P < 0.001$ ). No significant  
51  
52 340 differences between the dietary treatments were observed at any of the sampling times with  
53  
54 341 respect to the relative weights of the small intestinal segments and the caeca.  
55  
56 342 The relative weight of gizzard contents (Table 5) was likewise higher in birds receiving  
57  
58 343 silage as compared to the other two dietary groups, ( $P < 0.01$  and  $< 0.001$  at 23 and 53  
59  
60



1  
2  
3 344 weeks of age, respectively). There was also a difference between groups regarding the  
4  
5 345 relative weight of contents in the upper small intestine (duodenum/jejunum). At both  
6  
7 346 sampling times, birds fed with LD+MS had the highest, and birds fed solely with LD had  
8  
9 the lowest amounts of small intestinal contents ( $P < 0.05$  and  $< 0.01$  at 23 and 53 weeks of  
10  
11 347 age, respectively). Following intake of barley-pea silage, the relative weight of the contents  
12  
13 348 in the upper small intestine increased over time to reach values similar to maize silage at  
14  
15 349 53 week of age. A similar picture was observed considering the relative weight of ileal  
16  
17 350 contents (Table 5). At 23 weeks, the relative weight of caecal contents was similar in all  
18  
19 351 dietary groups, whereas at 53 weeks, the amount of caecal digesta tended to be lower ( $P =$   
20  
21 352  $0.07$ ) in hens receiving supplements of barley pea silage as compared to the other groups.  
22  
23 353 At both sampling times a difference between treatments regarding the percentage of dry  
24  
25 354 matter in the gizzard contents ( $P < 0.01$ ) was observed (Table 5). The dry matter content  
26  
27 355 was lowest in hens supplemented with carrots, and increased slightly when fed with barley  
28  
29 356 pea silage. The highest DM contents in gizzard contents was observed in birds receiving  
30  
31 357 maize silage and in birds fed solely LD (Table 5).  
32  
33 358 The pH in the contents of the gizzard (Table 6) was not different at 23 weeks of age,  
34  
35 359 whereas at 53 weeks of age a decrease of the pH about 0.7-0.9 units was found in birds fed  
36  
37 360 with both types of silages as compared to birds supplemented with carrots or birds  
38  
39 361 receiving no foraging material ( $P < 0.05$ ).  
40  
41 362  
42  
43  
44  
45  
46  
47  
48  
49

#### **Table 6**

50  
51 364 In the upper part of the small intestine, the pH was lower in birds receiving  
52  
53 365 maize silage and carrots as compared to the other groups ( $P < 0.01$ ). At 23 weeks, the pH  
54  
55 366 of the caecal content was not influenced significantly by the dietary treatment, however, at  
56  
57 367 53 weeks, the caecal pH value was about 0.4 units higher in birds fed the LD as compared  
58  
59 368 to the birds supplemented with the different foraging material ( $P < 0.05$ ).  
60

1  
2  
3 369 Counts of intestinal bacteria were conducted on two occasions (23 weeks and  
4  
5 370 53 weeks of age) and showed that differences between the dietary groups with respect to  
6  
7  
8 371 bacterial composition became more distinct with the progress of the experiment. Because  
9  
10 372 of the large amount of data, only the results obtained at the last sampling are presented  
11  
12 373 (Table 6). No difference between the diets was observed with respect to the number of  
13  
14 374 anaerobic bacteria in the different gastrointestinal segments. Coliform bacteria and lactose  
15  
16 375 negative enterobacteria in the small intestine were more numerous in hens fed with pellets  
17  
18 376 alone as compared to birds supplemented with forages, but these differences were not  
19  
20  
21 377 statistically significant.

22  
23  
24 378 Lactic acid bacteria in the contents of the small intestine tended to be fewer  
25  
26 379 in the group fed with barley pea silage compared to the other groups. The numbers of  
27  
28 380 enterococci in the gizzard and small intestine of hens receiving only LD were in the range  
29  
30 381 of 0.6 (gizzard) up to 2 log units (small intestine) higher than in birds supplemented with  
31  
32 382 forage ( $P < 0.05$ ).

33  
34  
35 383 The concentration of lactic acid (Table 7) was highest in gizzard contents of  
36  
37 384 birds fed both types of silages as compared to hens fed LD+C and LD ( $P < 0.01$ ).

38  
39  
40  
41 385 **Table 7**

42  
43 386 In the upper small intestine the highest lactic acid concentrations were found in  
44  
45 387 small intestinal contents of hens LD+MS followed by LD+BS, LD and LD+C ( $P <$   
46  
47 388 0.02). Birds fed with both types of silages had higher concentrations of acetic acid  
48  
49 389 in the gizzard contents than the other two groups ( $P < 0.05$ ). The amounts of acetic  
50  
51 390 acid found in the contents of the upper small intestine were very low in all groups  
52  
53 391 and did not exceed 1.5  $\mu\text{mol/g}$  contents. In ileal contents, acetic acid concentration  
54  
55 392 was lower in birds receiving supplements of carrots than in the other dietary  
56  
57 393 groups. In caecal contents, the concentrations of both acetic and propionic acid  
58  
59  
60

1  
2  
3 394 generally tended to be higher in birds fed with maize silage than in the other  
4  
5 395 dietary groups.

6  
7  
8 396 **Feather pecking and plumage condition**

9  
10 397 No effects of treatments were found at 24 and 38 weeks of age. At 53 weeks of age the  
11  
12 398 incidence of feather pecking bouts was significantly higher in the control compared to the  
13  
14 399 groups receiving foraging material (Table 8).

15  
16  
17  
18 400 **Table 8**

19  
20 401 The number of feather pecks was higher in the control compared to hens fed carrots and  
21  
22 402 with hens fed barley-pea silage and maize silage intermediate. These differences were  
23  
24 403 caused by more severe feather pecking in the control than in forage treatments. Gentle  
25  
26 404 feather pecking bouts were higher in control and barley than in the others, whereas gentle  
27  
28 405 feather pecking pecks did not differ significantly amongst treatments. Treatments did not  
29  
30 406 affect aggressive nor beak pecking. Plumage condition deteriorated over time and birds in  
31  
32 407 the control treatment had in general the worst quality, even though this was only  
33  
34 408 statistically significant for the combined score at 54 weeks of age (Table 8). As regards  
35  
36 409 quality of neck, the barley-pea group had a significantly better score at 54 weeks than  
37  
38 410 control hens. Skin quality was in general very good in all treatments and ages (data not  
39  
40 411 shown).

41  
42  
43  
44  
45  
46  
47  
48 413 **DISCUSSION**

49  
50 414 The results show beneficial effects of supplemental feeding of silage and carrots. The hens  
51  
52 415 given access to silages or carrots showed great interest in the supplements after few days of  
53  
54 416 introduction, and the intake of layer diet was reduced in the groups given carrots or barley-  
55  
56 417 pea silage compared to the control. The daily intake of both types of silage was close to 60  
57  
58 418 g per hen, and the intake of carrots exceeded 100 g per hen, illustrating that the supplement

1  
2  
3 419 constituted a large proportion of the total feed intake on “as feed” basis. The daily intake  
4  
5 420 of essential nutrients necessary for optimal egg-production was sufficient, in spite of a  
6  
7  
8 421 reduced intake of layer diet. Hvidsten and Herstad (1972) reported that silage of barley  
9  
10 422 grain or barley meal could be used to supplement a layer diet with good results on egg  
11  
12 423 production and feed utilisation. Tkalcic *et al.* (2000) fed ensiled maize grains mixed with a  
13  
14 424 diet based on soyabean meal, meat meal, wheat meal and limestone in order to study the  
15  
16 425 effect on egg-production. They concluded that groups fed diets with 60% of the dry matter  
17  
18 426 intake from ensiled maize silage grain had better laying performance than the control  
19  
20 427 group fed a complete commercial diet, indicating that inclusion of ensiled maize grains  
21  
22 428 instead of dry maize grains had a positive effect on production, in line with the present  
23  
24 429 experiment.

25  
26  
27  
28  
29 430 The daily protein intake from the layer diet was 22.2 g (LD), 20.3 g  
30  
31 431 (LD+MS), 18.3 g (LD+BS) and 19.5 g (LD+C) per hen. Protein from the supplements  
32  
33 432 contributed further with 1.8 g (LD+MS), 1.6 g (LD+BS) and 0.8 g (LD+C) daily per hen.  
34  
35 433 Based on the calculated methionine content in the layer diet (Table 1), the daily methionine  
36  
37 434 intake per day per hen from the layer diet ranged from 572 mg (LD) to 472 mg (LD+BS)  
38  
39 435 emphasising that methionine intake was sufficient for egg production, as the recommended  
40  
41 436 requirement is 300 mg/hen /day (NRC, 1994). As earlier reported by Hammershøj and  
42  
43 437 Steinfeldt (2005), the content of methionine in maize-silage and carrots is very low,  
44  
45 438 analysed to be 1.30 and 0.84 g/kg DM, respectively, suggesting that the contribution of  
46  
47 439 foraging material to the total dietary essential amino acid concentration is negligible. A  
48  
49 440 major concern in modern animal husbandry, both with regard to commercial and organic  
50  
51 441 productions, is the excretion of nitrogen and phosphorus to the environment. The N-  
52  
53 442 retention was less efficient when hens were fed with carrots as supplements compared to  
54  
55 443 the other groups. Since no difference in N-retention was found between hens fed LD,  
56  
57  
58  
59  
60

1  
2  
3 444 LD+MS and LD+BS at any ages, we conclude that the high intake of silages did not  
4  
5 445 increase excretion of nitrogen to the environment.  
6  
7

8 446 Dietary fibre (NSP+ lignin) in feed is a complex group of components  
9  
10 447 differing widely in chemical composition, physical properties and physiological activity  
11  
12 448 (Theander *et al.*, 1989; Bach Knudsen, 1997). The content of dietary fibre was high in the  
13  
14 449 foraging material used in the present experiment, especially in the two silages, where the  
15  
16 450 insoluble fibre constituted the main part (~ 90% of NSP). In carrots, the soluble fraction  
17  
18 451 makes up the bulk of the total NSP. It is known that high amounts of soluble NSP in diets  
19  
20 452 can increase intestinal viscosity, often having negative effects in broiler chickens (Bedford  
21  
22 453 and Classen, 1992; Choct and Annison, 1992b; Smits *et al.*, 1998; Steinfeldt, 2001).  
23  
24 454 However, adult birds seem to be less sensitive to increased intestinal viscosity created by  
25  
26 455 soluble NSP in cereal-based diets (Salih *et al.*, 1991; Lázaro *et al.*, 2003). The results from  
27  
28 456 the present digestibility experiment indicate that the hens are able to digest a part of the  
29  
30 457 total NSP from diets and supplements, in agreement with studies performed with adult  
31  
32 458 hens or cockerels (Longstaff and McNab, 1989; Carré *et al.*, 1990; Steinfeldt *et al.*, 1995;  
33  
34 459 Lázaro *et al.*, 2003). Furthermore, in the present study the soluble part of NSP was  
35  
36 460 digested to a higher extent than the insoluble fraction, in agreement with Carré *et al.*  
37  
38 461 (1990). It is believed that the main part of the soluble NSP from the supplements are  
39  
40 462 fermented during the passage through the intestinal system, especially in the caeca, and  
41  
42 463 may contribute with energy through production of short chain fatty acids. In an experiment  
43  
44 464 with broiler chickens Jørgensen *et al.* (1996), concluded that NSP fermentation contributed  
45  
46 465 3-4% of the metabolisable energy intake. Longstaff and McNab (1989) fed adult cockerels  
47  
48 466 with carrots (freeze-dried and tube-fed) and overall average digestibility of soluble uronic  
49  
50 467 acids to be 49.9%, whereas the corresponding figure for insoluble uronic acids was minus  
51  
52 468 111%. It was concluded that polysaccharide digestibility is very difficult to quantify, due  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 469 to differences in polysaccharide solubility, when different fibre sources pass through the  
4  
5  
6 470 intestinal system.

7  
8 471 The large amount of supplements eaten by the hens in the present experiment  
9  
10 472 had only minor effects on AME<sub>n</sub> and total tract apparent digestibility of the measured  
11  
12 473 nutrients. (Table 4). In some recent studies the inclusion of different insoluble fibre  
13  
14 474 sources (oat hulls, wood shavings, cellulose) in poultry diets had positive effects on starch  
15  
16 475 digestibility in both broilers and laying hens (Hetland and Svihus, 2001; Svihus and  
17  
18 476 Hetland, 2001; Hetland *et al.*, 2003).

19  
20  
21  
22 477 Hens fed with carrots had a higher final body weight than the other groups,  
23  
24 478 suggesting that large amounts of easily-fermented components as sugars and soluble NSP  
25  
26 479 contribute some energy to the hens, which confirm the results of Lázaro *et al.* (2003). In  
27  
28 480 the present study, hens fed with both types of silage developed larger gizzards than hens  
29  
30 481 receiving carrots or no supplementation, due to increased mechanical stimulation by the  
31  
32 482 increased amounts of dietary fibre retained in the gizzard and to a coarser feed structure  
33  
34 483 (Engberg *et al.*, 2002, 2004; Hetland *et al.*, 2003; Idi *et al.*, 2005). In addition, the pH of  
35  
36 484 the gizzard contents was lower in silage-fed birds, suggesting an increased secretion of  
37  
38 485 hydrochloric acid as a response to the coarse structure of the silage (Engberg, *et al.*, 2004;  
39  
40 486 Idi *et al.*, 2005). The relative weight of contents increased both in the gizzard and in the  
41  
42 487 upper part of the small intestine following intake of the silages, which might indicate a  
43  
44 488 prolonged passage time in these segments of the digestive tract. Hetland *et al.* (2004)  
45  
46 489 propose that accumulation of insoluble fibres in the gizzard results in a slower passage rate  
47  
48 490 of especially the fibre fraction. A full gizzard is likely to make the birds feel more satiated  
49  
50 491 resulting in birds appearing more calm, which, in turn, may contribute to a lower feather  
51  
52 492 pecking pressure.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 493 The composition of the microflora in the digestive tract of the hens was  
4  
5 494 influenced only marginally by the dietary treatments. Coliform bacteria and lactose  
6  
7 495 negative bacteria tended to decrease in small intestinal contents of forage fed birds  
8  
9  
10 496 compared to non-supplemented hens (Table 6). These results are in accordance with earlier  
11  
12 497 findings in broilers, where a coarser feed structure, such as feeding whole wheat, reduced  
13  
14 498 the counts of these gram-negative bacteria compared to pellet feeding (Engberg *et al.*,  
15  
16 499 2004, Bjerrum *et al.*, 2005). This may be explained by a lower gizzard pH and longer  
17  
18 500 retention time of feed in the gizzard, providing a barrier for acid labile bacteria, including  
19  
20 501 pathogens like *E. coli* and *Salmonella* (Bjerrum *et al.*, 2005).  
21  
22  
23

24 502 With respect to the lactic acid producing bacteria, only the enterococci were  
25  
26 503 significantly influenced in the entire digestive tract, with the highest counts in hens fed the  
27  
28 504 control diet without forage (Table 6). However, the increase of enterococcus counts in  
29  
30 505 control hens was not reflected by the lactic acid concentration, which was highest in the  
31  
32 506 upper digestive tract (gizzard and duodenum jejunum) of hens fed both types of silages.  
33  
34 507 This might be explained by the contribution of lactic acid in addition to the acetic acid  
35  
36 508 already present in both types of silage.  
37  
38  
39

40 509 The lower pH in the caecal contents of hens given foraging material,  
41  
42 510 compared to the controls, may indicate that forage provides some carbohydrate, which  
43  
44 511 escapes enzymatic hydrolysis in the small intestine and is fermented by the caecal  
45  
46 512 microflora. Although not significant, higher concentrations of acetic, propionic and butyric  
47  
48 513 acid in caecal contents of forage-fed hens (Table 7) may point in this direction. Whether  
49  
50 514 the increased caecal fermentation followed by a pH decrease has a significant beneficial  
51  
52 515 influence on gut health, or following absorption of short chain fatty acids contributes to the  
53  
54 516 energy supply of the bird, is a matter of speculation (Jozefiak *et al.*, 2004).  
55  
56  
57  
58  
59  
60



1  
2  
3 517 Feeding supplements dramatically reduced mortality in the groups given  
4  
5 518 foraging material (0.5-2.5% vs. 15% in the control). Approximately 50% of the mortality  
6  
7 519 recorded in the LD group was identified as cannibalism directed against the cloacal region.  
8  
9 520 Layer diets deficient in protein or methionine and cystine can increase feather pecking and  
10  
11 521 cannibalism (Ambrosen and Petersen, 1997). However, in the present experiment the layer  
12  
13 522 diet contained sufficient essential nutrients to cover requirements, meaning that deficiency  
14  
15 523 in the layer diet cannot explain the high incidence of this kind of cannibalism. Rather it  
16  
17 524 seems that access to supplemental foraging material decreased damaging pecking in  
18  
19 525 general (to feathers as well as skin/cloaca). This reduction in feather pecking was also seen  
20  
21 526 with pheasant chickens given access to green clover, beech branches or range area  
22  
23 527 (Hoffmeyer, 1969; Kjaer, 2004). In a number of studies with laying hens, access to  
24  
25 528 different kinds of foraging material or dietary insoluble fibre resulted in reduced feather  
26  
27 529 pecking and improved plumage condition (Blokhuys, 1986; Nørgaard-Nielsen *et al.*, 1993;  
28  
29 530 Wechsler and Huber-Eicher, 1998; Aerni *et al.*, 2000; Köhler *et al.*, 2001). In addition, El-  
30  
31 531 Lethey (2000) found improved immune response in groups of laying hens with access to  
32  
33 532 foraging material (long-cut straw) compared to controls fed diets without foraging  
34  
35 533 material, which might indicate lower stress. This corresponds well with the fact that  
36  
37 534 divergent selection on feather pecking (Kjaer *et al.*, 2001) seems to affect a range of  
38  
39 535 immune variables (Buitenhuis *et al.*, 2006). The type of feather pecking in this study was  
40  
41 536 predominantly of the severe kind causing damage to the plumage and even feathers pulled  
42  
43 537 out. Therefore it was quite surprising that so few hens had skin damage. Much pecking was  
44  
45 538 directed to the cloacal region and, in many cases, this led to cannibalism and death of the  
46  
47 539 hen prior to the records of pecking damages at 25, 39 and 54 weeks of age. Hughes and  
48  
49 540 Duncan (1972) suggested that there was a distinction between cannibalistic pecking at the  
50  
51 541 cloacal region (vent pecking) and at other parts of the body. All three types of forage  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 542 reduced severe feather pecking behaviour and increased the quality of the plumage,  
4  
5 543 whereas the effect on gentle, non damaging feather pecking was less clear. The  
6  
7  
8 544 significantly higher total feed consumption of the hens fed silages or carrots could indicate  
9  
10 545 that more time was spent on feeding compared to the control LD, suggesting less time was  
11  
12 546 spent feather pecking in these groups..

13  
14  
15 547 In conclusion, access to three types of foraging material, silages and carrots,  
16  
17 548 had beneficial effects on feather pecking behaviour and mortality. In addition, egg-  
18  
19 549 production was improved indicating that high fibre sources, like the silages and carrots,  
20  
21 550 have some nutritional value. Caution must be taken with regard to the use of carrots,  
22  
23 551 because a high intake increases the water content of the excreta, which in turn may result  
24  
25 552 in wet litter increasing the risk of infection with bacteria and parasites. However, moderate  
26  
27 553 amounts of carrots can be recommended. With regard to production, the use of maize  
28  
29 554 silage as supplement was overall a better choice than the barley-pea silage. The very high  
30  
31 555 content of mainly insoluble NSP, including cellulose, in barley-pea silage was probably the  
32  
33 556 reason for the limited feed intake. The results also showed that the influence of forage  
34  
35 557 supplementation on gastrointestinal characteristics depends on the type of foraging  
36  
37 558 material. In contrast to carrots, silages are fermented feeds and provide large amounts of  
38  
39 559 fibre that stimulate gizzard function. Both silage and carrots seem to stimulate caecal  
40  
41 560 fermentation. On the basis of the present results, the provision of foraging material to  
42  
43 561 laying hens can be recommended, and may benefit some aspects of animal welfare.  
44  
45  
46  
47  
48  
49  
50

#### 51 REFERENCES

- 52  
53  
54 563 AMBROSEN, T. & PETERSEN, V.E. (1997) The influence of protein level in the diet on  
55  
56 564 cannibalism and quality of plumage of layers. *Poultry Science*, **76**: 559–563.  
57  
58 565 AERNI, V., EL-LETHEY, H. & WECHSLER, B. (2000) Effect of foraging material and  
59  
60 566 food form on feather pecking in laying hens. *British Poultry Science*, **41**: 16-21.

- 1  
2  
3 567 ASSOCIATION OF OFFICIAL ANALYTICAL CHEMISTS (1990a) Method No. 984.13.  
4  
5 568 Chapter 4, p 74 in *Official methods of analysis*, 15th edn. AOAC, Arlington, VA. 1298  
6  
7 569 pp.  
8  
9  
10 570 ASSOCIATION OF OFFICIAL ANALYTICAL CHEMISTS (1990b) Method No.923.03  
11  
12 571 Chapter 32, p 777 in *Official Methods of Analysis*, 15th edn. AOAC, Arlington, VA. 1298  
13  
14 572 pp.  
15  
16  
17 573 BACH KNUDSEN, K.E. (1997) Carbohydrate and lignin contents of plant materials used in  
18  
19 574 animal feeding. *Animal Feed Science and Technology*, **67**: 319-338.  
20  
21  
22 575 BACH KNUDSEN, K.E. & LI, B. (1991) Determination of oligosaccharides in protein rich  
23  
24 576 feedstuffs by gas-liquid chromatography and high-performance liquid chromatography.  
25  
26 577 *Journal of Agricultural and Food Chemistry*, **39**: 689-694.  
27  
28  
29 578 BEDFORD, M.R. & CLASSEN, H.L. (1992) Reduction of intestinal viscosity through  
30  
31 579 manipulation of dietary rye and pentosanase concentration is effected through changes in  
32  
33 580 the carbohydrate composition of the intestinal aqueous phase and results in improved rates  
34  
35 581 and food conversion efficiency of broiler chicks. *Journal of Nutrition*, **122**: 560-569.  
36  
37  
38 582 BJERRUM, L., ENGBERG, R.M. & PEDERSEN, K. (2005) The influence of whole  
39  
40 583 wheat feeding on salmonella infection and gut flora composition in broilers. *Avian*  
41  
42 584 *Diseases*, **49**: 9-15.  
43  
44  
45 585 BLOKHUIS, H.J. (1986) Feather-pecking in poultry: its relation with ground-pecking.  
46  
47 586 *Applied Animal Behaviour Science*, **16 (1)**: 63-67.  
48  
49  
50 587 BUITENHUIS, B., KJAER, J.B. & JUUL-MADSEN, H.R. (2006) A comparison of  
51  
52 588 plasma serotonin and tryptophan levels, immunological and hematological parameters  
53  
54 589 between laying hen lines divergently selected on feather pecking behavior. *Poultry*  
55  
56 590 *Science*, **85**: 1722-1728.  
57  
58  
59  
60

- 1  
2  
3 591 CARRÉ, B., DEROUET, L. & LECLERCQ, B. (1990) The digestibility of cell wall  
4 polysaccharides from wheat (bran or whole grain), soya bean meal, and white lupin  
5 592  
6 meal in cockerels, muscovy ducks, and rats. *Poultry Science*, **69**: 623-633.  
7 593  
8  
9 594 CHOCT, M. & ANNISON, G. (1992a) The inhibition of nutrient digestion by wheat  
10 pentosans. *British Journal of Nutrition*, **67**: 123-132.  
11 595  
12  
13 596 CHOCT, M. & ANNISON, G. (1992b) Antinutritive effect of wheat pentosans in broiler  
14 chicken: Role of viscosity and gut microflora. *British Poultry Science*, **33**: 821-834.  
15 597  
16  
17 598 CHOCT, M., HUGHES, R.J., WANG, J., BEDFORD, M., MORGAN, A.J. & ANNISON, G.  
18 (1996) Increased small intestine fermentation is partly responsible for the anti-nutritive  
19 599  
20 activity of non-starch polysaccharides in chickens. *British Poultry Science*, **37**: 609-621.  
21 600  
22  
23 601 DER DANSKE FJERKRÆRÅD / DANISH POULTRY COUNCIL (2006) Beretning  
24 2006. pp 128 (In Danish).  
25 602  
26  
27 603 EL-LETHEY, H., AERNI, V., JUNGI, T.W. & WECHSLER, B. (2000) Stress and feather  
28 pecking in laying hens in relation to housing condition. *British Poultry Science*, **41**: 22-  
29 604  
30 28.  
31 605  
32  
33 606 ENGBERG, R.M., HEDEMANN, M.S. & JENSEN, B.B. (2002) The influence of grinding  
34 and pelleting of feed on the microbial composition and activity in the digestive tract of  
35 607  
36 broiler chickens. *British Poultry Science*, **43**: 569-579.  
37 608  
38  
39 609 ENGBERG, R.M., HEDEMANN, M.S., STEENFELDT, S. & JENSEN, B.B. (2004)  
40 Influence of whole wheat and xylanase on broiler performance and microbial  
41 610  
42 composition and activity in the digestive tract. *Poultry Science*, **83**: 925-938.  
43 611  
44  
45 612 GABRIEL, I., MALLET, S. & LECONTE, M. (2003) Differences in the digestive tract  
46 characteristics of broiler chickens fed on complete pelleted diet or on whole wheat  
47 613  
48 added to pelleted protein concentrate. *British Poultry Science*, **44**: 283-290.  
49 614  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 615 HAMMERSHØJ, M. & STEENFELDT, S. (2005) Effects of blue lupin (*Lupinus*  
4 *angustifolius*) in organic layer diets and supplementation with foraging material on egg  
5 616 *production and some egg quality parameters. Poultry Science, 84: 723-733.*  
6  
7  
8  
9  
10 618 HARTINI, S., M. CHOCT, HINCH, G.H., & NOLAN, J.V. (2003) Effect of diet  
11  
12 619 composition, gut microbial status and fibre forms on Cannibalism in Layers. A report  
13  
14 620 for the Australian Egg Corporation Limited. AECL Publication No 03/03, URL:  
15  
16 621 <http://www.aecl.org/images/UNE-72a.pdf> .  
17  
18  
19  
20 622 HETLAND, H. & SVIHUS, B. (2001) Effect of oat hulls on performance, gut capacity and  
21  
22 623 feed passage time in broiler chickens. *British Poultry Science, 42: 354 361.*  
23  
24 624 HETLAND, H., SVIHUS, B. & OLAISEN, V. (2002) Effect of feeding whole cereals on  
25  
26 625 performance, starch digestibility, and duodenal particle size distribution in broiler  
27  
28 626 chickens. *British Poultry Science, 43: 416 423.*  
29  
30  
31 627 HETLAND, H., SVIHUS, B. & KROGDAHL, Å. (2003) Effects of oat hulls and wood  
32  
33 628 shavings on digestion in broilers and layers fed diets based on whole or ground wheat.  
34  
35 629 *British Poultry Science, 44: 275-282.*  
36  
37  
38 630 HETLAND, H., CHOCT, M. & SVIHUS, B. (2004) Role of insoluble non-starch  
39  
40 631 polysaccharides in poultry nutrition. *World's Poultry Science Journal, 60 (4): 415-422.*  
41  
42  
43 632 HILL, F.W. & ANDERSON, D.L. (1958) Comparison of metabolizable energy and  
44  
45 633 productive energy determinations with growing chicks. *Journal of Nutrition, 64: 587-*  
46  
47 634 *603.*  
48  
49  
50 635 HOFFMEYER, I. (1969) Feather pecking in pheasants - an ethological approach to the  
51  
52 636 problem. *Danish Review Game Biology, 6: 1-36*  
53  
54 637 HUGHES, B.O. & DUNCAN, I.J.H. (1972) The influence of strain and environmental  
55  
56 638 factors upon feather pecking and cannibalism in fowls. *British Poultry Science, 13:*  
57  
58 639 *525-547*  
59  
60

- 1  
2  
3 640 HVIDSTEN, H. & HERSTAD, O. (1972) Ensiled and dried barley as feed for poultry.  
4  
5 641 *Acta Agriculturae Scandinavica*, **22**: 183-187.  
6  
7  
8 642 IDI, A., PERMIN, A., CHRISTENSEN, J.P., STEENFELDT, S., ENGBERG, R.M. &  
9  
10 643 FINK, M. (2005) Effect of carrots and maize silage on colonization of hens by  
11  
12 644 *Ascaridia galli* and *Salmonella enterica* serovar Enteritidis. *Helminthologia*, **42 (3)**:  
13  
14 645 121-131.  
15  
16  
17 646 JENSEN, M.T., COX, R.P. & JENSEN, B.B. (1995) Microbial production of skatole in the  
18  
19 647 hind gut of pigs given different diets and its relation to skatole deposition in backfat.  
20  
21 648 *Animal Science*, **61**: 293-304.  
22  
23  
24 649 JÓZEFIAK, D., RUTKOWSKI, A. & MARTIN, S.A. (2004) Carbohydrate fermentation in  
25  
26 650 the avian ceca-A review. *Animal Feed Science and Technology*, **113**: 1-15.  
27  
28  
29 651 JØRGENSEN, H., ZHAO, X., BACH KNUDSEN, K.E. & EGGUM, B.O. (1996) The  
30  
31 652 influence of dietary fibre source and level on the development of the gastrointestinal  
32  
33 653 tract, digestibility and energy metabolism in broiler chickens. *British Journal of*  
34  
35 654 *Nutrition*, **75**: 379-395.  
36  
37  
38 655 KJAER, J.B. (2004) Effect of stocking density and group size on feather and skin  
39  
40 656 condition of the skin and feathers of pheasant chicks. *Veterinary Record*, **154(18)**: 556-  
41  
42 657 558.  
43  
44  
45 658 KJAER, J.B., SØRENSEN, P. & SU, G. (2001) Divergent selection of feather pecking  
46  
47 659 behaviour in laying hens (*Gallus gallus domesticus*). *Applied Animal Behaviour*  
48  
49 660 *Science*, **71**: 29-239  
50  
51  
52 661 KJAER, J.B. & SØRENSEN, P. (2002) Feather pecking and cannibalism in free-range  
53  
54 662 laying hens as affected by genotype, level of dietary methionine + cystine, light  
55  
56 663 intensity during rearing and age at access to the range area. *Applied Animal Behaviour*  
57  
58 664 *Science*, **76**: 21-39.  
59  
60

- 1  
2  
3 665 KÖHLER, B., FÖHLISCH, J., STRUBE, J. & LANGE, K. (2001) Influences of green  
4  
5 666 forage and lighting conditions on egg quality and hen welfare. *Proceedings of the 6th*  
6  
7 667 *European Symposium on Poultry welfare*, Zollikofen, Switzerland, pp.309-311.  
8  
9  
10 668 LANGHOUT, D.J. (1999) The role of the intestinal flora as affected by NSP in broilers.  
11  
12 669 *Proceedings of the 12th European Symposium on Poultry Nutrition*, Veldhoven, The  
13  
14 670 Netherlands, pp. 203-212.  
15  
16  
17 671 LÁZARO, R., GARCÍA, M., ARANÍBAR, M.J. & MATEOS, G.G. (2003) Effects of  
18  
19 672 enzyme addition to wheat-, barley- and rye-based diets on nutrient digestibility and  
20  
21 673 performance of laying hens. *British Poultry Science*, **44**: 256-265.  
22  
23  
24 674 LONGSTAFF, M. & MCNAB, J.M. (1989) Digestion of fibre polysaccharides of pea  
25  
26 675 (*Pisum sativum*) hulls, carrot and cabbage by adult cockerels. *British Journal of*  
27  
28 676 *Nutrition*, **62**: 563-577.  
29  
30  
31 677 MASON, V.C., BECH-ANDERSEN, S. & RUDEMO, M. (1980) Hydrolysate preparation  
32  
33 678 for amino acid determination in feed constituents. *Journal of Animal Physiology and*  
34  
35 679 *Animal Nutrition*, **43**: 146-164.  
36  
37  
38 680 NATIONAL RESEARCH COUNCIL (NRC), (1994) Nutrient Requirements of Poultry.  
39  
40 681 (Ninth Revised Edition. Washington, D.C., National Academic Press, USA).  
41  
42  
43 682 NØRGAARD-NIELSEN, G., VESTERGAARD, K. & SIMONSEN, H.B. (1993) Effects  
44  
45 683 of rearing experience and stimulus enrichment on feather damage in laying hens.  
46  
47 684 *Applied Animal Behaviour Science*, **38**: 345-352.  
48  
49  
50 685 SALIH, M.E., CLASSEN, H.L. and CAMPBELL, G.L. (1991) Response of chickens fed on  
51  
52 686 hull-less barley to dietary  $\beta$ -glucanase of different ages. *Animal Feed Science and*  
53  
54 687 *Technology*, **33**: 139-149.  
55  
56  
57 688 SAS INSTITUTE INC. (1990) SAS/STAT Guide for personal computers. SAS Institute Inc.,  
58  
59 689 Cary, NC, USA.



- 1  
2  
3 690 SAVORY, C.J. (1995) Feather pecking and cannibalism. *World's Poultry Science Journal*,  
4  
5 691 **15 (3):** 281-286.  
6  
7  
8 692 SCOTT, M.J., NESHEIM, M.C. & YOUNG, R.J. (1982). Nutrition of the chicken, (3rd  
9  
10 693 edn. Cornell University, Ithaca, NY).  
11  
12 694 SMITS, C.H.M., VELDMAN, A., VERKADE, H.J. & BEYNEN, A.C. (1998) The inhibitory  
13  
14 695 effect of carboxymethylcellulose with high viscosity on lipid absorption in broiler  
15  
16 696 chickens coincides with reduced bile salt concentration and raised microbial numbers in  
17  
18 697 the small intestine. *Poultry Science*, **77:** 1534-1539.  
19  
20  
21 698 STEENFELDT, S. (2001) The dietary effect of different wheat cultivars for broiler chickens.  
22  
23 699 *British Poultry Science*, **42:** 595-609.  
24  
25  
26 700 STEENFELDT, S., BACH KNUDSEN, K.E., BØRSTING, C.F. & EGGUM, B.O. (1995)  
27  
28 701 The nutritive value of decorticated mill fractions of wheat. 2. Evaluations with raw and  
29  
30 702 enzyme treated fractions using adult cockerels. *Animal Feed Science and Technology*,  
31  
32 703 **54:** 249-265.  
33  
34  
35 704 STEENFELDT, S., HAMMERSHØJ, M., MÜLLERTZ, A. & JENSEN, J.F. (1998) Enzyme  
36  
37 705 supplementation to wheat-based diets for broilers. 2. Effect on apparent metabolisable  
38  
39 706 energy and nutrient digestibility. *Animal Feed Science and Technology*, **75:** 45-64.  
40  
41  
42 707 STOLDT, W. (1952) Vorschlag zur Vereinheitlichung der Fettbestimmung in Lebensmitteln  
43  
44 708 (Suggestion to standardise the determination of fat in foodstuffs). *Fetten, Seifen*,  
45  
46 709 *Anstrichmittel*, **54:** 206-207.  
47  
48  
49 710 SVIHUS, B. & HETLAND, H. (2001) Ileal starch digestibility in growing broiler chickens  
50  
51 711 fed on a wheat-based diets is improved by mash feeding, dilution with cellulose or  
52  
53 712 whole wheat inclusion. *British Poultry Science*, **42:** 633-637.  
54  
55  
56 713 TAUSON, R., KJAER, J.B., MARIA LEVRINO, G. & CEPERO BRIZ, R. (2005) Applied  
57  
58 714 scoring of integument and health in laying hens. *Proceedings of the 7th European*

- 1  
2  
3 715     *Symposium on Poultry Welfare, 15-19 June, Lublin, Poland. Polish Academy of*  
4  
5 716     *Sciences, 23(Supp 1): 153-159.*  
6  
7  
8 717     TKALCIC, E., STEKAR, J.M.A., KOVAC, M. & HOLCMAN, A. (2000) The effect of  
9  
10 718     ensiled maize grains on laying of hens. *Zbornik Biotehnske fakultete Univerze v*  
11  
12 719     *Ljubljani, Kmetijstvo Zootehnika, 76: 67-81.*  
13  
14  
15 720     THEANDER, O., WESTERLUND, E.A., ÅMAN, P. & GRAHAM, H. (1989) Plant cell  
16  
17 721     walls and monogastric diets. *Animal Feed Science and Technology, 23: 205-225.*  
18  
19  
20 722     THEANDER, O., ÅMAN, P., WESTERLUND, E.A. & GRAHAM, H. (1994)  
21  
22 723     Enzymatic/chemical analysis of dietary fiber. *Journal of AOAC International, 77: 703-*  
23  
24 724     709.  
25  
26  
27 725     WECHSLER, B. & HUBER-EICHER, B. (1998) The effect of foraging material and perch  
28  
29 726     height on feather pecking and feather damage in laying hens. *Applied Animal*  
30  
31 727     *Behaviour Science, 58: 131-141.*  
32  
33  
34 728     WENNRICH, G. (1975) Beziehungen zwischen ranghöhe, federpickaktivität und  
35  
36 729     aggressivität von haushennen (*Gallus domesticus*). *Archives für Geflügelkunde, 39:*  
37  
38 730     167-171.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Table 1.** *Composition of the layer diet (g/kg).*

Ingredients	Diet
Wheat	383
Barley	160
Oat	100
Soyabean meal	136
Fish meal	20
Meat-and-bone-meal	37
Animal fat	27
Soyabean oil	20
Calcium carbonate	70
Dicalcium phosphate	5
Sodium chloride	1
Sodium bicarbonate	2.5
Alfalfa meal	32
DL-Methionine (40%)	4
Vitamin and mineral premix <sup>1</sup>	2.5
Calculated content:	
ME, MJ per kg DM	11.5
Protein (Nx6.25, g/kg)	174
Lysine (g/kg)	8.3
Methionine (g/kg)	4.4
Methionine + cystine (g/kg)	7.2
Calcium (g/kg)	33.5
Total phosphorus (g/kg)	5.4

<sup>1</sup>The vitamin and mineral premix provided per kg of diet: Vitamin A 4.1 mg; Vitamin D<sub>3</sub> 0.08 mg; Vitamin E 30 mg; Vitamin K<sub>3</sub> 2.5 mg; Vitamin B<sub>1</sub>; 1 mg; Vitamin B<sub>2</sub>, 5 mg; Vitamin B<sub>6</sub>, 2.5 mg; D-pantothenic acid 9 mg; Niacin 30 mg; Betaine anhydrate 338 mg; Folic acid 1 mg; Biotin 0.05 mg; Vitamin B<sub>12</sub> 0.02 mg; Fe 80 mg; Zn 120 mg; Mn 100 mg; Cu 6 mg; I 0.5; Se 0.3 mg.

**Table 2.** Chemical composition of layer diet, maize<sup>1</sup>- and barley/pea silage<sup>1</sup> and carrots<sup>1</sup> (g/kg DM).

Constituents	Layer diet	Maize silage	Barley/pea silage	Carrots
Dry matter	914.4	317.1-328.8	212.1-230.4	83.7-125.2
Ash	126.6	42.4-37.8	73.3-73.2	63.4-53.8
Protein (Nx6.25)	186.9	98.8-89.4	122.5-126.9	72.8-65.0
HCL-fat	75.9	32.4	29.2-28.3	11.6-16.4
Gross energy (MJ/kg DM)	181.0	19.0-18.7	18.3-18.5	16.1-16.7
Starch	364.9	304.8-319.2	123.5-125.6	t
Sugars				
Glucose	0.4	t	t	150.0-155.5
Fructose	1.0	t	0.5	157.5-142.5
Sucrose	15.6	t	t	150.0-234.5
Raffinose	2.2	t	t	t
Stachyose	6.4	t	t	t
Verbascose	t	t	t	t
Cellulose	33	171-166	201-180	60-66
NCP <sup>2</sup>				
Rhamnose	1	t	3-1	4-5
Fucose	t	t	t	1
Arabinose	23	22-21	16	20-11
Xylose	38	11-98	86-77	3
Mannose	4	3	7	4
Galactose	10	8	9	31-16
Glucose	17	13-12	12	3-4
Uronic acid	9	16	56-58	70-94
NSP <sup>3</sup>	133(29) <sup>4</sup>	342(13)-323(28)	390(47)-361(42)	196(112)-203(112)
Lignin	23	75-80	93-103	30-35
DF (NSP+lignin)	156	417-403	483-464	226-238

<sup>1</sup>Values given represent analyses from the supplement samples taken at hen age 23 (I) and 54 (II) weeks.

<sup>2</sup>Non-cellulosic polysaccharides.

<sup>3</sup>Non-starch polysaccharides (NCP+cellulose).

<sup>4</sup>Values in parentheses are the soluble fraction of NCP.

t=trace.

**Table 3.** Production, body weight and mortality of laying hens fed a layer diet with or without supplements of maize silage, barley pea silage or carrots.

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM <sup>1</sup>	P-value
Egg production, per hen						
Number of eggs	214 <sup>ab</sup>	218 <sup>a</sup>	208 <sup>b</sup>	219 <sup>a</sup>	1.61	0.04
Egg mass, kg	13.2 <sup>ab</sup>	13.3 <sup>ab</sup>	12.8 <sup>b</sup>	13.6 <sup>a</sup>	0.10	0.03
Egg weight, g	61.5	61.1	61.5	61.9	0.16	0.36
Rate of lay, %	89.9 <sup>ab</sup>	91.4 <sup>a</sup>	87.2 <sup>b</sup>	92.0 <sup>a</sup>	0.67	0.04
Feed consumption, per hen						
Layer diet, kg	31.0 <sup>a</sup>	28.2 <sup>ab</sup>	25.5 <sup>b</sup>	27.2 <sup>b</sup>	0.63	0.004
Layer diet, g/day	130.1 <sup>a</sup>	118.5 <sup>ab</sup>	107.3 <sup>b</sup>	114.1 <sup>b</sup>	2.63	0.004
Silage/carrots, kg	-	14.1 <sup>b</sup>	13.8 <sup>b</sup>	25.6 <sup>a</sup>	1.66	0.0001
Silage/carrots, g/day	-	59.3 <sup>b</sup>	58.1 <sup>b</sup>	107.6 <sup>a</sup>	6.99	0.0001
LD + silage/carrots, g/day	130.1 <sup>a</sup>	177.7 <sup>b</sup>	165.4 <sup>b</sup>	221.7 <sup>a</sup>	8.64	0.0001
Silage/carrots, % of total feed intake	-	33.4 <sup>b</sup>	35.1 <sup>b</sup>	48.5 <sup>a</sup>	2.06	0.0001
Feed consumption, kg per kg egg						
Layer diet	2.35 <sup>a</sup>	2.12 <sup>ab</sup>	2.00 <sup>b</sup>	2.00 <sup>b</sup>	0.05	0.02
Silage/carrots	-	1.06 <sup>b</sup>	1.08 <sup>b</sup>	1.88 <sup>a</sup>	0.12	0.0001
LD + silage/carrots	2.35 <sup>a</sup>	3.18 <sup>b</sup>	3.08 <sup>b</sup>	3.88 <sup>a</sup>	0.15	0.0001
Body weight, g						
25 weeks	1750	1742	1718	1726	18.1	0.59
39 weeks	1889 <sup>a</sup>	1796 <sup>b</sup>	1795 <sup>b</sup>	1915 <sup>a</sup>	23.4	0.0001
54 weeks	1813 <sup>b</sup>	1787 <sup>b</sup>	1805 <sup>b</sup>	1917 <sup>a</sup>	24.8	0.001
Mortality, % of hens housed	15.3 <sup>a</sup>	1.5 <sup>b</sup>	2.5 <sup>b</sup>	0.5 <sup>b</sup>	2.10	0.02

Means in each row followed by different superscript letters <sup>a</sup> <sup>b</sup> differ significantly.

<sup>1</sup>Standard error of mean.

**Table 4.** Apparent metabolisable energy (MJ/kg DM), coefficients of total tract apparent digestibility and nitrogen retention in laying hens fed a layer diet with or without supplements of maize silage, barley pea silage or carrots.

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM <sup>4</sup>	P-value
23 weeks of age						
Organic matter	0.763	0.760	0.758	0.734	0.01	0.10
Fat	0.872	0.853	0.847	0.833	0.01	0.31
Starch	0.999 <sup>a</sup>	0.998 <sup>a</sup>	0.992 <sup>b</sup>	0.997 <sup>a</sup>	0.01	0.01
Total NSP <sup>1</sup>	0.309 <sup>a</sup>	0.276 <sup>ab</sup>	0.288 <sup>a</sup>	0.131 <sup>b</sup>	0.02	0.03
S-NSP <sup>2</sup>	0.414	0.429	0.353	0.381	0.02	0.87
I-NSP <sup>3</sup>	0.280 <sup>a</sup>	0.245 <sup>a</sup>	0.273 <sup>a</sup>	0.041 <sup>b</sup>	0.02	0.03
N-retention	0.472 <sup>ab</sup>	0.521 <sup>a</sup>	0.472 <sup>ab</sup>	0.412 <sup>b</sup>	0.02	0.005
AME <sub>n</sub>	13.39 <sup>a</sup>	13.41 <sup>a</sup>	13.14 <sup>ab</sup>	12.78 <sup>b</sup>	0.09	0.04
DM in excreta	22.39	17.86	19.88	19.14	0.69	0.11
53 weeks of age						
Organic matter	0.721 <sup>a</sup>	0.703 <sup>ab</sup>	0.689 <sup>b</sup>	0.706 <sup>ab</sup>	0.01	0.03
Fat	0.848 <sup>ab</sup>	0.861 <sup>a</sup>	0.848 <sup>ab</sup>	0.804 <sup>b</sup>	0.01	0.03
Starch	0.993	0.994	0.983	0.985	0.01	0.08
Total NSP <sup>1</sup>	0.220	0.239	0.264	0.244	0.01	0.70
S-NSP <sup>2</sup>	0.270	0.497	0.543	0.258	0.04	0.34
I-NSP <sup>3</sup>	0.206	0.185	0.201	0.240	0.02	0.60
N-retention	0.419	0.400	0.419	0.363	0.02	0.74
AME <sub>n</sub>	12.82	12.61	12.36	12.42	0.08	0.19
DM in excreta	22.31	20.06	21.06	17.00	0.82	0.11

Means in each row followed by different superscript letters <sup>a, b</sup> differ significantly.

<sup>1</sup>Non-starch polysaccharides (NCP+cellulose).

<sup>2</sup>Soluble NSP.

<sup>3</sup>Insoluble NSP.

<sup>4</sup>Standard error of mean.

**Table 5.** Relative weights of digestive tract segments and their contents and dry matter content in gastrointestinal contents of hens fed with or without supplements of maize silage, barley pea silage or carrots at 23 and 53 weeks of age.

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM <sup>1</sup>	P-value
<i>Relative segment weight, g/kg BW</i>						
23 weeks						
Gizzard	20.62 <sup>b</sup>	24.72 <sup>a</sup>	24.87 <sup>a</sup>	22.51 <sup>ab</sup>	1.06	0.04
Duodenum/jejunum	20.14	20.86	19.93	19.97	0.60	0.68
Ileum	7.56	8.53	8.60	8.67	0.52	0.42
Caeca	4.51	4.50	4.57	4.73	0.38	0.97
53 weeks						
Gizzard	16.3 <sup>b</sup>	30.5 <sup>a</sup>	33.0 <sup>a</sup>	19.0 <sup>b</sup>	1.21	0.001
Duodenum/jejunum	16.3	16.1	15.2	15.7	0.64	0.67
Ileum	6.50	7.98	7.35	7.05	0.44	0.17
Caeca	4.67	4.14	4.11	4.30	0.38	0.71
<i>Relative weight of contents, g/kg BW</i>						
23 weeks						
Gizzard	8.35 <sup>b</sup>	11.69 <sup>a</sup>	11.27 <sup>a</sup>	9.89 <sup>ab</sup>	0.60	0.01
Duodenum/jejunum	7.20 <sup>c</sup>	9.69 <sup>a</sup>	7.64 <sup>bc</sup>	9.26 <sup>ab</sup>	0.63	0.04
Ileum	5.64 <sup>b</sup>	6.71 <sup>ab</sup>	5.74 <sup>b</sup>	7.54 <sup>a</sup>	0.41	0.02
Caeca	3.26	3.24	3.64	3.40	0.40	0.89
53 weeks						
Gizzard	8.2 <sup>b</sup>	14.0 <sup>a</sup>	15.9 <sup>a</sup>	8.3 <sup>b</sup>	0.67	0.001
Duodenum/jejunum	9.8 <sup>c</sup>	14.9 <sup>a</sup>	13.5 <sup>ab</sup>	11.2 <sup>bc</sup>	0.76	0.004
Ileum	5.50	7.59	6.55	6.58	0.65	0.21
Caeca	3.80	3.18	2.21	3.78	0.43	0.07
<i>Dry matter content, %</i>						
23 weeks						
Gizzard	34.13 <sup>a</sup>	34.02 <sup>a</sup>	29.24 <sup>b</sup>	24.19 <sup>c</sup>	1.59	0.002
Duodenum/jejunum	19.00 <sup>a</sup>	18.72 <sup>a</sup>	18.39 <sup>a</sup>	15.81 <sup>b</sup>	0.70	0.03
Ileum	22.88	20.87	21.83	19.47	0.82	0.07
Caeca	22.79	22.53	23.68	21.53	0.95	0.48
Colon/rectum	25.36	24.11	23.80	23.60	0.75	0.38
53 weeks						
Gizzard	33.56 <sup>a</sup>	29.46 <sup>b</sup>	23.96 <sup>c</sup>	21.71 <sup>c</sup>	1.09	0.001
Duodenum/jejunum	21.29 <sup>a</sup>	18.66 <sup>b</sup>	18.54 <sup>b</sup>	19.20 <sup>b</sup>	0.49	0.01
Ileum	29.76	19.88	20.17	19.87	3.69	0.21
Caeca	28.56	26.14	25.02	24.16	1.31	0.15
Colon/rectum	22.44	20.98	19.23	20.32	0.91	0.14

Means in each row followed by different superscript letters <sup>a b c</sup> differ significantly.

<sup>1</sup>Standard error of mean.

**Table 6.** Counts of total anaerobic bacteria, coliform bacteria, lactose negative enterobacteria, lactic acid bacteria, enterococci (log cfu/g digesta ) and pH in the digestive tract of hens fed with or without supplements of maize silage, barley pea silage or carrots.

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM <sup>1</sup>	P-value
<b>Anaerobic bacteria</b>						
Gizzard	7.68	7.15	7.54	7.54	0.25	0.49
Duodenum/jejunum	8.52	8.06	8.23	7.97	0.26	0.50
Ileum	8.59	8.51	8.77	8.52	0.14	0.52
Caeca	10.06	9.79	9.95	10.21	0.13	0.21
Colon/rectum	9.15	8.76	8.96	8.84	0.18	0.48
<b>Coliform bacteria</b>						
Gizzard	4.54	3.72	4.22	4.14	0.32	0.38
Duodenum/jejunum	5.73	4.28	5.13	4.90	0.44	0.19
Ileum	7.29	7.07	6.59	6.42	0.27	0.17
Caeca	7.39	7.15	7.28	7.31	0.17	0.78
Colon/rectum	7.08	7.48	7.24	7.10	0.27	0.71
<b>Lactose negative enterobacteria</b>						
Gizzard	3.76	3.43	3.47	3.46	0.18	0.54
Duodenum/jejunum	4.67 <sup>a</sup>	3.06 <sup>b</sup>	3.85 <sup>a</sup>	3.85 <sup>a</sup>	0.34	0.04
Ileum	6.40	5.76	5.46	5.82	0.26	0.19
Caeca	6.06	5.92	5.52	6.08	0.28	0.50
Colon/rectum	6.18	5.91	5.65	6.08	0.25	0.47
<b>Lactic acid bacteria</b>						
Gizzard	7.63	7.49	7.49	7.69	0.41	0.98
Duodenum/jejunum	8.47	8.38	7.99	8.43	0.20	0.32
Ileum	8.96	8.73	8.46	8.79	0.16	0.23
Caeca	9.53	9.43	9.56	9.68	0.10	0.40
Colon/rectum	8.93	8.84	8.60	9.15	0.13	0.06
<b>Enterococci</b>						
Gizzard	5.64 <sup>a</sup>	3.89 <sup>b</sup>	4.04 <sup>b</sup>	5.05 <sup>a</sup>	0.27	0.002
Duodenum/jejunum	6.85 <sup>a</sup>	4.70 <sup>c</sup>	4.70 <sup>c</sup>	5.74 <sup>b</sup>	0.27	0.001
Ileum	7.72 <sup>a</sup>	6.49 <sup>b</sup>	6.01 <sup>b</sup>	6.55 <sup>b</sup>	0.32	0.02
Caeca	7.96	7.43	6.97	7.95	0.25	0.05
Colon/rectum	7.41	6.85	6.82	7.16	0.29	0.44
<b>pH</b>						
Gizzard	4.82 <sup>a</sup>	3.91 <sup>b</sup>	3.90 <sup>b</sup>	4.64 <sup>a</sup>	0.21	0.02
Duodenum/jejunum	6.27 <sup>a</sup>	5.92 <sup>b</sup>	6.22 <sup>a</sup>	6.08 <sup>ab</sup>	0.06	0.01
Ileum	7.48	7.44	7.57	7.17	0.18	0.47
Caeca	6.53 <sup>a</sup>	6.18 <sup>b</sup>	6.14 <sup>b</sup>	6.14 <sup>b</sup>	0.10	0.04
Colon/rectum	7.19	7.14	7.21	7.17	0.26	0.99

Means in each row followed by different superscript letters <sup>a</sup><sup>b</sup> differ significantly.

<sup>1</sup>Standard error of mean.

**Table 7.** Concentrations of lactic - and acetic acid in the digestive tract and caecal concentrations of butyric and propionic acid ( $\mu\text{mol/g}$  digesta) of hens fed with or without supplements of either maize silage, barley pea silage or carrots.

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM	P-value
Lactic acid						
Gizzard	4.7 <sup>b</sup>	16.9 <sup>a</sup>	14.3 <sup>a</sup>	1.6 <sup>b</sup>	2.81	0.01
Duodenum/jejunum	16.4 <sup>b</sup>	25.0 <sup>a</sup>	19.6 <sup>ab</sup>	13.5 <sup>b</sup>	2.08	0.01
Ileum	13.9	22.3	12.8	17.5	3.83	0.34
Colon/rectum	13.5	16.6	14.1	17.0	3.65	0.84
Acetic acid						
Gizzard	3.5 <sup>b</sup>	5.9 <sup>a</sup>	6.2 <sup>a</sup>	3.6 <sup>b</sup>	0.71	0.03
Duodenum/jejunum	1.5	0.2	1.3	0.7	0.65	0.41
Ileum	9.7 <sup>a</sup>	11.5 <sup>a</sup>	9.7 <sup>a</sup>	6.8 <sup>b</sup>	0.82	0.01
Colon/rectum	26.1	24.5	24.0	25.0	4.43	0.99
Caeca						
Acetic acid	70.0	91.5	77.0	74.3	7.30	0.24
Propionic acid	25.0	32.0	27.0	27.3	2.67	0.35
Butyric acid	13.6	17.1	16.5	18.8	2.06	0.39

Means in each row followed by different superscript letters<sup>a,b</sup> differ significantly.

<sup>1</sup>Standard error of mean.

**Table 8.** *Allo-pecking at 53 week of age (LSmeans of bouts/pecks per bird per hour) and plumage condition at 54 weeks of age of hens fed with or without supplements of either maize silage, barley pea silage or carrots.*

	Layer diet (LD)	Layer diet + maize silage (LD+MS)	Layer diet + barley/pea silage (LD+BS)	Layer diet + carrots (LD+C)	SEM <sup>3</sup>
Feather pecking					
-Total, bouts	0.92 <sup>a</sup>	0.21 <sup>b</sup>	0.44 <sup>b</sup>	0.20 <sup>b</sup>	0.09
-Total, pecks	1.58 <sup>a</sup>	0.57 <sup>ab</sup>	0.95 <sup>ab</sup>	0.36 <sup>b</sup>	0.15
-Severe, bouts	0.56 <sup>a</sup>	0.01 <sup>b</sup>	0.03 <sup>b</sup>	0.07 <sup>b</sup>	0.08
-Severe, pecks	0.60 <sup>a</sup>	0.03 <sup>b</sup>	0.04 <sup>b</sup>	0.09 <sup>b</sup>	0.08
-Gentle, bouts	0.36 <sup>a</sup>	0.20 <sup>b</sup>	0.42 <sup>a</sup>	0.13 <sup>b</sup>	0.04
-Gentle, pecks	0.98	0.54	0.91	0.27	0.15
Aggressive pecking					
-Bouts	0.21	0.18	0.19	0.23	0.02
-Pecks	0.22	0.21	0.21	0.24	0.02
Beak pecking					
-Bouts	0.25	0.23	0.10	0.17	0.04
-Pecks	0.98	0.78	0.32	0.51	0.17
Plumage condition <sup>1</sup>					
-Total score <sup>2</sup>	13.9 <sup>b</sup>	18.3 <sup>a</sup>	19.2 <sup>a</sup>	16.3 <sup>ab</sup>	1.24
Separate body parts					
-Neck	3.2	3.8	3.9	3.6	0.19
-Breast	2.4	3.1	3.5	3.1	0.26
-Back	2.4	3.8	4.0	3.0	0.38
-Wings	2.4	3.8	3.9	3.6	0.21
-Tail	2.6	3.8	3.9	3.1	0.33

Means in each row followed by different superscript letters <sup>a</sup><sup>b</sup> differ significantly ( $P < 0.05$ ).

<sup>1</sup>Maximum of 4 points for perfect plumage, minimum of 1 point for a very poor plumage.

<sup>2</sup>Total score is the sum of body part scores, max. 20 and min. 5 points.

<sup>3</sup>Standard error of mean.