HAL
open science

REMARKS ON SYMMETRIES OF 2 D-QUASICRYSTALS (SI - CMMSE - 2006)

Sergio Sánchez

To cite this version:

Sergio Sánchez. REMARKS ON SYMMETRIES OF 2 D-QUASICRYSTALS (SI - CMMSE - 2006). International Journal of Computer Mathematics, 2008, 85 (03-04), pp.319-328. 10.1080/00207160701429766 . hal-00545356

HAL Id: hal-00545356

https://hal.science/hal-00545356v1

Submitted on 10 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



REMARKS ON SYMMETRIES OF 2 D-QUASICRYSTALS (SI - CMMSE - 2006)

Journal:	*International Journal of Computer Mathematics*
Manuscript ID:	GCOM-2007-0229
Manuscript Type:	Original Article
Date Submitted by the Author:	23-Apr-2007
Complete List of Authors:	Sánchez, Sergio; Rey Juan Carlos, Applied Mathematics
Keywords:	symmetries, quasicrystals, a model set, a cut and project set, Abelian group

scholarONE™
Manuscript Central

REMARKS ON SYMMETRIES OF 2 D-QUASICRYSTALS (SI - CMMSE - 2006)

V.ARTAMONOV, S.SÀNCHEZ

Abstract. The paper presents a strict mathematical proof of the classifications of finite groups of symmetries of 2-dimensional quasicrystals given in [4].

For Peer Review Only

Introduction

There are various approaches to a construction of mathematical models of quasicrystals and a definition of their symmetries, see for example [1], [2, Chapter 6], [4], [5], [9]. We shall adopt the following model which is usually call *cut and project scheme* [9]. Nowadays the common definition of this method is the following one. Let V be an additive locally compact topological Abelian group, U an additive group of a real *physical* or *internal* vector space of dimension d and M a discrete subgroup in $E = U \oplus V$ such that E/M is compact and $M \cap V$ consists only of zero vector. The group E is often called a *hyperspace* or *embedding space*. Consider the diagram of projections of groups

$$\begin{array}{ccccc} U & \xleftarrow[\pi]{} & E = U \oplus V & \xrightarrow{\rho} & V \\ & & \bigcup & & \\ & & M & & \end{array}$$

Since the intersection **of** V and M consists only of zero element the projection π is injective on M. It is assumed that $\rho(M)$ is dense is in V.

Let us take a convex compact subset W in V called a *window*. Let us put $Q = \rho^{-1}(W) \cap M$. Then the set $\pi(Q)$ is called a *model set* or a *cut and project set*. Note that the set Q under the projection π is mapping injectively into U.

It is necessary to mention [8, Chapter 2, §3.2] that any locally compact Abelian group V is an inverse limit of direct product of additive groups $\mathbb{R}^m \times \mathbb{T}^s \times D$ where $\mathbb{T}$ is a one-dimensional torus and D is a

Date: September 29, 2006.
2000 *Mathematics Subject Classification.* Primary 52C23; Secondary 37L20.
Key words and phrases. symmetries, quasicrystals.
Research partially supported by grants RFBR 06-01-00037, NSh-5666.2006.1.

discrete group. So the case when V is a real vector space is in fact one of the most important.

Throughout the paper we shall consider the special case of the model when V is a normed real vector space of a finite dimension $n-d>0$. Then E is a vector space of dimension $n>d$ and M is a lattice in E. It means that M is discrete and $E=\mathbb{R}\otimes_{\mathbb{Z}} M$. Equivalently the rank of the free Abelian group M is equal to n. We shall assume that Q contains a basis of M as a free Abelian group.

Since all norms in E are equivalent without loss of generality we can assume that E is an Euclidean space.

According to [1] and [2, Chapter 6] a symmetry of the model Q in this case is the group of all affine transformations of E which map Q bijectively onto itself. Since Q contains a basis of M each symmetry leaves the lattice M invariant. We shall now call all these symmetries *proper*. **It is necessary to mention that there exist non-proper symmetries. For example for Fibonacci tilings powers of the matrix**

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

show examples of non-proper symmetries.

Suppose that M has also zero intersection with V. If f is a symmetry of Q and 0 is the origin then $f(0)\in M\cap U=0$. Hence each symmetry f is in fact a linear operator. As in [1] we can show that U is invariant under an action of f.

Generalizing this idea we shall denote by $\operatorname{Sym} Q$ the subgroup in the group $\mathrm{GL}(E)$ of all invertible linear operators in E such that both U and M are invariant under an action of each element from $\operatorname{Sym} Q$. The group of proper symmetries is a subgroup of $\operatorname{Sym} Q$. Each element of the group $\operatorname{Sym} Q$ in some special basis of E has the lower left zero block from (1). Moreover there exists an invertible matrix $Z\in \mathrm{GL}(n,\mathbb{R})$ such that $Z^{-1}\operatorname{Sym} Q\, Z\subset \mathrm{GL}(n,\mathbb{Z})$. These two conditions are equivalent to the definition of the group $\operatorname{Sym} Q$.

In the paper we shall consider the case when $\dim U=\dim V=2$ and G is a finite subgroup of $\operatorname{Sym} Q$. The main results of the paper are in Theorems 2.5, 2.11, 2.12, 2.13, 2.14, 2.15. The list of these finite groups is given in [4] without proof. Here we justify this classification and expose strict mathematical proofs based on group theory approach. The main results of the present paper are complementary to those in [4], [3] where realizations of some symmetries are presented. We give a list of group and in also in §3 their representation in 4-dimensional hyperspace.

We thank Prof. E.S. Golod for useful consultations concerning cyclotomic fields and the referee for his valuable comments and remarks.

For Peer Review Only

1. Matrix representations of symmetries

Suppose that $e_1, \dots, e_n$ is a basis of the lattice M. Then these elements form a basis of E. If $\mathcal{A} \in \operatorname{Sym} Q$ and A is a matrix of $\mathcal{A}$ is the basis $e_1, \dots, e_n$. Since M is invariant under $\mathcal{A}$ we can conclude that A is an integer matrix, $A \in \mathrm{GL}(n, \mathbb{Z})$.

Let $u_1, \dots, u_d$ be a basis of U and $u_{d+1}, \dots, u_n$ a basis of V. Suppose that

$$(u_1, \dots, u_n) = (e_1, \dots, e_n)C, \quad C \in \mathrm{GL}(n, \mathbb{R}).$$

Since U is invariant under $\mathcal{A}$ we have

$$C^{-1}AC = \begin{pmatrix} A_1 & A_2 \\ 0 & A_3 \end{pmatrix}, \qquad (1)$$
$$A_1 \in \mathrm{GL}(d, \mathbb{R}),\ A_3 \in \mathrm{GL}(n-d, \mathbb{R}),\ A_2 \in \mathrm{Mat}(d \times (n-d), \mathbb{R}).$$

We shall use throughout the paper these properties of matrices representing elements from $\operatorname{Sym} Q$.

2. 2D symmetries

In this section we shall assume that $n = 4$, $d = 2$ and G is a finite subgroup in $\operatorname{Sym} Q$. Recall that a dihedral group D_k is a group of orthogonal symmetries of a regular k-gon [2]. D_k is a subgroup of the group of orthogonal matrices $\mathrm{O}(2, \mathbb{R})$ generated by two matrices

$$a = \begin{pmatrix} \cos\frac{2\pi}{k} & -\sin\frac{2\pi}{k} \\ \sin\frac{2\pi}{k} & \cos\frac{2\pi}{k} \end{pmatrix}, \quad b = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \qquad (2)$$

It is known [2] that D_k consists of $2k$ elements

$$1,\ a, \dots, a^{k-1},\ b,\ ba,\ \dots, ba^{k-1},$$

where $a^k = (ba^l)^2 = 1$ for any l. In particular $ba^l = a^{-l}b$. Note that

$$a^l = \begin{pmatrix} \cos\frac{2\pi l}{k} & -\sin\frac{2\pi l}{k} \\ \sin\frac{2\pi l}{k} & \cos\frac{2\pi l}{k} \end{pmatrix}, \quad ba^l = \begin{pmatrix} \cos\frac{2\pi l}{k} & \sin\frac{2\pi l}{k} \\ \sin\frac{2\pi l}{k} & -\cos\frac{2\pi l}{k} \end{pmatrix} \qquad (3)$$

In order to classify finite subgroups in $\operatorname{Sym} Q$ we need some auxiliary statements. We exclude some proofs of known facts.

Proposition 2.1. *Let H be a finite subgroup of $\mathrm{O}(2, \mathbb{R})$. If $H \subset \mathrm{SO}(2, \mathbb{R})$ then H is a cyclic group. If $H \not\subseteq \mathrm{SO}(2, \mathbb{R})$, then $H = D_k$ for some k.*

Proposition 2.2. *Suppose that a finite group G is a subgroup of a direct product of two cyclic groups $H = \langle a_1 \rangle_{k_1} \times \langle a_2 \rangle_{k_2}$ of orders k_1 and k_2 respectively. Denote by $\nu_i : G \to \langle a_i \rangle_{k_i}$ the natural projection and suppose that ν_1, ν_2 are surjective. Then $(a_1^{s_1}, a_2), (a_1, a_2^{s_2}) \in G$ for some $s_i \in \mathbb{Z}$ and therefore $a_i^{s_1 s_2 - 1} \in G \cap \langle a_i \rangle_{k_i}$. If $d = (k_1, k_2)$ is the greatest common divisor of k_1, k_2 then $a_i^d \in G \cap \langle a_i \rangle_{k_i}$ for $i = 1, 2$. In particular $G \cap \langle a_i \rangle_{k_i}$ is a cyclic group $\langle a_i^{l_i} \rangle$, where $l_i \mid d$ and $l_i \mid (s_1 s_2 - 1)$.*

For Peer Review Only

Proof. Since ν_1 is surjective, then there exists an element $(a_1, b) \in G$ for some $b \in \langle a_2 \rangle$. Then $(a_1, b)^{k_2} = (a_1^{k_2}, b^{k_2}) = (a_1^{k_2}, 1) \in G \cap \langle a_1 \rangle$. Since the order of a_1 is k_1 by [2, Theorem 1.6, § 1.4] we can conclude that $a_1^d \in G \cap \langle a_1 \rangle$. Apply [2, Theorem 1.5, § 1.4]. Similarly if $(a_1^{s_1}, a_2), (a_1, a_2^{s_2}) \in G$, then

$$(a_1^{s_1}, a_2)^{s_2}(a_1, a_2^{s_2})^{-1} = (a_1^{s_1 s_2 - 1}, 1) \in G \cap \langle a_1 \rangle .$$

Similarly $a_2^{s_1 s_2 - 1} \in G \cap \langle a_2 \rangle$. □

Recall that the *Euler* function $\phi(m)$, $m \in \mathbb{Z}$, is the number of all integers $1 \leqslant k < m$ which are coprime with m. Recall that if $\zeta = \exp \frac{2\pi i}{m}$ then $\phi(m)$ coincides with the degree of the minimal integer cyclotomic polynomial $\Phi_m(X) \in \mathbb{Z}[X]$ such that $\Phi_m(\zeta) = 0$ and with the order of the Galois group $\mathrm{Gal}\,(\mathbb{Q}[\xi]/\mathbb{Q})$.

Proposition 2.3. *Let m be a positive integer such that $\phi(m) \leqslant 4$. Then m is one of the numbers*

m	10	5	12	6	3	8	4	2	1
$\phi(m)$	4	4	4	2	2	4	2	1	1

Proposition 2.4 ([2], Theorem 2.16, § 2.4)**.** *Let Δ be a finite subgroup in $\mathrm{SO}(2, \mathbb{R})$. Then $\Delta = \langle a \rangle_k$ is a cyclic group of order k, where*

$$a = \begin{pmatrix} \cos \frac{2\pi}{k} & -\sin \frac{2\pi}{k} \\ \sin \frac{2\pi}{k} & \cos \frac{2\pi}{k} \end{pmatrix} .$$

Moreover, the following are equivalent

1) *Δ is conjugate in $\mathrm{GL}(2, \mathbb{R})$ to a subgroup of $\mathrm{SL}(2, \mathbb{Z})$;*
2) *the trace $\mathrm{tr}\, a = 2 \cos \frac{2\pi}{k} \in \mathbb{Z}$;*
3) $k = 1, 2, 3, 4, 6,$
4) $\phi(k) \leqslant 2$.

Recall that a subgroup G of a direct product of groups $G_1 \times G_2$ is a *subdirect* product of G_1, G_2 if each projection $G \to G_i$, $i = 1, 2$, is surjective. In this case we also say that G is a subdirect subgroup of $G_1 \times G_2$.

Theorem 2.5. *Let G be a finite subgroup in $\mathrm{Sym}\, Q$ where $\dim U = 2 = \dim V$. Then G is a subdirect product of either $C_{k_1} \times C_{k_2}$, $C_{k_1} \times D_{k_2}$, $D_{k_1} \times D_{k_2}$ (C_k: cyclic group of order k) The integers k_1, k_2 satisfy one of the following conditions*

1) $k_1, k_2 = 1, 2, 3, 4, 6;$
2) (k_1, k_2) ***is either (5,10) or (10,5)***
3) $k_1 = k_2 = 12.$
4) $k_1 = k_2 = 8.$

Proof. By [2, Theorem 3.1, § 3.1] there exists a scalar product $[x, y]$ in E such that each linear operator from G is orthogonal with respect to the new product $[x, y]$. The physical space U is G-invariant. Let

For Peer Review Only

W be an orthogonal complement of U with respect to $[x, y]$. Then W is also G-invariant. it means that there exists an invertible matrix C such that (1) holds with $A_2 = 0$. In this case G is a subgroup of a direct product $G \subseteq \nu_1(G) \times \nu_2(G)$, where ν_1, ν_2 are projections and each $\nu_i(G)$ is a finite subgroup of $\mathrm{O}(2, R)$.

According to Proposition 2.1 each $\nu_j(G)$ is either a cyclic group $\langle a_j \rangle_{k_j}$ generated by the matrix a_j of order k_j from (2) or a dihedral group D_{k_j} generated by matrices a_j, b from (2). A subgroup $G \cap \langle a_j \rangle$ of a cyclic group $\langle a_j \rangle$ is cyclic generated by the element $a_j^{l_j}$ where l_j is the least exponent of elements from $G \cap \langle a_j \rangle$. Note that if, say $k_1 = 1$, then $G = G_2$ and applying Proposition 2.4 we deduce the statement of the theorem. Hence in what follows we can assume that $k_1, k_2 > 1$. Let $d = (k_1, k_2)$ be the greatest common divisor of k_1 and k_2.

Without loss of generality we can assume that ν_1, ν_2 are surjective that is G is a subdirect subgroup of $\nu_1(G) \times \nu_2(G)$ and one of the required cases takes place.

Lemma 2.6. $\phi(k_1)$, $\phi(k_2) \leqslant 4$.

Proof. Since G is a subdirect subgroup it contains a matrix

$$F = \begin{pmatrix} a_1 & 0 \\ 0 & b_2^i a_2^j \end{pmatrix}, \quad j \in \mathbb{Z},\ i = 0, 1.$$

where $a_1, b_2^i a_2^j$ are from (2), (3). Since F is conjugate in $\mathrm{GL}(4, \mathbb{R})$ to an integer matrix the characteristic polynomial $\det(F - tE)$ has integer coefficients.

Let $\zeta = \exp \frac{2\pi i}{k_1} \in \mathbb{C}$. Then ζ is a root of the integer polynomial $\det(F - tE)$ of degree 4 because

$$\det(F - tE) = \det(a_1 - tE) \det(b_2^i a_2^j - tE).$$

The minimal integer polynomial of $\exp \frac{2\pi i}{k_1}$ is the cyclotomic integral polynomial $\Phi_{k_1}(t)$ of degree $\phi(k_1)$. Hence $\Phi_{k_1}(t)$ divides $\det(F - tE)$ and therefore $\phi(k_1) \leqslant 4$. The case of k_2 is similar. □

Lemma 2.7. *If G contains a matrix*

$$F = \begin{pmatrix} a_1 & 0 \\ 0 & b_2 a_1^j \end{pmatrix}, \quad j \in \mathbb{Z}, \tag{4}$$

then $\phi(k_1) \leqslant 2$. Similarly if

$$F_1 = \begin{pmatrix} b_1 a_1^j & 0 \\ 0 & a_1 \end{pmatrix} \in G, \quad j \in \mathbb{Z},$$

then $\phi(k_2) \leqslant 2$.

Proof. Note that $\operatorname{tr} F = \operatorname{tr} a_1 + \operatorname{tr}(b_2 a_1^j) = \operatorname{tr} a_1 \in \mathbb{Z}$ by (3). Apply Proposition 2.4. □

Corollary 2.8. *One of the cases takes place:*

For Peer Review Only

1) $\phi(k_1)$, $\phi(k_2) \leqslant 2$;
2) $\phi(k_1) = \phi(k_2) = 4$.

Proof. Suppose that $\phi(k_2) \leqslant 2$. If G contains a matrix F_1 then we apply Lemma 2.7. Suppose that G contains a matrix

$$B = \begin{pmatrix} a_1 & 0 \\ 0 & a_2^{s_2} \end{pmatrix} = \begin{pmatrix} \cos\frac{2\pi}{k_1} & -\sin\frac{2\pi}{k_1} & 0 & 0 \\ \sin\frac{2\pi}{k_1} & \cos\frac{2\pi}{k_1} & 0 & \\ 0 & 0 & \cos\frac{2\pi s_2}{k_2} & -\sin\frac{2\pi s_2}{k_2} \\ 0 & 0 & \sin\frac{2\pi s_2}{k_2} & \cos\frac{2\pi s_2}{k_2} \end{pmatrix} \in G \subseteq G_1 \times G_2. \quad (5)$$

Then $\operatorname{tr} B = 2\left(\cos\frac{2\pi}{k_1} + \cos\frac{2\pi s_2}{k_2}\right) \in \mathbb{Z}$. By the assumption $2\cos\frac{2\pi}{k_1} \in \mathbb{Z}$. Apply Proposition 2.4. □

Starting from now we shall assume that $\phi(k_1) = \phi(k_2) = 4$. By Proposition 2.3 it means that $k_1, k_2 = 10, 5, 12, 8$ and by Lemma 2.7 the group G contains the matrix B a nd a matrix

$$B_2 = \begin{pmatrix} a_1^{s_1} & 0 \\ 0 & a_2 \end{pmatrix}. \quad (6)$$

Let $\xi_n = \exp\frac{2\pi i}{n} \in \mathbb{C}$. As in the proof of Lemma 2.6 we know that $\Phi_4(t) = \det(B - tE)$. Hence the roots of $\det(B - tE)$ are $\xi_{k_1}^r$, where $1 \leqslant r < k_1$ and $(r, k_1) = 1$. Similarly the roots of $\det(B - tE)$ are $\xi_{k_2}^{s_2 t}$ where $1 \leqslant t < \frac{k_2}{(s,k_2)}$ and $(t, \frac{k_2}{(s,k_2)}) = 1$ Hence $\xi_{k_2}^{s_2} = \xi_{k_1}^r$, $\xi_{k_1} = \xi_{k_2}^{s_2 t}$. In particular $\mathbb{Z}[\xi_{k_1}] = \mathbb{Z}[\xi_{k_2}^{s_2}] \subseteq \mathbb{Z}[\xi_{k_2}]$. Similarly $\mathbb{Z}[\xi_{k_2}] = \mathbb{Z}[\xi_{k_1}^{s_1}] \subseteq \mathbb{Z}[\xi_{k_1}]$. Hence

$$\mathbb{Z}[\xi_{k_1}] = \mathbb{Z}[\xi_{k_1}^{s_1}] = \mathbb{Z}[\xi_{k_2}] = \mathbb{Z}[\xi_{k_2}^{s_2}]. \quad (7)$$

Lemma 2.9. *The following cases take place*

a) k_1, $k_2 = 10, 5$ *and* $s_1 s_2 \equiv 1 \mod 5$;
b) $k_1 = k_2 = 12$ *and* $s_1, s_2 = 1, 5, 7, 11$;
c) $k_1 = k_2 = 8$ *and* $s_1, s_2 = 1, 3, 5, 7$.

Proof. Recall that $\mathbb{Z}[\xi_{10}] = \mathbb{Z}[\xi_5]$ and

$$\mathbb{Z}[\xi_{10}] \neq \mathbb{Z}[\xi_{12}] \neq \mathbb{Z}[\xi_8] \neq \mathbb{Z}[\xi_{10}].$$

Apply (7). Note also that $a_i^{s_1 s_2 - 1} \in G$ by Proposition 2.2. Hence by Proposition 2.4 $s_1 s_2 - 1$ in the first case should be divisible by 5. □

By Lemma 2.7 we complete the proof of Theorem 2.5. □

In order to present more detailed classification of groups in Theorem 2.5 we need

For Peer Review Only

Proposition 2.10. *Let G, k_1, k_2 be from Theorem 2.5 and $\phi(k_1) = \phi(k_2) = 4$. Suppose that the group G contains a matrix*

$$F = \begin{pmatrix} a_1^t & 0 \\ 0 & b_2 \end{pmatrix}.$$

Then G contains

$$a_1^2 = \begin{pmatrix} a_1^2 & 0 \\ 0 & 1 \end{pmatrix}$$

and a matrix F with $t = 0$, that is b_2. In this case $G \cap (\langle a_1 \rangle \times D_{k_2})$ is a semidirect product of a normal subgroup $\langle B \rangle \times \langle a_2^2 \rangle$ by a cyclic group $\langle F \rangle$ of order 2 with $t = 0$, where B is from (5) with $s_2 = 1$.

Proof. We know that G contains a matrix B from (5). Then

$$U = BFBF^{-1} = a_1^2 \in G.$$

Multiplying F and B_2 from (6) by powers of U we can assume that $t = 0, 1$ and $s_1 = 1$. The case $t = 1$ is impossible because otherwise $\operatorname{tr} a_1^2 \in \mathbb{Z}$ which contradicts Proposition 2.4. Thus we can assume that $B_2 = B$ and $t = 0$.

Note that $G_1 = \langle B \rangle \times \langle a_2^2 \rangle$ is a subgroup in G containing a_1^2 and therefore a_2^2. Hence the index of G_1 in $H = \langle a_1 \rangle \times \langle a_2 \rangle$ is dividing 4. Since $a_1, a_2 \notin G$ we can conclude that $G_1 = G \cap H$. Now the proof follows. □

Theorem 2.11. *Let a group G be a subdirect product from Theorem 2.5 where $k_1 = k_2 = 10$. Then G is one of the following groups.*

a) *If G is a subdirect product of two cyclic groups $\langle a_1 \rangle_{10} \times \langle a_2 \rangle_{10}$ then G is direct product of two cyclic groups*

$$\langle B \rangle \times \langle a_1^l \rangle \tag{8}$$

where B is from (5) with $1 \leqslant s_2 \leqslant 9$, $s_2 \neq 5$ and $l = 0, 5$. If s_2 is even then $l = 5$.

b) *G is a subdirect subgroup of $D_{10} \times D_{10}$ and a semidirect product of the normal group (8) by a cyclic group $\langle W \rangle$ of order 2, where*

$$W = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 a_2^j \end{pmatrix}, \quad j = 0, \ldots, 9. \tag{9}$$

Proof. Case **a)**. We know that some matrices B, B_2 from (5), (6) belong to G where s_i are from Lemma 2.9, case a). The group G contains the cyclic group $\langle B \rangle$ of order 10 which projects onto $\langle a_1 \rangle_{10}$. Thus $G = \langle B \rangle \times (G \cap \langle a_2 \rangle_{10})$. But by Proposition 2.4 the order of the group $G \cap \langle a_2 \rangle_{10}$ is equal to 1,2,3,4 or 6. Hence $G \cap \langle a_2 \rangle_{10} = \langle a_1^l \rangle$ where $l = 0$ or 5. Suppose that s_2 is even and $l = 0$. Then the cyclic group $\langle B \rangle$ does not contain B_2 which is impossible.

Suppose that G is a subdirect product of $\langle a_1 \rangle_{10} \times D_{10}$. Then $G \cap \langle a_1 \rangle_{10} \times \langle a_2 \rangle_{10}$ is a group (8). Also G contains the matrix F from

For Peer Review Only

Proposition 2.10 which contradicts Proposition 2.4. So this case is impossible.

Case **b)**. Using previous argument we can show that G does not contain the matrix F from Proposition 2.10. Thus $G \cap (\langle a_1 \rangle_{10} \times \langle a_2 \rangle_{10})$ is a subdirect product and therefore as in the case **a)** it has the form (8). As above G does not contain the matrix F from Proposition 2.10. Hence G contains a matrix $W \in G$ as in (9). Then $WgW^{-1} = g^{-1}$ for all $g \in \langle a_1 \rangle_{10} \times \langle a_2 \rangle_{10}$ and $W^2 = E$. Thus G has the required form. □

Theorem 2.12. *Let $k_1 = 10$, $k_2 = 5$. Then there are only two cases.*

a) *If G is a subdirect product $\langle a_1 \rangle_{10} \times \langle a_2 \rangle_5$ then $G = \langle B \rangle$ where $s_2 = 1, 2, 3, 4$.*

b) *If G is a subdirect product of $D_{10} \times D_5$ then G is generated by B and by W from (9), where $j = 0, 1, 2, 3, 4$. Thus $G \simeq D_{10}$.*

Proof. The group G contains a matrix B from (5) where s_2 is invertible modulo 5 by Lemma 2.9. In the case **a)** either G is generated by B or G has a nontrivial intersection with $\langle a_2 \rangle$ which is impossible by Proposition 2.4.

The case when G is a subdirect product $\langle a_1 \rangle_{10} \times D_5$ is impossible by Proposition 2.10.

The proof in the case **b)** is similar to those in the previous Theorem. □

The proof of the next Theorem is similar.

Theorem 2.13. *Let $k_1 = 5$, $k_2 = 10$. Then there are only two cases.*

a) *If G is a subdirect product $\langle a_1 \rangle_5 \times \langle a_2 \rangle_{10}$ then $G = \langle B_2 \rangle$ from (6) where $s_1 = 1, 2, 3, 4$.*

b) *If G is a subdirect product of $D_5 \times D_{10}$ then G is generated by B_2 and by W from (9). Thus $G \simeq D_{10}$.*

Theorem 2.14. *Let group G be a subdirect product from Theorem 2.5, where $k_1 = k_2 = 8$.*

A) *If G is a subdirect product of cyclic groups $\langle a_1 \rangle_8 \times \langle a_2 \rangle_8$, then G is a direct product of two cyclic groups (8), where $l = 0, 2, 4$, and s_2 being odd in B from (5).*

B) *Suppose that G is a subdirect product of $\langle a_1 \rangle \times D_8$. Then G is a semidirect product of the normal subgroup (8) and a cyclic group $\langle b_2 \rangle$ of order 2, where $l = 0, 2, 4$, and s_2 being odd in B from (5).*

C) *Suppose that G is a subdirect subgroup of $D_8 \times D_8$. Then G is one of the groups*

- **a)** *a semidirect product of a normal subgroup (8) with $l = 2$ and s_2 odd by a direct product $\langle b_1 \rangle_2 \times \langle b_2 \rangle_2$.*
- **b)** *G is semidirect product of the normal subgroup (8) and a cyclic group $\langle W \rangle_2$, where $l = 0, 2, 4$, s_2 being odd in B from (5) and W is from (9).*

For Peer Review Only

Proof. In the case **A)** the group G contains a matrix B from (5) with odd s_2. If $G \neq \langle B \rangle$ then $G = \langle B \rangle \times (G \cap \langle a_2 \rangle)$, where $G \cap \langle a_2 \rangle = \langle a_2^l \rangle$ and $l = 0,\ 2,\ 4$ by Proposition 2.4.

The case **B)** follows from Proposition 2.10.

In the case **C)** by Lemma 2.9 the group G contains a matrix B from (5) with odd s_2. Hence $G \cap (\langle a_1 \rangle \times \langle a_2 \rangle)$ is a subdirect product from the case **A)**. If F from Proposition 2.10 belongs to G then $t = 0$ and

$$G \cap (\langle a_1 \rangle \times \langle a_2 \rangle) = \langle B \rangle \times \langle a_2^2 \rangle .$$

Symmetrically G contains a matrix

$$b_1 = \begin{pmatrix} b_1 & 0 \\ 0 & 1 \end{pmatrix} .$$

Hence G has the form Ca).

Suppose that $U, U' \notin G$. Then G contains a matrix W from (9) where $j = 0, \ldots, 8$. Also

$$G \cap (\langle a_1 \rangle \times \langle a_2 \rangle) = \langle B \rangle \times \langle a_2^l \rangle , \quad l = 0,\ 2,\ 4.$$

Then G has the form **Cb)**. □

Note that the case B) with $l = 0$ and $s_2 = 3$ corresponds to perfect 8-fold dihedral symmetry in a certain member of the LI-class of Ammann-Beenker tilings [4]**. The generating matrices of the group are**

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & -1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} .$$

Using the same argument one can prove

Theorem 2.15. *Let $k_1 = k_2 = 12$. Then G is one of the following groups.*

A) *If G is subdirect product of $\langle a_1 \rangle_{12} \times \langle a_2 \rangle_{12}$ then G is the group*

$$G = \langle B \rangle \times \langle a_2^l \rangle \tag{10}$$

where B is from (5) ***with*** *$s_2 = 1, 5, 7, 11$ and $l = 0,\ 2,\ 3,\ 4,\ 6$.*

B) *If G is a subdirect product of $\langle a_1 \rangle_{12} \times D_{12}$ then G is a semidirect product of the normal subgroup* (10) *with $l = 2$ and $\langle b_2 \rangle_2$.*

C) *If G is a subdirect product of $D_{12} \times D_{12}$ then G is one of groups:*

- **a)** *a semidirect product of a normal subgroup* (10) *with $l = 2$ and s_2 odd by a direct product $\langle b_1 \rangle_2 \times \langle b_2 \rangle_2$.*
- **b)** *a semidirect product of the normal subgroup* (10) *and a cyclic group $\langle W \rangle_2$, where $l = 0,\ 2,\ 4$, $s_2 = 1,\ 5, 7,\ 11$ in B from* (5) *and where W is from* (9).

For Peer Review Only

Applying previous results to the window W which is a closure of a projection of A into V we obtain

Corollary 2.16. *The symmetry group of the window of any plane module set with* $2\dim$ *internal space is one of the following:* C_i *or* D_i, $i = 1, 2, 3, 4, 5, 6, 8, 10, 12$.

3. Realization

In this section we shall show that two exceptional cases 2), 3) in Theorem 2.5 can be realized. The idea of this realization comes from [10], [7], [4]. Let $m \geqslant 3$ be a positive integer. Take the real vector space $E = \mathbb{R} \otimes_{\mathbb{Z}} \mathbb{Z}[\xi]$ of dimension $\phi(m)$ where $\xi = \exp\left(\frac{2\pi i}{m}\right)$. Then **basis** of E consists of $\phi(m)$ vectors $e_j = 1 \otimes \xi^j$, $0 \leqslant j < \phi(m)$, [6, Chapter IV, § 1]. Define in in E two linear operators

$$a(1 \otimes \xi^j) = 1 \otimes \xi^{j+1}, \quad b(1 \otimes \xi^j) = 1 \otimes \xi^{-j}$$

for all $j \in \mathbb{Z}$. These definitions are correct since $\mathbb{Z}[\xi]$ is a left $\mathbb{Z}[\xi]$-module and the map b is a Galois automorphism of $\mathbb{Z}[\xi]$. It is easy to see that $b^2 = (ba)^2 = 1$. Hence the group of invertible operators generated by a, b is the dihedral group D_m and we have a representation of D_m in E of dimension $\phi(m)$. It is decomposed into direct sum of irreducible ones. Since dimensions of irreducible representations of dihedral groups are 1 or 2 [2, chapter 3], it suffices to show that ± 1 are not eigenvalues of operators a and b. Note that the characteristic polynomial of the operator a is equal to $\Phi_m(t)$, and **in fact** ± 1 are not roots of $\Phi_m(t)$, provided $m \geqslant 3$.

We have proved

Theorem 3.1. *Let* $m \geqslant 3$ *be an integer. Then the dihedral group* D_m *is a subgroup of a symmetry group* $\operatorname{Sym} Q$ *of some 2D-quasicrystal* Q *which can be constructed by a cut and project method using a hyperspace* E *of dimension* $\phi(m) \geqslant 2$.

There is another way of constructing symmetries of quasicrystals based on algebraic integers. Let $\xi = \exp\left(\frac{2\pi i}{m}\right)$ be as above. Then again $\mathbb{Z}[\xi]$ is a left $\mathbb{Z}[\xi]$-module and therefore the Abelian group $\mathbb{Z}[\xi]^*$ of invertible elements of the ring $\mathbb{Z}[\xi]$ acts on $E = \mathbb{R} \otimes_{\mathbb{Z}} \mathbb{Z}[\xi]$. By [2, Theorem 3.3, Chapter3] the complex space $\mathbb{C} \otimes_{\mathbb{R}} E$ contains one-dimensional subspace which is invariant under the action of the **Abelian** group $\mathbb{Z}[\xi]^*$. Hence E contains 2-dimensional real invariant subspace U. So taking U as a physical space and the lattice $\mathbb{Z}[\xi]$ in E as M we can construct a quasicrystal Q in which the symmetry group $\operatorname{Sym} Q$ contains $\mathbb{Z}[\xi]^*$ as a subgroup. By [3, page 561] the Abelian group $\mathbb{Z}[\xi]^*$ has rank $\frac{\phi(m)}{2} - 1$. Applying Proposition 2.3 as in [7] we obtain

Theorem 3.2. *Let* m *be an integer and either* $m = 5$ *or* $m \geqslant 7$. *Consider 2D-quasicrystal* Q *constructed by a cut and project method*

For Peer Review Only

using a hyperspace E of dimension $\phi(m) \geqslant 2$ and $M = \mathbb{Z}[\xi]$ as a lattice. The symmetry group $\operatorname{Sym} Q$ *contains an Abelian group isomorphic to* $\mathbb{Z}[\xi]^*$ *and having rank* $\frac{\phi(m)}{2} - 1 \geqslant 1$. *In particular* $\operatorname{Sym} Q$ *contains elements of infinite order.*

For example take $E = \mathbb{R} \otimes \mathbb{Z}[\xi]$, $\xi = \frac{1+i}{\sqrt{2}}$. Then $\dim E = \phi(8) = 4$ and E is a left module over $\mathbb{Z}[\xi]$. Therefore the group $\mathbb{Z}[\xi]^*$ of invertible elements of the ring $\mathbb{Z}[\xi]$ acts on E. The rank of the group $\mathbb{Z}[\xi]^*$ is equal to $\frac{\phi(8)}{2} - 1 = 1$ [3, page 561] and therefore it contains an element g of infinite order.

References

[1] Artamonov V.A., On symmetries of quasicrystals. Contemp. Math. v. 376. Algebraic Structures and Their Representations: XV Colloquium Latinoamericano de Álgebra, Cocoyoc, Morelos, Mexico, July 20-26, 2003, Jose A. de la Pena, Ernesto Vallejo, and Natig Atakishiyev ed., AMS, 2005, 175-188.

[2] Artamonov V.A., Slovokhotov Yu.L., Groups and their applications in physics, chemistry and crystallography. Publ. center Academia, Moscow, 2005, P.512.

[3] Hasse H. Number theory, Springer-Verlag Berlin, Heidelberg, New York, 1980.

[4] Hermisson J., Richard Ch., Baake V., A Guide to the symmetry structure of quasiperiodic tiling classes, J. Phys. I France, 7 (1997), 1003-1018.

[5] Baake M., Hermisson J., Pleasants A.B., The torus parametrization of quasiperiodic LI-classes, J. Phys A, 30 (1997), 3029-3056.

[6] Lang S. Algebraic number theory. Addison-Wesley Publ. Company, 1970, 354.

[7] **Niizeki K., Self-similarity of quasilattices in two dimensions. I The n-gonal quasilattice. Journal of Physics A: Mathematical and General. 22(1989), 193-204.**

[8] Skornyakov L.A. Ed., General algebra, v.1, Moscow: Nauka, Fizmatlit, 1990.

[9] **Moody R.V. Model sets: a survey. In the book: From quasicrystals to more Complex Systems. Springer, Berlin, pp. 145-166, arXiv:math.MG/0002020 v1**

[10] **Pleasants P.A.B., Designer quasicrystals: cut-and-project sets with pre-assigned properties. In the book: Directions in Mathematical Quasicrystals (M. Baake and R. V. Moody, eds.), CRM Monograph Series, AMS, Providence, Rhode Island, 93-138.**

Department of Algebra, Faculty of Mechanics and Mathematics, Moscow State University

Department of Applied Mathematics, University Rey Juan Carlos, Madrid

E-mail address: artamon@mech.math.msu.su, sergio.sanchez@urjc.es

For Peer Review Only