



**HAL**  
open science

## **Toxoplasmosis in Rodents: Ecological Survey and First Evidences in Thailand.**

Sathaporn Jittapalapong, Nachai Sarataphan, Soichi Maruyama, Jean-Pierre Hugot, Serge Morand, Vincent Herbreteau

► **To cite this version:**

Sathaporn Jittapalapong, Nachai Sarataphan, Soichi Maruyama, Jean-Pierre Hugot, Serge Morand, et al.. Toxoplasmosis in Rodents: Ecological Survey and First Evidences in Thailand.. Vector-Borne and Zoonotic Diseases, 2010, 11 (3), pp.231-237. 10.1089/vbz.2009.0238 . hal-00545606v1

**HAL Id: hal-00545606**

**<https://hal.science/hal-00545606v1>**

Submitted on 4 Jul 2012 (v1), last revised 4 Jul 2012 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Toxoplasmosis in rodents: ecological survey and first evidences in Thailand**

Sathaporn Jittapalapong,<sup>1</sup> Nachai Sarataphan,<sup>2</sup> Soichi Maruyama,<sup>3</sup> Jean-Pierre Hugot,<sup>4</sup> Serge Morand,<sup>5,6</sup> and Vincent Herbreteau,<sup>6,7</sup>

<sup>1</sup> Department of Parasitology, Faculty of Veterinary Medicine, Kasetsart University, Bangkok, Thailand.

<sup>2</sup> Bureau of Biotechnology for Livestock Production, Department of Livestock Development, Ministry of Agriculture and Cooperatives, Pathumthani, Thailand.

<sup>3</sup> Laboratory of Veterinary Public Health, Department of Veterinary Medicine, College of Bioresource Sciences, Nihon University, Kanagawa, Japan.

<sup>4</sup> Muséum National d'Histoire Naturelle, OSEB, UMR 7205, Paris, France.

<sup>5</sup> CNRS-UM2, Institut des Sciences de l'Evolution (ISEM), Université Montpellier 2, Montpellier, France.

<sup>6</sup> CIRAD, UR AGIRs (Animal et Gestion Intégrée des Risques), Campus International de Baillarguet, Montpellier, France.

<sup>7</sup> IRD Research Unit 178, Center for Vectors and Vector-borne Diseases, Faculty of Sciences, Mahidol University, Bangkok, Thailand.

Corresponding Author: Vincent Herbreteau

Email: vincent.herbreteau@teledetection.fr

Address: CIRAD, UR AGIRs, Maison de la télédétection, 500, rue Jean-François Breton, 34093 Montpellier Cedex 5, France

Running title: Toxoplasmosis of rodents in Thailand

Key words: toxoplasmosis, *T. gondii*, rodent, Murinae, *Rattus*, ecology

## **Abstract**

Domestic and wild rodents known as the most abundant and diversified order of mammals have a key role in the ecological food chain and also in the transmission of parasites and pathogens to other animals. While foraging on the ground, they can get infected by *Toxoplasma gondii*, a protozoan parasite, which is the causative agent of toxoplasmosis. Therefore they serve as intermediate hosts of *T. gondii* that they can transmit to their predators. To assess their role in the maintenance of *T. gondii* lifecycle in Thailand, we sampled rodents in a range of biotopes representative of the high biodiversity and conducted serological survey with latex agglutination test (LAT) to detect anti-*T. gondii* antibodies. Overall 21 of 461 (4.6%) rodents had diagnostically significant antibody titers (cut-off, 1:64). Every species with at least 37 individuals captured tested positive, confirming the wide range of potential mammalian hosts of toxoplasmosis. None of the ecological traits (sex, maturity, morphology, season or habitat) was found significant to predict the susceptibility to *T. gondii* both univariately and in a multivariate analysis. However high prevalences were reported either in forested or anthropized areas. This survey constitutes the first confirmed serological investigation of *T. gondii* in rodents in Thailand. The rarity of both domestic and wild felids in Thailand emphasizes the importance of rodents in maintaining *T. gondii*, and questions the involvement of other carnivores in the life cycle.

## **Introduction**

More than one hundred years ago, a protozoan parasite *Toxoplasma gondii* was described from an African rodent, *Ctenodactylus gundi* (Nicolle and Manceaux 1908), and simultaneously from a rabbit in Brasil (Splendore 1908). *T. gondii* is the unique species in the genus *Toxoplasma* of the Coccidian family and has the widest host range and distribution, being potentially carried by any warm-blooded animal (mammals and birds) with cats as definitive hosts (Dubey 2008, Tenter, et al. 2000). As a consequence, toxoplasmosis is one of the most common parasitic disease worldwide, probably infecting about one third of humans (Tenter, et al. 2000, Kim and Weiss 2008). *T. gondii* is mainly transmitted congenitally or by ingestion either of bradizoites in infected tissues, or of oocysts spread in the environment. The risk of infection by ingestion for rodents is related to the complexity of environmental factors that determine the presence of other mammals, especially felids, and oocysts. Climate also conditions the dispersal, density and survival of oocysts, since their resistance depends on humidity and temperature (Dubey and Frenkel 1998).

Researches on rodents have been limited even if they might play a significant role in the epidemiology of toxoplasmosis, by being potential intermediate hosts before the transmission to pigs, dogs or cats (Dubey and Frenkel 1998). A major review of this work was realized by Dubey and Frenkel in 1998, who analyzed investigations of *T. gondii* in rats worldwide and proposed interesting information about the implication of rats in the life cycle of this parasite. The authors report a low average prevalence of antibodies to *T. gondii* in rats, generally under 5–10%, and also correctly point out the difficulties to compare various serological screenings conducted with different methods. Nevertheless, we considered this review to be a fundamental source of

information about rodent species also occurring in Thailand, according to the reference book Mammal species of the world (Carleton and Musser 2005).

Furthermore, the implication of rodents in the life cycle of *T. gondii* is still misunderstood while considering the transmission between different hosts, as well as the influence of ecological factors (Tenter, et al. 2000, Kim and Weiss 2008). Few studies have considered how ecological traits of the animal host can influence the seroprevalence of toxoplasmosis. In the Philippines, Salibay and Claveria (2005) found no relationship between *T. gondii* prevalence and rodent species (*Rattus norvegicus* and *R. rattus mindanensis*) and no correlation between the prevalences in their three collection sites. They also reported a higher but not statistically significant seroprevalence in males (Salibay and Claveria 2005). In USA, Dabritz *et al.* (2008) found no relationship between *T. gondii* prevalence and the association of rodent habitat with water. Nevertheless, they could associate a greater distance to houses (>200 meters) and the maturity (adult) with a higher prevalence (Dabritz, et al. 2008). In French Guyana, de Thoisy *et al.* found that terrestrial mammals are significantly more exposed to *T. gondii* than other mammals, and noticed discordant results for the agouti and the acouchy, because of their feeding behaviour (they are granivorous and highly selective foragers). They finally recommended further analysis regarding the influence of habitat and behavior (de Thoisy, et al. 2003).

In Thailand, whereas toxoplasmosis has never been investigated in rodents, recent studies have shown widespread cases of infection in humans (Maruyama, et al. 2000), dogs and cats (Jittapalpong, et al. 2007), swine (Thiptara, et al. 2006), goats (Jittapalpong, et al. 2005), elephants (Tuntasuvan, et al. 2001) and tigers (Thiangtum, et al. 2006). Therefore, this study was first designed to determine the presence and prevalence of *T. gondii* in wild rodents in Thailand, then to enlighten from an

ecological point of view their importance in the transmission of this parasite either to other animals or to humans.

## **Materials and Methods**

### *Study sites and animal trapping*

From 2003 to 2006, rodents were captured in thirteen provinces throughout Thailand for studying their implication in the transmission of different pathogens to humans (Figure 1). Trapping locations were chosen to sample animals in a range of biotopes representative of the high biodiversity peculiar to Thailand and South-East Asia. In each site, twenty locally-made cages were used during five nights to catch animals alive. This field work was repeated during the dry and wet seasons. Geographical coordinates of the trapping places were systematically recorded. Trapping and processing were conducted according to established safety recommendations (Mills, et al. 1995). Animals were collected early in the morning and transferred to a field laboratory, where they were dissected to collect organs for different studies of rodent-borne pathogens. Blood was collected by cardiac puncture and later separated by centrifugation. Serum samples were stored in liquid nitrogen then kept in -80°C freezer before testing. Species were identified based on morphological criteria following regional taxonomic identification keys (Marshall 1988, Askins 1988, Aplin, et al. 2003, Corbet and Hill 1992).

### *Serological assays*

*T. gondii* antibodies were detected using a latex agglutination test (LAT) kit (Toxocheck-MT, Eiken Chemical Co., Ltd. Japan). This test has been evaluated as a suitable screening serological method to assess the presence of toxoplasmosis in all

species of animals with a high sensitivity and specificity (Tilahun, et al. 1998, Mazumder, et al. 1988). The manufacturer's protocol with a cut-off titer at 1:64 was respected: 25  $\mu$ l latex agglutination buffer was first introduced to each well of a U-shaped 96 wells microtitration plate; sera were diluted serially starting with 1:8 dilution then twofold dilutions and the final 25  $\mu$ l discarded; then 25  $\mu$ l of *T. gondii*-antigen-coated latex beads were added to each well. The plate was incubated at room temperature overnight. Antibody titers were determined by the last dilution where agglutination pattern could be read in comparison with the positive control.

### *Statistical analyses*

During rodent trapping and processing were collected a set of variables that could possibly explain the infection by *T. gondii*, and cause intra or interspecific differences in prevalences. These variables are phylogeny (at family, genus or species levels), sex, maturity (determined after external and internal observation of sexual organs), season of trapping (dry or wet), morphology and habitat. Several standard body measurements were taken in the field laboratory: weight, head and body length, tail length, hind foot length, ear length and skull length. Whereas a precise environmental description of each trapping site was recorded in the field, the variable habitat refers here to a global classification of the landscape from wild to anthropized biotopes, coded as follows: forests (1), fields at the edge of forests (2), dry fields (3), wet ricefields (4) and domestic areas (5) (in proximity to human settlements, in houses, villages or cities). Each variable was first analyzed univariately to measure the relation with *T. gondii* prevalence through Spearman correlations for continuous variables and Fischer's exact test (or Pearson  $\chi^2$  tests where appropriate) for proportions of categorical ones. Then we tried to model the relationship between *T.*

*gondii* seroprevalence and these variables using generalized linear and non-linear models with prevalence as the dependent variable; habitat, season, sex, and maturity as categorical variables and body measurements as continuous variables. These models with binomial error and a logistic link function have proved to be appropriate statistical tool to assess the influence of explanatory variables on infection (Crawley 1993).

## Results

461 rodents were collected during the field study, including 408 rats and mice (Muridae) and 53 squirrels (Sciuridae) with a total of 18 species (Table 1). Only two mice were trapped because of the large size of the cages used, which are not suitable for small-sized rodents. Most of the rodent species occurring in Thailand can be easily caught, whether they live in wild habitats or close to human settlements.

The overall prevalence of *T. gondii* infection in rodents was 4.6% (21 positive out of 461 tested). The number of seropositive animals for each species is significantly correlated with the number of animals tested (0.901,  $p < 0.05$ ). All species with more than 37 animals tested were found positive for at least one individual (Table 1): *Rattus tanezumi* (7.1%, 11/156), *Rattus exulans* (1.3%, 1/79), *Menetes berdmorei* (7.7%, 4/52), *Maxomys surifer* (5.3%, 2/38) and *Bandicota indica* (2.7%, 1/37). This attests the wide spectrum of potential mammalian hosts of toxoplasmosis. In addition two *Leopoldamys sabanus* individuals were found positive from sixteen captured (12.5%), which is the highest prevalence in rats. All *Rattus norvegicus* individuals tested negative for *T. gondii* antibodies, even if a relatively high number of individuals (34) were screened. No positives were found from the eleven other species that were captured in low numbers, such as *Bandicota savilei*, *Berylmys bowersi*, *Berylmys*



*berdmorei*, *Leopoldamys edwardsi*, *Niviventer fulvescens*, *Rattus andamanensis*, *R. losea*, *R. tiomanicus*, *Mus cervicolor*, *M. cookii* and *Callosciurus sp.* For these species, the number of animals tested is not sufficient to conclude that they might not get involved in the transmission of *T. gondii*. Among positive animals, the highest titer (1:512) was found in *Bandicota indica* and *Rattus tanezumi*.

The seroprevalence was higher in Sciuridae (7.5%) than in Muridae (4.2%) but not significantly different (two-tailed Fisher's exact test,  $p=0.285$ ,  $p>0.05$ ). Rodent females also presented a higher seroprevalence (5.7%) than males (3.3%) but not significantly different (two-tailed Fisher's exact test,  $p=0.265$ ,  $p>0.05$ ). This was also verified for the 156 *Rattus tanezumi* tested (8.3% for females and 5.7% for males) without statistical significance ( $p=0.755$ ). Moreover, even if the prevalence of adult rodents (4.7%) was nearly twice the prevalence of young rodents (2.6%), the effect of maturity on infection could not be statistically attested (two-tailed Fisher's exact test,  $p=1$ ,  $p>0.05$ ). Finally, there was no relation between seroprevalences and body measurements either standardized or not.

According to the habitat, the number of seropositive animals is significantly correlated to the number of animals tested (0.979,  $p<0.05$ ). However, according to the level of anthropization of each habitat, the seroprevalence is not correlated to the anthropization index (Figure 2). The highest seroprevalences are either observed in the less anthropized areas (forests: 6.2%, 8/130) or the highly anthropized ones, near human settlements (domestic areas: 4.5%, 12/269).

From the multivariate analysis, only few models could be considered to estimate the presence or absence of infection. The best model (logit model) with two effects is obtained with the log-transformed weight and a three-way interaction between season, sex and maturity ( $p<0.01$ ). A one-effect model with the three-way interaction between

season, sex and maturity is also relevant to predict infection ( $p < 0.005$ ). However, only categorical variables (maturity, sex and season) entered at  $p < 0.001$ , while all continuous variables did not enter the model. Therefore the model obtained is not robust and presents high deviance corresponding to overdispersion (residual deviance / residual df = 0.37).

## **Discussion**

To compare our results with other investigations of *T. gondii* in rodents, we analyzed the review realized by Dubey and Frenkel in 1998 and completed available information with more recent papers (Supplemental data). In this review, only three species have been described in Thailand, namely *Rattus norvegicus* (the urban brown rat), *R. rattus* (the black rat) and *R. exulans* (the Polynesian rat). *R. norvegicus* and *R. rattus* have a worldwide distribution and have been frequently studied in several countries, while there is less information on *R. exulans* present from mainland Asia to New Guinea and the Pacific region (Marshall 1988, Aplin, et al. 2003). We could find out further research papers investigating the presence of *T. gondii* in rats also known to occur in Thailand. Four studies have tested *R. norvegicus* (Salibay and Claveria 2005, Lubroth, et al. 1983, Dubey, et al. 2006, Kijlstra, et al. 2008) and three studies have examined a fourth species, *Mus musculus*, the house mouse (Lubroth, et al. 1983, Kijlstra, et al. 2008, Murphy, et al. 2008). Most of these studies have detected the presence of antibodies to *T. gondii* using various serological tests or *T. gondii* DNA by molecular method (Polymerase Chain Reaction, PCR). The use of different techniques makes a global comparison of results hazardous if we do not consider their variability of specificity and sensitivity. The dye-test has been the first technique developed in 1948 and the most used (Ferguson 2009). It has proved to be highly

specific and sensitive (Dubey 2008) and was then considered as a reference to evaluate other methods. Investigations that used the dye test have shown the highest prevalences in *R. norvegicus* (12.8%, 310/2,426) then *R. rattus* (8.3%, 78/936) and *R. exulans* (3.8%, 19/505). When compiling all serological results, rates are slightly higher for every species: *R. norvegicus* (15.6%, 699/4,482), *R. rattus* (12.2%, 169/1,390) and *R. exulans* (3.8%, 19/505) and a very high prevalence was shown for *Mus musculus* (60.0%, 12/20) in a single study by indirect immuno-fluorescent test. Recent studies have used PCR in the Netherlands and the United Kingdom also demonstrating a very high prevalence in *Mus musculus* (51.9%, 120/231) and lower in *R. norvegicus* (10.3%, 4/39) (Kijlstra, et al. 2008, Murphy, et al. 2008). Despite the heterogeneity of this summarized data, results from serology and molecular diagnostic are relatively concordant.

This literature review also reveals a limited geographical range of *T. gondii* in *M. musculus*, *R. norvegicus* and *R. rattus*, despite the worldwide distribution of these three rodent species. Indeed, the map of the worldwide prevalence of *T. gondii* in *R. norvegicus* shows a biased image of the infection, with the highest prevalences in North America, Western Europe and Egypt (Figure 3). For the four species that we have selected, most of investigations have been conducted in developed countries located in the Northern hemisphere (except Australia) where toxoplasmosis is a concern for human health and considered during pregnancy. There is no data from Africa, except from Egypt (Rifaat, et al. 1971) and another regional study (de Roever-Bonnet 1972), no data from East and South-East Asia, except the Philippines (Salibay and Claveria 2005) and Japan (Dubey and Frenkel 1998), and no data from South America.

These four species known in Thailand and cited in the literature are also the most common rodent species, living close to humans and potentially depending on humans as commensal. The high *T. gondii* prevalences reported in these species reflect their implication in a transmission cycle close to humans. Contrary to other investigations on rodents, our survey with no positive *Rattus norvegicus* questions its implication in the transmission of *T. gondii* in Bangkok metropolitan area where it lives. On one hand, the high prevalence (15.5%) reported in the literature suggests that we might have missed positive animals or endemic areas. On the other hand, we can assess that *R. norvegicus* has possibly a minor role in the transmission cycle in Bangkok metropolitan area. Indeed, this species occupies different spaces than the increasing number of cats and stray dogs that are designated to be the main actors of this transmission (Jittapalapong, et al. 2007).

Global prevalences reported in the literature for *R. rattus* (12.2%) and *R. exulans* (3.8%) are slightly higher than those observed in our study for *R. tanezumi* (7.1%) and *R. exulans* (1.3%). It is surprising to find low prevalences of *T. gondii* in *R. exulans* both in the literature and in our study, since this species exclusively lives inside and around houses and can share its biotope with domestic cats, which are susceptible to shed oocysts in their feces that can infect rats. However, despite a low prevalence, *R. exulans* has an important role in the transmission cycle by maintaining parasites close to humans. *R. rattus* cited in these articles can refer to different species since it is now considered as a species group including an Oceanian or European type referred as *R. rattus* and an Asian type referred as *R. tanezumi* (Carleton and Musser 2005). Indeed the complexity of this species group taxonomy makes hazardous any comparison of our results with other investigations.

Therefore, wild species described in Thailand have never been investigated and their importance in the transmission of *T. gondii* was not considered. Even if they do not live in contact with humans, *Leopoldamys* spp., *Maxomys* spp. and *Menetes* spp. may play a major role in maintaining tachyzoites and bradizoites in forested ecosystems and increasing the probability of infection of wild and domestic felids. This consideration is reinforced by the importance of the species of these two genera in wild habitats where other mammal species have become rare under a high hunting pressure in non-protected areas in Thailand. Furthermore, it was unexpected to find such high prevalences in wild rodents, when considering the rarity of wild felids in Thai forests that may theoretically result in a low circulation of oocysts. This questions the exclusive role of felids in hosting the sexual part of the life cycle of *T. gondii*. According to the IUCN (International Union for Conservation of Nature and Natural Resources) Red List of Threatened Species, only two species of wild felids are common in Thailand: *Prionailurus bengalensis* (Leopard Cat) and *Felis chaus* (jungle cat). Other felid species are now threatened or extinct. This assessment suggests that the sexual part of the life cycle of *T. gondii* might also be hosted by other carnivores considering their diversity and relative importance in the ecosystems, even if it has never been demonstrated in any countries. Common species are limited to one mongoose (Herpestidae), *Herpestes javanicus* (small Asian mongoose), two mustelids: *Martes flavigula* (Yellow-throated Marten) and *Melogale personata* (Burmese Ferret-badger), and a few viverrids including: *Paradoxurus hermaphroditus* (Common palm civet), *Paguma larvata* (Masked palm civet), *Viverricula indica* (Small Indian Civet). Further investigations should be conducted to verify these observations.

Finally our results attest the low specificity of *T. gondii* that can potentially infect any murine rodent species, whatever their sex, age or morphology. Differences in prevalences related to the sex and maturity are not clearly observed and may vary from one species to another one according to specific social or feeding behaviour that can determine the exposure to the parasite. Nevertheless, the prevalence measured in young rodents (2.6%), which is more than half the prevalence in adults (4.7%), is relatively high when considering that youngest have less chance to get infected by free oocysts, as they are fed by their mothers. These high levels of infection may confirm a role for vertical transmission in maintaining infections within rodent populations, as it was demonstrated experimentally in *Mus musculus* and *Apodemus sylvaticus* (Owen and Trees 1998). However, it seems that the efficiency of this vertical transmission is less in wild rodents than in experimental condition using captive-bred mice, with respectively 82.7% (n=83) and 85.0% (n=207) in pups. Furthermore, high prevalences are observed in different habitats and suggest the existence of two cycles of transmission, a wild sylvatic cycle in forested habitats and a domestic cycle within human settlements, which might be rarely connected. This hypothesis should be better supported by molecular data. Rodents as well as non-rodent species may serve as such intermediate hosts and accidentally bring parasites horizontally from wild to highly anthropized biotopes or vice versa. While rodent distribution is generally delimited in space by different biotopes, a few species such as *R. tanezumi* can inhabit a variety of biotopes. We trapped this species, also known as the Asian black rat, in the roof of houses in villages, in gardens, in ricefields but also in secondary forests. If we consider its high prevalence (7.1%) compared with other species, we can conclude that *R. tanezumi* is a main actor of a horizontal transmission between biotopes.

Finally *T. gondii* prevalence in rodents is influenced by environmental conditions that determine in particular the food and shelter availability and therefore the abundance and distribution of rodents and their predators (Meerburg, et al. 2009). Indeed, rodent species have a high capacity for adaptation to rapid environmental changes. Rodents are usually the first mammals to invade deforested areas or agricultural fields, benefiting from a high reproductive potential and a short period of maturation (Aplin, et al. 2003). The deterioration of biotopes consequently contributes to a horizontal transmission of toxoplasmosis by bringing healthy populations in new spaces where other wild mammals may have previously occurred and potentially spread oocysts.

In addition, the consumption of infected rats may increase the risk of a direct transmission from rats to humans, as it has been postulated for some populations in Nigeria with a significantly higher prevalence for those handling or eating rats (Dubey and Frenkel 1998, Uneke, et al. 2007, Olusi, et al. 1994). In Thailand, some rural populations also hunt wild rats as food and may be consequently exposed to infection by *T. gondii*. The transmission can occur by ingestion of blood when handling dead animals or when preparing them before cooking. Proper hygiene is recommended to avoid any contamination of food.

### **Acknowledgments**

Laboratory work was conducted at the Department of Parasitology, Faculty of Veterinary Medicine, at Kasetsart University in Thailand. We thank Kasetsart University Research Development Institution (KURDI) for partial funding. Field work was part of the “Eco-epidemiology of Hantaviruses in Thailand” (ANR, Program 00121 05) and the “CERoPath project” (Community ecology of rodents and their pathogens in South-East Asia: effects of biodiversity changes and implications in

health ecology / ANR 07 BDIV 012) funded by the French National Agency for Research. We particularly thank all those who have helped us hardly in the field and the anonymous reviewers for insightful remarks.

## References

- Nicolle C, Manceaux L. Sur une infection à corps de Leishman (ou organismes voisins) du gundi. Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences 1908;147:763–766.
- Splendore A. Un nuovo parassita deconigli incontrato nelle lesioni anatomiche d'una malattia che ricorda in molti punti il Kala-azar dell'uomo. Nota preliminare pel. Rev. Soc. Sci. Sao Paulo 1908;3:109-112.
- Dubey JP. The History of *Toxoplasma gondii*-The First 100 Years. J. Eukaryot. Microbiol. 2008;55:467-475.
- Tenter AM, Heckeroth AR, Weiss LM. *Toxoplasma gondii*: from animals to humans. International Journal for Parasitology 2000;30:1217-1258.
- Kim K, Weiss LM. *Toxoplasma*: the next 100 years. Microbes and Infection 2008;10:978-984.
- Dubey JP, Frenkel JK. Toxoplasmosis of rats: a review, with considerations of their value as an animal model and their possible role in epidemiology. Veterinary Parasitology 1998;77:1-32.
- Carleton MD, Musser GG. Superfamily Muroidea. In: Wilson DE, Reeder DM, eds. *Mammal species of the World, a taxonomic and geographic reference*. Baltimore: The Johns Hopkins University Press; 2005:2142.
- Salibay CC, Claveria FG. Serologic detection of *Toxoplasma gondii* infection in *Rattus* spp collected from three different sites in Dasmariñas, Cavite , Philippines Southeast Asian Journal of Tropical Medicine and Public Health 2005;36:46-49.
- Dabritz HA, Miller MA, Gardner IA, Packham AE, Atwill ER, Conrad PA. Risk factors for *Toxoplasma gondii* infection in wild rodents from central coastal California and a review of *T. gondii* prevalence in rodents. Journal of Parasitology 2008;94:675-683.
- de Thoisy B, Demar M, Aznar C, Carme B. Ecologic correlates of *Toxoplasma gondii* exposure in free-ranging neotropical mammals. Journal of wildlife diseases 2003;39:456-459.
- Maruyama S, Boonmar S, Morita Y, Sakai T, Tanaka S, Yamaguchi F, Kabeya H, Katsube Y. Seroprevalence of *Bartonella henselae* and *Toxoplasma gondii* among Healthy Individuals in Thailand. Journal of Veterinary Medical Science 2000;62:635-637.
- Jittapalapong S, Nimsupan B, Pinyopanuwat N, Chimnoi W, Kabeya H, Maruyama S. Seroprevalence of *Toxoplasma gondii* antibodies in stray cats and dogs in the Bangkok metropolitan area, Thailand. Veterinary Parasitology 2007;145:138-141.
- Thiptara A, Kongkaew W, Bilmad U, Bhumibhamon T, Anan S. Toxoplasmosis in Piglets. Annals of the New York Academy of Sciences 2006;1081:336-338.



- Jittapalapong S, Sangvaranond A, Pinyopanuwat N, Chimnoi W, Khachaeram W, Koizumi S, Maruyama S. Seroprevalence of *Toxoplasma gondii* infection in domestic goats in Satun Province, Thailand. *Veterinary Parasitology* 2005;127:17-22.
- Tuntasuvan D, Mohkaew K, Dubey JP. Seroprevalence of *Toxoplasma gondii* in Elephants (*Elephas maximus indicus*) in Thailand. *Journal of Parasitology* 2001;87:229-230.
- Thiangtum K, Nimsuphun B, Pinyopanuwat N, Chimnoi W, Tunwattana W, Tongthainan D, Jittapalapong S, Rukkwamsuk T, Maruyama S. Seroprevalence of *Toxoplasma gondii* in captive felids in Thailand *Veterinary Parasitology* 2006;136:351-355.
- Mills JN, Childs JE, Ksiazek TG, Peters CJ, Velleca WM. Methods for trapping and sampling small mammals for virologic testing. Atlanta: U.S. Department of Health and Human Services - Centers for Disease Control and Prevention; 1995.
- Marshall JT. Family Muridae: Rats and mice. In: Lekagul B, Mc Neely JA, eds. *Mammals of Thailand*. Bangkok: Association for the Conservation of Wildlife; 1988:397-487.
- Askins R. Family Sciuridae: squirrels. In: Lekagul B, Mc Neely JA, eds. *Mammals of Thailand*. Bangkok: Association for the Conservation of Wildlife; 1988:337-387.
- Aplin KP, Brown PR, Jacob J, Krebs CJ, Singleton GR. Field methods for rodent studies in Asia and the Indo-Pacific. Canberra: Australian Center for International Agricultural Research; 2003.
- Corbet GB, Hill JE. Mammals of the Indomalayan Region: a systematic review. London and Oxford: Natural History Museum Publications and Oxford University Press; 1992.
- Tilahun W, Arnaud LF, Tefera S, Henk G, Tsehaynesh M, Tobias FRdW, Hailu Y, Roel AC, Tom Van G. Evaluation of the Eiken latex agglutination test for anti-*Toxoplasma* antibodies and seroprevalence of *Toxoplasma* infection among factory workers in Addis Ababa, Ethiopia. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1998;92:401-403.
- Mazumder P, Chuang HY, Wentz MW, Wiedbrauk DL. Latex agglutination test for detection of antibodies to *Toxoplasma gondii*. *Journal of Clinical Microbiology* 1988;26:2444-2446.
- Crawley MJ. GLIM for Ecologists. Oxford: Blackwell Science; 1993.
- Lubroth J-S, Dreesen DW, Ridenhour RA. The role of rodents and other wildlife in the epidemiology of swine toxoplasmosis. *Preventive Veterinary Medicine* 1983;1:169-178.
- Dubey JP, Bhaiyat MI, Macpherson CNL, de Allie C, Chikweto A, Kwok OCH, Sharma RN. Prevalence of *Toxoplasma gondii* in Rats (*Rattus norvegicus*) in Grenada, West Indies. *Journal of Parasitology* 2006;92:1107-1108.
- Kijlstra A, Meerburg B, Cornelissen J, De Craeye S, Vereijken P, Jongert E. The role of rodents and shrews in the transmission of *Toxoplasma gondii* to pigs. *Veterinary Parasitology* 2008;156:183-190.
- Murphy RG, Williams RH, Hughes JM, Hide G, Ford NJ, Oldbury DJ. The urban house mouse (*Mus domesticus*) as a reservoir of infection for the human parasite *Toxoplasma gondii*: an unrecognised public health issue? *International Journal of Environmental Health Research* 2008;18:177-185.

- Ferguson DJP. *Toxoplasma gondii*: 1908-2008, homage to Nicolle, Manceaux and Splendore. *Memórias do Instituto Oswaldo Cruz* 2009;104:133-148.
- Rifaat MA, Mahdi AH, Arafa MS, Nasr NT, Sadek MS. Isolation of *Toxoplasma* from *Rattus norvegicus* in Egypt. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1971;65:788-9.
- de Roever-Bonnet H. Toxoplasmosis in tropical Africa. *Tropical and geographical medicine* 1972;24:7-13.
- Owen MR, Trees AJ. Vertical transmission of *Toxoplasma gondii* from chronically infected house (*Mus musculus*) and field (*Apodemus sylvaticus*) mice determined by polymerase chain reaction. *Parasitology* 1998;116:299-304.
- Meerburg BG, Singleton GR, Kijlstra A. Rodent-borne diseases and their risks for public health. *Critical Reviews in Microbiology* 2009;35:221-270.
- Uneke C, Duhlińska D, Ngwu B, Njoku M. Seroprevalence of *Toxoplasma gondii* infection in Kwal, a rural district of Plateau-Nigeria. *African journal of medicine and medical sciences* 2007;36:109-113.
- Olusi T, Ajaya J, Makinde A. Antibodies to *Toxoplasma gondii* in a rat-eating population on Benue State, Nigeria. *Annals of tropical medicine and parasitology* 1994;88:217-218.

**Address reprint requests to:**

Dr. Vincent Herbreteau

CIRAD, UR AGIRs, Maison de la télédétection,

500, rue Jean-François Breton

34093 Montpellier Cedex 5

France

Email: [vincent.herbreteau@teledetection.fr](mailto:vincent.herbreteau@teledetection.fr)

TABLE 1: PREVALENCE OF *T GONDII* ANTIBODIES IN RODENTS TRAPPED IN THAILAND

<b>Species</b>	<b>No. of positive / tested (%)</b>	<b>Provinces (See figure 1)</b>
<b>Murinae</b>		
<i>Bandicota indica</i>	1/37 (2.7)	1, 2, 3, 4, 8, 12
<i>Bandicota savilei</i>	0/11 (0)	1
<i>Berylmys berdmorei</i>	0/4 (0)	2, 3
<i>Berylmys bowersi</i>	0/5 (0)	2, 3
<i>Leopoldamys edwardsi</i>	0/3 (0)	1, 2
<i>Leopoldamys sabanus</i>	2/16 (12.5)	3
<i>Maxomys surifer</i>	2/38 (5.3)	2, 3, 5
<i>Mus cervicolor</i>	0/1 (0)	8
<i>Mus cookii</i>	0/1 (0)	1
<i>Niviventer fulvescens</i>	0/13 (0)	3, 12
<i>Rattus andamanensis</i>	0/7 (0)	2, 3
<i>Rattus exulans</i>	1/79 (1.3)	1, 2, 3, 5, 6, 7, 8, 10, 11, 12
<i>Rattus losea</i>	0/1 (0)	2
<i>Rattus norvegicus</i>	0/34 (0)	8
<i>Rattus tanezumi</i>	11/156 (7.1)	2, 3, 5, 6, 8, 9, 10, 12, 13
<i>Rattus tiomanicus</i>	0/2 (0)	13
<b>Total Muridae</b>	<b>17/408 (4.2)</b>	
<b>Sciuridae</b>		
<i>Callosciurus sp.</i>	0/1 (0)	3
<i>Menetes berdmorei</i>	4/52 (7.7)	3
<b>Total Sciuridae</b>	<b>4/53 (7.5)</b>	
<b>Total</b>	<b>21/461 (4.6)</b>	

FIGURE 1: LOCATION OF SAMPLING SITES IN 13 THAI PROVINCES




FIGURE 2: PREVALENCE OF *T GONDII* IN RELATION TO A GRADIENT OF ANTRHOPIZATION

IN HABITATS





FIGURE 3: WORLDWIDE PREVALENCE OF *T. GONDII* IN *R. NORVEGICUS*



SUPPLEMENTAL DATA: LITERATURE REVIEW OF THE SEROPREVALENCE OF *T. GONDII* IN

*RODENTS (KNOWN TO OCCUR IN THAILAND)*

Species	Country	Total tested	Positive	Prevalence (%)	Method	Reference
<i>Mus musculus</i>	Netherlands	31	2	6,5	PCR	Kijlstra et al., 2008
	USA	20	12	60,0	IIF	Lubroth et al., 1983
	UK	200	118	59,0	PCR	Murphy et al., 2008
	<b>Total</b>	<b>251</b>	<b>132</b>	<b>52,6</b>		
<i>Rattus exulans</i>	Micronesia	420	14	3,3	DT	Wallace et al., 1972
	USA	85	5	5,9	DT	Wallace, 1973
	<b>Total</b>	<b>505</b>	<b>19</b>	<b>3,8</b>		
<i>Rattus norvegicus</i>	Australia	41	3	7,3	CF	Cook and Pope, 1959
	Costa Rica	23	7	30,4	DT	Chinchilla, 1978
	Costa Rica	107	22	20,6	DT	Ruiz and Frenkel, 1980
	Czech Republic	519	49	9,4	DT	Zàstera et al., 1966
	Egypt	60	8	13,3	IHA	El Nahal et al., 1982
	Egypt	100	34	34,0	DT	Rifaat et al., 1971
	Finland	108	13	12,0	DT	Grönroos and Salminen, 1955
	France	4	1	25,0	DT	Doby et al., 1974
	Germany	34	5	14,7	DT	Laven and Westphal, 1950
	Grenada	238	2	0,8	MAT	Dubey et al., 2006
	India	186	0	0,0	IHA	Mir et al., 1982
	Italy	152	34	22,4	IFA	Zardi et al., 1983
	Italy	20	14	70,0	IFA	Genchi et al., 1991
	Japan	116	7	6,0	DT	Izutani, 1958
	Japan	100	12	12,0	DT	Jyono, 1960
	Japan	300	25	8,3	HA	Iseki et al., 1972
	Japan	15	0	0,0	LAT	Murata, 1987
	Kuwait	142	4	2,8	HA	Bezjak and Thorburn, 1982
	Mexico	360	83	23,1	DT	Varela, 1955
	Netherlands	39	4	10,3	PCR	Kijlstra et al., 2008
	Panama	226	81	35,8	MAT	Frenkel et al., 1995
	Philippines	83	50	60,2	LAT	Salibay and Claveria, 2005
	Poland	646	58	9,0	DT	Slowakiewicz et al., 1967
	Saudi Arabia	142	20	14,1	IHA	Morsy et al., 1994
	UK	235	82	34,9	LAT	Webster, 1994
	UK	46	5	10,9	DT	Hay et al., 1983
	UK	65	5	7,7	DT	Jackson et al., 1986
	USA	8	2	25,0	IHA	Burridge et al., 1979
	USA	2	2	100,0	IHA	Lubroth et al., 1983
	USA	73	1	1,4	DT	Wallace, 1973
	USA	95	6	6,3	MAT	Dubey et al., 1995
	USA	109	54	49,5	IFA	Childs and Seegar, 1986
USA	25	0	0,0	DT	Eyles et al., 1959	
USA	100	8	8,0	DT	Eyles, 1952	
USA	2	2	100,0	IIF	Lubroth et al., 1983	
<b>Total</b>	<b>4521</b>	<b>703</b>	<b>15,5</b>			
<i>Rattus rattus</i>	Africa	61	5	8,2	DT	de Roever-Bonnet, 1972
	Australia	117	4	3,4	CF	Cook and Pope, 1959
	France	40	1	2,5	DT	Doby et al., 1974
	Italy	249	71	28,5	IFA	Zardi et al., 1983
	Japan	42	2	4,8	DT	Izutani, 1958
	Japan	40	0	0,0	DT	Jyono, 1960
	Japan	26	4	15,4	HA	Iseki et al., 1972
	Mexico	39	27	69,2	DT	Varela, 1955
	Micronesia	238	7	2,9	DT	Wallace et al., 1972
	Saudi Arabia	24	7	29,2	IHA	Morsy et al., 1994



USA	38	5	13,2	IHA	Burridge et al., 1979
USA	476	36	7,6	DT	Wallace, 1973
<b>Total</b>	<b>1390</b>	<b>169</b>	<b>12,2</b>		

CF	Complement Fixation
DT	Dye Test
HA	Hemagglutination
IFA	Immuno-Fluorescence Assay
IHA	Indirect Hemagglutination
IIF	Indirect Immuno-Fluorescent
LAT	Latex Agglutination Test
MAT	Modified Agglutination Test
PCR	Polymerase Chain Reaction

### Supplementary references:

- Bezjak V, Thorburn H. Survey of rats (*Rattus norvegicus*) in Kuwait for the presence of some human pathogens: 2. Results of *Toxoplasma* and *Trichinella* investigation. Journal of the Kuwait Medical Association 1982;16:11-15.
- Burridge MJ, Bigler WJ, Forrester DJ, Hennemann JM. Serologic survey for *Toxoplasma gondii* in wild animals in Florida. Journal of the American Veterinary Medical Association 1979;175:964-967.
- Childs JE, Seegar WS. Epidemiologic observations on infection with *Toxoplasma gondii* in three species of urban mammals from Baltimore, Maryland, USA. International Journal of Zoonoses 1986;13:249-261.
- Chinchilla M. Epidemiologia de la toxoplasmosis en Costa Rica: importancia de los roedores domésticos. Revista de Biología Tropical 1978;26:113-124.
- Cook I, Pope JH. *Toxoplasma* in Queensland: III. A preliminary survey of animal hosts. Australian Journal of Experimental Biology and Medical Sciences 1959;37:253-262.
- El Nahal HS, Morsy TA, Bassili WR, El Missiry AG, Saleh MSM. Antibodies against three parasites of medical importance in *Rattus* sp. collected in Giza Governorate, Egypt. Journal of the Egyptian Society of Parasitology 1982;12:287-293.
- Doby JM, Desmots G, Beaucournu JC, Akinchina GT. Recherche immunologique systématique de toxoplasmose chez les petits mammifères sauvages en France. Folia Parasitologica 1974;21:289-300.
- Dubey JP, Weigel RM, Siegel AM, Thulliez P, Kitron UD, Mitchell MA, Mannelli A, Mateus-Pinilla NE, Shen SK, Kwok OCH and others. Sources and reservoirs of *Toxoplasma gondii* infection on 47 swine farms in Illinois. Journal of Parasitology 1995;81:723-729.
- Eyles DE. *Toxoplasma* in the Norway rat. Journal of Parasitology 1952;38:226-229.
- Eyles DE, Gibson CL, Coleman N, Smith CS, Jumper JR, Jones FE. The prevalence of toxoplasmosis in wild and domesticated animals of the Memphis region. American Journal of Tropical Medicine and Hygiene 1959;8:505-510.
- Frenkel JK, Hassanein KM, Hassanein RS, Brown E, Thulliez P, Quintero-Nunez R. Transmission of *Toxoplasma gondii* in Panama City, Panama: a five-year prospective cohort study of children, cats, rodents, birds, and soil. American Journal of Tropical Medicine and Hygiene 1995;53:458-468.
- Genchi G, Polidori GA, Zaghini L, Lanfranchi P. Aspetti epidemiologici della toxoplasmosi nell'allevamento intensivo del suino. Archivio veterinario italiano 1991;42:105-111.

- Grönroos P, Salminen A. Toxoplasmosis in Norway rats in Helsinki. *Annales Medicinæ Experimentalis et Biologiæ Fenniae* 1955;33:141-144.
- Hay J, Hutchison WM, Jackson MH, Siim JC. Prevalence of *Toxoplasma* infection in a wild rodent population from central Scotland. *Annals of Tropical Medicine and Parasitology* 1983;77:653-654.
- Iseki M, Nishibayashi M, Sano R, Ogo T, Inamoto T, Shibuya J. Prevalence of *Toxoplasma* antibodies in rats in Osaka. [In Japanese]. *Japanese Journal of Parasitology* 1972;21:39-44.
- Izutani T. *Toxoplasma* antibody maintained by house rats, cattle, horses, fowls, and sparrows. [In Japanese]. *Journal of the Osaka City Medical Center* 1958;7:181-192.
- Jackson MH, Hutchison WM, Siim JC. Toxoplasmosis in a wild rodent population of central Scotland and a possible explanation of the mode of transmission. *Journal of Zoology London* 1986;A 209:549-557.
- Jyono K. Surveys of *Toxoplasma* antibodies in groups of man and rats, and on suitable conditions of dye test. [In Japanese]. *Journal of the Osaka City Medical Center* 1960;9:1907-1925.
- Laven H, Westphal A. Die UÜbertragung von *Toxoplasma gondii* unter besonderer berücksichtigung des blutes als infektionsquelle. *Z. Tropenmed. Parasitol.* 1950;2:221-235.
- Mir NA, Chhabra MB, Bhardwaj RM, Gautam OP. *Toxoplasma* infection and some other protozoan parasites of the wild rat in India. *Indian Veterinary Journal* 1982;59:60-63.
- Morsy TA, Sabry AHA, Habib KSM, Arafa MAS, El Bahrawy AFA, Al Dakhil MM. Antibodies against *Toxoplasma* in commensal rodents trapped in Riyadh region, Saudi Arabia. *Journal of the Egyptian Society of Parasitology* 1994;24:279-284.
- Murata K. Antibodies against *Toxoplasma gondii* in wild animals wounded or diseased and caught in the Kobe area. [In Japanese]. *Journal of the Japanese Veterinary Medical Association* 1987;40:846-849.
- Ruiz A, Frenkel JK. Intermediate and transport hosts of *Toxoplasma gondii* in Costa Rica. *American Journal of Tropical Medicine and Hygiene* 1980;29:1161-1166.
- Slowakiewicz E, Starzyk JR, Siuda K, Drozd A. Badania nad czestoscia wystepowania pierwotniaka *Toxoplasma gondii* u dzikich zwierzat. [Investigations on the frequency of appearance of the protozoon *Toxoplasma gondii* in wild animals]. *Acta Biologica Cracoviensia* 1967;10:267-271.
- Varela G. Toxoplasmosis en la Republica Mexicana. *An. Soc. Mexico Oftalmol* 1955;28:227-239.
- Wallace GD, Marshall L, Marshall MAC. Cats, rats, and toxoplasmosis on a small Pacific island. *American Journal of Epidemiology* 1972;95:475-482.
- Wallace GD. Intermediate and transport hosts in the natural history of *Toxoplasma gondii*. *American Journal of Tropical Medicine and Hygiene* 1973;22:456-464.
- Webster JP. Prevalence and transmission of *Toxoplasma gondii* in wild brown rats, *Rattus norvegicus*. *Parasitology* 1994;108:407-411.
- Zardi O, Adorisio E, La Commare F, Milana G, Pozio E, Gramiccia M, Gradoni L. *Toxoplasma gondii* in roditori e cani dell'Italia centrale. *Rev. Parasitol.* 1983;44:147-149.

Zástěra M, Hubner J, Pokorný J, Seeman J. Králci jako možný zdroj toxoplazmické infekce člověka. (Rabbits as a possible source of *Toxoplasma* infection for man). Ceska Epidemiol. Mikrobiol. Imunol. 1966;15:172-177.