



HAL
open science

Regulation of mitochondrial biogenesis during myogenesis

A.H.V. Remels, R.C.J. Langen, P. Schrauwen, G. Schaart, A.M.W.J. Schols,
H.R. Gosker

► **To cite this version:**

A.H.V. Remels, R.C.J. Langen, P. Schrauwen, G. Schaart, A.M.W.J. Schols, et al.. Regulation of mitochondrial biogenesis during myogenesis. *Molecular and Cellular Endocrinology*, 2009, 315 (1-2), pp.113. 10.1016/j.mce.2009.09.029 . hal-00547658

HAL Id: hal-00547658

<https://hal.science/hal-00547658v1>

Submitted on 17 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Regulation of mitochondrial biogenesis during myogenesis

Authors: A.H.V. Remels, R.C.J. Langen, P. Schrauwen, G. Schaart, A.M.W.J. Schols, H.R. Gosker



PII: S0303-7207(09)00526-7
DOI: doi:10.1016/j.mce.2009.09.029
Reference: MCE 7341

To appear in: *Molecular and Cellular Endocrinology*

Received date: 27-4-2009
Revised date: 27-9-2009
Accepted date: 28-9-2009

Please cite this article as: Remels, A.H.V., Langen, R.C.J., Schrauwen, P., Schaart, G., Schols, A.M.W.J., Gosker, H.R., Regulation of mitochondrial biogenesis during myogenesis, *Molecular and Cellular Endocrinology* (2008), doi:10.1016/j.mce.2009.09.029

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Regulation of mitochondrial biogenesis during myogenesis

Remels AHV₁, Langen RCJ₁, Schrauwen P₂, Schaart G₂, Schols AMWJ₁ and Gosker HR₁

1. Department of Respiratory Medicine, Maastricht University Medical Centre +, the Netherlands
2. Department of Human Biology, Maastricht University Medical Centre +, the Netherlands

Correspondence and requests for reprints

Alexander H. Remels

Department of Respiratory Medicine

Maastricht University Medical Centre +, P.O. Box 5800, 6202 AZ Maastricht, the Netherlands

E-mail: a.remels@pul.unimaas.nl

Fax: +31-43-3875051; Telephone: +31-43-3881563

Abstract

Pathways involved in mitochondrial biogenesis associated with myogenic differentiation are poorly defined. Therefore, C₂C₁₂ myoblasts were differentiated into multi-nucleated myotubes and parameters/regulators of mitochondrial biogenesis were investigated. Mitochondrial respiration, citrate synthase- and β -hydroxyacyl-CoA dehydrogenase activity as well as protein content of complexes I, II, III and V of the mitochondrial respiratory chain increased 4-8 fold during differentiation. Additionally, an increase in the ratio of myosin heavy chain (MyHC) slow vs MyHC fast protein content was observed. PPAR transcriptional activity and transcript levels of PPAR- α , the PPAR co-activator PGC-1 α , mitochondrial transcription factor A and nuclear respiratory factor 1 increased during differentiation while expression levels of PPAR- γ decreased. In conclusion, expression and activity levels of genes known for their regulatory role in skeletal muscle oxidative capabilities parallel increases in oxidative parameters during the myogenic program. In particular, PGC-1 α and PPAR- α may be involved in the regulation of mitochondrial biogenesis during myogenesis.

Keywords: peroxisome proliferator-activated receptors, myogenesis, mitochondrial transcription factor A, nuclear respiratory factor 1, skeletal muscle

Introduction

Intrinsically, skeletal muscle is a tissue characterized by a high level of plasticity. The potential to alter muscle size, metabolic properties and/or protein isoform expression gives skeletal muscle the ability to adapt to the different challenges that may be placed upon it [Baldwin and Haddad, 2002]. Events triggering skeletal muscle adaptive changes include endurance- and strength training, overload, aging, disuse, starvation, chronic illness and degenerative disorders [Fluck, 2006, Narici et al., 2004, Gosker et al., 2002]. Growth and repair (regeneration) of skeletal muscle tissue is inextricably linked to the action of a group of myogenic precursor cells, called satellite cells. Upon activation, these cells proliferate, differentiate and ultimately fuse with existing myofibers [Anderson, 2006]. The irreversible transition from the proliferation competent myoblast stage into fused multi-nucleated myotubes is known as myogenic differentiation and recapitulates developmental and regenerative myogenesis [Le Grand and Rudnicki, 2007].

Oxidative phenotype (OXPHEN) of a muscle depends on its oxidative capacity (determined by the activity of mitochondrial enzymes involved in oxidative substrate metabolism) and fiber type composition, (e.g. ratio type I oxidative (slow-twitch) vs type IIB/X glycolytic (fast-twitch) fibers). The peroxisome proliferator-activated receptors (PPARs), especially the PPAR- δ and PPAR- α isoforms, and related co-activator molecules as PGC-1 α are highly implicated in regulation of skeletal muscle OXPHEN [Luquet et al., 2003, Koves et al., 2005]. Accordingly, muscle-specific over-expression of PPAR- δ or PGC-1 α in mice potently up-regulated oxidative capabilities of the muscle and mediated a fiber type shift towards an increased OXPHEN [Lin et al., 2002, Wang et al., 2004]. Moreover, PGC-1 α and the PPARs govern mitochondrial biogenesis through control over other regulatory proteins including nuclear respiratory factor 1 (NRF-1) and the master

mitochondrial regulator mitochondrial transcription factor A (Tfam) [Gleyzer et al., 2005, Wu et al., 1999]. An improved muscle OXPHEN may be established during the process of myogenesis (i.e. the formation, repair or hypertrophy of muscle), or through alterations within the pre-existing muscle fibers, independent of myogenic differentiation. The regulation of oxidative profile in existing muscle fibers has been described quite extensively in the literature [Lin et al., 2002, Wang et al., 2004, Canto et al., 2007] whereas knowledge regarding the development of muscle oxidative profile and the involvement of key regulators herein during myogenic differentiation is rather limited.

Studying the molecular events governing regulation of mitochondrial biogenesis during myogenesis is of clinical relevance for a number of pathological conditions including chronic inflammatory myopathies and degenerative muscle conditions as e.g. Duchenne's muscular dystrophy [Oexle and Kohlschutter, 2001, Wallace and McNally, 2009], but also in conditions characterized by loss of OXPHEN in skeletal muscle, such as COPD [Gosker et al., 2002].

It has been shown that mitochondria play a mechanistic role not only in myoblast proliferation [Duguez et al., 2004, Rochard et al., 2000], but also in myogenesis by targeting key regulators of myogenic differentiation as myogenin [Rochard et al., 2000] and cellular oncogens as c-myc [Seyer et al., 2006]. Moreover, blocking mitochondrial biogenesis potently inhibits myogenic differentiation [Herzberg et al., 1993, Rochard et al., 2000]. This demonstrates that mitochondria display a retrograde signaling to regulate their own biogenesis as well as cell cycle progression [Jahnke et al., 2009], which subsequently implies that the investigation of myogenic differentiation is inevitably coupled to studying the process of mitochondrial biogenesis.

Although markers of OXPHEN (including mitochondrial biogenesis) during myogenesis have been described previously [Moyes et al., 1997, Rochard et al., 2000, Duguez et al., 2004, Duguez et al., 2002, Kraft et al., 2006, Jahnke et al., 2009], information about the expression profile of key regulators during the entire myogenic program is incomplete. Therefore, the aim of the present study was to chart the development of multiple aspects of mitochondrial biogenesis during the full length of the differentiation process of skeletal muscle cells in order to increase our understanding of the key regulatory molecules of OXPHEN during myogenesis.

Materials and methods

Cell culture

The murine skeletal muscle cell line C₂C₁₂ was obtained from the American Type Culture Collection (ATCC CRL1772; Manassas, VA, USA). These cells are able to undergo differentiation into spontaneously contracting myotubes after growth factor withdrawal [Yaffe and Saxel, 1977]. Myoblasts were cultured in growth medium (GM) composed of low-glucose Dulbecco's Modified Eagle's medium (DMEM) containing antibiotics (50 U/ml penicillin and 50 µg/ml streptomycin) and 9% (v/v) foetal bovine serum (FBS) (all from GIBCO, Rockville, MD, USA). Cells were plated at 10⁴ cells/cm² and cultured in GM for 24 h. To induce differentiation, cells were washed in Hank's Balanced Salt solution (HBSS) and GM was replaced with DM, which contained DMEM with 0.5% heat-inactivated FBS and antibiotics. C₂C₁₂ cells were grown on Matrigel (BD Biosciences, Bedford, MA, USA) (1:50 in DMEM) (Becton Dickinson Labware, Bedford, MA, USA), which contains constituents of the natural extracellular matrix of skeletal muscle fibers including collagen IV, laminin, heparin sulfate proteoglycan and entactin.

Transfections and Plasmids

For assessment of PPAR-dependent transcriptional activity, C₂C₁₂ cells were stably transfected with a PPAR-sensitive luciferase reporter plasmid containing 3 PPAR responsive elements in tandem derived from the promoter region of the acyl CoA oxidase (ACO) gene (pSG5, Stratagene, La Jolla, CA, USA). Cells were grown to 70% confluency and transfected by Lipofectamine according to manufacturer's instructions (Invitrogen, Carlsbad, CA, USA). Transfection was performed in the presence of 3.75 µg ACO luciferase plasmid DNA (pSG5, Stratagene) and 0.25 µg of a plasmid containing the neomycin resistance gene (pSV2-Neo, Stratagene). For selection of positive clones, cells were cultured in GM containing 800 µg/ml G418 (Calbiochem, Nottingham, United Kingdom).

May-Grunwald Giemsa staining

As a morphological parameter of differentiation, the myogenic index was defined as the number of nuclei residing in cells containing three or more nuclei divided by the total number of nuclei in May-Grunwald Giemsa stained cells. Cells were grown on Matrigel-coated 60 mm dishes and differentiation was induced. After 0, 3 or 7 days of differentiation cells were washed 2x in phosphate buffered saline (PBS) (room temperature), fixed in methanol and stained in May-Grunwald Giemsa (Sigma, Saint Louis, MO, USA) according to manufacturer's instructions.

Reporter assays

The PPAR-sensitive reporter cell line was plated in 35 mm dishes and allowed to grow to 70% confluency. After induction of differentiation (day 0), cells were differentiated into myotubes for 3, 5 or 7 days after which cells were harvested. Cells were washed 2x with cold PBS and subsequently lysed by adding (100 µl) 1x Reporter Lysis Buffer (Promega, Madison,

WI, USA) and incubation on ice for 10 min. Cell lysates were centrifuged (13 000 g, 1 min), and supernatants were snap-frozen and stored at -80 °C for later analysis. Luciferase activity was measured according to manufacturer's instructions (Promega) and corrected for total protein content (Biorad, Hercules, CA, USA).

Real time quantitative PCR (QPCR)

Total RNA was extracted using the acid guanidium thiocyanate-phenol-chloroform extraction method (Ambion Ltd., IJssel, The Netherlands). RNA concentration was determined using a spectrophotometer. One µg of RNA per sample was reverse transcribed into cDNA according to the manufacturer's instructions (Reverse-iT™ 1st strand synthesis, Abgene, Leusden, The Netherlands). cDNA was PCR amplified with SYBR Green Fluorescein Mix (Westburg, Leusden, The Netherlands) on a Biorad PCR apparatus (Biorad). Specificity of the amplification was verified by melt curve analysis and evaluation of efficiency of PCR amplification. Primers were designed to generate a PCR amplification product of 100 to 150 bp (Table 1). Transcript levels for the constitutive housekeeping gene products cyclophilinA, Hypoxanthine phosphoribosyltransferase (HPRT), β -actin and Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) were quantitatively measured in each sample to control for sample-to-sample differences in RNA concentration. Additionally, geNorm software was applied to calculate a normalization factor based on the expression levels of these 4 housekeepers. geNorm calculates the gene expression stability measure M for a reference gene as well as the average pairwise variation V for that gene with all other tested reference genes. Stepwise exclusion of the gene with the highest M value allows ranking of the tested genes according to their expression stability. The underlying principles and calculations are described in Vandesompele et al. [Vandesompele et al., 2002]. Gene

expression was quantified and expressed as arbitrary units (AU). **Please include Table 1 here.**

Western blotting

Abundance of proteins involved in oxidative phosphorylation (OXPHOS) as well as myosin heavy chain (MyHC) protein content was evaluated by Western blotting. Cells were washed in PBS and whole cell lysates were prepared by the addition of lysis buffer (20 mM Tris, 150 mM NaCl, 1% (v/v) Nonidet P-40, 1 mM DTT, 1 mM Na₃VO₄, 1 mM PMSF, 10 µg/ml leupeptin, and 1% (v/v) aprotinin). Lysates were incubated on ice for 30 min, followed by 30 min of centrifugation at 13 000 g. A fraction of the supernatant was saved for protein determination and 4 x Laemmli sample buffer (0.25 M Tris·HCl; pH 6.8, 8% (w/v) SDS, 40% (v/v) glycerol, 0.4 M DTT, and 0.04% (w/v) bromophenol blue) was added, followed by boiling of the samples for 5 min and storage at -20 °C. Total protein was assessed using the Biorad DC protein assay kit (Biorad) according to the manufacturer's instructions. 25 µg or 2 µg of protein (OXPHOS and MyHC respectively) was loaded per lane and separated on a 10% polyacrylamide gel (Mini Protean 3 system; Biorad), followed by transfer to a 0.45 µm nitrocellulose membrane (Biorad) by electro blotting. The membrane was blocked for 1 h at room temperature in 5% (w/v) non-fat dry milk. Nitrocellulose blots were washed in PBS-Tween 20 (0.05%), followed by overnight incubation (4 °C) with a polyclonal antibody specific for OXPHOS proteins (Cell Signaling, Beverly, MA, USA) or MyHC fast or slow (Sigma). After three wash steps of 20 min each, the blots were probed with a peroxidase-conjugated secondary antibody (Vector Laboratories, Burlingame, CA, USA) and visualized using SuperSignal West Pico chemiluminescent substrate (Pierce Biotechnology, Rockford, IL, USA) according to the manufacturer's instructions.

Enzyme activity assay

Cells were washed twice in ice-cold PBS and subsequently lysed by 10 min incubation with 200 μ l of 0.1% Triton-X100 dissolved in PBS. Cells were harvested by scraping, using rubber policemen, and transferred to an Eppendorf tube containing 45 μ l 5% bovine serum albumine (BSA). After vortexing for 10 sec, samples were centrifuged (10 min, 13 000 g, 4 °C), and the supernatant was used for spectrophotometrical assessment of the following enzyme activities: HAD; EC 1.1.1.35; see Ref. [Gosker et al., 2005], CS; EC 2.3.3.1 ;see Ref. [Gosker et al., 2005] and CK; EC2.7.3.2 see Ref [Langen et al., 2003]) (Multiskan Spectrum; Thermo Labsystems, Breda, The Netherlands). Enzyme activities were corrected for total protein content (Biorad).

Mitochondrial respiration

C₂C₁₂ myoblasts or differentiated myotubes were harvested in trypsin, washed twice with MIRO5 respiration medium (37 °C) (prepared according to MiPNet protocols) (Oroboros Instruments, Innsbruck, Austria) and, after a 5 min centrifugation at 500 g, resuspended in 2.0 ml of MIRO5 respiration medium. Protein content was determined in dishes that were cultured in a parallel experiment. Measurement of respiration was performed by a polarographic oxygen sensor in a 2 ml glass chamber of the respirometer Oxygraph 2K (Oroboros Instruments). The amplified signal from the oxygen sensor was recorded on a computer at sampling intervals of 2 seconds using DatLab acquisition software (Oroboros Instruments). The respiration medium was equilibrated with air at 37 °C and stirred until a stable signal was obtained for calibration at air saturation. After calibration the medium was replaced by aerated cell suspensions at a volume of 2.0 ml. After insertion, cells were permeabilized using 1 μ l of digitonin (0.5 μ g/ml final concentration) and substrates were added in chronological order (volume; final concentration): malate (5 μ l; 2 mM), followed by

glutamate (10 μ l; 10 mM) or pyruvate (5 μ l; 5 mM) or palmitoyl-carnitine (10 μ l; 50 μ M) followed by ADP (20 μ l; 5 mM). At the end of the substrate trace a titration with the uncoupling agent FCCP was performed (1 μ l; 0.5 μ M titration). Oxygen consumption (pmol/ml.s) was calculated as a negative time derivative of oxygen concentration (nmol/ml). Respiration rates were corrected for total protein content to correct for variability in cellular input.

Mitochondrial DNA copy number

Mitochondrial DNA (mtDNA) copy number was determined as a marker for mitochondrial density using QPCR. Total DNA was isolated from the cells according to manufacturer's instructions (Nucleospin, Machery Nagel, Ede, The Netherlands). A serial dilution standard curve was prepared from a pool of all the samples. Real-time PCR was carried out at the following parameters: 15 min denaturation step at 95 °C, followed by 40 cycles at 95 °C for 15 sec and 60 °C for 45 sec, 30 sec at 95 °C and 30 sec at 60 °C followed by a stepwise increase of 0.5 °C until 95 °C was reached (melt curve analysis). Mitochondrial DNA copy number was calculated from the ratio COXII/cyclophilinA or COXIII/cyclophilinA.

Statistical analysis

Data was analyzed according to the guidelines of Altman *et al.* using SPSS (Statistical Package for the Social sciences, SPSS Inc., Chicago, IL, USA) [Altman et al., 1983]. Kruskal-Wallis tests were applied for mean comparisons as a function of time and non-parametric tests were used to detect significant differences between individual time-points. Data are represented as the mean \pm SD. A two-tailed probability value of less than 0.05 was considered to be significant.

Results

Markers of myogenic differentiation

In order to characterize myogenic differentiation of C₂C₁₂ cells we compared biochemical and structural markers of differentiation in myoblasts vs differentiating/differentiated myotubes. Results are summarized in Table 2 and are illustrative of good morphological and biochemical differentiation. **Please include Table 2 here.**

Mitochondrial respiration in myoblasts and myotubes

To investigate the development of oxidative capacity during C₂C₁₂ myogenesis, we measured mitochondrial O₂ consumption in proliferating C₂C₁₂ myoblasts and differentiated myotubes (differentiated 7 days) in response to different substrates. As depicted in Fig. 1, basal respiratory rate was higher in myotubes when compared to myoblasts in the presence of endogenous substrates (basal respiration) (p=0.050). FCCP-induced respiration (maximal respiration), bypassing respiratory control by ATPase and thus reflecting maximal respiratory capacity, was 4-fold higher in myotubes compared to myoblasts (p=0.050). Shown in Fig. 1 is the maximal respiratory capacity with palmitoyl-carnitine as a substrate. Differences in maximal respiratory capacity between myoblasts and myotubes were similar when other substrates were used (glutamate and pyruvate respectively, data not shown). **Please include Figure 1 here.**

Mitochondrial markers during myogenesis

To gain insight in mitochondrial functioning and mitochondrial biogenesis during differentiation of C₂C₁₂ cells into mature myotubes, we measured several mitochondrial markers. These included activity levels of enzymes involved in substrate oxidation, to assess

functionality of the citric acid cycle and fatty acid β -oxidation pathway, structural components of the respiratory chain (OXPHOS) and mtDNA copy number as a general marker for mitochondrial content. Additionally, we also investigated mRNA transcript levels of cytochrome *c* oxidase (COX) subunits II and III (mitochondrial-encoded) and subunits Va and VIIc (nuclear-encoded). Activity levels of enzymes involved in oxidative metabolism, CS (Krebs cycle) and HAD (β -oxidation), increased approximately 4-fold during myogenesis ($p=0.016$) (Fig. 2A). Accordingly, protein levels of all investigated complexes of the respiratory chain increased up to 8-fold ($p=0.019$, 0.023 , 0.021 and 0.016 for complex I, II, III and V respectively) (Fig. 2B) while mtDNA copy number was 4-fold higher in fully differentiated myotubes compared to myoblasts ($p=0.025$) (Fig. 2C). As illustrated in Fig. 2D, transcript levels of the mitochondrial-encoded subunits COX II and COX III as well as the nuclear-encoded subunits Va and VIIc increased during differentiation ($p=0.023$ and 0.026 respectively). **Please include Figure 2 here.**

Myosin heavy chain isoform protein content during myogenesis

To assess the distribution of MyHC isoforms during myogenesis, we investigated protein levels of MyHC fast (type II) and MyHC slow (type I) isoforms in C₂C₁₂ cells. MyHC protein levels are expressed as the ratio between the slow and fast isoform. Myoblasts expressed extremely low levels of both MyHC fast and slow isoforms whereas differentiated myotubes expressed significant amounts of both isoforms. During differentiation a relative higher increase in MyHC slow isoform was observed compared to MyHC fast isoform expression ($p=0.025$) (Fig. 3). **Please include Figure 3 here.**

PPARs and related factors during differentiation

We determined mRNA transcript levels of PGC-1 α and all PPAR isotypes at different time points during the differentiation process. As illustrated in Fig. 4, PGC-1 α and PPAR- α mRNA expression levels increased ($p=0.063$ and 0.024 respectively) while PPAR- γ showed decreased mRNA expression levels during differentiation ($p=0.016$). PPAR- δ mRNA expression levels were stable during the full myogenic program. Data was corrected for a normalization factor derived from a geNorm analysis of 4 housekeepers (cyclophilinA, β -actin, GAPDH and HPRT). The average expression stability (M-value) for cyclophilinA, β -actin, GAPDH and HPRT was 0.473, 0.987, 0.688 and 0.470 respectively. Pairwise comparison for inclusion of a fourth housekeeping gene (V3/4) was 0.244 indicating preferable use of 4 housekeeping genes. It must be mentioned that expression levels of all 4 investigated housekeeping genes decreased approximately 2-fold during differentiation of C₂C₁₂ cells (data not shown). However, similar patterns of expression of all genes of interest were observed when data was corrected for any of the 4 individual housekeepers or when correction based on the geNorm normalization factor was applied. In addition to the PPARs and their co-activator PGC-1 α , mRNA levels of 2 key regulators of mitochondrial biogenesis downstream of PGC-1 α (NRF-1 and Tfam), were also determined during the myogenic program of C₂C₁₂ cells. Tfam and NRF-1 mRNA expression levels increased from day 0 until day 3 of the differentiation process and remained increased for the duration of the differentiation process ($p=0.096$ and 0.024 respectively) (Fig. 5). Data corrected for the housekeeper cyclophilinA is shown as a representative housekeeper, but similar results were obtained with all other individual housekeeping genes as well as with the geNorm normalization factor. **Please include Figure 4 and 5 here.**

PPAR-dependent transcriptional activity

The PPARs as well as PGC-1 α are highly implicated in the regulation of skeletal muscle oxidative capacity [Lin et al., 2002,Wang et al., 2004,Canto et al., 2007]. Therefore, we investigated PPAR-dependent transcriptional activity during differentiation of C₂C₁₂ cells into multi-nucleated myotubes. As depicted in Fig. 6, PPAR-dependent transcriptional activity increased during differentiation (p=0.022). **Please include Figure 6 here.**

Discussion

In the present paper we show that multiple key determinants of OXPHEN develop during myogenic differentiation, which coincides with the initiation of mitochondrial biogenesis and increased expression and activity levels of key regulators of muscle OXPHEN.

Constituents of OXPHEN

During the process of myogenic differentiation a remarkable morphological transition from myoblasts into multi-nucleated myotubes is observed. In addition, myogenesis is accompanied by an increase in oxidative metabolism as we demonstrate in a detailed manner in the present study. We show an increased mitochondrial respiratory capacity, as well as increased protein content of the complexes of the inner mitochondrial membrane respiratory chain, which is the site of O₂ consumption [Mathieu-Costello and Hepple, 2002]. In addition, we observed increased enzyme activity levels of HAD and CS and more oxidative type 1 MyHC slow protein in myotubes as compared to myoblasts. This is in concordance with previous studies and is illustrative of development of OXPHEN during myogenesis [Duguez et al., 2002,Leary et al., 1998]. The observed increase in the activity level of CS (established marker of mitochondrial volume) as well as increased mtDNA copy number and increases in different COX mRNA levels confirm increased mitochondrial biogenesis [Kim et al., 1995].

This data collectively shows that myogenic differentiation of C₂C₁₂ cells is obviously associated with mitochondrial biogenesis. It can however not be excluded that, in addition to mitochondrial biogenesis, alterations in efficiency of mitochondria contribute to an increased mitochondrial capacity during myogenesis.

Regulators of OXPHEN

Coordination of the nuclear and mitochondrial genome during mitochondrial biogenesis is regulated by a network of transcription factors including PGC-1 α , PPARs, NRF-1 and Tfam [Butow and Bahassi, 1999]. In this signaling pathway, PGC-1 α is the master regulator coordinating PPAR activity levels as well as mitochondrial biogenesis through activation of NRF-1-mediated transcription of Tfam [Baar, 2004, Scarpulla, 2002]. In the present study, PGC-1 α expression increased during myogenesis. The role of PGC-1 α in positive regulation of oxidative capacity and fiber type composition of existing muscle is well established [Lin et al., 2002]. Our data now also suggests a role for PGC-1 α in the development of OXPHEN during myogenic differentiation. In line with this notion, PGC-1 α mRNA levels increased during mitochondrial biogenesis in a mouse model of skeletal muscle regeneration, although these authors did not demonstrate that the induction of PGC-1 α expression was associated with myogenic differentiation or took place in pre-existing muscle fibers [Duguez et al., 2002].

The PPARs are known to be strongly involved in the control of genes encoding enzymes of the fatty acid oxidation pathway as e.g. HAD [Wang et al., 2003]. In concordance, we observed a clear induction of PPAR transcriptional activity and HAD activity during the initiation of myogenic differentiation. To the best of our knowledge, this is the first study investigating PPAR transcriptional activity during skeletal muscle cell differentiation.

Typically, PPAR- α and PPAR- δ are the 2 isoforms most implicated in positive regulation of fatty acid oxidative capacity of skeletal muscle, whereas PPAR- γ is the isoform implicated in fatty-acid uptake and storage [Singh et al., 2007, Fredenrich and Grimaldi, 2004, Lefebvre et al., 2006]. Therefore, it is conceivable that the increase in PPAR transcriptional activity is mainly due to increased PPAR- α and/or PPAR- δ transcriptional activity.

In concordance with this suggestion, we found that PPAR- γ mRNA levels markedly decreased during differentiation. It has been shown that constitutive high expression levels of PPAR- γ in skeletal muscle cells can result in trans-differentiation into adipocytes [Hu et al., 1995]. On the other hand, low critical levels of PPAR- γ expression are required to maintain the ability to differentiate properly [Singh et al., 2007]. Since C₂C₁₂ cell metabolism during differentiation is directed towards a more OXPHEN and PPAR- γ is implicated in storage of fatty acids, reduction of PPAR- γ expression may be a regulatory mechanism to ensure an adequate development of the oxidative capacity of the cell. In contrast to PPAR- γ , PPAR- α expression levels increased dramatically in our study. PPAR- δ mRNA levels on the other hand remained constant during the entire differentiation process despite its clearly established role in regulation and development of skeletal muscle oxidative capacity [Luquet et al., 2005, Luquet et al., 2003]. Kraft *et al.* previously showed a 10-fold increase in PPAR- α expression levels in differentiating C₂C₁₂ cells, although they only investigated the myogenic program up to 3 days of differentiation [Kraft et al., 2006]. We now show that, in line with enzyme activity data, PPAR- α mRNA expression levels increase until day 5 of differentiation reaching a 40-fold higher level in myotubes compared to myoblasts. To date, no other studies have previously investigated PPAR- δ expression levels during skeletal muscle differentiation.

The increase in expression levels of PGC-1 α and PPAR- α during differentiation parallels the observed increase in markers of OXPHEN. In addition to their well established role as positive regulators of oxidative capacity of existing muscle fibers [Lin et al., 2002, Wang et al., 2004, Canto et al., 2007], this therefore also suggests a role for these regulators in the development of oxidative capacity during myogenesis. Although whole body knockout PGC-1 α animals only display a 20-30% reduction in mitochondrial content [Leick et al., 2009, Leick et al., 2008], muscle-specific over-expression of PGC-1 α does result in an increase in basal OXPHEN [Lin et al., 2002]. This implies that PGC-1 α is a potent stimulator of mitochondrial biogenesis but may not always be required. Possibly, other co-factors in the PGC-1 α family as PRC and PGC-1 β compensate for the loss of PGC-1 α in the PGC-1 α knockouts [Scarpulla, 2008, Ishii et al., 2009]. It must also be noted that the PGC-1 α knockout mouse is characterized by physical hyperactivity which may cause exercise-induced mitochondrial biogenesis and potentially obscures any loss of baseline mitochondrial biogenesis [Lin et al., 2004]. In addition to PGC-1 α , our data also suggests a role for PPAR- α in mitochondrial biogenesis. It has been demonstrated that PPAR- α knockout mice display reduced activity levels of fatty acid oxidation enzymes as HAD, however no loss of other mitochondrial enzymes was observed [Bedu et al., 2007]. Future studies are therefore needed to address the requirement and the exact role of PPAR- α in skeletal muscle mitochondrial biogenesis. In contrast to PGC-1 α and PPAR- α , our data suggests that differential expression levels of PPAR- δ protein are not implicated in the process of mitochondrial biogenesis during myogenic differentiation. This is in line with a recent paper from Kleiner et al. who demonstrated that PPAR- δ agonism activates fatty acid oxidation, which is confirmed in mice with skeletal muscle-specific PPAR- δ over expression [Luquet et al., 2003], but fails to induce mitochondrial gene expression and function [Kleiner et al., 2009]. It can be speculated that PPAR- δ activation directs the metabolic profile of existing fibers towards a more

developed OXPHEN while PGC1 α -mediated PPAR- α activation may regulate development of oxidative capacity in new muscle fibers during myogenesis. Nevertheless, as PPAR- δ remains a specific regulator of fatty acid oxidation, which is part of the mitochondrial machinery, it still contributes to muscle OXPHEN. In line with a positive role in determining skeletal muscle OXPHEN, increased PGC-1 α and PPAR- α mRNA expression levels during differentiation may mediate the observed increase in transcription of nuclear-encoded constituents of the mitochondrial respiratory chain. Next to being a co-activator for PPARs, PGC-1 α also functions as a co-activator for NRF-1. Subsequently, PGC-1 α /NRF-1 induces the expression of Tfam which is the master regulator of mitochondrial biogenesis [Gleyzer et al., 2005]. In the present study, Tfam and NRF-1 expression levels increased at the onset of differentiation, which is in line with the start of mitochondrial biogenesis in these cells at the beginning of the myogenic program. Since Tfam is a master regulator of the transcription and replication of mtDNA [Gleyzer et al., 2005], its observed up-regulation likely mediated the increased mtDNA content during C₂C₁₂ myogenic differentiation. Additionally, we observed increased mRNA levels of mitochondrial-encoded COX subunits which may also result from increased transcription of the mitochondrial genome by Tfam.

In conclusion, in the present paper we describe temporal profiles of a number of key constituents of OXPHEN and mitochondrial capacity during the full myogenic program of C₂C₁₂ myoblasts. We show that expression and activity levels of genes known for their regulatory role in skeletal muscle oxidative capabilities parallel the observed increase in oxidative parameters. Based on our results we propose that increased PGC-1 α expression, either directly or through increased PPAR- α expression levels and PPAR transcriptional activity, mediates mitochondrial biogenesis and concurrent up regulation of skeletal muscle OXPHEN during myogenic differentiation.

Acknowledgements

The authors like to thank E. Phielix for excellent technical support in measurements of cellular respiration and M. Kelders for technical assistance in QPCR analysis.

Funding

The research of Dr. H. Gosker is supported by a grant from the Netherlands Asthma Foundation (NAF 3.4.05.038). The research of Dr. P. Schrauwen has been made possible by fellowships of the Royal Netherlands Academy of Arts and Sciences. Dr. Langen was sponsored by a VENI grant (VENI 916.56.112). A. Remels is supported by a grant from Numico Research.

Disclosures

Funding sources had no involvement in collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication. The authors would like to disclose that there are no advisory board affiliations or financial or personal interests in any organization sponsoring the research and accept full responsibility for conduct of the study and the decision to publish the present work. We declare that the manuscript, including related data, figures and tables, has not been previously reported or published, that the manuscript is not under consideration elsewhere, and that it will not be submitted elsewhere while under review by the journal of molecular cellular endocrinology.

Figure legends

Figure 1: **Mitochondrial respiration in C₂C₁₂ myoblasts vs differentiated myotubes.** Mitochondrial respiration was measured in proliferating myoblasts and in myotubes which were differentiated for 7 days in DM. Cells were harvested and oxygen consumption was measured. Basal respiration (on endogenous substrates) was determined after adding the cells to the chambers. After permeabilization with digitonin (1 μ l; 0.5 μ g/ml), the following substrates were added to the cells: malate (5 μ l; 2 mM), glutamate (10 μ l; 10 mM) or pyruvate (5 μ l; 5 mM) or palmitoyl-carnitine (10 μ l; 50 μ M) followed by ADP (20 μ l; 5 mM) and oligomycin (1 μ l; 1.66 μ g/ml). Maximal respiration was determined after titration with FCCP (1 μ l; 0.5 μ M titration). Respiration rates were corrected for total protein content. Data are represented as mean \pm SD for 3 independent experiments. Oxygen consumption is given in pmol/mg·s. Significance of difference: * $p \leq 0.050$

Figure 2: **Mitochondrial content during myogenesis.** C₂C₁₂ cells were harvested at different time points during differentiation (day 0, 3, 5 and 7) and processed for Western blotting, enzyme activity measurements, mRNA abundance and mtDNA copy number determination. Data are represented as mean \pm SD from triplicate samples from 3 independent experiments. A) Enzyme activity levels, B) Protein levels of the different complexes of the respiratory chain are given as arbitrary units (AU) in % of levels in myoblasts. C) mitochondrial copy number and D) expression levels of mitochondrial- (COX II and COX III) and nuclear- (COX Va and COX VIIc) encoded subunits of COX are expressed as arbitrary units (AU) after correction for the housekeeping gene cyclophilinA. Significance of difference compared to myoblasts (day 0): * $p \leq 0.050$.

Figure 3: **Myosin heavy chain (MyHC) distribution in C₂C₁₂ cells during differentiation.** C₂C₁₂ cells were harvested at different time points during differentiation (day 0, 3, 5 and 7) and MyHC isoform protein levels were quantified by Western blot analysis. Protein levels of MyHC isoforms were corrected for total protein content. Data are represented as mean \pm SD from triplicate samples from 3 independent experiments. Protein levels are given as arbitrary units (AU). Quantified MyHC protein levels are expressed as the ratio between the slow and fast isoform. Significance of difference compared to myoblasts (day 0): * $p \leq 0.050$.

Figure 4: **Expression of PGC-1 α , PPAR- α , PPAR- δ , and PPAR- γ mRNA in C₂C₁₂ skeletal muscle cells during differentiation.** C₂C₁₂ cells were differentiated in DM for 0, 3, 5, or 7 days and harvested. Expression levels of PGC-1 α (A), PPAR- α (B), PPAR- δ (C) and PPAR- γ (D) were corrected for cyclophilinA and expressed as arbitrary units (AU). Data are represented as mean \pm SD from triplicate samples from 3 independent experiments. Similar results were observed when corrected for other individual housekeeping genes as HPRT, β -actin or GAPDH or when the geNorm normalization factor (based on 4 pooled housekeepers) was used. Significance of difference compared to myoblasts: * $p \leq 0.050$.

Figure 5: **Expression of Tfam and NRF-1 mRNA in C₂C₁₂ skeletal muscle cells during differentiation.** Cells were differentiated in DM for 0, 3, 5, or 7 days and harvested. Expression levels of Tfam (A) and NRF-1 (B) were corrected for cyclophilinA as a representative housekeeper. Expression levels are expressed as % of day 0. Data are represented as mean \pm SD from triplicate samples from 3 independent experiments. Similar results were observed when corrected for individual housekeeping genes as HPRT, β -actin or

GAPDH or when the geNorm normalization factor based on 4 different housekeepers was used. Significance of difference compared to myoblasts: * $p \leq 0.050$.

Figure 6: **PPAR-dependent transcriptional activity in C₂C₁₂ skeletal muscle cells during differentiation.** C₂C₁₂ cells stably containing a PPAR sensitive reporter construct were differentiated in DM for 0, 3, 5, or 7 days and harvested. Cells were lysed; luciferase activity was determined and normalized to total protein content. Significance of difference compared to myoblasts: * $p \leq 0.050$. Error bars indicate the SD. Shown are the mean \pm SD of 3 independent experiments.

Table legends

Table 1: Peroxisome proliferator-activated receptor: PPAR; PPAR- γ co-activator 1 alpha: PGC-1 α ; Mitochondrial transcription factor A: Tfam; Nuclear respiratory factor 1: NRF-1; Cytochrome *c* oxidase: COX; Glyceraldehyde 3-phosphate dehydrogenase: GAPDH; Hypoxanthine phosphoribosyltransferase: HPRT.

Table 2: Values are expressed as mean \pm SD from 3 independent experiments. Significance of difference compared to myoblasts: * $p < 0.001$. MyHC: myosin heavy chain; CK: creatine kinase; Myogenic index: the number of nuclei residing in cells containing three or more nuclei divided by the total number of nuclei in May-Grunwald Giemsa stained cells.

References

- [1] D.G. Altman, S.M. Gore, M.J. Gardner and S.J. Pocock, Statistical guidelines for contributors to medical journals, *Br Med J (Clin Res Ed)* **286** (1983), 1489-93.
- [2] J.E. Anderson, The satellite cell as a companion in skeletal muscle plasticity: currency, conveyance, clue, connector and colander, *J Exp Biol* **209** (2006), 2276-92.
- [3] K. Baar, Involvement of PPAR gamma co-activator-1, nuclear respiratory factors 1 and 2, and PPAR alpha in the adaptive response to endurance exercise, *Proc Nutr Soc* **63** (2004), 269-73.
- [4] K.M. Baldwin and F. Haddad, Skeletal muscle plasticity: cellular and molecular responses to altered physical activity paradigms, *Am J Phys Med Rehabil* **81** (2002), S40-51.
- [5] E. Bedu, D. Desplanches, J. Pequignot, B. Bordier and B. Desvergne, Double gene deletion reveals the lack of cooperation between PPARalpha and PPARbeta in skeletal muscle, *Biochem Biophys Res Commun* **357** (2007), 877-81.
- [6] R.A. Butow and E.M. Bahassi, Adaptive thermogenesis: orchestrating mitochondrial biogenesis, *Curr Biol* **9** (1999), R767-9.
- [7] C. Canto, S. Pich, J.C. Paz, R. Sanches, V. Martinez, M. Orpinell, M. Palacin, A. Zorzano and A. Guma, Neuregulins increase mitochondrial oxidative capacity and insulin sensitivity in skeletal muscle cells, *Diabetes* **56** (2007), 2185-93.
- [8] S. Duguez, L. Feasson, C. Denis and D. Freyssenet, Mitochondrial biogenesis during skeletal muscle regeneration, *Am J Physiol Endocrinol Metab* **282** (2002), E802-9.
- [9] S. Duguez, O. Sabido and D. Freyssenet, Mitochondrial-dependent regulation of myoblast proliferation, *Exp Cell Res* **299** (2004), 27-35.
- [10] M. Fluck, Functional, structural and molecular plasticity of mammalian skeletal muscle in response to exercise stimuli, *J Exp Biol* **209** (2006), 2239-48.

- [11] A. Fredenrich and P.A. Grimaldi, Roles of peroxisome proliferator-activated receptor delta in skeletal muscle function and adaptation, *Curr Opin Clin Nutr Metab Care* **7** (2004), 377-81.
- [12] N. Gleyzer, K. Vercauteren and R.C. Scarpulla, Control of mitochondrial transcription specificity factors (TFB1M and TFB2M) by nuclear respiratory factors (NRF-1 and NRF-2) and PGC-1 family coactivators, *Mol Cell Biol* **25** (2005), 1354-66.
- [13] H.R. Gosker, P. Schrauwen, R. Broekhuizen, M.K. Hesselink, E. Moonen-Kornips, K.A. Ward, F.F. Franssen, E.F. Wouters and A.M. Schols, Exercise training restores uncoupling protein-3 content in limb muscles of patients with chronic obstructive pulmonary disease, *Am J Physiol Endocrinol Metab* (2005).
- [14] H.R. Gosker, H. van Mameren, P.J. van Dijk, M.P. Engelen, G.J. van der Vusse, E.F. Wouters and A.M. Schols, Skeletal muscle fibre-type shifting and metabolic profile in patients with chronic obstructive pulmonary disease, *Eur Respir J* **19** (2002), 617-25.
- [15] N.H. Herzberg, R. Zwart, R.A. Wolterman, J.P. Ruiter, R.J. Wanders, P.A. Bolhuis and C. van den Bogert, Differentiation and proliferation of respiration-deficient human myoblasts, *Biochim Biophys Acta* **1181** (1993), 63-7.
- [16] E. Hu, P. Tontonoz and B.M. Spiegelman, Transdifferentiation of myoblasts by the adipogenic transcription factors PPAR gamma and C/EBP alpha, *Proc Natl Acad Sci U S A* **92** (1995), 9856-60.
- [17] K.A. Ishii, T. Fumoto, K. Iwai, S. Takeshita, M. Ito, N. Shimohata, H. Aburatani, S. Taketani, C.J. Lelliott, A. Vidal-Puig and K. Ikeda, Coordination of PGC-1beta and iron uptake in mitochondrial biogenesis and osteoclast activation, *Nat Med* **15** (2009), 259-66.

- [18] V.E. Jahnke, O. Sabido and D. Freyssenet, Control of mitochondrial biogenesis, ROS level, and cytosolic Ca²⁺ concentration during the cell cycle and the onset of differentiation in L6E9 myoblasts, *Am J Physiol Cell Physiol* **296** (2009), C1185-94.
- [19] K. Kim, A. Lecordier and L.H. Bowman, Both nuclear and mitochondrial cytochrome c oxidase mRNA levels increase dramatically during mouse postnatal development, *Biochem J* **306** (Pt 2) (1995), 353-8.
- [20] S. Kleiner, V. Nguyen-Tran, O. Bare, X. Huang, B. Spiegelman and Z. Wu, PPAR{delta} Agonism Activates Fatty Acid Oxidation via PGC-1{alpha} but Does Not Increase Mitochondrial Gene Expression and Function, *J Biol Chem* **284** (2009), 18624-33.
- [21] T.R. Koves, P. Li, J. An, T. Akimoto, D. Slentz, O. Ilkayeva, G.L. Dohm, Z. Yan, C.B. Newgard and D.M. Muoio, Peroxisome proliferator-activated receptor-gamma co-activator 1alpha-mediated metabolic remodeling of skeletal myocytes mimics exercise training and reverses lipid-induced mitochondrial inefficiency, *J Biol Chem* **280** (2005), 33588-98.
- [22] C.S. Kraft, C.M. LeMoine, C.N. Lyons, D. Michaud, C.R. Mueller and C.D. Moyes, Control of mitochondrial biogenesis during myogenesis, *Am J Physiol Cell Physiol* **290** (2006), C1119-27.
- [23] R.C. Langen, A.M. Schols, M.C. Kelders, E.F. Wouters and Y.M. Janssen-Heininger, Enhanced myogenic differentiation by extracellular matrix is regulated at the early stages of myogenesis, *In Vitro Cell Dev Biol Anim* **39** (2003), 163-9.
- [24] F. Le Grand and M.A. Rudnicki, Skeletal muscle satellite cells and adult myogenesis, *Curr Opin Cell Biol* **19** (2007), 628-33.

- [25] S.C. Leary, B.J. Battersby, R.G. Hansford and C.D. Moyes, Interactions between bioenergetics and mitochondrial biogenesis, *Biochim Biophys Acta* **1365** (1998), 522-30.
- [26] P. Lefebvre, G. Chinetti, J.C. Fruchart and B. Staels, Sorting out the roles of PPAR alpha in energy metabolism and vascular homeostasis, *J Clin Invest* **116** (2006), 571-80.
- [27] L. Leick, Y. Hellsten, J. Fentz, S.S. Lyngby, J.F. Wojtaszewski, J. Hidalgo and H. Pilegaard, PGC-1{alpha} mediates exercise-induced skeletal muscle VEGF expression in mice, *Am J Physiol Endocrinol Metab* **297** (2009), E92-E103.
- [28] L. Leick, J.F. Wojtaszewski, S.T. Johansen, K. Kiilerich, G. Comes, Y. Hellsten, J. Hidalgo and H. Pilegaard, PGC-1alpha is not mandatory for exercise- and training-induced adaptive gene responses in mouse skeletal muscle, *Am J Physiol Endocrinol Metab* **294** (2008), E463-74.
- [29] J. Lin, H. Wu, P.T. Tarr, C.Y. Zhang, Z. Wu, O. Boss, L.F. Michael, P. Puigserver, E. Isotani, E.N. Olson, B.B. Lowell, R. Bassel-Duby and B.M. Spiegelman, Transcriptional co-activator PGC-1 alpha drives the formation of slow-twitch muscle fibres, *Nature* **418** (2002), 797-801.
- [30] J. Lin, P.H. Wu, P.T. Tarr, K.S. Lindenberg, J. St-Pierre, C.Y. Zhang, V.K. Mootha, S. Jager, C.R. Vianna, R.M. Reznick, L. Cui, M. Manieri, M.X. Donovan, Z. Wu, M.P. Cooper, M.C. Fan, L.M. Rohas, A.M. Zavacki, S. Cinti, G.I. Shulman, B.B. Lowell, D. Krainc and B.M. Spiegelman, Defects in adaptive energy metabolism with CNS-linked hyperactivity in PGC-1alpha null mice, *Cell* **119** (2004), 121-35.
- [31] S. Luquet, C. Gaudel, D. Holst, J. Lopez-Soriano, C. Jehl-Pietri, A. Fredenrich and P.A. Grimaldi, Roles of PPAR delta in lipid absorption and metabolism: a new target for the treatment of type 2 diabetes, *Biochim Biophys Acta* **1740** (2005), 313-7.

- [32] S. Luquet, J. Lopez-Soriano, D. Holst, A. Fredenrich, J. Melki, M. Rassoulzadegan and P.A. Grimaldi, Peroxisome proliferator-activated receptor delta controls muscle development and oxidative capability, *Faseb J* **17** (2003), 2299-301.
- [33] O. Mathieu-Costello and R.T. Hepple, Muscle structural capacity for oxygen flux from capillary to fiber mitochondria, *Exerc Sport Sci Rev* **30** (2002), 80-4.
- [34] C.D. Moyes, O.A. Mathieu-Costello, N. Tsuchiya, C. Filburn and R.G. Hansford, Mitochondrial biogenesis during cellular differentiation, *Am J Physiol* **272** (1997), C1345-51.
- [35] M.V. Narici, N.D. Reeves, C.I. Morse and C.N. Maganaris, Muscular adaptations to resistance exercise in the elderly, *J Musculoskelet Neuronal Interact* **4** (2004), 161-4.
- [36] K. Oexle and A. Kohlschutter, Cause of progression in Duchenne muscular dystrophy: impaired differentiation more probable than replicative aging, *Neuropediatrics* **32** (2001), 123-9.
- [37] P. Rochard, A. Rodier, F. Casas, I. Cassar-Malek, S. Marchal-Victorion, L. Daury, C. Wrutniak and G. Cabello, Mitochondrial activity is involved in the regulation of myoblast differentiation through myogenin expression and activity of myogenic factors, *J Biol Chem* **275** (2000), 2733-44.
- [38] R.C. Scarpulla, Nuclear control of respiratory chain expression by nuclear respiratory factors and PGC-1-related coactivator, *Ann N Y Acad Sci* **1147** (2008), 321-34.
- [39] R.C. Scarpulla, Transcriptional activators and coactivators in the nuclear control of mitochondrial function in mammalian cells, *Gene* **286** (2002), 81-9.
- [40] P. Seyer, S. Grandemange, M. Busson, A. Carazo, F. Gamaleri, L. Pessemesse, F. Casas, G. Cabello and C. Wrutniak-Cabello, Mitochondrial activity regulates myoblast differentiation by control of c-Myc expression, *J Cell Physiol* **207** (2006), 75-86.

- [41] J. Singh, N.K. Verma, S.M. Kansagra, B.N. Kate and C.S. Dey, Altered PPARgamma expression inhibits myogenic differentiation in C2C12 skeletal muscle cells, *Mol Cell Biochem* **294** (2007), 163-71.
- [42] J. Vandesompele, K. De Preter, F. Pattyn, B. Poppe, N. Van Roy, A. De Paepe and F. Speleman, Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes, *Genome Biol* **3** (2002), RESEARCH0034.
- [43] G.Q. Wallace and E.M. McNally, Mechanisms of muscle degeneration, regeneration, and repair in the muscular dystrophies, *Annu Rev Physiol* **71** (2009), 37-57.
- [44] Y.X. Wang, C.H. Lee, S. Tjep, R.T. Yu, J. Ham, H. Kang and R.M. Evans, Peroxisome-proliferator-activated receptor delta activates fat metabolism to prevent obesity, *Cell* **113** (2003), 159-70.
- [45] Y.X. Wang, C.L. Zhang, R.T. Yu, H.K. Cho, M.C. Nelson, C.R. Bayuga-Ocampo, J. Ham, H. Kang and R.M. Evans, Regulation of muscle fiber type and running endurance by PPARdelta, *PLoS Biol* **2** (2004), e294.
- [46] Z. Wu, P. Puigserver, U. Andersson, C. Zhang, G. Adelmant, V. Mootha, A. Troy, S. Cinti, B. Lowell, R.C. Scarpulla and B.M. Spiegelman, Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1, *Cell* **98** (1999), 115-24.
- [47] D. Yaffe and O. Saxel, Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle, *Nature* **270** (1977), 725-7.


Table 1: Primer details


QPCR primer sequences		
Primer	Forward 5'-3'	Reverse 5'-3'
PPAR- α	AGTGCCCTGAACATCGAGTGT	AAGCCCTTACAGCCTTCACATG
PPAR- δ	GAGCAAGTGGCCACGGGTGAC	CCACCTGAGGCCCATCACAG
PPAR- γ	TGGGGATGTCTCACAAATGCCA	TTCCTGTCAAGATCGCCCTCG
PGC-1 α	AAACCACACCCACAGGATCAG	TCTTCGCTTTATTGCTCCATGA
Tfam	CGGAGTTCACGCTGG	CCGTCCGGTTCGTCTCAC
NRF-1	CCTCCAAACCTGCATCTCAC	TCGCACCACATTCTCCAAAG
COXVa	TGCGAGCATGTAGACGGTAAAT	GAGGCCTGCTTTGTCCTTAACA
COXVIIc	GGCCAGAGTATCCGGAGGTT	CCACTGAAAATGGCAAATTCTTC
COXII	CCATCCCAGCCCGACTAA	ATTCAGAGCATTGGCCATAGAA
COXIII	GGTCTCGGAAGTATTTTTCTTTGC	CAGCAGCCTCCTAGATCATGTG
CyclophilinA	TTCCTCCTTTCACAGAATTATTCCA	CCGCCAGTGCCATTATGG
GAPDH	CAACTCACTCAAGATTGTCAGCAA	TGGCAGTGATGGCATGGA
HPRT	TGGATATGCCCTTGACTATAATGAGTAC	AGGACTCCTCGTATTTGCAGATTC
β -actin	CTGAATGGCCCAGGTCTGA	CCCTCCCAGGGAGACCAA

Table 2: Parameters of myogenic differentiation

	Myoblasts	Myotubes (day 3)	Myotubes (day 7)
MyHC fast (AU)	0	1.7 ± 0.4 *	3.9 ± 0.6 *
CK (U/mg protein)	0.010 ± 0.003	0.081 ± 0.023 *	0.734 ± 0.087 *
Myogenic index	0	0.22 ± 0.04*	0.51 ± 0.08*

trip





trip

