



HAL
open science

First Stars XIV. Sulphur abundances in extremely metal-poor (EMP) stars

Monique Spite, E. Caffau, S.M. Andrievsky, S. A. Korotin, E. Depagne, F. Spite, P. Bonifacio, H.-G. Ludwig, R. Cayrel, P. Francois, et al.

► **To cite this version:**

Monique Spite, E. Caffau, S.M. Andrievsky, S. A. Korotin, E. Depagne, et al.. First Stars XIV. Sulphur abundances in extremely metal-poor (EMP) stars. 2010. hal-00548834v1

HAL Id: hal-00548834

<https://hal.science/hal-00548834v1>

Preprint submitted on 20 Dec 2010 (v1), last revised 7 Jan 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First Stars XIV. Sulphur abundances in extremely metal-poor stars [★]

M. Spite¹, E. Caffau^{2,1}, S.M. Andrievsky^{1,3}, S.A. Korotin³, E. Depagne⁴, F. Spite¹, P. Bonifacio¹, H.-G. Ludwig², R. Cayrel¹, P. François¹, V. Hill⁵, B. Plez⁶, J. Andersen^{7,8}, B. Barbuy⁹, T. C. Beers¹⁰, P. Molaro¹¹, B. Nordström⁷, and F. Primas¹²

¹ GEPI Observatoire de Paris, CNRS, Université Paris Diderot, F-92195 Meudon Cedex France e-mail : monique.spite@obspm.fr

² Zentrum für Astronomie der Universität Heidelberg, Landessternwarte, Königstuhl 12, 69117 Heidelberg, Germany

³ Department of Astronomy and Astronomical Observatory, Odessa National University, and Isaac Newton Institute of Chile, Odessa branch, Shevchenko Park, 65014, Odessa, Ukraine

⁴ Las Cumbres Observatory, Goleta, CA 93117, USA

⁵ Observatoire de la Côte d'Azur, CNRS UMR6202, BP4229, 06304 Nice Cedex 4, France

⁶ GRAAL, Université de Montpellier II, F-34095 Montpellier Cedex 05, France

⁷ The Niels Bohr Institute, Astronomy, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

⁸ Nordic Optical Telescope, Apartado 474, E-38700 Santa Cruz de La Palma, Spain

⁹ Universidade de São Paulo, Departamento de Astronomia, Rua do Matão 1226, 05508-900 São Paulo, Brazil

¹⁰ Department of Physics & Astronomy and JINA: Joint Institute for Nuclear Astrophysics, Michigan State University, East Lansing, MI 48824, USA

¹¹ Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34143 Trieste, Italy

¹² European Southern Observatory (ESO), Karl-Schwarzschild-Str. 2, D-85748 Garching b. München, Germany

ABSTRACT

Context. Sulphur is important in the study of the early chemical evolution of the Galaxy, because the site of its formation remains uncertain, since at low metallicity the trend of this α -element against $[\text{Fe}/\text{H}]$ is controversial. Different authors find $[\text{S}/\text{Fe}]$ to remain constant or decrease with $[\text{Fe}/\text{H}]$ below $[\text{Fe}/\text{H}]=-2.0$, depending on the multiplet used in the analysis. Moreover, although sulphur is not significantly bound in dust grains in the ISM, it seems to behave differently in DLAs and in old metal-poor stars.

Aims. We aim here to perform a precise measurement of the S abundance in a sample of extremely metal-poor stars observed with the ESO VLT equipped with UVES, taking into account NLTE and 3D effects.

Methods. NLTE profiles of the lines of multiplet 1 of S I have been computed with a version of the program MULTI (including opacity sources from ATLAS9, and based on a new model atom for S), and fitted to the observed spectra.

Results. We find sulphur in EMP stars to behave like the other α -elements, with $[\text{S}/\text{Fe}]$ remaining approximately constant below $[\text{Fe}/\text{H}]=-3$. However, $[\text{S}/\text{Mg}]$ seems to decrease slightly as a function of $[\text{Mg}/\text{H}]$. The overall abundance patterns of O, Na, Mg, Al, S, and K are best matched by the SN model yields by Heger & Woosley. The $[\text{S}/\text{Zn}]$ ratio in EMP stars is solar, as found also in DLAs. We obtain an upper limit on the abundance of sulphur, $[\text{S}/\text{Fe}] < +0.5$, for the ultra metal-poor star CS 22949-037. This, along with a previous reported measurement of zinc, argues against the conjecture that the light-element abundances pattern in this star, and, by analogy, the hyper metal-poor stars HE 0107-5240 and HE 1327-2326, are due to dust depletion.

Key words. Galaxy: abundances – Galaxy: halo – Galaxy: evolution – Stars: abundances – Stars: Supernovae

1. Introduction

The determination of the abundance of sulphur in the atmosphere of old Galactic stars is of interest for two primary reasons:

* Based on observations obtained with the ESO Very Large Telescope at Paranal Observatory, Chile (Large Programme "First Stars", ID 165,N-0276, P.I.: R. Cayrel.

-First, since unlike most other heavy elements, sulphur is generally considered not to be significantly bound in interstellar dust grains, it is a good indicator of the chemical composition of the ISM and thus allows a direct comparison between the abundances in the DLA's (Damped Ly α clouds) and in early Galactic matter.

-Secondly, the formation of sulphur is still controversial. Sulphur is generally considered as an α -capture element like

magnesium and calcium, formed preferentially in massive Type II supernovae, including those of the first generations of stars. In analogy with other α -elements, [S/Fe] is expected to be constant and positive at low metallicity (see e.g. François, 1987, 1988). Yet, the trend of [S/Fe] vs. [Fe/H] in the early Galaxy is still debated: Israelian & Rebolo (2001) and Takada-Hidai et al. (2002) found [S/Fe] to rise with decreasing [Fe/H], while Ryde & Lambert (2004) and Nissen et al. (2004, 2008) found [S/Fe] to remain flat. Finally, Caffau et al. (2005, 2010a), combining their own data with a large sample from the literature covering multiplets 1, 3, 6, and 8 of S I (see Table 1), suggest a bimodal distribution of [S/Fe] for [Fe/H] < -1.0 to explain the high value of [S/Fe] found in several metal-poor stars.

Table 1. Parameters of the sulphur lines

Mult	wavelength (nm) air	Transition	log gf	χ_{ex} (eV)
1	921.2863	4s-4p $^5S_2^0 - ^5P_3$	0.42	6.525
1	922.8093	$^5S_2^0 - ^5P_2$	0.26	6.525
1	923.7538	$^5S_2^0 - ^5P_1$	0.04	6.525
3	1045.5449	$^3S_1^0 - ^3P_2$	0.26	6.860
3	1045.6757	$^3S_1^0 - ^3P_0$	-0.43	6.860
3	1045.9406	$^3S_1^0 - ^3P_1$	0.04	6.860
6	869.3931	4p-4d $^5P_3 - ^5D_3^0$	-0.51	7.870
6	869.4626	$^5P_3 - ^5D_4^0$	0.08	7.870
8	675.6851	4p-5d $^5P_3 - ^5D_2^0$	-1.76	7.870
8	675.7007	$^5P_3 - ^5D_3^0$	-0.90	7.870
8	675.7171	$^5P_3 - ^5D_4^0$	-0.31	7.870
1F	1082.1176	$^3P - ^1D$	-8.62	0.000

A rise of [S/Fe] with decreasing [Fe/H] has no simple explanation. It has been suggested that SNe with large explosion energies (hypernovae) could have made a substantial contribution to the nucleosynthesis of elements in the early Galaxy (Israelian & Rebolo 2001, Nakamura et al. 2001), or even that it could be related to a delayed deposition of the supernova-synthesized products into the interstellar medium (Ramaty et al. 2000).

On the other hand, Takeda et al. (2005) noted that, even taking into account non-LTE effects on the formation of the S lines, the result depended on the multiplet used for the determination of the sulphur abundance. There was, in particular, an apparent discrepancy between the abundance trends deduced from multiplets 6 and 1.

Taking advantage of a new model atom of sulphur (Korotin 2008, 2009), based on new radiative photoionization rates, we here report new S abundances for 33 very metal-poor stars, including 21 stars with [Fe/H] < -2.9 (Extremely Metal-Poor or EMP stars). Our study is based on a precise high-resolution NLTE analysis of the lines of the Mult. 1 of S I at $\lambda\lambda$ 921.29, 922.81, and 923.75 nm (see Table 1).

2. Observations and Reduction

All stars were observed as part of the ESO large programme “First Stars”, using the high-resolution spectrograph UVES (Dekker et al. 2000) at the ESO-VLT. The spectra have been discussed in detail in Cayrel et al. (2004) and Bonifacio et al. (2007, 2009). The resolving power of the spectra is $R \approx 43\,000$ with 5 pixels per spectral resolution element. The spectra were reduced using the UVES context within MIDAS (Ballester et al. 2000). The S/N ratio per pixel in the region of the S I lines is typically ~ 180 for the giants, ~ 90 for the dwarfs (or 400 and 200 per resolution element, see Cayrel et al., 2004).

Three Ba-poor giants (McWilliam et al. 1995) were also observed as part of the programme, under similar conditions as the other metal-poor stars (see also Andrievsky et al. 2011, to be subm. to A&A). The S abundance could be determined for one of them (CS 22949-048), which has been added to the sample.

3. Choice of sulphur lines and determination of fundamental atmospheric parameters

In cool stars, five multiplets of S I can be observed in the red and near-infrared spectral region (Table 1): Mult. 1 at about 922 nm, Mult. 3 at 1045 nm, Mult. 6 at 869 nm, and Mult. 8 at 675 nm (Caffau et al. 2005, 2010a), and Mult. 1F, the forbidden line at 1082.1 nm (Ryde, 2006, Caffau & Ludwig, 2007). However, in EMP stars, the lines of multiplets 6 and 8 are too weak for precise analysis (see Fig. 1). The lines of Mult. 3 and Mult. 1F, are non detectable in stars with [Fe/H] < -2.9, and are in any case outside the range of our spectra. The S abun-




Fig. 1. Spectra of BD+17 3248 and HD 2796, showing the line at 869.46 nm in Mult. 6 of S I. We detect the line in BD+17 3248 ([Fe/H]=-2.1), but not in HD 2796 ([Fe/H]=-2.5). A synthetic spectrum (full line) corresponding to the equivalent width of 0.26 pm found by Israelian & Rebolo for this line has been superimposed on our spectrum of HD 2796.

dances reported in this paper are therefore derived from the lines of Mult. 1 at 921.286, 922.809, and 923.754 nm.




Fig. 2. Spectrum of the giant star CS 29491-053 ($[\text{Fe}/\text{H}] = -3.0$) in the region of the S I lines of multiplet 1 (wavelength in nm). The synthetic spectrum (red line) has been computed for the adopted abundance (Table 3), and for an abundance 2.5 times smaller (thin blue line). The S line at 922.8 nm lies in the wing of the hydrogen line $P\zeta$; all other lines in this region are of telluric origin. In our analysis, all S lines blended with telluric lines have been rejected.

The line at 922.809 nm is located in the wing of the Paschen ζ line, and the absorption from the hydrogen line has been taken into account in the computations (see Fig. 2). This spectral region is also crowded with telluric absorption lines of H_2O . Since no hot star was observed together with the programme stars, we have not tried to remove these telluric lines, but we have carefully computed their position relative to the stellar lines and only used S I lines free of contamination. Generally, at least one or two lines in each spectrum, are free from blending.

Table 2 provides equivalent widths of the S I lines, since they were not published in Cayrel et al. (2004) or Bonifacio et al. (2009). In this spectral region the observed spectra are affected by fringing, which is not completely compensated for by division by a flat field (Nissen et al. 2008). The level of the continuum is therefore often not well defined, and we estimate that the error on the equivalent width can reach 0.4 pm for giants and 0.5 pm for dwarfs. Since in all the cases the sulphur lines in Table 2 are “weak”, the abundances, as a first approx-

Table 2. Equivalent widths (pm) of the S I lines of multiplet 1.

Wavelength (nm)	921.29	922.81	923.75
χ_{ex} (eV)	6.525	6.525	6.525
$\log gf$	0.42	0.26	0.04
	EW (pm)	EW (pm)	EW (pm)
Turnoff stars			
BS 16968-061	1.68	-	-
BS 17570-063	1.30	-	-
CS 22965-054	1.30	-	-
CS 29499-060	1.80	-	-
CS 29506-007	1.98	-	-
CS 29506-090	1.63	-	-
CS 30301-024	1.95	-	-
Giants			
HD 2796	4.80	3.20	2.76
HD 122563	-	1.92	1.29
HD 186478	-	2.70	1.68
BD+17 3248	-	4.88	-
BD-18 5550	1.65	-	-
BS 17569-049	-	1.20	1.20
CS 22169-035	-	1.00	(0.76) †
CS 22186-025	(2.09)	1.52	0.79
CS 22189-009	0.72	-	0.32
CS 22873-055	2.37	-	-
CS 22873-166	2.00	1.24	1.08
CS 22878-101	-	-	0.72
CS 22891-209	1.28	(1.00)	-
CS 22892-052	1.18	-	-
CS 22896-154	1.98	-	0.79
CS 22897-008	0.90	-	-
CS 22948-066	-	1.20	-
CS 22953-003	1.43	-	0.74
CS 22956-050	1.08	-	-
CS 22966-057	2.49	1.04	1.15
CS 29491-053	1.53	1.02	0.88
CS 29502-042	0.83	-	-
CS 29518-051	2.30	1.34	-
CS 30325-094	1.33	0.98	-
CS 31082-001	2.19	1.90	0.97
CS 22949-048	-	0.55	0.45

† Equivalent widths in parenthesis are very uncertain

imation, are affected proportionally to the EWs of the lines: $\Delta(\log ab) = \Delta(\log EW)$. However, these equivalent widths have not been used in the computations since we have directly compared synthetic and observed spectra; they are given here only for comparison.

In total, we could thus determine S abundances in 26 giants and 7 turnoff stars from our original sample.

4. NLTE calculations and 3D correction

Atmospheric parameters of the stars (T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$) are adopted from Cayrel et al. (2004), Hill et al. (2002) and Bonifacio et al. (2007); for the Ba-poor star: from Andrievsky et al. (2011). Briefly, the temperature of the giants has been deduced from their colours, adopting the calibration of Alonso et al. (1996), and the temperature of the turnoff stars from the

Table 3. Adopted atmospheric parameters and NLTE S abundances for our sample of EMP stars. The solar abundance of sulphur has been taken from Caffau et al.: $\log\epsilon(\text{S})_{\odot} = 7.16$, the NLTE Mg abundance (column 6) is from Andrievsky et al., (2010). The error (column 8) represents the uncertainty on the spectral fitting. The last column gives the number of sulphur lines used in the determination

Star name	T_{eff}	$\log g$	v_t	[Fe/H]	NLTE [Mg/H]	LTE $\log\epsilon(\text{S})$	err. \pm	NLTE $\log\epsilon(\text{S})$	3D corr.	3D+ NLTE [S/H]	3D+ NLTE [S/Fe]	3D+ NLTE [S/Mg]	N
Turnoff stars													
BS 16968–061	6040	3.8	1.5	-3.05	-2.46	4.90	0.20	4.33	0.14	-2.69	0.36	-0.23	1
BS 17570–063	6240	4.8	0.5	-2.92	-2.52	5.05	0.20	4.61	0.12	-2.43	0.49	0.09	1
CS 22965–054	6090	3.8	1.4	-3.04	-2.42	4.74	0.20	4.21	0.13	-2.82	0.22	-0.40	1
CS 29499–060	6320	4.0	1.5	-2.70	-2.14	4.90	0.20	4.50	0.12	-2.54	0.16	-0.40	1
CS 29506–007	6270	4.0	1.7	-2.91	-2.25	5.03	0.20	4.62	0.12	-2.44	0.47	-0.19	1
CS 29506–090	6300	4.3	1.4	-2.83	-2.22	4.98	0.20	4.56	0.12	-2.48	0.35	-0.26	1
CS 30301–024	6330	4.0	1.6	-2.75	-2.25	4.98	0.20	4.60	0.12	-2.44	0.31	-0.19	1
Giants													
HD 2796	4950	1.5	2.1	-2.47	-1.84	5.25	0.10	4.97	0.07	-2.12	0.35	-0.28	3
HD 122563	4600	1.1	2.0	-2.82	-2.19	4.87	0.10	4.70	0.06	-2.40	0.42	-0.21	2
HD 186478	4700	1.3	2.0	-2.59	-1.86	5.07	0.08	4.84	0.06	-2.26	0.33	-0.40	2
BD+17 3248	5250	1.4	1.5	-2.07	-1.39	5.57	0.10	5.11	0.10	-1.95	0.12	-0.56	2
BD-18 5550	4750	1.4	1.8	-3.06	-2.44	4.65	0.12	4.41	0.07	-2.68	0.38	-0.24	1
BS 17569–049	4700	1.2	1.9	-2.88	-2.11	4.70	0.20	4.40	0.06	-2.70	0.18	-0.59	2
CS 22169–035	4700	1.2	2.2	-3.04	-2.66	4.50	0.11	4.20	0.05	-2.91	0.13	-0.25	1
CS 22186–025	4900	1.5	2.0	-3.00	-2.39	4.70	0.10	4.43	0.06	-2.67	0.33	-0.28	2
CS 22189–009	4900	1.7	1.9	-3.49	-3.11	4.25	0.20	3.97	0.07	-3.12	0.37	-0.01	2
CS 22873–055	4550	0.7	2.2	-2.99	-2.34	4.72	0.10	4.52	0.10	-2.54	0.45	-0.20	1
CS 22873–166	4550	0.9	2.1	-2.97	-2.14	4.70	0.10	4.55	0.07	-2.54	0.43	-0.40	3
CS 22878–101	4800	1.3	2.0	-3.25	-2.56	4.55	0.16	4.36	0.05	-2.75	0.50	-0.19	1
CS 22891–209	4700	1.0	2.1	-3.29	-2.69	4.40	0.15	4.21	0.06	-2.89	0.40	-0.20	1
CS 22892–052*	4850	1.6	1.9	-3.03	-2.52	4.48	0.10	4.29	0.05	-2.82	0.21	-0.30	1
CS 22896–154	5250	2.7	1.2	-2.69	-2.06	5.00	0.08	4.56	0.07	-2.53	0.16	-0.47	2
CS 22897–008	4900	1.7	2.0	-3.41	-2.83	4.35	0.18	4.10	0.06	-3.00	0.41	-0.17	1
CS 22948–066	5100	1.8	2.0	-3.14	-2.59	4.65	0.14	4.26	0.06	-2.84	0.30	-0.25	1
CS 22949–037*	4900	1.5	1.8	-3.97	-2.42	< 3.9	-	< 3.6	0.07	< 3.7	< 0.5	< -1.1	2
CS 22953–003	5100	2.3	1.7	-2.84	-2.34	4.72	0.10	4.41	0.06	-2.69	0.15	-0.35	2
CS 22956–050	4900	1.7	1.8	-3.33	-2.59	4.45	0.20	4.18	0.06	-2.92	0.41	-0.33	1
CS 22966–057	5300	2.2	1.4	-2.62	-1.89	5.12	0.10	4.62	0.08	-2.46	0.16	-0.57	3
CS 29491–053	4700	1.3	2.0	-3.04	-2.34	4.60	0.10	4.45	0.05	-2.66	0.38	-0.32	3
CS 29502–042	5100	2.5	1.5	-3.19	-2.52	4.50	0.14	4.13	0.06	-2.97	0.22	-0.45	1
CS 29518–051	5200	2.6	1.4	-2.69	-2.02	4.95	0.10	4.50	0.07	-2.59	0.10	-0.57	2
CS 30325–094	4950	2.0	1.5	-3.30	-2.54	4.64	0.18	4.19	0.06	-2.91	0.39	-0.37	2
CS 31082–001	4825	1.5	1.8	-2.91	-2.09	4.84	0.10	4.54	0.07	-2.55	0.36	-0.46	3
CS 22949–048	4800	1.5	2.0	-3.25	-2.68	4.42	0.15	4.07	0.05	-3.04	0.21	-0.39	2

* An asterisk after the name of the star means that the star is carbon-rich.

wings of H_{α} . The random error on the temperature is about 80 K (1σ). For a given stellar temperature, the ionization equilibrium provides an estimate of the stellar gravity with an internal accuracy of about 0.1 dex in $\log g$, and the microturbulence velocity v_t can be constrained within 0.2 kms^{-1} . The adopted parameters are given in Table 3. Table 4 lists the uncertainty of the sulphur abundance arising from the errors on T_{eff} , $\log g$ and v_t . Because gravity is determined from the ionization equilibrium, a variation of T_{eff} will change $\log g$ and also sometimes slightly influences v_t . Hence, the total error budget is not the quadratic sum of the various sources of uncertainties, but con-

tains significant covariance terms (see Cayrel et al. 2004 for more details).

Our $\log g$ value may be affected by NLTE effects (overionization) and by uncertainties in the oscillator strengths of the Fe and Ti lines. This has been discussed in Cayrel et al. (2004), section 3.1. It has been shown that even for giants the spectroscopic $\log g$ and the $\log g$ value deduced from isochrones should not differ by more than about 0.3 dex.

For the non-LTE computations of the sulphur abundance, Kurucz models without overshooting were used (Castelli et al., 1997). These models have been shown to provide LTE abun-

Table 4. Abundance uncertainties linked to stellar parameters. Case of a typical dwarf CS 29506-090, and a typical giant HD 122563.

CS 29506-090				
A: $T_{\text{eff}}=6300\text{K}$, $\log g=4.3$ dex, $vt=1.4$ km s $^{-1}$				
B: $T_{\text{eff}}=6300\text{K}$, $\log g=4.2$ dex, $vt=1.4$ km s $^{-1}$				
C: $T_{\text{eff}}=6300\text{K}$, $\log g=4.3$ dex, $vt=1.2$ km s $^{-1}$				
D: $T_{\text{eff}}=6200\text{K}$, $\log g=4.3$ dex, $vt=1.4$ km s $^{-1}$				
E: $T_{\text{eff}}=6200\text{K}$, $\log g=4.1$ dex, $vt=1.3$ km s $^{-1}$				
El.	Δ_{B-A}	Δ_{C-A}	Δ_{D-A}	Δ_{E-A}
[Fe/H]	-0.03	0.03	-0.06	-0.07
[S I/Fe]	0.00	-0.03	0.09	0.04
[Fe I/Fe]	0.02	0.01	-0.04	0.01
[Fe II/Fe]	-0.01	-0.01	0.04	-0.01
HD 122563				
A: $T_{\text{eff}}=4600\text{K}$, $\log g=1.0$ dex, $vt=2.0$ km s $^{-1}$				
B: $T_{\text{eff}}=4600\text{K}$, $\log g=0.9$ dex, $vt=2.0$ km s $^{-1}$				
C: $T_{\text{eff}}=4600\text{K}$, $\log g=1.0$ dex, $vt=1.8$ km s $^{-1}$				
D: $T_{\text{eff}}=4500\text{K}$, $\log g=1.0$ dex, $vt=2.0$ km s $^{-1}$				
E: $T_{\text{eff}}=4500\text{K}$, $\log g=0.6$ dex, $vt=1.8$ km s $^{-1}$				
El.	Δ_{B-A}	Δ_{C-A}	Δ_{D-A}	Δ_{E-A}
[Fe/H]	-0.00	0.06	-0.09	-0.06
[S I/Fe]	-0.02	-0.02	0.12	0.04
[Fe I/Fe]	0.03	0.03	-0.11	0.03
[Fe II/Fe]	-0.03	-0.04	0.11	-0.03

dances very similar (within 0.05 dex) to those of the MARCS models used by Cayrel et al. (2004) and Bonifacio et al. (2009).

It has been shown (see Takeda et al. 2005, Korotin 2008) that Mult. 1 of SI is affected by a significant negative non-LTE correction. To compute NLTE profiles of the SI lines, we have used the model atom of Korotin (2008), which contains 64 levels of SI and the ground level of SII. The radiative photoionization rates for all the levels taken into account are based on the new detailed ionization cross sections listed in the Opacity Project TopBase (<http://cdsweb.u-strasbg.fr/topbase/topbase.html>). The oscillator strengths of the lines are also taken from TopBase. The line profiles have been computed from a modified version of the program MULTI (Carlsson 1986, Korotin et al. 1999). This version includes, in particular, opacity sources from ATLAS9 (Kurucz 1992) which modify the continuum opacity and the intensity distribution in the UV region. Such sources are very important for determination of the radiative rates of the bound-bound transitions in the sulphur atom. In Fig. 3 we present the new non-LTE correction computed for metal-poor stars for the 921.2 nm sulphur line of the Mult. 1. This correction is sometimes rather different from the correction computed by Takeda et al. (2005). For example, for turnoff stars with a metallicity of about -3.0 and a temperature of 6000 K, the correction is found to be close to -0.6 dex, whereas it was only about -0.2 dex following Takeda et al. (2005).

We have also computed the 3D correction (as explained in Caffau et al. 2010b or Caffau & Ludwig 2007). This correction is small, but positive (Table 3), and has been computed and applied separately.

4.1. Test of coherence

With the model atom adopted here, there is good agreement (inside 0.1 dex) in the Sun and in Procyon between the S abundances derived from the different multiplets when 3D models are used.

–For Procyon we found $\epsilon(\text{S}) = 7.23 \pm 0.03$ from the 3D, LTE computations of Caffau et al. (2007) with a Korotin’s NLTE correction (private communication).

–For the Sun the mean sulfur abundance derived from these computations is $\log \epsilon(\text{S}) = 7.16 \pm 0.05$, for $\log \epsilon(\text{H}) = 12.0$ (Caffau et al. (2010b)). This value is very close to the meteoritic value (7.17 ± 0.02 , following Lodders et al., 2009). The solar abundance $\log \epsilon(\text{S}) = 7.16$ has been adopted as a reference in our computations.

Moreover, for the most metal-rich star of our sample, BD+17 3248, the S abundance could be determined from the lines of the Mult. 1 and the main line of Mult. 6 (Fig. 1a). The agreement between the NLTE S abundances derived from multiplets 6 and 1 is very good when the 3D correction is also included (-0.02 for Mult. 6, $+0.10$ for Mult. 1): $\epsilon(\text{S}) = 5.28 \pm 0.2$ from Mult. 6 and $\epsilon(\text{S}) = 5.22 \pm 0.1$ from Mult. 1.




Fig. 3. NLTE correction for the 921.2nm sulphur line in metal-poor stars as a function of the temperature, the gravity and the metallicity for $[\text{Fe}/\text{H}]=-2$, $[\text{Fe}/\text{H}]=-3$ (dashed lines) and $[\text{Fe}/\text{H}]=-4$ (solid line). This correction has been computed for $[\text{S}/\text{H}]=[\text{Fe}/\text{H}]$.

5. Results

In Table 3 we present the result of the computations for our sample of stars. The S abundances have been derived from synthetic spectra fits (Fig. 2). The sulphur abundances could not be determined for the most metal-poor stars ($[\text{Fe}/\text{H}] < -3.6$ for giants, $[\text{Fe}/\text{H}] < -3.2$ for dwarfs), because the lines were too weak to be measured reliably. However, an upper limit of the sulphur abundance has been given for CS 22949-037 because this upper limit can help explaining the chemical anomalies of this star (see section 5.3). The error on the sulphur abundance, given in Table 3, represents the uncertainty on the spectral fitting. This error strongly dominates the total error on $[\text{S}/\text{Fe}]$ or $[\text{S}/\text{Mg}]$.




Fig. 4. Final NLTE $[S/Fe]$ ratios for our EMP stars. Solid symbols show the turnoff stars, open symbols the giants.

5.1. $[S/Fe]$ vs. $[Fe/H]$ and comparison to previous studies

Fig. 4 shows $[S/Fe]$ as a function of $[Fe/H]$ for our sample of EMP stars. Below $[Fe/H] = -2.9$, the slope of the relation is very small: $a = -0.012 \pm 0.006$; $[S/Fe]$ is thus practically constant with a mean of $\overline{[S/Fe]} = +0.35 \pm 0.10$. The level of this plateau is in good agreement with the value found by Nissen et al. (2004, 2008) and Caffau et al. (2005): $\overline{[S/Fe]} \approx +0.35$ for the most metal-poor stars of their sample.




Fig. 5. Spectrum of BD-18 5550 in the region of Mult. 1 of S I. Only the main line can be measured in our spectrum, as the others are severely blended with telluric lines. The thick line shows the theoretical profile computed with our value of the S abundance (Table 3). The profile computed with the abundance adopted by Takeda & Takada-Hidai (2010) (thin line), is not compatible with the observed spectrum.

• Behavior of stars with $-2.5 < [Fe/H]$

It is interesting that none of the stars in Fig. 4 is found to be “S-rich” ($[S/Fe] > +0.7$), in contrast to several stars in the interval $-2 < [Fe/H] < -1$ (Israelian & Rebolo 2001, Takada-Hidai et al. 2002, Takeda et al. 2005, and Caffau et al. 2005).

The most S-rich star in Israelian & Rebolo (2001) is HD 2796, with $[S/Fe] = +0.8$. HD 2796 is included in our sample, but its very high S abundance is not confirmed by our measurement of the lines of Mult. 1; we find $[S/Fe] = +0.3$. Israelian & Rebolo found $EW = 0.26$ pm for the main line of Mult. 6 in this star, but although our spectrum has very high S/N in this

region (over 500/pix at 869 nm), we do not detect the line (Fig. 1b). Its equivalent width is clearly less than 0.26 pm, and it is probable that the error of the equivalent width has been underestimated by Israelian & Rebolo. Ryde & Lambert (2004), re-analysed three other S-rich stars from the list of Israelian & Rebolo with $0.55 < [S/Fe] < 0.7$ and for all of them they found also a normal sulphur abundance.

In Caffau et al. (2005) three other stars have been found to be S-rich (with $[S/Fe] > +0.7$): BD+02 263, BD+05 3640 and CD-45 13178 (HD 181743), these three stars have metallicities in the interval $-2.2 < [Fe/H] < -1.1$ and are thus outside the range of metallicity explored in our paper. Moreover, in HD 181743, Nissen et al. (2008) found from the Mult. 1 a normal abundance of sulphur ($[S/Fe] = +0.34$), but on the other hand, in BD+05 3640 the high sulphur abundance has been confirmed (but at a lower level) by Caffau et al. (2010a) by the analysis of the Mult. 3.

We conclude that the existence of S-rich stars in the interval $-2.5 < [Fe/H] < -1.1$ remains uncertain.

• Behavior of stars with $[Fe/H] < -2.5$

Recently, Takeda & Takada-Hidai (2010) have measured Mult. 3 of S I in metal-poor stars and find $[S/Fe]$ to be constant in the interval $-2.5 < [Fe/H] < -1.1$, but to increase suddenly for $[Fe/H] < -2.6$. They have only five stars with $[Fe/H] < -2.6$, and in one of them the S line cannot be detected. For G64-37, the uncertainty is so high that the detection is in doubt (see their Fig. 13). Of the three remaining stars, two of them are in common with our sample: HD 122563 and BD-18 5550. For these stars we find “normal” S abundances of $[S/Fe] = +0.42$ and $+0.38$ from Mult. 1 (Table 3).

To understand the cause of this discrepancy, we have re-measured the equivalent widths of the 1045.5449 nm line in the spectra of Takeda & Takada-Hidai (2010), neglecting the other two lines, which are even fainter. (The lines of Mult. 3 used by Takeda & Takada-Hidai are weaker than those of Mult. 1 and their spectra have only half the resolution of ours).

– For HD 122563 we find $EW = 1.5$ pm, corresponding to $[S/Fe] = +0.45$, in excellent agreement with our determination. (Since the sulphur lines are weak the value of $[S/Fe]$ found by Takeda & Takada-Hidai has been simply corrected by $\Delta[S/Fe] = \log(EW_2/EW_1)$ where EW_1 and EW_2 represent the old and the new equivalent widths.)

– BD-18 5550 is the most S-rich star in the sample of Takeda & Takada-Hidai with $[S/Fe] = +0.76$, against our value of $+0.38$ from Mult. 1 – see Fig. 5. For this star, the S/N of the spectrum of Takeda & Takada-Hidai seems to be rather good, but we note that the full width at half maximum (FWHM) of the lines is much larger than those in HD 122563, while the FWHM of the lines in these stars is about the same in our UVES spectra. If we fix the FWHM of the lines in BD-18 5550 at the same value as in HD 122563, we find the equivalent width of the 1045.5449 nm line to be about 1.2 pm and $[S/Fe] = +0.62$. Adopting the Cayrel formula (Cayrel 1988) to estimate the precision of this measurement (with $S/N = 150$), we find the error of this equivalent width to be close to 0.5 pm. Hence, the error in the abundance ratio should be about 0.2 dex, which makes their result compatible with ours.




Fig. 6. $[S/Mg]$ vs. $[Fe/H]$ (top) and $[Mg/H]$ (bottom). Symbols as in Fig. 4.

5.2. $[S/Mg]$ vs. predicted supernova yields

It is often noted that magnesium should be a better reference element than iron, because Mg is only formed in massive SN II and its production is less affected by explosive nucleosynthesis, mixing, and “fallback”. As a consequence, the predictions of $[S/Mg]$ should be more robust than for $[S/Fe]$ (Woosley & Weaver, 1995, Shigeyama & Tsujimoto 1998).

The NLTE abundance of Mg for our sample of stars has been computed by Andrievsky et al. (2010). Fig. 6 shows the observed value of $[S/Mg]$ as a function of $[Fe/H]$ and $[Mg/H]$. The scatter is small, but two stars seem to have a high $[S/Mg]$ ratio: BS 17570–063 ($[Fe/H]=-2.92$) and CS 22189–009 ($[Fe/H]=-3.49$). In both cases the high $[S/Mg]$ ratio seems to be real, and not the consequence of an uncertainty on the magnesium abundance (see Fig. 6). In fact the full spread of NLTE $[Mg/Fe]$ of the metal-poor stars below $[Fe/H] < -1$ is definitely larger than expected from the error estimate of the analysis itself (see Gehren et al., 2006, Andrievsky et al., 2010, and also Cayrel et al., 2004) and the anomalies of the magnesium abundances are not clearly correlated with the other elements.

In the interval $-3.4 < [Fe/H] < -2.5$ we can define a mean value of $[S/Mg]$ as was done for O, Na, Al and K by Andrievsky et al. (2010): $[S/Mg] = -0.32 \pm 0.14$. In the same metallicity interval, the scatter of $[S/Fe]$ around the mean is only 0.12 dex (see Fig 4). It appears that, as already observed in LTE (Cayrel et al. 2004), the scatter of the abundance ratios is slightly larger when Mg rather than Fe is used as the reference element. This is surprising, considering that Fe forms in processes quite dif-

ferent from those forming Mg and S (explosive burning for Fe, and mainly hydrostatic C, Ne, and O burning for Mg and S).

For $[Mg/H] < -2.7$, a slight increase of $[Na/Mg]$, $[Al/Mg]$, and $[K/Mg]$ as $[Mg/H]$ decreases was seen by Andrievsky et al. (2010). A slight increase of $[S/Mg]$ is also possible (Fig. 6), but there are very few stars with $[Mg/H] < -2.7$ in our sample. We plan to discuss the evolution of $[S/Mg]$ in the Galaxy in more detail when we have completed a new precise analysis of disk stars, including NLTE and 3D corrections.




Fig. 7. $[X/H]$ abundances in CS 22949-037 against the solar condensation temperatures (Lodders, 2003). There is no correlation between T_{cond} and the abundance of the elements, in particular S and Zn have a condensation temperature lower than Na and they are more deficient, at variance with the abundances in λ Boo stars. (The filled circles represent the abundances corrected or not sensitive to NLTE effects, the open symbols are abundances which could not be corrected for NLTE effects)

5.3. The case of the peculiar C-rich star CS 22949-037

CS 22949-037 is a very peculiar giant: it is extremely metal-poor ($[Fe/H] \approx -4.0$), and its unusual chemical composition was first pointed out by McWilliam et al. (1995) and later studied in more detail by Norris et al. (2001) and Depagne et al. (2002). This star exhibits light element (C-N-O, Na, Mg) enhancements that are quite similar to those associated with the hyper metal-poor stars HE 0107-5240 and HE 1327-2326 with $[Fe/H] < -5$ (see Frebel et al., 2005, 2008). It is generally admitted that these stars are the product of specific zero-metal supernovae. However, Venn & Lambert (2008) have argued that this abundance pattern might be associated with the same peculiarity that is used to account for λ Boo stars (presumably related to dust depletion). If this were so, the abundance of the elements should be smoothly correlated with the condensation temperature T_{cond} . The abundances of S and Zn could help to choose between the two hypotheses, because their temperature are intermediate between those of O and Na, both of which are strongly enhanced in CS 22949-037. If dust deple-

tion is responsible for the pattern of CS 22949-037 we would expect that sulphur and zinc be also strongly enhanced in this star. The abundance of zinc has been measured in CS 22949-037 by Depagne et al. (2002). The sulphur lines are not visible in this star but (Table 3) an upper limit can be estimated as $[S/H] < -3.5$ or $[S/Fe] < +0.5$. In Fig. 7 we have plotted $[X/H]$ vs T_{cond} . The abundances $[X/H]$ of C, N, O, Ca, Fe, and Zn are taken from Depagne et al. (2002); for the other elements Na, Mg, Al, S, Sr, and Ba, since the NLTE abundances have been computed (Andrievsky et al., 2007, 2008, 2009, 2010, 2011) they have been preferred. (The oxygen abundance determined from the forbidden line is insensitive to the NLTE effects). It can be seen that CS 22949-037 does not show any clear signature of dust depletion.

5.4. Comparison with predicted SN yields




Fig. 8. Comparison of the most recent O, Na, Mg, Al, S, and K abundances in EMP stars to the predicted yields of metal-poor SNe and hypernovae. The solid lines represent the predictions of different models, and the dots the mean values of the observed abundances in the interval $-3.0 < [Mg/H] < -2.0$

In Fig. 8 we compare the abundance ratios of the light metals to some predicted abundances in the ejecta of metal-poor SNe or hypernovae following Woosley & Weaver (1995), Kobayashi et al. (2006) and Heger & Woosley (2010). The mean values of the observed abundances of the different elements are computed in the interval $-3.0 < [Mg/H] < -2.0$, their uncertainty should be of the order of 0.1 dex. The amount of ejected S seems generally overestimated by the models. If we consider all the elements from O to K, the best agreement is obtained with the predictions by the models of Heger & Woosley 2010 (hypotheses B or D, with a low mass cutoff at $10M_{\odot}$), mainly because other simulations show larger disagreement for the K abundance.

5.5. $[S/Zn]$ in metal-poor stars and damped $\text{Ly}\alpha$ systems

Nissen et al. (2004) suggested that the $[S/Zn]$ ratio is higher in stars than in damped $\text{Ly}\alpha$ systems. Neither S nor Zn shows much affinity for dust, although MgS has been recently mentioned as a possible grain constituent by Zhang et al. (2009). Thus, depletion onto dust in DLAs probably can not explain




Fig. 9. $[S/Zn]$ vs. $[Zn/H]$ in EMP stars (circles) and in metal-poor DLAs (triangles).

Table 5. $[Zn/H]$ and $[S/Zn]$ in the most metal-poor DLAs.

Ident	$[Zn/H]$	$[S/Zn]$	reference S	reference Zn
0010-0012	-1.35	0.24	Srianand,05	Ledoux,03
0058-2914	-1.52	0.22	Srianand,05	Pettini,00
0100+130	-1.55	0.15	Molaro,98	Dessauges-,04
0102-1902	-1.88	0.06	Srianand,05	Ledoux,03
0112+029	-1.01	-0.18	Ledoux,03	Pettini,94
0347-383	-1.15	0.06	Srianand,05	Ledoux,03
0405-443	-1.34	-0.09	Lopez,03	Ledoux,06
0405-443	-1.06	0.04	Lopez,03	Ledoux,06
0528-2505	-1.47	0.07	Centurion,03	Ledoux,06
0841+129	-1.45	0.10	Ledoux,06	Centurion,03
0841+129	-1.74	0.32	Centurion,03	Dessauges-,06
1223+178	-1.63	0.25	Dessauges-,07	Ledoux,03
1331+170	-1.25	0.07	Dessauges-,04	Dessauges-,04
2138-4427	-1.64	0.04	Srianand,05	Ledoux,03
2314-409	-1.03	0.11	Ellison,01	Ledoux,06
2343+1232	-1.30	-0.06	Noterdaeme,07	Ledoux,06

this difference. Fig. 9 shows $[S/Zn]$ vs. $[Zn/H]$ for our sample of metal-poor stars and for the most metal-poor DLAs with both Zn and S abundance determinations (Table 5). Since the Zn line is very weak in turnoff stars, we could only measure Zn abundances in our giant stars (Cayrel et al., 2004). Following Takeda et al. (2005), non-LTE effects for Zn are not very important in these stars ($\leq +0.1$ dex), but we have interpolated a correction in their tables and applied it.

Fig. 9 shows that, when non-LTE effects are taken into account, the sulphur abundance in metal-poor stars is rather similar to that in DLAs – if anything slightly lower. However, the chemical evolution of the Galactic halo might still be different from the average of DLAs.

In EMP stars as in DLAs the $[S/Zn]$ ratio is approximately solar. This is due to positive values of both $[S/Fe]$ (Fig. 4) and $[Zn/Fe]$ at low metallicity (Cayrel et al. 2004; see also Primas et al. 2000 and Nissen et al. 2004). We will discuss the evolution of the $[S/Zn]$ ratio in the Galaxy in more detail when we have completed a new precise analysis of S abundances in less metal-poor stars.

6. Conclusions

We have determined S abundances in 33 metal-poor stars, including 21 with $[Fe/H] < -2.9$, and find $[S/Fe]$ to be constant:

$[S/Fe] = 0.35 \pm 0.10$. However, a slight increase of $[S/Mg]$ when $[Mg/H]$ decreases cannot be excluded (Fig. 6). The same effect is observed also for Na, Al, and K (Spite et al. 2009, Andrievsky et al. 2010).

We do not confirm earlier claims of a significant rise in $[S/Fe]$ with decreasing $[Fe/H]$, but corroborate the findings of Nissen et al. (2004, 2008) and Ryde & Lambert (2004) that S behaves like the other α -elements, also at very low metallicity.

The low sulphur abundance (along with a low abundance of zinc) in CS 22949-037 suggests that the abundance pattern in this star (and thus by analogy in the hyper metal-poor stars HE 0107-5240 and HE 1327-2326) is the result of specific zero-metal supernovae and is not related to dust depletion as suggested by Venn & Lambert (2008).

When the NLTE effects are taken into account the pattern of the light elements from O to K relative to magnesium, is rather well represented by ejecta of zero-metal supernovae (Heger & Woosley 2010).

The ratio $[S/Zn]$ is found to be approximately solar as it is in DLAs since $[S/Fe]$ and $[Zn/Fe]$ are both equally positive at low metallicity.

Acknowledgements. SMA gratefully acknowledges the support and hospitality of Observatoire de Paris-Meudon. MS, EC, FS, PB, RC, PF and VH acknowledge the support of CNRS (PNCG and PNPS). BB acknowledges grants from FAPESP and CNPq. BN and JA acknowledge the support of the Carlsberg Foundation and the Danish Natural Science Research Council and TCB acknowledges partial funding of this work from grants PHY 02-16783 and PHY 08-22648: Physics Frontiers Center/Joint Institute for Nuclear Astrophysics (JINA), awarded by the U.S. National Science Foundation. We also thank the anonymous referee for his helpful comments.

References

- Alonso A., Arribas S., Martinez-Roger C., 1996, A&A 313, 873
- Andrievsky S. M., Spite M., Korotin S. A., Spite F., Bonifacio P., Cayrel R., Hill V., François P., 2007, A&A 464, 1081
- Andrievsky S. M., Spite M., Korotin S. A., Spite F., Bonifacio P., Cayrel R., Hill V., François P., 2008, A&A 481, 481
- Andrievsky S. M., Spite M., Korotin S.A., Spite F., Bonifacio P., Cayrel R., Hill V., François P., 2009, A&A 494, 1083
- Andrievsky S. M., Spite M., Korotin S. A., Spite F., Bonifacio P., Cayrel R., François P., Hill V., 2010, A&A 509, 88
- Andrievsky S. M., Spite F., Korotin S. A., Spite M. et al., 2011 to be submitted to A&A
- Ballester P., Modigliani A., Boitquin O., Cristiani S., Hanuschik R., et al., 2000 ESO Messenger 101 31
- Bonifacio P., Molaro P., Sivarani T., et al., 2007 A&A 462 851 (“**First Stars VII**”)
- Bonifacio P., Spite M., Cayrel R., Hill V., et al., 2009, A&A 501, 519 (“**First Stars XII**”)
- Caffau E., Bonifacio P., Faraggiana R., François P., Gratton R.G., Barbieri M., 2005, A&A 441, 533
- Caffau E., Ludwig H.-G., 2007, A&A 467, L11
- Caffau E., Faraggiana R., Bonifacio P., Ludwig H.-G., Steffen M., 2007, A&A 470, 699
- Caffau E., Sbordone L., Faraggiana R., Ludwig H.-G., Bonifacio P., Spite M., 2010a, AN 331, 725
- Caffau, E., Ludwig, H.-G., Steffen, M., Freytag, B., & Bonifacio, P. 2010b, *Solar Phys.*, 66
- Carlsson M., 1986, Uppsala Obs. Rep. 33
- Castelli F., Gratton R. G., Kurucz, R. L. 1997, A & A 318, 841
- Cayrel R., 1988, Proceedings of the IAU symposium 132 “The impact of very high S/N spectroscopy on stellar physics”, eds. G. Cayrel de Strobel and M. Spite, Kluwer Academic Publishers, Dordrecht, p.345
- Cayrel R., Depagne E., Spite M., Hill V., Spite F., François P., Plez B., Beers T.C., Primas F., Andersen J., Barbuy B., Bonifacio P., Molaro P., Nordström B., 2004 A&A 416 1117 (“**First Stars V**”)
- Centurión M., Molaro P., Vladilo G., 2003, A&A, 403, 55
- Dekker H., D’Odorico S., Kaufer A., et al., 2000 in Optical and IR Telescopes Instrumentation and Detectors eds I., Masanori & A..F., Morwood Proc., SPIE 4008 534
- Depagne E., Hill V., Spite M., Spite F., Plez B., Beers T.C., Barbuy B., Cayrel R., Andersen J., Bonifacio P., François P., Nordström B., Primas F., 2004 A&A 416 1117 (“**First Stars II**”)
- Dessauges-Zavadsky, M.; Calura, F.; Prochaska, J. X.; D’Odorico, S.; Matteucci, F., 2004, A&A 416, 79
- Dessauges-Zavadsky, M.; Prochaska, J. X.; D’Odorico, S.; Calura, F.; Matteucci, F., 2006, A&A 445, 93
- Dessauges-Zavadsky, M.; Calura, F.; Prochaska, J. X.; D’Odorico, S.; Matteucci, F., 2007, A&A 470, 431
- Ellison S. L., Lopez S., 2001, A&A, 380, 117
- François P., 1987, A&A 176, 294
- François P., 1988, A&A 195, 226
- Frebel A., Aoki W., Christlieb N., Ando H., Asplund M., Barklem P. S., Beers T. C., Eriksson K., Fechner C., Fujimoto M. Y., 2005, Nature 434, 871
- Frebel A., Collet R., Eriksson K. et al., 2008, ApJ 684, 588
- Gehren T., Shi J.R., Zhang H.W. et al. 2006, A&A 451, 1065
- Heger A., Woosley S.E., 2010, ApJ 724, 341
- Hill V., Plez B., Cayrel R., Beers T.C., Nordström B., Andersen J., Spite M., Spite F., Barbuy B., Bonifacio P., Depagne E., François P., Primas F., 2002, A&A 387, 560 (“**First Stars I**”)
- Israeli G., Rebolo R., 2001, ApJ 557, L43
- Kobayashi C., Umeda H., Nomoto K., Tominaga N., and Ohkubo T. 2006, ApJ 653, 1145
- Korotin S.A., 2008, Odessa Astronomical Pub. 21, 42
- Korotin S.A., 2009, Astronomicheskii Zhurnal, 2009 86, 702 and 2009, Astr. Rep. 53, 651
- Korotin S.A., Andrievsky S. M., Luck R.E., 1999, A&A 351, 168
- Kurucz R.L., 1992, RMxAA 23, 181
- Ledoux, C., Petitjean P., Srianand, R., 2003, MNRAS, 346, 209L
- Ledoux C., Petitjean P., Fynbo J. P. U., Miller P., Srianand R., 2006, A&A 457, 71L
- Lodders K., 2003, ApJ 591, 1220
- Lodders K., Palme H., Gail H.-P., 2009, Landolt-Börnstein, New Series, Astronomy and Astrophysics, Springer Verlag, Berlin
- Lopez S., Reimers D., D’Odorico S., Prochaska J.X., 2002, A&A, 385, 778
- Lopez S., Ellison S. L., 2003, A&A, 403, 573
- McWilliam A., Preston G. W., Sneden C., Searle L., 1995, AJ, 109, 2757
- Molaro P., Vladilo G., Centurion M., 1998, MNRAS 293, L37
- Nakamura T., Umeda H., Iwamoto K., Nomoto K., Hashimoto M., Hix W.R., & Thielemann F.-K. 2001, ApJ, 555, 880
- Nissen P., Chen Y.Q., Asplund M., Pettini M., 2004, A&A 415, 993
- Nissen P., Akerman C., Asplund M., Fabbian D., Pettini M., 2008, psa Conf, 51
- Norris J.E., Ryan S.G., Beers T.C., 2001, ApJ 561, 1034
- Noterdaeme P., Ledoux C., Petitjean P., Le Petit F., Srianand R., Smette A., 2007, A&A 474, 393
- Pettini M., Smith L. J., Hunstead R. W., King, D. L., 1994, ApJ 426, 79

- Pettini M., Ellison S. L., Steidel C. C., Shapley A. E., Bowen D. V., 2000, *ApJ* 532, 65
- Primas F., Brugamyer E., Sneden C., King J.R., Beers T.C., Boesgaard A.M., Deliyannis C.P., 2000, “The First Stars”: Proceedings of the MPA/ESO Workshop, ESO ASTROPHYSICS SYMPOSIA., Eds. A. Weiss, T.G. Abel, and V. Hill. Springer-Verlag, 2000, p. 51
- Prochaska J. X., Wolfe A. M., Tytler D., et al. 2001, *ApJS*, 137, 21
- Ramaty R., Scully S., Lingenfelter R., Kozlovsky B. 2000, *ApJ*, 534, 747
- Ryde N., Lambert D.L., 2004, *A&A* 415, 559
- Ryde N., 2006, *A&A* 455, L13
- Shigeyama, T., Tsujimoto, T. 1998, *ApJ*, 507, L135
- Spite M., Spite F., Bonifacio P., Hill V., Andrievsky S., Cayrel R., François P., Korotin S., 2009 “Chemical Abundances in the Universe: Connecting First Stars to Planets”, Proceedings of the IAU Symposium 265, Eds. K. Cunha, M. Spite & B. Barbuy, p. 380
- Srianand R., Petitjean P., Ledoux C., Ferland G., Shaw G., 2005, *MNRAS*, 362, 549
- Takada-Hidai, M., Takeda, Y., Sato, S., et al. 2002, *ApJ*, 573, 614
- Takeda Y., Hashimoto O., Taguchi H., Yoshioka K. Takada-Hidai M., Saito Y., Honda S., 2005, *PASJ* 57, 751
- Takeda Y., Takada-Hidai M., 2010, ArXiv 1009.0824 (to be published in PASJ)
- Venn K.A., Lambert D.L., 2008, *ApJ*, 677, 572
- Woosley S. E., Weaver T. A. 1995, *ApJS*, 101, 181
- Zhang K., Jiang B.W., Li A., 2009, *ApJ* 702, 680