



HAL
open science

DENSE ORBITS FOR ABELIAN AFFINE GROUPS

Adlene Ayadi

► **To cite this version:**

| Adlene Ayadi. DENSE ORBITS FOR ABELIAN AFFINE GROUPS. 2010. hal-00549263v1

HAL Id: hal-00549263

<https://hal.science/hal-00549263v1>

Preprint submitted on 21 Dec 2010 (v1), last revised 23 Mar 2011 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DENSE ORBITS FOR ABELIAN AFFINE GROUPS

ADLENE AYADI

ABSTRACT. In this paper, we give a characterization of existence of a dense orbit for the action on \mathbb{C}^n of an abelian affine subgroup \mathcal{G} of $GA(n, \mathbb{C})$. We prove in particular that if \mathcal{G} has a dense orbit then the closure $\overline{\mathcal{G}}$ is an affine space of dimension n . If \mathcal{G} is finitely generated, this characterization is explicit.

1. Introduction

Let $M_n(\mathbb{C})$ be the set of all square matrices of order $n \geq 1$ with entries in \mathbb{C} and $GL(n, \mathbb{C})$ be the group of all invertible matrices of $M_n(\mathbb{C})$. A map $f : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is called an affine map if there exist $A \in M_n(\mathbb{C})$ and $a \in \mathbb{C}^n$ such that $f(x) = Ax + a$, $x \in \mathbb{C}^n$. We denote $f = (A, a)$, we call A the *linear part* of f . Denote by $MA(n, \mathbb{C})$ the set of all affine maps and $GA(n, \mathbb{C})$ the set of all invertible affine maps of $MA(n, \mathbb{C})$. $MA(n, \mathbb{C})$ is a vector space and for composition of maps, $GA(n, \mathbb{C})$ is a group.

Let \mathcal{G} be an abelian affine subgroup of $GA(n, \mathbb{C})$. For a vector $v \in \mathbb{C}^n$, we consider the orbit of \mathcal{G} through v : $\mathcal{G}(v) = \{f(v) : f \in \mathcal{G}\} \subset \mathbb{C}^n$. A subset $E \subset \mathbb{C}^n$ is called \mathcal{G} -invariant if $f(E) \subset E$ for any $f \in \mathcal{G}$; that is E is a union of orbits. Before stating our main results, we introduce the following notions:

A subset \mathcal{H} of \mathbb{C}^n is called an *affine subspace* of \mathbb{C}^n if there exist a vector subspace H of \mathbb{C}^n and $a \in \mathbb{C}^n$ such that $\mathcal{H} = H + a$. For $a \in \mathbb{C}^n$, denote by $T_a : \mathbb{C}^n \rightarrow \mathbb{C}^n$; $x \mapsto x + a$ the translation map by vector a , so $\mathcal{H} = T_a(H)$. We say that \mathcal{H} has dimension p ($0 \leq p \leq n$), denoted $\dim(\mathcal{H}) = p$, if H has dimension p .

Denote by \overline{A} the closure of a subset $A \subset \mathbb{C}^n$. A subset E of \mathbb{C}^n is called a *minimal set* of \mathcal{G} if E is closed in \mathbb{C}^n , non empty, \mathcal{G} -invariant and has no proper subset with these properties. It is equivalent to say that E is a \mathcal{G} -invariant set such that every orbit contained in E is dense in it.

If V is a \mathcal{G} -invariant open set in \mathbb{C}^n , we say that E is a *minimal set in V* if it is a minimal set of the restriction $\mathcal{G}|_V$ of \mathcal{G} to V .

Define the map

$$\Phi : GA(n, \mathbb{C}) \rightarrow \Phi(GA(n, \mathbb{C})) \subset GL(n+1, \mathbb{C})$$

2000 *Mathematics Subject Classification.* 37C85.

Key words and phrases. affine action, dense, orbit, group, abelian, transitive.

This work is supported by the research unit: systèmes dynamiques et combinatoire: 99UR15-15.

$$f = (A, a) \mapsto \begin{bmatrix} 1 & 0 \\ a & A \end{bmatrix}$$

We have the following composition formula

$$\begin{bmatrix} 1 & 0 \\ a & A \end{bmatrix} \begin{bmatrix} 1 & 0 \\ b & B \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ Ab + a & AB \end{bmatrix}.$$

Then Φ is a homomorphism of groups.

Let \mathcal{G} be an abelian affine subgroup of $GA(n, \mathbb{C})$. Then $\Phi(\mathcal{G})$ is an abelian subgroup of $GL(n+1, \mathbb{C})$.

Let $\pi : \mathbb{C}^n \rightarrow \{1\} \times \mathbb{C}^n$ be the map defined by $\pi(y) = (1, y)$. Clearly π is an invertible affine maps and we have for every $x \in \mathbb{C}^n$,

$$\pi(\mathcal{G}(x)) = \Phi(\mathcal{G})(1, x) = \{1\} \times \mathcal{G}(x) \subset \{1\} \times \mathbb{C}^n.$$

Let $n \in \mathbb{N}_0$ be fixed. For each $m = 1, 2, \dots, n+1$, denote by:

- $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ and $\mathbb{N}_0 = \mathbb{N} \setminus \{0\}$.
- $\mathcal{B}_0 = (e_1, \dots, e_{n+1})$ the canonical basis of \mathbb{C}^{n+1} and I_{n+1} the identity matrix of $GL(n+1, \mathbb{C})$.
- $\mathbb{T}_m(\mathbb{C})$ the set of matrices over \mathbb{C} of the form

$$\begin{bmatrix} \mu & & & 0 \\ a_{2,1} & \mu & & \\ \vdots & \ddots & \ddots & \\ a_{m,1} & \dots & a_{m,m-1} & \mu \end{bmatrix} \quad (1)$$

- $\mathbb{T}_m^*(\mathbb{C})$ the group of matrices of the form (1) with $\mu \neq 0$.

Let $r \in \mathbb{N}$ and $\eta = (n_1, \dots, n_r)$ be a sequence of positive integers such that $n_1 + \dots + n_r = n+1$. In particular, $r \leq n+1$.

Write

- $\mathcal{K}_{\eta,r}(\mathbb{C}) := \mathbb{T}_{n_1}(\mathbb{C}) \oplus \dots \oplus \mathbb{T}_{n_r}(\mathbb{C})$. In particular if $r = 1$, then $\mathcal{K}_{\eta,1}(\mathbb{C}) = \mathbb{T}_{n+1}(\mathbb{C})$ and $\eta = (n+1)$.
- $\mathcal{K}_{\eta,r}^*(\mathbb{C}) := \mathcal{K}_{\eta,r}(\mathbb{C}) \cap GL(n+1, \mathbb{C})$.

Define the map

$$\Psi : MA(n, \mathbb{C}) \rightarrow \Psi(GA(n, \mathbb{C})) \subset M_{n+1}(\mathbb{C})$$

$$f = (A, a) \mapsto \begin{bmatrix} 0 & 0 \\ a & A \end{bmatrix}$$

We have Ψ is an isomorphism. Remark that $\Phi(f) = \Psi(f) + D_1$, where $D_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \in M_{n+1}(\mathbb{C})$, $f \in GA(n, \mathbb{C})$, so Φ is an affine map.

Denote by:

- $\mathcal{F}_{n+1} = \left\{ \begin{bmatrix} 0 & 0 \\ b & B \end{bmatrix} \in M_{n+1}(\mathbb{C}), B \in M_n(\mathbb{C}), b \in \mathbb{C}^n \right\} = \Psi(MA(n, \mathbb{C}))$.

- $\exp : \mathbb{M}_{n+1}(\mathbb{C}) \longrightarrow \mathrm{GL}(n+1, \mathbb{C})$ is the matrix exponential map; set $\exp(M) = e^M$.

There always exists a $P \in \Phi(\mathrm{GA}(n, \mathbb{C}))$ and a partition η of $n+1$ such that $G' = P^{-1}GP \subset \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(\mathrm{GA}(n, \mathbb{C}))$ (see Proposition 2.3). For such a choice of matrix P , we let

- $\mathfrak{g} = \exp^{-1}(G) \cap (P(\mathcal{K}_{\eta,r}(\mathbb{C}))P^{-1}) \cap \mathcal{F}_{n+1}$. If $G \subset \mathcal{K}_{\eta,r}^*(\mathbb{C})$, we have $\mathfrak{g} = \exp^{-1}(G) \cap \mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}$.
- $\mathfrak{g}_u = \{Bu : B \in \mathfrak{g}\}$, $u \in \mathbb{C}^{n+1}$.
- $\mathfrak{g} = \Psi^{-1}(\mathfrak{g})$.
- $\mathfrak{g}_v = \{f(v), f \in \mathfrak{g}\}$, $v \in \mathbb{C}^n$.

Let $\tilde{\pi} : \mathbb{C}^n \longrightarrow \{0\} \times \mathbb{C}^n$ be the map defined by $\tilde{\pi}(y) = (0, y)$. Clearly $\tilde{\pi}$ is an isomorphism.

Denote by:

- $u_0 = [e_{1,1}, \dots, e_{r,1}]^T \in \mathbb{C}^{n+1}$ where $e_{k,1} = [1, 0, \dots, 0]^T \in \mathbb{C}^{n_k}$, for $k = 1, \dots, r$. One has $u_0 \in \{1\} \times \mathbb{C}^n$.
- $w_0 = \pi^{-1}(Pu_0) \in \mathbb{C}^n$. As $P \in \Phi(\mathrm{GA}(n, \mathbb{C}))$, $Pu_0 \in \{1\} \times \mathbb{C}^n$. So $\pi(w_0) = (1, w_0) = Pu_0$.
- $e^{(k)} = [e_1^{(k)}, \dots, e_r^{(k)}]^T \in \mathbb{C}^{n+1}$ where

$$e_j^{(k)} = \begin{cases} 0 \in \mathbb{C}^{n_j} & \text{if } j \neq k \\ e_{k,1} & \text{if } j = k \end{cases} \quad \text{for every } 1 \leq j, k \leq r.$$

- $p_2 : \mathbb{C}^{n+1} \longrightarrow \mathbb{C}^n$ the projection defined by $p_2(x_1, \dots, x_{n+1}) = (x_2, \dots, x_{n+1})$. Notice that π^{-1} is the restriction of p_2 to $\{1\} \times \mathbb{C}^n$.
- M^T is the transpose of any $M \in M_m(\mathbb{C})$, $m \leq n+1$.
- $\overline{\mathcal{G}}$ is the closure of \mathcal{G} in $MA(n, \mathbb{C})$, which is considered as a vector space with the norm defined by $\|f\| = \max(\|A\|, \|a\|)$ the maximum of $\|A\|$ and $\|a\|$, for every $f = (A, a) \in MA(n, \mathbb{C})$.

For a *finitely generated* subgroup $\mathcal{G} \subset \mathrm{GA}(n, \mathbb{R})$, let introduce the following property. Consider the following rank condition on a collection of affine maps $f_1, \dots, f_p \in \mathrm{GA}(n, \mathbb{C})$, where $f'_1, \dots, f'_p \in \mathfrak{g}$ such that $e^{\Psi(f'_k)} = \Phi(f_k)$, $k = 1, \dots, p$.

We say that f_1, \dots, f_p satisfy *property D* if for every $(s_1, \dots, s_p; t_1, \dots, t_r) \in \mathbb{Z}^{p+r} \setminus \{0\}$:

$$\mathrm{rank} \begin{bmatrix} \mathrm{Re}(f'_1(w_0)) & \dots & \mathrm{Re}(f'_p(w_0)) & 0 & \dots & 0 \\ \mathrm{Im}(f'_1(w_0)) & \dots & \mathrm{Im}(f'_p(w_0)) & 2\pi e^{(1)} & \dots & 2\pi e^{(r)} \\ s_1 & \dots & s_p & t_1 & \dots & t_r \end{bmatrix} = n+1.$$

For a vector $v \in \mathbb{C}^n$, we write $v = \mathrm{Re}(v) + i\mathrm{Im}(v)$ where $\mathrm{Re}(v)$ and $\mathrm{Im}(v) \in \mathbb{R}^n$.

For groups of affine maps on \mathbb{K}^n ($\mathbb{K} = \mathbb{R}$ or \mathbb{C}), their dynamics were recently initiated for some classes in different point of view, (see for instance, [3], [4], [5],[6]). The purpose here is to give analogous results of that theorem for linear abelian subgroup of $GL(n, \mathbb{C})$ proved in [1]. Our main results are the following:

Theorem 1.1. *Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$. The following are equivalent:*

- (i) \mathcal{G} has a dense orbit in \mathbb{C}^n
- (ii) The orbit $\mathcal{G}(w_0)$ is dense in \mathbb{C}^n
- (iii) \mathfrak{g}_{w_0} is an additive subgroup dense in \mathbb{C}^n

Corollary 1.2. *If \mathcal{G} has a dense orbit then $\overline{\mathcal{G}}$ is an affine subspace of $GA(n, \mathbb{C})$ of dimension n .*

For a finitely generated subgroup \mathcal{G} , the Theorem can be stated as follows:

Theorem 1.3. *Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$ generated by f_1, \dots, f_p and let $f'_1, \dots, f'_p \in \mathfrak{g}$ such that $e^{\Psi(f'_1)} = \Phi(f_1), \dots, e^{\Psi(f'_p)} = \Phi(f_p)$. Then the following are equivalent:*

- (i) \mathcal{G} has a dense orbit in \mathbb{C}^n
- (ii) the maps f_1, \dots, f_p satisfy property \mathcal{D}
- (iii) $\mathfrak{g}_{w_0} = \sum_{k=1}^p \mathbb{Z}f'_k(w_0) + 2i\pi \sum_{k=1}^r \mathbb{Z}(p_2 \circ Pe^{(k)})$ is an additive group dense in \mathbb{C}^n .

Corollary 1.4. *If \mathcal{G} is of finite type p with $p \leq 2n - r$, then it has no dense orbit.*

Corollary 1.5. *If \mathcal{G} is of finite type p with $p \leq n$, then it has no dense orbit.*

2. Notations and Lemmas

Denote by $\mathcal{L}_{\mathcal{G}}$ the set of the linear parts of all elements of \mathcal{G} and $\text{vect}(F)$ is the vector space generated by a subset $F \subset \mathbb{C}^{n+1}$.

Lemma 2.1. *Let $P \in \Phi(GA(n, \mathbb{C}))$ and $h : M_{n+1}(\mathbb{C}) \rightarrow M_{n+1}(\mathbb{C})$ the isomorphism defined by $h(A) = PAP^{-1}$. Then $h(\mathcal{F}_{n+1}) = \mathcal{F}_{n+1}$ and $h(\Phi(GA(n, \mathbb{C}))) = \Phi(GA(n, \mathbb{C}))$.*

Proof. Let $f = (A, a) \in MA(n, \mathbb{C})$. As $P \in \Phi(GA(n, \mathbb{C}))$, write

$$P = \begin{bmatrix} 1 & 0 \\ d & C \end{bmatrix}, \quad C \in GL(n, \mathbb{C}), \quad d \in \mathbb{C}^n.$$

Then

$$\begin{aligned} h(\Psi(f)) &= \begin{bmatrix} 1 & 0 \\ d & C \end{bmatrix} \begin{bmatrix} 0 & 0 \\ a & A \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -C^{-1}d & C^{-1} \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ d + Ca - CAC^{-1}d & CAC^{-1} \end{bmatrix}, \end{aligned}$$

so $h(\Psi(f)) \in \mathcal{F}_{n+1}$. It follows that $h(\mathcal{F}_{n+1}) \subset \mathcal{F}_{n+1}$. Conversely, let $h' = h^{-1}$, so $h' : M_{n+1}(\mathbb{C}) \rightarrow M_{n+1}(\mathbb{C})$ is the isomorphism defined by $h(B) = P^{-1}BP$. By replacing h by h' in above, we have $h'(\mathcal{F}_{n+1}) \subset \mathcal{F}_{n+1}$. It follows that

$$\mathcal{F}_{n+1} = h \circ h'(\mathcal{F}_{n+1}) \subset h(\mathcal{F}_{n+1}).$$

The same proof is used to show $h(\Phi(GA(n, \mathbb{C}))) = \Phi(GA(n, \mathbb{C}))$. \square

Proposition 2.2. ([1], Proposition 2.3) *Let G be an abelian subgroup of $GL(n, \mathbb{C})$. Then there exists $P \in GL(n, \mathbb{C})$ such that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta', r'}(\mathbb{C})$, for some $r' \leq n$ and $\eta' = (n'_1, \dots, n'_{r'}) \in \mathbb{N}_0^{r'}$.*

Proposition 2.3. *Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$ and $G = \Phi(\mathcal{G})$. Then there exists $P \in \Phi(GA(n, \mathbb{C}))$ such that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta, r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$, for some $r \leq n + 1$ and $\eta = (n_1, \dots, n_r) \in \mathbb{N}_0^r$. In particular, $Pu_0 \in \{1\} \times \mathbb{C}^n$.*

Proof. We have $\mathcal{L}_{\mathcal{G}}$ is an abelian subgroup of $GL(n, \mathbb{C})$. By Proposition 2.2, there exists $P_1 \in GL(n, \mathbb{C})$ such that $P_1^{-1}GP_1$ is a subgroup of $\mathcal{K}_{\eta', r'}^*(\mathbb{C})$ for some $r' \leq n$ and $\eta' = (n'_1, \dots, n'_{r'}) \in \mathbb{N}_0^{r'}$ such that $n'_1 + \dots + n'_{r'} = n$. Write for every $A \in \mathcal{L}_{\mathcal{G}}$, $P_1^{-1}AP_1 = \text{diag}(A_1, \dots, A_{r'})$ with $A_k \in \mathbb{T}_{n'_k}^*$ and μ_{A_k} is the only eigenvalue of A_k , $k = 1, \dots, r'$. Let $J = \{k \in \{1, \dots, r'\}, \mu_{A_k} = 1, \forall A \in \mathcal{L}_{\mathcal{G}}\}$. There are two cases:

- *Case1:* Suppose that $J = \{k_1, \dots, k_s\}$ for some $s \leq r'$. We can take $J = \{1, \dots, s\}$, otherwise, we replace P_1 by RP_1 for some permutation matrix R of $GL(n, \mathbb{C})$. Let $P = \text{diag}(1, P_1)$, so $P \in \Phi(GA(n, \mathbb{C}))$ and

$$P^{-1}\Phi(f)P = \begin{bmatrix} 1 & 0 \\ P_1^{-1}a & P_1^{-1}AP_1 \end{bmatrix}.$$

Write $P(\mathcal{B}_0) = (e_1, \mathcal{C}_1, \dots, \mathcal{C}_{r'})$, where $\mathcal{C}_k = (e'_{k,1}, \dots, e'_{k,n'_k})$ is a basis of $\mathbb{C}^{n'_k}$, $k = 1, \dots, r'$. Then $H = \text{vect}(e_1, \mathcal{C}_1, \dots, \mathcal{C}_s)$ is G -invariant and the restriction $\Phi(f)_{/H}$ has 1 as only eigenvalue for every $f \in \mathcal{G}$. Therefore for every $f = (A, a) \in \mathcal{G}$ we have

$$P^{-1}\Phi(f)P = \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix},$$

where $A_1 \in \mathbb{T}_q^*(\mathbb{C})$, $q = 1 + n'_1 + \dots + n'_s$ and $A_2 \in \mathcal{K}_{\eta'', r'-s}(\mathbb{C})$ with $\eta'' = (n'_{s+1}, \dots, n'_{r'})$. As $\mu_{A_1} = 1$ is the only eigenvalue of A_1 , so $P^{-1}\Phi(f)P \in \Phi(GA(n, \mathbb{C}))$. It follows that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta, r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$, where $r = r' - s + 1$ and $\eta = (q, n'_{s+1}, \dots, n'_{r'})$.

- *Case:2* Suppose that $J = \emptyset$, then 1 is the commune eigenvalue of all element of G and let v be the commune eigenvector of all element of G associated to 1. Since $\{0\} \times \mathbb{C}^n$ is G -invariant, one has $v \in \mathbb{C}^* \times \mathbb{C}^n$, so we can take $v = (1, v_1)$, $v_1 \in \mathbb{C}^n$, otherwise, if $v = (\lambda, v'_1)$ we replace v by $\frac{1}{\lambda}v$. As $\Phi(f)(1, v_1) = (1, f(v_1)) = (1, v_1)$

so $f(v_1) = \pi^{-1}(\Phi(f)(1, v_1)) = \pi^{-1}(1, v_1) = v_1$, hence $f(v_1) = Av_1 + a = v_1$. Let $P = \begin{bmatrix} 1 & 0 \\ v_1 & P_1 \end{bmatrix} \in \Phi(GA(n, \mathbb{C}))$, so for every $f = (A, a) \in \mathcal{G}$

$$\begin{aligned} P^{-1}\Phi(f)P &= \begin{bmatrix} 1 & 0 \\ -P_1^{-1}v_1 & P_1^{-1} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ a & A \end{bmatrix} \begin{bmatrix} 1 & 0 \\ v_1 & P_1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ P_1^{-1}(Av_1 + a - v_1) & P_1^{-1}AP_1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & P_1^{-1}AP_1 \end{bmatrix}. \end{aligned}$$

It follows that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta, r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$, where $r = r' + 1$ and $\eta = (1, n'_1, \dots, n'_{r'})$.

Since $u_0 \in \{1\} \times \mathbb{C}^n$ and $P \in \Phi(GA(n, \mathbb{C}))$, so $Pu_0 \in \{1\} \times \mathbb{C}^n$. \square

Denote by:

- $G' = P^{-1}GP$.
- $\mathfrak{g}' = \exp^{-1}(G') \cap \mathcal{K}_{\eta, r}(\mathbb{C}) \cap \mathcal{F}_{n+1}$.
- $\mathcal{C}(G') = \{B \in \mathcal{K}_{\eta, r}(\mathbb{C}), A'B = BA', A' \in G'\}$.
- $\mathcal{C}(G) = P\mathcal{C}(G')P^{-1}$.

Proposition 2.4. (Under above notation), we have $\exp(\mathfrak{g}) = G$. In particular, $\exp(\mathfrak{g}') = G'$.

The proof of Proposition 2.4 uses the following Lemmas:

Lemma 2.5. ([1], Lemma 4.2) Let G be an abelian subgroup of $GL(n+1, \mathbb{C})$ and $P \in GL(n+1, \mathbb{C})$ such that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta, r}(\mathbb{C})$. If $\mathfrak{g}_1 = \exp^{-1}(G) \cap (P\mathcal{K}_{\eta, r}(\mathbb{C})P^{-1})$ then:

- (i) $\exp(\mathfrak{g}_1) = G$.
- (ii) $\mathfrak{g}_1 \subset \mathcal{C}(G)$.

Lemma 2.6. ([1], Proposition 3.2) $\exp(\mathcal{K}_{\eta, r}(\mathbb{C})) = \mathcal{K}_{\eta, r}^*(\mathbb{C})$.

Lemma 2.7. (i) If $N \in P\mathcal{K}_{\eta, r}(\mathbb{C})P^{-1}$ such that $e^N \in \Phi(GA(n, \mathbb{C}))$, so $N - 2ik\pi I_{n+1} \in \mathcal{F}_{n+1}$, for some $k \in \mathbb{Z}$.

(ii) $\exp(\mathcal{F}_{n+1}) = \Phi(GA(n, \mathbb{C}))$.

(iii) $\exp(\mathcal{F}_{n+1} \cap \mathcal{K}_{\eta, r}(\mathbb{C})) = \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta, r}^*(\mathbb{C})$.

Proof. (i) Let $N' = P^{-1}NP \in \mathcal{K}_{\eta, r}(\mathbb{C})$, $M = e^N$ and $M' = P^{-1}MP$. We have $e^{N'} = M'$ and by Lemma 2.6, $M' \in \mathcal{K}_{\eta, r}^*(\mathbb{C})$. Write $M' = \text{diag}(M'_1, \dots, M'_r)$ and $N' = \text{diag}(N'_1, \dots, N'_r)$, $M'_k, N'_k \in \mathbb{T}_{n_k}(\mathbb{C})$, $k = 1, \dots, r$. Then $e^{N'} = \text{diag}(e^{N'_1}, \dots, e^{N'_r})$, so $e^{N'_1} = M'_1$. As 1 is the only eigenvalue of M'_1 , N'_1 has an eigenvalue $\mu \in \mathbb{C}$ such that $e^\mu = 1$. Thus $\mu = 2ik\pi$ for some $k \in \mathbb{Z}$. Therefore, $N'' = N' - 2ik\pi I_{n+1} \in \mathcal{F}_{n+1}$ and satisfying $e^{N''} = e^{-2ik\pi} e^{N'} = M'$. It follows by Lemma 2.1, that $N - 2ik\pi I_{n+1} = PN''P^{-1} \in P\mathcal{F}_{n+1}P^{-1} = \mathcal{F}_{n+1}$, since $P \in \Phi(GA(n, \mathbb{C}))$.

(ii) let $N = \begin{bmatrix} 0 & 0 \\ b & B \end{bmatrix} \in \mathcal{F}_{n+1}$, we have $N^k = \begin{bmatrix} 0 & 0 \\ B^{k-1}b & B^k \end{bmatrix} \in \mathcal{F}_{n+1}$ for every $k \in \mathbb{N}_0$, so

$$\begin{aligned} e^N &= I_{n+1} + \sum_{k \in \mathbb{N}_0} \frac{1}{k!} N^k \\ &= \begin{bmatrix} 1 & 0 \\ \sum_{k \in \mathbb{N}_0} \frac{1}{k!} B^{k-1}b & I_n + \sum_{k \in \mathbb{N}_0} \frac{1}{k!} B^k \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ \sum_{k \in \mathbb{N}_0} \frac{1}{k!} B^{k-1}b & e^B \end{bmatrix}. \end{aligned}$$

Then $e^N \in \Phi(GA(n, \mathbb{C}))$. It follows that $\exp(\mathcal{F}_{n+1}) \subset \Phi(GA(n, \mathbb{C}))$.

Conversely, let $M = \begin{bmatrix} 1 & 0 \\ a & A \end{bmatrix} \in \Phi(GA(n, \mathbb{C}))$. By applying Proposition 2.3 on the group generated by M , there exists $P \in \Phi(GA(n, \mathbb{C}))$ such that $M' = P^{-1}MP \in \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. By Lemma 2.6, there exists $N' \in \mathcal{K}_{\eta,r}(\mathbb{C})$ such that $e^{N'} = M'$. By (i), $N'' = N' - 2ik\pi I_{n+1} \in \mathcal{F}_{n+1}$ for some $k \in \mathbb{Z}$ and satisfying $e^{N''} = M'$. Hence, $N = PN'P^{-1} \in P\mathcal{F}_{n+1}P^{-1}$ and satisfying $e^N = Pe^{N''}P^{-1} = PM'P^{-1} = M$. By Lemma 2.1, $N \in \mathcal{F}_{n+1}$. It follows that $\Phi(GA(n, \mathbb{C})) \subset \exp(\mathcal{F}_{n+1})$.

(iii) By Lemma 2.6 and (ii) we have $\exp(\mathcal{F}_{n+1}) \cap \mathcal{K}_{\eta,r}(\mathbb{C}) \subset \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$.

Conversely, let $M = \begin{bmatrix} 1 & 0 \\ a & A \end{bmatrix} \in \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$. By Lemma 2.6, there exists $N' \in \mathcal{K}_{\eta,r}(\mathbb{C})$ such that $e^{N'} = M$. By (i), $N = N' - 2ik\pi I_{n+1} \in \mathcal{F}_{n+1}$ for some $k \in \mathbb{Z}$ and satisfying $e^N = e^{-2ik\pi} e^{N'} = M$. It follows that $\Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}) \subset \exp(\mathcal{F}_{n+1})$. The proof is complete. \square

Proof of Proposition 2.4. By Lemma 2.5.(i), we have $\exp(\mathfrak{g}_1) = G$. Since $\mathfrak{g} = \mathfrak{g}_1 \cap \mathcal{F}_{n+1}$ then

$$\exp(\mathfrak{g}) \subset \exp(\mathfrak{g}_1) = G.$$

Conversely, let $M \in G$. By lemma 2.5.(i) we have $\exp(\mathfrak{g}_1) = G$ then there exists $N \in \mathfrak{g}_1 \subset \mathcal{K}_{\eta,r}(\mathbb{C})$ such that $e^N = M$. By Lemma 2.7.(i), there exists $k \in \mathbb{Z}$ such that $N' = N - 2ik\pi I_{n+1} \in \mathcal{F}_{n+1}$. Since N and $2ik\pi I_{n+1} \in P\mathcal{K}_{\eta,r}(\mathbb{C})P^{-1}$, $N' \in \mathfrak{g}_1 \cap \mathcal{F}_{n+1} = \mathfrak{g}$. It follows that $M = e^{N'} \in \exp(\mathfrak{g})$. The proof is complete. In particular if $P = I_{n+1}$, we have $\exp(\mathfrak{g}') = G'$. \square

3. Parametrization

Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$ and $G = \Phi(\mathcal{G})$. In all the following, P denotes the matrix defined in Proposition 2.3 (i.e. $P \in \Phi(GA(n, \mathbb{C}))$) such that $P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta, r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. Recall that $G' = P^{-1}GP$ and $\mathfrak{g}' = \exp^{-1}(G') \cap \mathcal{K}_{\eta, r}(\mathbb{C}) \cap \mathcal{F}_{n+1}$, so every $A' \in G'$ is in the form

$$A' = \text{diag}(A'_1, \dots, A'_r) \in \mathcal{K}_{\eta, r}^*(\mathbb{C}), \quad A'_k \in \mathbb{T}_{n_k}^*(\mathbb{C}), \quad k = 1, \dots, r$$

and every $B' \in \mathfrak{g}'$ is in the form

$$B' = \text{diag}(B'_1, \dots, B'_r) \in \mathcal{K}_{\eta, r}(\mathbb{C}), \quad B'_k \in \mathbb{T}_{n_k}(\mathbb{C}), \quad k = 1, \dots, r$$

For every $k = 1, \dots, r$, denote by:

- $G'_k = \{A'_k, \quad A' \in G'\}$.
- $\mathfrak{g}'_k = \{B'_k, \quad B' \in \mathfrak{g}'\}$.
- $F_{G'_k} = \left\{ (A'_k - \mu_{A'_k} I_{n_k}) e_{k,i}, \quad A' \in G', \quad 1 \leq i \leq n_k - 1 \right\}$.
- $F_{\mathfrak{g}'_k} = \left\{ (B'_k - \mu_{B'_k} I_{n_k}) e_{k,i}, \quad B' \in \mathfrak{g}', \quad 1 \leq i \leq n_k - 1 \right\}$.

where $\mu_{A'_k}$ (resp. $\mu_{B'_k}$) is the only eigenvalue of A'_k . (resp. B'_k) and $(e_{k,1}, \dots, e_{k,n_k})$ is the canonical basis of \mathbb{C}^{n_k} .

- Let $h : M_{n+1}(\mathbb{C}) \rightarrow M_{n+1}(\mathbb{C})$ be the isomorphism defined by $h(B) = PBP^{-1}$. one has $h(G') = G$.

See that π is an affine map, because $\pi = (D_0, e_1)$, where $D_0 = \text{diag}(0, I_n)$. Remark that $\pi \notin GA(n, \mathbb{C})$, however $\pi : \mathbb{C}^n \rightarrow \{1\} \times \mathbb{C}^n$ is invertible.

Denote by :

- $U = \prod_{k=1}^r \mathbb{C}^* \times \mathbb{C}^{n_k-1}$, we have U is a G' -invariant open set.
- $V = \pi^{-1} \circ P(U \cap (\{1\} \times \mathbb{C}^n))$.

We have V is an open subset of \mathbb{C}^n , because $V = \pi^{-1}(P(U) \cap P(\{1\} \times \mathbb{C}^n))$, π is continuous and $P(U) \cap P(\{1\} \times \mathbb{C}^n) = P(U) \cap (\{1\} \times \mathbb{C}^n)$ is an open set of $\{1\} \times \mathbb{C}^n$ (since $P \in \Phi(GA(n, \mathbb{C}))$ and so $P(\{1\} \times \mathbb{C}^n) = \{1\} \times \mathbb{C}^n$).

- $\mathcal{C}(\mathfrak{g}') = \{B \in \mathcal{K}_{\eta, r}(\mathbb{C}), \quad A'B = BA', \quad A' \in \mathfrak{g}'\}$.
- $\mathcal{C}(\mathfrak{g}) = P\mathcal{C}(\mathfrak{g}')P^{-1}$ (i.e. $\mathcal{C}(\mathfrak{g}) = h(\mathcal{C}(\mathfrak{g}'))$).
- $\mathcal{C}(\mathcal{G}) = \Phi^{-1}(\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C})))$.
- $\mathcal{C}(\mathfrak{g}) = \Psi^{-1}(\mathcal{C}(\mathfrak{g}) \cap \mathcal{F}_{n+1})$.

Notice that by construct, $\mathcal{C}(\mathcal{G})$ is an affine subspace of $MA(n, \mathbb{C})$, since Φ is affine map. However, $\mathcal{C}(\mathfrak{g})$ is a vector subspace of $MA(n, \mathbb{C})$, since Ψ is a linear map.

Recall the following results proved in [1] for an abelian subgroup G' of $\mathcal{K}_{\eta, r}^*(\mathbb{C})$.

Lemma 3.1. ([1], Lemma 4.2) *We have $\mathcal{C}(G') = \mathcal{C}(\mathfrak{g}')$.*

By Lemma 3.1 we have $\mathcal{C}(G') = \mathcal{C}(\mathfrak{g}')$ and so $\mathcal{C}(G) = \mathcal{C}(\mathfrak{g})$. By construct, this identification can not be true in the affine case (i.e. $\mathcal{C}(\mathcal{G}) \neq \mathcal{C}(\mathfrak{g})$). Because $f_0 = (0, 0) \notin \mathcal{C}(\mathcal{G})$ and $\mathcal{C}(\mathfrak{g})$ is a vector space.

Proposition 3.2. ([1], Theorem 5.2 and Corollary 5.4) *Let G' be an abelian subgroup of $\mathcal{K}_{\eta,r}^*(\mathbb{C})$. If $\text{rank}(F_{G'_k}) = n_k - 1$ (resp. $\text{rank}(F_{\mathfrak{g}'_k}) = n_k - 1$), $k = 1, \dots, r$, then there exists a linear injective map $\varphi : \mathbb{C}^{n+1} \rightarrow \mathcal{K}_{\eta,r}(\mathbb{C})$ (resp. $\psi : \mathbb{C}^{n+1} \rightarrow \mathcal{K}_{\eta,r}(\mathbb{C})$) such that:*

- (i) $\mathcal{C}(G') \subset \varphi(\mathbb{C}^{n+1})$, (resp. $\mathcal{C}(G') \subset \psi(\mathbb{C}^{n+1})$).
- (ii) For every $v \in \mathbb{C}^{n+1}$, $\varphi(v)u_0 = v$, (resp. $\psi(v)u_0 = v$). In particular $\varphi(G'(u_0)) = G'$ and $\varphi(\mathfrak{g}'_{u_0}) = \mathfrak{g}'$ (resp. $\psi(G'(u_0)) = G'$ and $\psi(\mathfrak{g}'_{u_0}) = \mathfrak{g}'$).

Lemma 3.3. ([1], Corollaries 2.7 and 3.2.(ii)) *The linear injective map $\psi : \mathbb{C}^{n+1} \rightarrow \mathcal{K}_{\eta,r}(\mathbb{C})$ defined in Proposition 3.2 satisfies $\exp(\mathcal{C}(\mathfrak{g}')) = \mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$.*

Corollary 3.4. ([1], Proposition 6.2) *If $\overline{G'(u_0)} = \mathbb{C}^{n+1}$ (resp. $\overline{\mathfrak{g}'_{u_0}} = \mathbb{C}^{n+1}$) then φ (resp. ψ) is an isomorphism of \mathbb{C}^{n+1} on $\mathcal{C}(G')$.*

Corollary 3.5. ([1], Corollary 3.7 and Propositions 3.6, 3.2) *The restriction $\exp|_{\mathcal{K}_{\eta,r}(\mathbb{C})} : \mathcal{K}_{\eta,r}(\mathbb{C}) \rightarrow \mathcal{K}_{\eta,r}^*(\mathbb{C})$ is a local diffeomorphism.*

Lemma 3.6. *The restriction $\exp|_{(\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1})} : \mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1} \rightarrow \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$ is a local diffeomorphism.*

Proof. The proof results from Corollary 3.5 and Lemma 2.7.(iii). □

Recall the following Proposition:

Proposition 3.7. ([1], Proposition 6.6) *U is G' -invariant dense open set in \mathbb{C}^{n+1} and all orbits of G' in U are minimal in U .*

Lemma 3.8. *Under notation of Proposition 3.2, we have:*

- (i) $h(\mathfrak{g}') = \mathfrak{g}$ and $P^{-1}(\mathfrak{g}_{(1, w_0)}) = \mathfrak{g}'_{u_0}$.
- (ii) $\varphi(\mathbb{C}^{n+1}) \cap \Phi(GA(n, \mathbb{C})) \subset \varphi(\{1\} \times \mathbb{C}^n)$ and $\varphi(\mathbb{C}^{n+1}) \cap \mathcal{F}_{n+1} \subset \varphi(\{0\} \times \mathbb{C}^n)$.
- (iii) $h(\mathcal{C}(G')) = \mathcal{C}(G)$.
- (iv) $\tilde{\pi}(f'(u)) = \Psi(f')(1, u)$, for every $f' \in \mathfrak{g}$ and $u \in \mathbb{C}^n$. In particular, $\tilde{\pi}(\mathfrak{g}_u) = \mathfrak{g}_{(1, u)}$.

Proof. (i) Let $B' \in \mathfrak{g}' = \exp^{-1}(G) \cap \mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}$, so $e^{h(B')} = h(e^B) \in h(G') = G$ and since $h(\mathcal{F}_{n+1}) = \mathcal{F}_{n+1}$ so $h(B) \in \mathcal{F}_{n+1}$, then

$$h(B) \in \exp^{-1}(G) \cap h(\mathcal{K}_{\eta,r}(\mathbb{C})) \cap \mathcal{F}_{n+1} = \mathfrak{g}.$$

Conversely, we use the same proof by replacing h by h^{-1} .
 Since $h(g') = \mathfrak{g}$, so $\mathfrak{g} = h^{-1}(g') = P^{-1}g'P$. Then

$$\begin{aligned} P^{-1}(\mathfrak{g}_{(1,w_0)}) &= \{P^{-1}B(1, w_0) : B \in \mathfrak{g}\} \\ &= \{P^{-1}BPu_0 : B \in \mathfrak{g}\} \\ &= \{B'u_0 : B' \in \mathfrak{g}'\} \\ &= \mathfrak{g}'_{u_0} \end{aligned}$$

(ii) Let $v = [x_1, \dots, x_{n+1}]^T \in \mathbb{C}^{n+1}$ such that $\varphi(v) \in \Phi(GA(n, \mathbb{C}))$, so by Proposition 3.2.(ii), $\varphi(v)u_0 = v$. It follows that $x_1 = 1$ and so $v \in \{1\} \times \mathbb{C}^n$.

(iii) Since $h(G') = G$ and $AB = BA$ if and only if $h(A)h(B) = h(B)h(A)$, for every $A, B \in M_{n+1}(\mathbb{C})$, it follows that $h(\mathcal{C}(G')) = \mathcal{C}(G)$.

(iv) Let $f' = (B, b) \in \mathfrak{g}$ and $u \in \mathbb{C}^n$, then

$$\Psi(f')(1, u) = \begin{bmatrix} 0 & 0 \\ b & B \end{bmatrix} \begin{bmatrix} 1 \\ u \end{bmatrix} = (0, f'(u)) = \tilde{\pi}(f'(u)),$$

so

$$\tilde{\pi}(\mathfrak{g}_u) = \Psi(\mathfrak{g})_{(1,u)} = \mathfrak{g}_{(1,u)} = \{0\} \times \mathfrak{g}_u \subset \{0\} \times \mathbb{C}^n.$$

□

Proposition 3.9. *Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$, $G = \Phi(\mathcal{G})$ and $P \in \Phi(GA(n, \mathbb{C}))$ so that $G' = P^{-1}GP$ is a subgroup of $\mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. If $\text{rank}(F_{G'_k}) = n_k - 1$ (resp. $\text{rank}(F_{\mathfrak{g}'_k}) = n_k - 1$), $k = 1, \dots, r - 1$, then there exists two injective affine maps $\gamma : \mathbb{C}^n \rightarrow GA(n, \mathbb{C})$ and $\tilde{\gamma} : \mathbb{C}^n \rightarrow MA(n, \mathbb{C})$ (resp. $\delta : \mathbb{C}^n \rightarrow GA(n, \mathbb{C})$ and $\tilde{\delta} : \mathbb{C}^n \rightarrow MA(n, \mathbb{C})$) such that :*

(i) $\mathcal{C}(\mathcal{G}) \subset \gamma(\mathbb{C}^n)$ and $\mathcal{C}(\mathfrak{g}) \subset \tilde{\gamma}(\mathbb{C}^n)$ (resp. $\mathcal{C}(\mathcal{G}) \subset \delta(\mathbb{C}^n)$ and $\mathcal{C}(\mathfrak{g}) \subset \tilde{\delta}(\mathbb{C}^n)$).

(ii) $\gamma(\mathcal{G}(w_0)) = \mathcal{G}$ and $\tilde{\gamma}(\mathfrak{g}_{w_0}) = \mathfrak{g}$. (resp. $\delta(\mathcal{G}(w_0)) = \mathcal{G}$ and $\tilde{\delta}(\mathfrak{g}_{w_0}) = \mathfrak{g}$).

Proof. Let's prove the Proposition for $\text{rank}(F_{G_k}) = n_k - 1$. The proof for $\text{rank}(F_{\mathfrak{g}_k}) = n_k - 1$, $k = 1, \dots, r$ is analogous.

(ii) As $G(1, w_0) = \{1\} \times \mathcal{G}(w_0)$ we have $\pi(\mathcal{G}(w_0)) = G(1, w_0)$. Then

$$P^{-1} \circ \pi(\mathcal{G}(w_0)) = P^{-1}G(1, w_0) = P^{-1}GP(u_0) = G'(u_0) \quad (2)$$

By proposition 3.2, there exists a linear injective map $\varphi : \mathbb{C}^{n+1} \rightarrow \mathcal{K}_{\eta,r}(\mathbb{C})$ such that $\varphi(G'(u_0)) = G'$. Then by (2) we have

$$h \circ \varphi(P^{-1} \circ \pi(\mathcal{G}(w_0))) = h \circ \varphi(G'(u_0)) = h(G') = G. \quad (3)$$

Since $\Phi^{-1}(G) = \mathcal{G}$ then by (3), we have

$$\mathcal{G} = \Phi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \pi(\mathcal{G}(w_0)).$$

It follows that $\gamma = \Phi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \pi$ is an injective affine map from \mathbb{C}^n to $GA(n, \mathbb{C})$ and satisfying $\gamma(\mathcal{G}(w_0)) = \mathcal{G}$. (Because π and Φ are invertible affine maps; h, P are isomorphisms and φ is an injective linear map)

Let $\tilde{\gamma} = \Psi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \tilde{\pi}$, so $\tilde{\gamma}$ is an injective affine map from \mathbb{C}^n to $MA(n, \mathbb{C})$ and satisfying:

$$\begin{aligned}
\tilde{\gamma}(\mathfrak{g}_{w_0}) &= \Psi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \tilde{\pi}(\mathfrak{g}_{w_0}) \\
&= \Psi^{-1} \circ h \circ \varphi \circ P^{-1}(\mathfrak{g}_{(1, w_0)}) && \text{(by Lemma 3.8.(iv))} \\
&= \Psi^{-1} \circ h \circ \varphi(\mathfrak{g}'_{u_0}) && \text{(by Lemma 3.8.(i))} \\
&= \Psi^{-1} \circ h(\mathfrak{g}') \\
&= \Psi^{-1}(\mathfrak{g}) && \text{(by Lemma 3.8.(i))} \\
&= \mathfrak{g}
\end{aligned}$$

By analogous proof, we replace φ by ψ in the construction of γ and $\tilde{\gamma}$ and we find

$$\delta = \Phi^{-1} \circ h \circ \psi \circ P^{-1} \circ \pi \quad \text{and} \quad \tilde{\delta} = \Psi^{-1} \circ h \circ \psi \circ P^{-1} \circ \tilde{\pi} \quad (4)$$

(i) By Lemma 3.8.(iii), $h(\mathcal{C}(G')) = \mathcal{C}(G)$ and by proposition 3.2, $\mathcal{C}(G') \subset \varphi(\mathbb{C}^{n+1})$. Then we have:

$$\begin{aligned}
\mathcal{C}(\mathcal{G}) &= \Phi^{-1}(\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C}))) \\
&= \Phi^{-1}(h(\mathcal{C}(G')) \cap \Phi(GA(n, \mathbb{C}))) \\
&\subset \Phi^{-1} \circ h(\varphi(\mathbb{C}^{n+1}) \cap h^{-1}(\Phi(GA(n, \mathbb{C})))) \\
&\subset \Phi^{-1} \circ h(\varphi(\mathbb{C}^{n+1}) \cap \Phi(GA(n, \mathbb{C}))) && \text{(by Lemma 2.1)} \\
&\subset \Phi^{-1} \circ h \circ \varphi(\{1\} \times \mathbb{C}^n) && \text{(by Lemma 3.8.(ii))} \quad (5)
\end{aligned}$$

On the other hand, we have $P^{-1}(\{1\} \times \mathbb{C}^n) = \{1\} \times \mathbb{C}^n$, because $P \in \Phi(GA(n, \mathbb{C}))$. As $\{1\} \times \mathbb{C}^n = \pi(\mathbb{C}^n)$, so $\{1\} \times \mathbb{C}^n = P^{-1}(\{1\} \times \mathbb{C}^n) = P^{-1} \circ \pi(\mathbb{C}^n)$. By (5), we have

$$\mathcal{C}(\mathcal{G}) \subset \Phi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \pi(\mathbb{C}^n) = \gamma(\mathbb{C}^n).$$

By Lemma 3.1 and Proposition 3.2, $\mathcal{C}(\mathfrak{g}') = \mathcal{C}(G') \subset \varphi(\mathbb{C}^{n+1})$. By Lemma 3.8.(iii), $h(\mathcal{C}(\mathfrak{g}')) = \mathcal{C}(\mathfrak{g})$, then we have:

$$\begin{aligned}
\mathcal{C}(\mathfrak{g}) &= \Psi^{-1}(\mathcal{C}(\mathfrak{g}') \cap \mathcal{F}_{n+1}) \\
&= \Psi^{-1}(h(\mathcal{C}(\mathfrak{g}')) \cap h(\mathcal{F}_{n+1})) && \text{(by Lemma 2.1)} \\
&\subset \Psi^{-1} \circ h(\varphi(\mathbb{C}^{n+1}) \cap \mathcal{F}_{n+1}) \\
&\subset \Psi^{-1} \circ h \circ \varphi(\{0\} \times \mathbb{C}^n) && \text{(by Lemma 3.8.(i))} \quad (6)
\end{aligned}$$

On the other hand, we have $P^{-1}(\{0\} \times \mathbb{C}^n) = \{0\} \times \mathbb{C}^n$, because $P \in \Phi(GA(n, \mathbb{C}))$. As $\{0\} \times \mathbb{C}^n = \tilde{\pi}(\mathbb{C}^n)$, so $\{0\} \times \mathbb{C}^n = P^{-1}(\{0\} \times \mathbb{C}^n) = P^{-1} \circ \tilde{\pi}(\mathbb{C}^n)$. By (6), we have

$$\mathcal{C}(\mathfrak{g}) \subset \Psi^{-1} \circ h \circ \varphi \circ P^{-1} \circ \tilde{\pi}(\mathbb{C}^n) = \tilde{\gamma}(\mathbb{C}^n).$$

This completes the proof. \square

Corollary 3.10. *If $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$ (resp. $\overline{\mathfrak{g}_{w_0}} = \mathbb{C}^n$) then $\gamma : \mathbb{C}^n \rightarrow \mathcal{C}(\mathcal{G})$ and $\tilde{\gamma} : \mathbb{C}^n \rightarrow \mathcal{C}(\mathfrak{g})$ (resp. $\delta : \mathbb{C}^n \rightarrow \mathcal{C}(\mathcal{G})$ and $\tilde{\delta} : \mathbb{C}^n \rightarrow \mathcal{C}(\mathfrak{g})$) are homeomorphisms.*

Proof. Suppose that $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$. Then by Proposition 3.9 there exists an injective affine map $\gamma : \mathbb{C}^n \rightarrow GA(n, \mathbb{C})$ such that $\mathcal{C}(\mathcal{G}) \subset \gamma(\mathbb{C}^n)$. Let's prove that $\gamma(\mathbb{C}^n) \subset \mathcal{C}(\mathcal{G})$:

By proposition 3.9.(ii), we have $\gamma(\mathcal{G}(w_0)) = \mathcal{G}$. Then since γ is continuous, we have:

$$\gamma(\mathbb{C}^n) = \gamma(\overline{\mathcal{G}(w_0)}) \subset \overline{\gamma(\mathcal{G}(w_0))} = \overline{\mathcal{G}}.$$

By construction, $\mathcal{C}(\mathcal{G})$ is an affine subspace of \mathbb{C}^n then it is closed and so $\overline{\mathcal{G}} \subset \mathcal{C}(\mathcal{G})$. Hence $\gamma(\mathbb{C}^n) \subset \mathcal{C}(\mathcal{G})$. We conclude that γ is an homeomorphism of \mathbb{C}^n on $\mathcal{C}(\mathcal{G})$.

The same proof is given for $\tilde{\gamma}$, δ and $\tilde{\delta}$. □

Corollary 3.11. *If $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$ then the map $\alpha : \mathbb{C}^n \rightarrow \mathbb{C}^n$ defined by $\alpha = \gamma^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}) \circ h^{-1} \circ \Psi \circ \tilde{\gamma}$ satisfies:*

- (i) α is an open map
- (ii) $\alpha^{-1}(\mathcal{G}(w_0)) = \mathfrak{g}_{w_0}$.

Proof. If $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$ then φ is invertible so are γ and $\tilde{\gamma}$. Hence α is well defined.

- (i) Since $\exp/\mathcal{K}_{\eta,r}(\mathbb{C})$ is a local diffeomorphism (Corollary 3.5) then α is open.
- (ii) By Proposition 3.9.(ii), we have $\tilde{\gamma}^{-1}(\mathfrak{g}) = \mathfrak{g}_{w_0}$ and $\gamma(\mathcal{G}(w_0)) = \mathcal{G}$. Then

$$\begin{aligned} \alpha^{-1}(\mathcal{G}(w_0)) &= \tilde{\gamma}^{-1} \circ \Psi^{-1} \circ h \circ (\exp/(\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}))^{-1} \circ h^{-1} \circ \Phi \circ \gamma(\mathcal{G}(w_0)) \\ &= \tilde{\gamma}^{-1} \circ \Psi^{-1} \circ h \circ (\exp/(\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}))^{-1} \circ h^{-1} \circ \Phi(\mathcal{G}) \\ &= \tilde{\gamma}^{-1} \circ \Psi^{-1} \circ h \circ (\exp/(\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}))^{-1} \circ h^{-1}(G) \\ &= \tilde{\gamma}^{-1} \circ \Psi^{-1} \circ h \circ (\exp/(\mathcal{K}_{\eta,r}(\mathbb{C}) \cap \mathcal{F}_{n+1}))^{-1}(G') \\ &= \tilde{\gamma}^{-1} \circ \Psi^{-1} \circ h(g') && \text{(by definition of } g') \\ &= \tilde{\gamma}^{-1} \circ \Psi^{-1}(g) && \text{(by Lemma 3.8.(i))} \\ &= \tilde{\gamma}^{-1}(\mathfrak{g}) \\ &= \mathfrak{g}_{w_0}. \end{aligned}$$

□

Lemma 3.12. *Let $v \in \mathbb{C}^{n+1}$. Then $v \in U \cap (\{1\} \times \mathbb{C}^n)$ if and only if $\psi(v) \in \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. Moreover, $\psi^{-1}(\mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))) = U \cap (\{1\} \times \mathbb{C}^n)$.*

Proof. Let $x = [x_{1,1}, \dots, x_{1,n_1}; \dots; x_{r,1}, \dots, x_{r,n_r}]^T \in \mathbb{C}^{n+1}$. By Proposition 3.2.(i), $\psi(x) = \text{diag}(B_1, \dots, B_r) \in \mathcal{K}_{\eta,r}(\mathbb{C})$ with $B_k \in \mathbb{T}_{n_k}(\mathbb{C})$, $k = 1, \dots, r$. By Proposition 3.2.(ii), $x = \psi(x)u_0 = [B_1 e_{1,1}, \dots, B_r e_{r,1}]^T$. Therefore $B_1 e_{1,1} = [1, x_{1,2}, \dots, x_{1,n_1}]^T$ and $B_k e_{k,1} = [x_{k,1}, \dots, x_{k,n_k}]^T$, $k = 2, \dots, r$. The proof of Lemma 3.12 follows, since B_k has $x_{k,1}$ as only eigenvalue, for every $k = 1, \dots, r$. □

Lemma 3.13. ([1], Proposition 2.3) *If $A, B \in \mathcal{K}_{\eta,r}(\mathbb{C})$ satisfy $e^A e^B = e^B e^A$ then $AB = BA$.*

Lemma 3.14. *Under notation of Proposition 3.9. If $\overline{\mathfrak{g}_{w_0}} = \mathbb{C}^n$ then:*

- (i) $U \cap (\{1\} \times \mathbb{C}^n) = \psi^{-1}(\mathcal{C}(G') \cap \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}))$.
- (ii) $V = \delta^{-1}(\mathcal{C}(\mathcal{G}) \cap \Phi^{-1} \circ h(\mathcal{K}_{\eta,r}^*(\mathbb{C})))$.
- (iii) $\exp(\mathcal{C}(g') \cap \mathcal{F}_{n+1}) = \mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$.

Proof. (i) By Lemma 3.12, we have

$$\psi^{-1}(\mathcal{C}(G') \cap \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})) \subset \psi^{-1}(\mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))) = U \cap (\{1\} \times \mathbb{C}^n).$$

Conversely, let $x \in U \cap (\{1\} \times \mathbb{C}^n)$, so $x_{k,1} \neq 0$, $k = 2, \dots, r$. Then $\psi(x) \in \psi(\mathbb{C}^n) = \mathcal{C}(G')$ (Corollary 3.4). By Lemma 3.12, $\psi(x) \in \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. It follows that $\psi(x) \in \mathcal{C}(G') \cap \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$.

(ii) Recall that $\mathcal{C}(\mathcal{G}) = \Phi^{-1}(\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C})))$, by Lemma 2.1, $h(\Phi(GA(n, \mathbb{C}))) = \Phi(GA(n, \mathbb{C}))$ and by (4), $\delta = \Phi^{-1} \circ h \circ \psi \circ P^{-1} \circ \pi$. Then we have

$$\begin{aligned} V &= \pi^{-1} \circ P(U \cap (\{1\} \times \mathbb{C}^n)) \\ &= \pi^{-1} \circ P[\psi^{-1}(\mathcal{C}(G') \cap \Phi(GA(n, \mathbb{C})) \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}))] && \text{(by (i))} \\ &= \pi^{-1} \circ P \circ \psi^{-1} \circ h^{-1}[\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C})) \cap h(\mathcal{K}_{\eta,r}^*(\mathbb{C}))] && \text{(by Lemma 3.8.(iii))} \\ &= (\pi^{-1} \circ P \circ \psi^{-1} \circ h^{-1} \circ \Phi) (\Phi^{-1}[\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C}))] \cap \Phi^{-1} \circ h(\mathcal{K}_{\eta,r}^*(\mathbb{C}))) \\ &= \delta^{-1}(\mathcal{C}(\mathcal{G}) \cap \Phi^{-1} \circ h(\mathcal{K}_{\eta,r}^*(\mathbb{C}))) \end{aligned}$$

(iii) By Lemma 3.3.(i), $\exp(\mathcal{C}(g')) = \mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C})$ and by Lemma 2.7.(i), $\exp(\mathcal{F}_{n+1}) = \Phi(GA(n, \mathbb{C}))$. So $\exp(\mathcal{C}(g') \cap \mathcal{F}_{n+1}) \subset \mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. Conversely, let $M \in \mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))$. By Lemma 2.7.(iii), there exists $N \in \mathcal{F}_{n+1} \cap \mathcal{K}_{\eta,r}(\mathbb{C})$ such that $e^N = M$. As $M \in \mathcal{C}(G') = \mathcal{C}(g')$ (Lemma 3.1), so $MA = AM$ for every $A \in g'$ thus $Me^A = e^A M$ then $e^N e^A = e^A e^N$. Since $A, N \in \mathcal{K}_{\eta,r}(\mathbb{C})$, then by Lemma 3.13, $NA = AN$, so $N \in \mathcal{C}(g')$. It follows that $M = e^N \in \exp(\mathcal{C}(g') \cap \mathcal{F}_{n+1})$. \square

Corollary 3.15. *If $\overline{\mathfrak{g}_{w_0}} = \mathbb{C}^n$ then the map $\beta : \mathbb{C}^n \rightarrow \mathbb{C}^n$ defined by*

$$\beta = \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp|_{\mathcal{K}_{\eta,r}(\mathbb{C})}) \circ h^{-1} \circ \Psi \circ \tilde{\delta}$$

- (i) $\beta(\mathfrak{g}_{w_0}) = \mathcal{G}(w_0)$.
- (ii) $\beta(\mathbb{C}^n) = V$.

Proof. If $\overline{\mathfrak{g}_{w_0}} = \mathbb{C}^n$ then by Corollary 3.10, $\delta : \mathbb{C}^n \rightarrow \mathcal{C}(\mathcal{G})$ and $\tilde{\delta} : \mathbb{C}^n \rightarrow \mathcal{C}(\mathfrak{g})$ are homeomorphisms, so $\tilde{\delta}(\mathbb{C}^n) = \mathcal{C}(\mathfrak{g})$.

(i) By proposition 2.4, we have $\exp(\mathfrak{g}') = G'$. By Proposition 3.9, we have $\tilde{\delta}(\mathfrak{g}_{w_0}) = \mathfrak{g}$ and $\delta(\mathcal{G}(w_0)) = \mathcal{G}$. As δ is invertible, then $\delta^{-1}(\mathcal{G}) = \mathcal{G}(w_0)$. Therefore :

$$\begin{aligned}
\beta(\mathfrak{g}_{w_0}) &= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1} \circ \Psi \circ \tilde{\delta}(\mathfrak{g}_{w_0}) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1} \circ \Psi(\mathfrak{g}) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1}(\mathfrak{g}) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) (\mathfrak{g}') \\
&= \delta^{-1} \circ \Phi^{-1} \circ h(G') \\
&= \delta^{-1} \circ \Phi^{-1}(G) \\
&= \delta^{-1}(\mathcal{G}) \\
&= \mathcal{G}(w_0).
\end{aligned}$$

(ii) By Lemma 2.1, $h(\mathcal{F}_{n+1}) = \mathcal{F}_{n+1}$ and $h(\Phi(GA(n, \mathbb{C}))) = \Phi(GA(n, \mathbb{C}))$ and by Lemma 3.14, we have:

$$\begin{aligned}
\beta(\mathbb{C}^n) &= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1} \circ \Psi \circ \tilde{\delta}(\mathbb{C}^n) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1} \circ \Psi(\mathcal{C}(\mathfrak{g})) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) \circ h^{-1}(\mathcal{C}(\mathfrak{g}) \cap \mathcal{F}_{n+1}) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) (\mathcal{C}(\mathfrak{g}') \cap h^{-1}(\mathcal{F}_{n+1})) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h \circ (\exp/\mathcal{K}_{\eta,r}(\mathbb{C})) (\mathcal{C}(\mathfrak{g}') \cap \mathcal{F}_{n+1}) \\
&= \delta^{-1} \circ \Phi^{-1} \circ h(\mathcal{C}(G') \cap \mathcal{K}_{\eta,r}^*(\mathbb{C}) \cap \Phi(GA(n, \mathbb{C}))) \\
&= \delta^{-1} \circ \Phi^{-1}(\mathcal{C}(G) \cap \Phi(GA(n, \mathbb{C})) \cap h(\mathcal{K}_{\eta,r}^*(\mathbb{C}))) \\
&= \delta^{-1}(\mathcal{C}(\mathcal{G}) \cap \Phi^{-1} \circ h(\mathcal{K}_{\eta,r}^*(\mathbb{C}))) \\
&= V.
\end{aligned}$$

□

Proposition 3.16. *V is \mathcal{G} -invariant dense open set of \mathbb{C}^n and every orbit of \mathcal{G} , contained in V is minimal in V.*

Proof. Let $x \in V$ and $f = (A, a) \in \mathcal{G}$. Then $\pi(x) = (1, x) \in P(U)$ and $\Phi(f) \in G$. By Proposition 3.7, U is G' -invariant so $P(U)$ is G -invariant. Then $\Phi(f)(1, x) = (1, f(x)) \in P(U)$, so $f(x) = \pi^{-1}(1, f(x)) \in \pi^{-1} \circ P(U) = V$. It follows that V is \mathcal{G} -invariant.

By proposition 3.7, we have all orbits of $G' = P^{-1}GP$ and contained in U are minimal in U . Let $x \in V$ and $y \in \overline{\mathcal{G}(x)} \cap V$, so $P^{-1} \circ \pi(x) = P^{-1}(1, x) \in U$ and also $P^{-1}(1, y) \in U$. We have

$$\{1\} \times \overline{\mathcal{G}(x)} = \overline{G(1, x)} = P \left(\overline{G'(P^{-1}(1, x))} \right). \quad (7)$$

As $P^{-1}(1, x), P^{-1}(1, y) \in U$, then

$$\overline{G'(P^{-1}(1, x))} \cap U = \overline{G'(P^{-1}(1, y))} \cap U.$$

Therefore

$$\begin{aligned} (\{1\} \times \overline{\mathcal{G}(x)}) \cap P(U) &= \overline{G(1, x)} \cap P(U) \\ &= P(\overline{G'(P^{-1}(1, x))} \cap U) \\ &= P(\overline{G'(P^{-1}(1, y))} \cap U) \\ &= \overline{G(1, y)} \cap P(U) \\ &= (\{1\} \times \overline{\mathcal{G}(y)}) \cap P(U) \quad (\text{by (7)}) \end{aligned}$$

Thus

$$(\{1\} \times \overline{\mathcal{G}(x)}) \cap P(U \cap (\{1\} \times \mathbb{C}^n)) = (\{1\} \times \overline{\mathcal{G}(y)}) \cap P(U \cap (\{1\} \times \mathbb{C}^n))$$

Then

$$\overline{\mathcal{G}(x)} \cap [\pi^{-1} \circ P(U \cap (\{1\} \times \mathbb{C}^n))] = \overline{\mathcal{G}(y)} \cap [\pi^{-1} \circ P(U \cap (\{1\} \times \mathbb{C}^n))].$$

So

$$\overline{\mathcal{G}(x)} \cap V = \overline{\mathcal{G}(y)} \cap V.$$

The proof is complete. \square

Lemma 3.17. $\mathbb{C}^n \setminus V$ is union of at most n , \mathcal{G} -invariant affine spaces with dimension less to $n - 1$.

Proof. We have $\mathbb{C}^{n+1} \setminus U = \bigcup_{k=1}^r H_k$, where

$$H_k = \{u = (u_1, \dots, u_r) \in \mathbb{C}^{n+1} : u_k \in \{0\} \times \mathbb{C}^{n_k-1}, u_i \in \mathbb{C}^* \times \mathbb{C}^{n_i-1}, 1 \leq i \leq r, i \neq k\}$$

is a vector space G' -invariant with dimension n , $k = 1, \dots, r$. Since $r \leq n + 1$, $\mathbb{C}^{n+1} \setminus U$ is a union of at most $n + 1$, G' -invariant vector spaces with dimension n . Then $\mathbb{C}^{n+1} \setminus P(U)$ is a union of at most $n + 1$, G -invariant vector spaces with dimension n . As $H_1 \cap (\{1\} \times \mathbb{C}^n) = \emptyset$, so

$$\begin{aligned} (\{1\} \times \mathbb{C}^n) \setminus (U \cap (\{1\} \times \mathbb{C}^n)) &= (\mathbb{C}^{n+1} \setminus U) \cap (\{1\} \times \mathbb{C}^n) \\ &= \bigcup_{k=2}^r (H_k \cap (\{1\} \times \mathbb{C}^n)) \quad (8). \end{aligned}$$

Since $P(\{1\} \times \mathbb{C}^n) = \{1\} \times \mathbb{C}^n$, then by (8), we have

$$\begin{aligned} \mathbb{C}^n \setminus V &= \pi^{-1}(\{1\} \times \mathbb{C}^n) \setminus \pi^{-1} \circ P(U \cap (\{1\} \times \mathbb{C}^n)) \\ &= \pi^{-1} \circ P((\{1\} \times \mathbb{C}^n) \setminus (U \cap (\{1\} \times \mathbb{C}^n))) \\ &= \pi^{-1} \circ P\left(\bigcup_{k=2}^r (H_k \cap (\{1\} \times \mathbb{C}^n))\right) \\ &= \bigcup_{k=2}^r E_k, \end{aligned}$$

where $E_k = \pi^{-1} \circ P(H_k \cap (\{1\} \times \mathbb{C}^n))$ is \mathcal{G} -invariant affine space for every $2 \leq k \leq r$. As $H_k \neq \{1\} \times \mathbb{C}^n$, then $\dim(H_k \cap (\{1\} \times \mathbb{C}^n)) \leq n-1$, so $\dim(E_k) \leq n-1$. It follows that $\mathbb{C}^n \setminus V$ is union of at most n spaces E_k , since $r \leq n+1$. \square

Corollary 3.18. *If \mathcal{G} has a dense orbit in \mathbb{C}^n then all orbits of V are dense in \mathbb{C}^n .*

Proof. Let $u \in \mathbb{C}^n$ such that $\overline{\mathcal{G}(u)} = \mathbb{C}^n$. By Lemma 3.17, $\mathbb{C}^n \setminus V$ is union of at most n , \mathcal{G} -invariant affine spaces with dimension less to $n-1$, so $u \in V$. Let $v \in V$. Since $V = \overline{\mathcal{G}(u)} \cap V$ then by Proposition 3.16, $\overline{\mathcal{G}(v)} \cap V = \overline{\mathcal{G}(u)} \cap V = V$. As $\overline{V} = \mathbb{C}^n$, one has $\overline{\mathcal{G}(v)} = \mathbb{C}^n$. \square

4. Proof of Theorem 1.1 and corollary 1.2

4.1. Proof of Theorem 1.1. Let \mathcal{G} be an abelian affine subgroup of $GA(n, \mathbb{C})$ and let $P \in \Phi(GA(n, \mathbb{C}))$ defined in Proposition 2.3. Recall that $G = \Phi(\mathcal{G})$ and $\mathfrak{g} = \exp^{-1}(G) \cap (PK_{\eta,r}(\mathbb{C})P^{-1}) \cap \mathcal{F}_{n+1}$.

ii) \implies i) is clear.

i) \implies ii) This follows from Corollary 3.18, since $u_0 \in U$ and so $w_0 = \pi^{-1} \circ P(u_0) \in V$.

iii) \implies ii) : Suppose that $\overline{\mathfrak{g}_{w_0}} = \mathbb{C}^n$. Then by Corollary 3.15 there exists a continuous map $\beta : \mathbb{C}^n \longrightarrow \mathbb{C}^n$ such that $\beta(\mathfrak{g}_{w_0}) = \mathcal{G}(w_0)$ and $\beta(\mathbb{C}^n) = V$. By continuity, one has:

$$V = \beta(\mathbb{C}^n) = \beta(\overline{\mathfrak{g}_{w_0}}) \subset \overline{\beta(\mathfrak{g}_{w_0})} = \overline{\mathcal{G}(w_0)}$$

Since V is a dense open set in \mathbb{C}^n then $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$.

ii) \implies iii) : Suppose that $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$.

By Corollary 3.11, there exists an open map $\alpha : \mathbb{C}^n \longrightarrow \mathbb{C}^n$ such that $\mathfrak{g}_{w_0} = \alpha^{-1}(\mathcal{G}(w_0))$. Then:

$$\mathbb{C}^n = \alpha^{-1}(\mathbb{C}^n) = \alpha^{-1}(\overline{\mathcal{G}(w_0)}) \subset \overline{\alpha^{-1}(\mathcal{G}(w_0))} = \overline{\mathfrak{g}_{w_0}}$$

\square

4.2. Proof of Corollary 1.2. Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$ having a dense orbit. By Theorem 1.1, $\overline{\mathcal{G}(w_0)} = \mathbb{C}^n$. Then by Proposition 3.2 and corollary 3.10, there exists an invertible affine map $\gamma : \mathbb{C}^n \rightarrow \mathcal{C}(\mathcal{G})$ such that $\gamma(\mathcal{G}(w_0)) = \mathcal{G}$. Therefore:

$$\gamma(\mathbb{C}^n) = \gamma(\overline{\mathcal{G}(w_0)}) = \overline{\gamma(\mathcal{G}(w_0))} = \overline{\mathcal{G}}.$$

It follows that $\overline{\mathcal{G}}$ is an affine space of dimension n .

5. Finitely generated subgroups

5.1. Proof of Theorem 1.3.

Denote by $v_0 = Pu_0 = (1, w_0)$. So $w_0 = p_2(v_0)$.

Lemma 5.1. *We have $p_2(\Psi(f')v_0) = f'(w_0)$, for every $f' \in \mathfrak{g}$. In particular, $\mathfrak{g}_{w_0} = p_2(\mathfrak{g}_{v_0})$.*

Proof. Let $f' = (B, b) \in \mathfrak{g}$, then

$$\begin{aligned} \Psi(f')v_0 &= \begin{bmatrix} 0 & 0 \\ b & B \end{bmatrix} \begin{bmatrix} 1 \\ w_0 \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ b + Bv_0 \end{bmatrix}. \end{aligned}$$

So $p_2(\Psi(f')v_0) = b + Bv_0 = f'(w_0)$. \square

Proposition 5.2. ([1], Proposition 8.1) *Let G be a subgroup of $GL(n+1, \mathbb{C})$ generated by A_1, \dots, A_p . Let $B_1, \dots, B_p \in \mathfrak{g}$ such that $A_k = e^{B_k}$, $k = 1, \dots, p$. Then:*

$$\mathfrak{g}_{v_0} = \sum_{k=1}^p \mathbb{Z}B_k v_0 + \sum_{k=1}^r 2i\pi \mathbb{Z}P e^{(k)}.$$

Proposition 5.3. (Under notations of Proposition 2.3) *Let \mathcal{G} be an abelian subgroup of $GA(n, \mathbb{C})$ generated by f_1, \dots, f_p and let $f'_1, \dots, f'_p \in \mathfrak{g}$ such that $\Phi(f_k) = e^{\Psi(f'_k)}$, $k = 1, \dots, p$. Then:*

$$\mathfrak{g}_{w_0} = \sum_{k=1}^p \mathbb{Z}f'_k(w_0) + \sum_{k=1}^r 2i\pi \mathbb{Z}(p_2 \circ P e^{(k)}).$$

Proof. Let $G = \Phi(\mathcal{G})$. Then G is generated by $\Phi(f_1), \dots, \Phi(f_p)$ and \mathfrak{g} is the additive group generated by $\Psi(f'_1), \dots, \Psi(f'_p)$. By proposition 5.2 we have

$$\mathfrak{g}_{v_0} = \sum_{k=1}^p \mathbb{Z}\Psi(f'_k)v_0 + \sum_{k=1}^r 2i\pi \mathbb{Z}P e^{(k)}.$$

By Lemma 5.1, we have $\mathfrak{g}_{w_0} = p_2(\mathfrak{g}_{v_0})$ and $p_2(\Psi(f'_k)v_0) = f'_k(w_0)$. then

$$\begin{aligned} \mathfrak{g}_{w_0} &= p_2 \left(\sum_{k=1}^p \mathbb{Z} \Psi(f'_k)v_0 + \sum_{k=1}^r 2i\pi \mathbb{Z} P e^{(k)} \right). \\ &= \sum_{k=1}^p \mathbb{Z} f'_k(w_0) + \sum_{k=1}^r 2i\pi \mathbb{Z} (p_2 \circ P e^{(k)}). \end{aligned}$$

□

Recall the following Proposition which was proven in [2]:

Proposition 5.4. (cf. [2], page 35). *Let $H = \mathbb{Z}u_1 + \dots + \mathbb{Z}u_p$ with $u_k = (u_{k,1}, \dots, u_{k,n}) \in \mathbb{C}^n$ and $u_{k,i} = \operatorname{Re}(u_{k,i}) + i\operatorname{Im}(u_{k,i})$, $k = 1, \dots, p$, $i = 1, \dots, n$. Then H is dense in \mathbb{C}^n if and only if for every $(s_1, \dots, s_p) \in \mathbb{Z}^p \setminus \{0\}$:*

$$\operatorname{rank} \begin{bmatrix} \operatorname{Re}(u_{1,1}) & \dots & \dots & \operatorname{Re}(u_{p,1}) \\ \vdots & \vdots & \vdots & \vdots \\ \operatorname{Re}(u_{1,n}) & \dots & \dots & \operatorname{Re}(u_{p,n}) \\ \operatorname{Im}(u_{1,1}) & \dots & \dots & \operatorname{Im}(u_{p,1}) \\ \vdots & \vdots & \vdots & \vdots \\ \operatorname{Im}(u_{1,n}) & \dots & \dots & \operatorname{Im}(u_{p,n}) \\ s_1 & \dots & \dots & s_p \end{bmatrix} = 2n + 1.$$

Proof of Theorem 1.3: This follows directly from Theorem 1.1, Propositions 5.3 and 5.4.

5.2. Proof of Corollaries 1.4 and 1.5.

Proof of Corollary 1.4: We show first that if $H = \mathbb{Z}u_1 + \dots + \mathbb{Z}u_m$, $u_k \in \mathbb{C}^n$ with $m \leq 2n$, then H can not be dense: Write $u_k \in \mathbb{C}^n$, $u_k = \operatorname{Re}(u_k) + i\operatorname{Im}(u_k)$ and $v_k = [\operatorname{Re}(u_k); \operatorname{Im}(u_k); s_k]^T \in \mathbb{R}^{2n+1}$, $1 \leq k \leq m$. Since $m \leq 2n$, it follows that $\operatorname{rank}(v_1, \dots, v_m) \leq 2n$, and so H is not dense in \mathbb{C}^n by Proposition 5.4.

Now, by applying Theorem 1.3 and the fact that $m = p + r \leq 2n$, the Corollary 1.4 follows. □

Proof of Corollary 1.5: Since $p \leq n$ and $r \leq n$ then $p + r \leq 2n$. Corollary 1.5 follows from Corollary 1.4. □

5.3. The case: $n = 1$.

Let \mathcal{G} be a subgroup of $GA(1, \mathbb{C})$, generated by $f_1 = (a_1, \lambda_1), \dots, f_p = (a_p, \lambda_p)$, where $a_k = \rho_k e^{i\theta_k}$, $\lambda_k = r_k e^{i\alpha_k}$, $\rho_k, r_k > 0$, $\theta_k, \alpha_k \in \mathbb{R}$, $1 \leq k \leq p$. Therefore $G = \Phi(\mathcal{G})$ is an abelian subgroup of $GL(2, \mathbb{C})$. Let $f'_1, \dots, f'_p \in \mathfrak{g}$ such that $e^{\Psi(f'_k)} = \Phi(f_k)$, $1 \leq k \leq p$. Since \mathcal{G} is abelian then there are two cases:

◇ *Case 1: \mathcal{G} is a group of translations.*

In this case $\mathcal{G}(x) = T_x \left(\sum_{k=1}^p \lambda_k \mathbb{Z} \right)$. Then \mathcal{G} has a dense orbit if and only if $\mathcal{G}(0) =$

$\sum_{k=1}^p \lambda_k \mathbb{Z}$ has a dense orbit, which is equivalent, by Proposition 5.4, to:

For every $(s_1, \dots, s_p, t) \in \mathbb{Z}^{p+1} \setminus \{0\}$:

$$\text{rank} \begin{bmatrix} \log(r_1) & \dots & \log(r_p) & 0 \\ \alpha_1 & \dots & \alpha_p & 2\pi \\ s_1 & \dots & s_p & t \end{bmatrix} = 3.$$

◇ *Case 2 : \mathcal{G} contains no translation:*

In this case $w_0 = (1, 1)$, $a_k \neq 1$ for every $k = 1, \dots, p$ and $f'_k = (B_k, b_k)$ is defined by:

$$B_k = \begin{bmatrix} 0 & 0 \\ b_k & \mu_k \end{bmatrix}, \quad \text{where } b_k = \log(r_k) + i\theta_k \quad \text{and} \quad \mu_k = \frac{\lambda_k(\log(r_k) + i\theta_k)}{(a_k - 1)}.$$

Since $w_0 = 1$ then

$$\begin{aligned} \mathfrak{g}_{w_0} &= f'_1(w_0)\mathbb{Z} + \dots + f'_p(w_0)\mathbb{Z} + 2i\pi\mathbb{Z} \\ &= (b_1 + \mu_1)\mathbb{Z} + \dots + (b_p + \mu_p)\mathbb{Z} + 2i\pi\mathbb{Z} \end{aligned}$$

Moreover, the property \mathcal{D} in this cases is as follow:

f_1, \dots, f_p satisfy property \mathcal{D} if and only if for every $(s_1, \dots, s_p, t) \in \mathbb{Z}^{p+1} - \{0\}$:

$$\text{rank} \begin{bmatrix} \log|b_1 + \mu_1| & \dots & \log|b_p + \mu_p| & 0 \\ \arg(b_1 + \mu_1) & \dots & \arg(b_p + \mu_p) & 2\pi \\ s_1 & \dots & s_p & t \end{bmatrix} = 3.$$

where $\arg(z) = \theta \in [0, 2\pi[$ for every $z = re^{i\theta} \in \mathbb{C}$, $r > 0$. Therefore, from Theorem 1.3, we obtain:

Proposition 5.5. *If \mathcal{G} is a subgroup of $GA(1, \mathbb{C})$ generated by $f_1 = (a_1, \lambda_1), \dots, f_p = (a_p, \lambda_p)$, where $a_k = \rho_k e^{i\theta_k} \neq 1$, $\lambda_k = r_k e^{i\alpha_k}$, $\rho_k, r_k > 0$, $\theta_k, \alpha_k \in \mathbb{R}$, $1 \leq k \leq p$. Then the following are equivalent:*

(i) G is minimal (i.e. every orbit of \mathcal{G} is dense in \mathbb{C}).

(ii) G has a dense orbit

(iii) $\sum_{k=1}^p (b_k + \mu_k)\mathbb{Z} + 2i\pi\mathbb{Z}$ is dense in \mathbb{C} , where $b_k = \log(r_k) + i\theta_k$ and $\mu_k = (\log \rho_k + i\theta_k) \left(1 + \frac{\lambda_k}{a_k - 1} \right)$.

(iv) for every $(s_1, \dots, s_p, t) \in \mathbb{Z}^{p+1} \setminus \{0\}$:

$$\text{rank} \begin{bmatrix} \log|b_1 + \mu_1| & \dots & \log|b_p + \mu_p| & 0 \\ \arg(b_1 + \mu_1) & \dots & \arg(b_p + \mu_p) & 2\pi \\ s_1 & \dots & s_p & t \end{bmatrix} = 3.$$

6. Examples

Example 6.1. Let $a_k = e^{\pi(s_k + it_k)}$, $s_k, t_k \in \mathbb{Q}$, $1 \leq k \leq p$ and let \mathcal{G} be the subgroup of translations of \mathbb{C} generated by T_{a_1}, \dots, T_{a_p} . Then \mathcal{G} has no dense orbit.

Proof. We can write $s_k = \frac{m_k}{q}$ and $t_k = \frac{p_k}{q}$, $q \in \mathbb{N}^*$ and $m_k, p_k \in \mathbb{Z}$ for every $1 \leq k \leq p$. One has: $\mathfrak{g}_x = T_x(\mathfrak{g}_0)$, where

$$\mathfrak{g}_0 = \frac{\pi}{q} \left(\sum_{k=1}^p (m_k + ip_k)\mathbb{Z} + 2iq\mathbb{Z} \right) \subset \frac{\pi}{q} (\mathbb{Z} + i\mathbb{Z}).$$

So $\overline{\mathfrak{g}_0} \neq \mathbb{C}$. By Theorem 1.1, $\overline{\mathcal{G}(0)} \neq \mathbb{C}$. It follows by Proposition 5.5, that $\overline{\mathcal{G}(x)} \neq \mathbb{C}$. \square

Example 6.2. Let \mathcal{G} the group generated by $f_1 = (A_1, a_1)$, $f_2 = (A_2, a_2)$ and $f_3 = (A_3, a_3)$, where:

$$A_1 = \text{diag}(e^i, 1), \quad a_1 = (i(1 - e^i), 0), \quad A_2 = \text{diag}(1, e^i), \quad a_2 = (0, 2i(e^{-i} - 1)).$$

$$A_3 = \text{diag} \left(e^{\frac{-\sqrt{3}}{2\pi} + i\left(\frac{\sqrt{5}}{2} - \frac{\sqrt{3}}{2\pi}\right)}, e^{\frac{-\sqrt{2}}{\pi} + i\left(\frac{\sqrt{2}}{2\pi} - \frac{\sqrt{7}}{2}\right)} \right) \quad \text{and} \quad a_3 = (0, 0).$$

Then every orbit in $V = \mathbb{C}^* \times \mathbb{C}^*$ is dense in \mathbb{C}^2 .

Proof. Denote by $G = \Phi(\mathcal{G})$. Then G generated by

$$A'_1 = \Phi(f_1) = \begin{bmatrix} 1 & 0 & 0 \\ i(1 - e^i) & e^i & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A'_2 = \Phi(f_2) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2i(e^{-i} - 1) & 0 & e^i \end{bmatrix},$$

and

$$A'_3 = \Phi(f_3) = \text{diag} \left(1, e^{\frac{-\sqrt{3}}{2\pi} + i\left(\frac{\sqrt{5}}{2} - \frac{\sqrt{3}}{2\pi}\right)}, e^{\frac{-\sqrt{2}}{\pi} + i\left(\frac{\sqrt{2}}{2\pi} - \frac{\sqrt{7}}{2}\right)} \right).$$

Let $f'_1 = (B_1, b_1)$, $f'_2 = (B_2, b_2)$ and $f'_3 = (B_3, b_3)$ such that $e^{\Psi(f'_k)} = A'_k$, $k = 1, 2, 3$. We have

$$B_1 = \text{diag}(i, 0), \quad B_2 = \text{diag}(0, i),$$

$$B_3 = \text{diag} \left(\frac{-\sqrt{3}}{2\pi} + i \left(\frac{\sqrt{5}}{2} - \frac{\sqrt{3}}{2\pi} \right), \frac{-\sqrt{2}}{\pi} + i \left(\frac{\sqrt{2}}{2\pi} - \frac{\sqrt{7}}{2} \right) \right),$$

$$b_1 = (1, 0), \quad b_2 = (0, -2), \quad \text{and} \quad b_3 = (0, 0).$$

Here, we have:

- $P = I_3$, $U = (\mathbb{C}^*)^3$, $u_0 = (1, 1, 1)$.
- $V = \mathbb{C}^* \times \mathbb{C}^*$, $w_0 = (1, 1)$.

The additive group $\mathfrak{g} = \exp^{-1}(G) \cap (\mathcal{K}_{(1,1,1),3}(\mathbb{C})) \cap \mathcal{F}_3$ is generated by N_1, N_2 and N_3 , where

$$N_k = \Psi(f'_k) = \begin{bmatrix} 0 & 0 \\ b_k & B_k \end{bmatrix}, \quad 1 \leq k \leq 3$$

In the other hand, for every $(s_1, s_2, s_3, t_1, t_2) \in \mathbb{Z}^5 \setminus \{0\}$, one has the determinant:

$$\begin{aligned} \Delta &= \begin{vmatrix} \operatorname{Re}(B_1 w_0 + b_1) & \operatorname{Re}(B_2 w_0 + b_2) & \operatorname{Re}(B_3 w_0 + b_3) & 0 & 0 \\ \operatorname{Im}(B_1 w_0 + b_1) & \operatorname{Im}(B_2 w_0 + b_2) & \operatorname{Im}(B_3 w_0 + b_3) & 2\pi e_1 & 2\pi e_2 \\ & s_1 & s_2 & s_3 & t_1 & t_2 \end{vmatrix} \\ &= \begin{vmatrix} -\frac{\sqrt{3}}{2\pi} & 1 & 0 & 0 & 0 \\ -\frac{\sqrt{2}}{\pi} & 0 & -2 & 0 & 0 \\ \frac{\sqrt{5}}{2} - \frac{\sqrt{3}}{2\pi} & 1 & 0 & 2\pi & 0 \\ \frac{\sqrt{2}}{2\pi} - \frac{\sqrt{7}}{2} & 0 & 1 & 0 & 2\pi \\ s_1 & s_2 & s_3 & t_1 & t_2 \end{vmatrix} \\ &= -2\pi \left((4s_1)\pi - (2s_3)\sqrt{2} + (2s_2)\sqrt{3} + t_1\sqrt{5} + t_2\sqrt{7} \right). \end{aligned}$$

Since $\pi, \sqrt{2}, \sqrt{3}, \sqrt{5}$ and $\sqrt{7}$ are rationally independent then $\Delta \neq 0$ for every $(s_1, s_2, s_3, t_1, t_2) \in \mathbb{Z}^5 \setminus \{0\}$. It follows that:

$$\operatorname{rank} \begin{bmatrix} -\frac{\sqrt{3}}{2\pi} & 1 & 0 & 0 & 0 \\ -\frac{\sqrt{2}}{\pi} & 0 & -2 & 0 & 0 \\ \frac{\sqrt{5}}{2} - \frac{\sqrt{3}}{2\pi} & 1 & 0 & 2\pi & 0 \\ \frac{\sqrt{2}}{2\pi} - \frac{\sqrt{7}}{2} & 0 & 1 & 0 & 2\pi \\ s_1 & s_2 & s_3 & t_1 & t_2 \end{bmatrix} = 5$$

and by Theorem 1.3, \mathcal{G} has a dense orbit and every orbit of V is dense in \mathbb{C}^2 . \square

REFERENCES

1. A. Ayadi and H. Marzougui, *Dense orbits for abelian subgroups of $GL(n, \mathbb{C})$* , *Foliations 2005*: 47–69. World Scientific, Hackensack, NJ, 2006.
2. Waldschmidt, M., *Topologie des points rationnels.*, Cours de troisième Cycle, Université P. et M. Curie (Paris VI), 1994/95.
3. M. Javaheri, *A generalization of Dirichlet approximation theorem for the affine actions on real line*, *Journal of Number Theory* **128** (2008) 1146–1156.
4. R.S. Kulkarni, *Dynamics of linear and affine maps*, arXiv:0709.1353v1, 2007.
5. V. Bergelson, M. Misiurewicz and S. Senti, *Affine actions of a free semigroup on the real line* *Ergod. Th. and Dynam. Sys.* **26**, (2006), 1285–1305.
6. T. V. Budnytska, *Classification of topologically conjugate affine mapping*, *Ukrainian Mathematical Journal*, **61**, No. 1, (2009) 164–170.

ADLENE AYADI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES OF GAFSA, UNIVERSITY OF GAFSA, GAFSA, TUNISIA
E-mail address: adleneso@yahoo.fr