

Geometric Hyperplanes: Desargues Encodes Doily Metod Saniga

▶ To cite this version:

| Metod Saniga. Geometric Hyperplanes: Desargues Encodes Doily. 2010. hal-00550447

HAL Id: hal-00550447 https://hal.science/hal-00550447

Preprint submitted on 28 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geometric Hyperplanes: Desargues Encodes Doily

Metod Saniga

¹Astronomical Institute, Slovak Academy of Sciences, SK-05960 Tatranská Lomnica Slovak Republic (msaniga@astro.sk)

Abstract

It is shown that the structure of the generalized quadrangle of order two is fully encoded in the properties of the Desargues configuration. A point of the quadrangle is represented by a geometric hyperplane of the Desargues configuration and its line by a set of three hyperplanes such that one of them is the complement of the symmetric difference of the remaining two and they all share a pair of non-collinear points.

Keywords: Geometric Hyperplane – Desargues Configuration – Generalized Quadrangle of Order Two

Certain small v_3 -configurations exhibit a lot of remarkable properties and seem to be intricately linked with each other. A special relation in this respect is that where one configuration fully determines or, so to speak, encodes the properties of the other. A particularly nice illustration of such a link is the Fano plane (i. e., the unique 7_3 -configuration) encoding the structure of the split Cayley hexagon of order two [1]. The purpose of this research note is to point out — also in view of possible physical applications — a similar relation between other two well-known v_3 -configurations, the Desargues configuration and the generalized quadrangle of order two.

The Desargues configuration is one of the most important point-line incidence structures. It consists of 10 points and 10 lines, with three points on a line and three lines through a point. There exist altogether ten such 10_3 -configurations (see, for example, [2]). The Desargues configuration is, unlike the others, flag-transitive and the only one where for *each* of its points the three points that are not collinear with it lie on a line. This configuration is depicted in Figure 1, left, in a form showing its automorphism of order three (after Polster, [3]). The generalized quadrangle of order two, GQ(2, 2), is another prominent point-line geometry. It has 15 points/lines, also with three points/lines on a line/per point. And although there are as many as $245\,342$ such 15_3 -configurations, GQ(2, 2), apart from being flag-transitive, is the only one that is *triangle-free* (see, for example, [4, 5]). Its well-known picture, showing an automorphism of order five and often dubbed the doily, is given in Figure 1, right.

These two configurations can be linked in a number of ways. The one we shall briefly describe here, believing it is novel, relies on the concept of a geometric hyperplane [6]. A geometric hyperplane of a point-line incidence geometry is a point-subset of the geometry such that each line of the geometry meets the subset in one or all points. Employing Figure 1, left, it is quite straightforward to verify that the Desargues configuration is endowed with 15 distinct geometric hyperplanes. They form two different classes and are sketched in Figure 2; any of the first ten (Nos. 1 to 10) comprises four points, namely a point and the three points non-collinear with it, whilst each of the remaining five hyperplanes (Nos. 11 to 15) features six points on three lines forming a triangle and a fourth line cutting the three in triangle's "midpoints." Let us now construct the point-line incidence geometry where a point is a geometric hyperplane of the Desargues configuration and a line comprises three hyperplanes such that any of them is the complement of the symmetric difference of the remaining two and they all have a pair of non-collinear points in common; a brief inspection of Figure 3 confirms that this geometry is indeed isomorphic to GQ(2, 2).

Acknowledgement

This work was partially supported by the VEGA grant agency projects 2/0092/09 and 2/0098/10.




Figure 1: Left: — A picture of the Desargues configuration: small circles stand for its points, whereas its lines are represented by triples of points on common straight segments (6), arcs of circles (3) and a big circle. Right: — A picture of the generalized quadrangle of order two, the doily: as in the preceding case, circles stand for its points, while its lines are given by point triples joined by the same segments (10) and/or arcs of circles (5).




Figure 2: The 15 geometric hyperplanes (filled circles) of the Desargues configuration.




Figure 3: A diagrammatical illustration of the representation of GQ(2, 2) in terms of geometric hyperplanes of the Desargues configuration. (The reader familiar with the structure of the doily will recognize that, e.g., the hyperplanes 11 to 15 form an ovoid.)

References

- [1] Polster B, Schroth AE, and Van Maldeghem H. Generalized flatland. Math Intelligencer 2001:23:33–47.
- [2] Grünbaum B. Configurations of Points and Lines. Graduate Studies in Mathematics 103, American Mathematical Society: Providence; 2009.
- [3] Polster B. A Geometrical Picture Book. Springer: New York; 1998.
- [4] Betten A, Brinkmann G, and Pisanski T. Counting symmetric configurations v_3 . Discr Applied Math 2000;99:331–338.
- [5] Boben M, Grünbaum B, Pisanski T, and Žitnik A. Small triangle-free configurations of points and lines. Discr Comput Geometry 2006;35:405–427.
- [6] Ronan MA. Embeddings and hyperplanes of discrete geometries. European J Combin 1987;8:179–185.