



HAL
open science

HIV testing trends among gay men in Scotland, UK (1996-2005): Implications for HIV testing policies and prevention

Lisa M Williamson, Paul Flowers, Christina Knussen, Graham J Hart

► **To cite this version:**

Lisa M Williamson, Paul Flowers, Christina Knussen, Graham J Hart. HIV testing trends among gay men in Scotland, UK (1996-2005): Implications for HIV testing policies and prevention. *Sexually Transmitted Infections*, 2009, 85 (7), pp.550. 10.1136/sti.2008.033886 . hal-00552798

HAL Id: hal-00552798

<https://hal.science/hal-00552798>

Submitted on 6 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIV testing trends among gay men in Scotland, UK (1996-2005): Implications for HIV testing policies and prevention

Running head: HIV testing among gay men in Scotland

Lisa M Williamson^a

Paul Flowers^b

Christina Knussen^b

Graham J Hart^c

^a **MRC Social and Public Health Sciences Unit, Glasgow**

^b **Division of Psychology, School of Life Sciences, Glasgow Caledonian
University, Glasgow**

^c **Centre for Sexual Health and HIV Research, Research Department of Infection
and Population Health, University College London, London**

**Correspondence to: Lisa M Williamson, MRC Social and Public Health Sciences
Unit, 4 Lilybank Gardens, Glasgow, G12 8RZ, UK.**

Tel: +44 (0) 141 357 3949. Fax: +44 (0) 141 337 2389.

Email: Lisa@sphsu.mrc.ac.uk

**Key words: HIV testing; gay men; community-based surveys; sexual risk
behaviour**

ABSTRACT

Objective: To examine trends in the HIV testing behaviour of gay men in Scotland over a ten-year period.

Methods: Seven cross-sectional surveys in commercial gay venues in Glasgow and Edinburgh (1996-2005). 9613 men completed anonymous, self-complete questionnaires (70% average response rate).

Results: Among 8305 respondents included in these analyses, HIV testing increased between 1996 and 2005, from 49.7% to 57.8% ($P < 0.001$). The proportion of men who had tested recently (in the calendar year of, or immediately prior to, the survey) increased from 28.4% in 1996 to 33.2% in 2005, when compared with those who have tested but not recently, and those who have never tested (AOR 1.31, 95% CI 1.13-1.52). However, among ever testers, there was no increase in rates of recent testing. Recent testing decreased with age: 31.3% of the <25, 30.3% of the 25-34, 23.2% of the 35-44, and 21.2% of the >44 years age groups had tested recently. Among men reporting two or more unprotected anal intercourse partners in the previous year, only 41.4% had tested recently.

Conclusions: HIV testing among gay men in Scotland increased between 1996 and 2005, and corresponds with the Scottish Government policy change to routine, opt-out testing in genitourinary medicine clinics. However, testing rates remain low and compare unfavourably to near universal testing levels elsewhere. The limited change, and decline across age groups, in recent HIV testing rates suggest few men test repeatedly or regularly. Additional, innovative efforts are required to increase the uptake of regular HIV testing among gay men.

KEY MESSAGES

- HIV testing increased among gay men in Scotland between 1996 and 2005, from 49.7% to 57.8%, with recent testing increasing from 28.4% to 33.2%.
- The increase corresponds with the policy change to routine, opt-out testing in GUM clinics but rates still compare unfavourably to near universal levels elsewhere.
- Recent testing rates did not increase among ever testers and decreased with age, suggesting few men test repeatedly or regularly.
- Further efforts to increase regular HIV testing among gay men are essential, and must also challenge HIV stigma and 'normalise' testing at the community-level.

INTRODUCTION

Increases in HIV incidence among men who have sex with men (MSM) have been reported,¹ and MSM remain the group most at risk of acquiring HIV in the United Kingdom (UK).² In Scotland, MSM account for 36% of HIV diagnoses, and current HIV prevalence in this group is 4.3%.³ In 2005, community-based surveys of gay men in Scotland found 42% of HIV positive men were undiagnosed,⁴ compared with 30% of MSM genitourinary medicine (GUM) clinic attenders, but recent estimates suggest less than 10% remain undiagnosed following a clinic attendance.³ This has been credited to the introduction of routine, opt-out testing in GUM clinics, where by all patients should be offered an HIV test regardless of symptoms or risk factors. This was implemented as part of the Scottish Sexual Health Strategy,⁵ which was distinct from the Strategy for England and Wales⁶. This policy is now recommended throughout the UK.⁷

The promotion of regular and frequent HIV antibody testing as a means of identifying, and therefore limiting, the potential onward transmission of infection is a core component of prevention efforts in the United States (US) and the Centers for Disease Control and Prevention (CDC) recently recommended routine HIV testing should be performed in all health care settings.⁸ In the UK, HIV testing has traditionally played a lesser part in prevention efforts but its promotion has been central to recent campaigns (see for example <http://www.hivcomebacktour.co.uk/>). The UK Chief Medical Officers recently advocated extension of this to all health care settings,⁹ and the 2008 UK National Guidelines for HIV Testing are designed to facilitate this.⁷ However, HIV testing rates among gay men in Scotland have traditionally been lower than among men in similar surveys elsewhere in the UK and

Europe,^{10 11} and compare unfavourably with the near universal rates reported in the US and Australia.^{12 13} In this paper, we examine trends in the HIV testing behaviour of gay men in Scotland over a ten-year period and discuss whether we are doing enough to promote testing in the group most at risk of acquiring HIV in Scotland.

METHODS

We conducted seven cross-sectional surveys between 1996 and 2005 in commercial gay venues in Glasgow and Edinburgh (Table 1).^{4 14-16} Time and location sampling was used to recruit representative samples. All seven surveys utilised anonymous, self-complete questionnaires and respondents were asked whether they had ever had an HIV test and for the date of their most recent test. Those tested in the year of the survey, or the year immediately prior to it, were categorised as recent HIV testers (e.g. for the 2005 survey, men testing in 2005 or 2004 were categorised as recent testers).

[TABLE 1]

Ethical approval was granted for the 1996, 1999, 2002^a, and 2005 surveys by the University of Glasgow Ethics Committees for Non-clinical Research Involving Human Subjects, and for the 2000 survey by the Psychology Ethics Committee at Glasgow Caledonian University. The 2002^b and 2003 surveys were exact replications of the 2000 study design and methodology. Consequently no further ethical approval for these surveys was sought.

The data were analysed by chi-squared tests and logistic regression. The analysis was conducted using SPSS 16.0 for Mac.

RESULTS

Questionnaires were obtained from 9613 men with an average response rate of 70% (Table 1). Men responding to the question on HIV testing, and who also provided information on age, date of last HIV test, and numbers of unprotected anal intercourse (UAI) partners in the previous year, are included in the analysis (N=8305). The age breakdown of each survey is shown in Table 2.

[TABLE 2]

Overall, 4370 respondents (52.6%) reported that they had had an HIV test. The proportion reporting they had ever had an HIV test increased between 1996 and 2005, from 49.7% to 57.8% ($P < 0.001$). Table 3 shows the proportions who reported having a recent test (in the calendar year of, or immediately prior to, the survey), having ever, but not recently, tested (within >1-5 and 6+ years ago), and having never tested by age group and year of survey. Although the proportion who reported recent testing increased from 28.4% in 1996 to 33.2% in 2005 (reaching their highest level of 40.4% in 2003), the proportion tested >1-5 years ago varied slightly over the seven surveys but remained at 15% in 1996 and 2005, and the proportion tested 6+ years ago increased from 6.2% in 1996 to 9.6% in 2005. The odds ratios for recent testing (compared with those who have tested but not recently, and those who have never tested), adjusted for age, are shown in Table 3. Compared with 1996, the odds ratios associated with recent testing for the 1999 and 2002^a surveys were somewhat lower ($P=0.003$ and $P=0.038$ respectively), while the odds ratio for the 2000 survey did not differ significantly from that associated with 1996. However, the odds ratios for the

2002^b, 2003 and 2005 surveys were all significantly higher (all $P < 0.01$), indicating that recent testing rates in these years were higher than those recorded in 1996.

[TABLE 3]

However, recency of testing among those who reported having had an HIV test at some point in their lives varied but there was no real increase over time, falling from 57.1% (595/1042) in 1996 to 48.2% (369/766) in 2002^a and rising to 57.5% (488/850) in 2005. The adjusted odds ratios for recent testing compared with those who had ever, but not recently, tested (excluding those who have never tested) were also calculated. Compared with 1996, the odds ratios were significantly different (lower) in the 1999 (AOR= 0.79, 95% CI =0.62-0.89) and 2002^a (AOR = 0.69, 95% CI =0.57-0.84) surveys. However, the odds ratios were not significantly different in the 2002^b (AOR=1.16, 95% CI = 0.81-1.65), 2003 (AOR=1.17, 95% CI = 0.83-1.65) or 2005 (AOR=1.10, 95% CI = 0.91-1.33) surveys. This indicates that among men who had had an HIV test, there was no increase in rates of *recent* testing between 1996 and 2005.

Although the proportion of men who had never had an HIV test decreased with age between the <25, 25-34, and 35-44 years age groups, the proportion tested recently also decreased with age. On average, 31.3% of the <25, 30.3% of the 25-34, 23.2% of the 35-44, and 21.2% of the >44 years age groups had tested recently. Table 3 shows the pattern of change over time for each age group. Among the <25 years age group, recent testing fell from 33.9% in 1996 to 26.9% in 2002^a and rose to 34.2% in 2005 (with a high of 51.1% in the Edinburgh only 2003 survey). Among the 25-34

years age group, recent testing increased from 29.7% in 1996 to 41.4% in 2005. In the 35-44 years age group, recent testing rose from 22.4% in 1996 to 39.0% in 2003 and fell to 26.0% in 2005. In the >44 years age group, the proportions tested recently varied over time, rising from 20.4% in 1996 to 24.4% in 2005.

Aggregate rates of recent HIV testing were consistently higher among men who reported two or more UAI partners in the previous year than among men who reported none or one partner (AOR = 1.93, 95% CI = 1.68-2.23) (Table 4). Overall, 41.4% of men reporting UAI with two or more partners had tested recently. Among men who reported UAI with two or more partners, recent testing was lowest in the 35-44 year age group, with only 34.2% having had a recent test.

[TABLE 4]

DISCUSSION

Rates of self-reported HIV testing increased among gay men in Scotland surveyed in commercial venues between 1996 and 2005 and corresponds with the Scottish Government policy change to routine, opt-out testing in genitourinary medicine clinics, which was introduced in the Scottish Sexual Health Strategy.⁵ However, with increasing testing rates over time, and a corresponding increase in recent testing, one might have expected to find higher proportions of those who have had a test, to have had one recently, but this was not the case. This, combined with the consistent trend that older age is associated with decreased likelihood of recent testing, is consistent with HIV testing being a one-off event. It is not indicative of HIV testing being a 'routine' part of a sexual health check up.

There are some limitations to consider when interpreting our results. This was a bar-based sample so only men who visit the venues surveyed had the opportunity to participate and caution should be taken when generalising to the wider population of gay men. The analyses are also limited to the available variables included in all seven surveys; preventing the exploration of other potential confounding factors.

Furthermore, our measure of recent testing would not necessarily pick up all those who had had their first test between studies, given that the gaps between the bigger surveys are at least two years. However, our findings still have considerable implications for HIV testing and prevention policies.

UK guidelines recommend routine, opt-out HIV testing in genitourinary medicine (GUM) settings,^{7 17} and the offer of HIV testing to MSM in UK clinics increased by 35% between 2003 and 2006, with uptake reported to be 85%.² However, at the community-level in Scotland these changes have not yet achieved the near universal testing seen in the US and Australia.^{12 13} It is possible that the increase in uptake in GUM clinics reflects more frequent testing in these settings rather than an increase in the absolute numbers of MSM being tested. This is appropriate given that in the community-based surveys of HIV prevalence the majority of men with undiagnosed HIV had previously tested negative, while levels of HIV were low among men who have never tested.¹⁸ Specialist sexual health services are ideally placed to target and recall men who test negative, but report high risk sexual behaviour, for repeat testing. However, community, or scene-based, testing initiatives should also be trialled to increase testing uptake among men not accessing mainstream services.^{19 20}

HIV testing levels among gay men in Scotland could also be limited by factors specific to the Scottish context. In-depth research on the HIV testing behaviour of gay men in Scotland has demonstrated that fear of a positive result, along with HIV-related stigma and discrimination within the gay community, discouraged testing.^{14 21}

²² In fact, within this particular cultural context the anticipated burden of coping with a positive diagnosis was understood to be more important in shaping testing decisions than accessing appropriate treatment and care.²³

Our findings suggest that few men become regular or repeat testers. As would be expected, older men are more likely to have tested within their lifespan, but they are less likely than their younger counterparts to report recent testing. This may indicate that men are making informed decisions about the need to test based on their perceived risk or that there are age-related patterns in undiagnosed infection. Recent testing rates were higher among men who reported high risk sexual behaviour but even here, under half of men (and only one third of men aged 35-44 years) who reported UAI with multiple partners in the previous year had had a recent test. The limitations of serosorting (only having UAI with partners of the same status) as a risk reduction strategy for HIV negative and untested men have been widely reported,^{18 24-27} and men who have never, or not recently, tested cannot claim to accurately know their status if, as described here, they also report sexual risk behaviour. Receptive unprotected anal intercourse, particularly with partners of unknown HIV status, is one of the main sexual behaviours identified as a risk factor for HIV seroconversion,^{28 29} and therefore accurate knowledge of HIV status is essential if men are to avoid unknowingly putting others at risk of HIV.

Knowledge of HIV status is particularly important when levels of undiagnosed HIV infection are high. Recent data (measured through oral fluid samples collected and anonymously tested for HIV antibodies) show 48.1% of HIV-positive men in bar-based surveys in Glasgow, and 36.4% in Edinburgh, are undiagnosed.¹⁸ The proportion undiagnosed decreases with age, but with higher overall prevalence in older men, the prevalence of undiagnosed HIV is twice as high in older than younger men in community-based surveys (average of 4.5% in the 26-35 years and 1.9% in the 15-25 years age group).¹⁸ Given that among UK MSM, HIV diagnoses remains highest in the 25-34 and 35-44 year age groups,² there is a clear need to further promote regular HIV testing.

Testing and treatment decisions are complex,³⁰ but early diagnosis is essential to ensure timely access to treatments. In 2006, 20% of MSM diagnosed with HIV had a CD4 count less than 200 cells/mm³ (which at the time was the recommended threshold for treatment to commence), and men diagnosed late are fourteen times more likely to die within a year of diagnosis than those diagnosed earlier.² Nevertheless, efforts to increase regular HIV testing have to promote the positive benefits of knowing your HIV status beyond simply having access to treatments.

Accurate knowledge of status allows men to make effective use of primary and secondary prevention strategies, such as serosorting and post-exposure prophylaxis. It should also allow them to make informed sexual risk decisions and be better able to negotiate sex through effective status disclosure, to avoid unknowingly putting others at risk of HIV. However, we have previously demonstrated that gay men in Scotland exhibit undue confidence that their partners share the same assumed negative status as

themselves,¹⁵ and, as previously noted in this paper, there are continuing issues around HIV stigma and discrimination within the gay community.^{14 21} These remain significant barriers to efforts to further increase HIV testing among gay men in Scotland.

Increasing testing may not be enough in itself to reduce HIV infections among UK gay men, when sexual risk is reportedly higher among men living with diagnosed HIV,¹⁸ but the promotion of regular HIV testing should be an essential component of HIV prevention, and additional, innovative efforts are required to increase its uptake. These efforts must do as much to challenge HIV stigma and ‘normalise’ regular testing at the community-level, as they do to increase uptake at the individual-level.

Word Count: 2348

ACKNOWLEDGEMENTS

We thank the survey staff and fieldworkers in each city, the venue managers, their staff, and the men who agreed to participate in the survey.

COMPETING INTERESTS

None.

FUNDING

The UK Medical Research Council funded the 1996, 1999, 2002a and 2005 surveys as part of the Sexual and Reproductive Health Programme (WBS U.1300.00.005) at the Social and Public Health Sciences Unit. Healthy Gay Scotland funded the 2000, 2002b and 2003 surveys.

AUTHOR CONTRIBUTIONS

PF had the initial idea for the paper and all authors contributed to the interpretation of the data and development of the paper. CK conducted the analyses and LW wrote the first draft. All authors contributed to subsequent drafts and approved the final version of the manuscript. The 1996 and 1999 surveys were designed and conducted by GH and PF, the 2002a and 2005 surveys by GH and LW, and the 2000, 2002b and 2003 surveys by PF.

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in STI and any other BMJPG

products to exploit all subsidiary rights, as set out in our licence

<http://sextrans.bmj.com/ifora/licence.pdf>."

REFERENCES

1. **Elford J.** Changing patterns of sexual behaviour in the era of highly active antiretroviral therapy. *Curr Opin Infect Dis* 2006;**19**:26-32.
2. **The UK Collaborative Group for HIV and STI Surveillance.** Testing Times: HIV and other Sexually Transmitted Infections in the United Kingdom: 2007. London: Health Protection Agency, Centre for Infections, 2007.
3. **Health Protection Scotland.** HIV Infection and AIDS: Quarterly report to 30 September 2008 / HIV prevalence among noninjecting drug using heterosexuals, injecting drug users (IDUs), and men who have sex with men (MSM) who undergo attributable HIV testing in Scotland: 2007 update / Unlinked anonymous HIV testing of genito-urinary medicine clinic attendees in Glasgow: data to 2007. *HPS Wkly Rep* 2008;**42**:403-416.
4. **Williamson LM, Hart GJ.** HIV prevalence and undiagnosed infection among a community sample of gay men in Scotland. *J Acquir Immune Defic Syndr* 2007;**45**:224-230.
5. **Scottish Executive.** Respect and Responsibility: Strategy and Action Plan for Improving Sexual Health. Edinburgh: Scottish Executive, 2005.
6. **Department of Health.** The National Strategy for Sexual Health: Better prevention, better services, better sexual health. London: Department of Health, 2001.
7. **British HIV Association, British Association for Sexual Health and HIV, British Infection Society.** UK National Guidelines for HIV Testing 2008. London: British HIV Association, 2008.

8. **Centers for Disease Control and Prevention.** Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Morb Mort Wkly Rep* 2006;**55**:1-16.
9. **Chief Medical Officer.** Improving the detection and diagnosis of HIV in non-HIV specialities including primary care. Available at <http://www.info.doh.gov.uk/doh/embroadcast.nsf/vwDiscussionAll/EE0FA479BAA64A1B80257355003DFB47> [Accessed 12 August 2008].
10. **Bochow MJ,** Chiarotti F, Davies P, Dubois-Arber F, Dur W, Fouchard J, et al. Sexual behaviour of gay and bisexual men in eight European countries. *AIDS Care* 1994;**6**:533-549.
11. **Williamson LM,** Dodds JP, Mercey DE, Johnson AM, Hart GJ. Increases in HIV-related sexual risk behaviour among community samples of gay men in London and Glasgow: how do they compare? *J Acquir Immune Defic Syndr* 2006;**42**:238-241.
12. **Centers for Disease Control and Prevention.** Human immunodeficiency virus (HIV) risk, prevention, and testing behaviours - United States, National HIV Behavioural Surveillance System: men who have sex with men, November 2003-April 2005. *MMWR Morb Mort Wkly Rep* 2006;**55**:1-16.
13. **Prestage G,** Jin F, Zablotska IB, Imrie J, Grulich AE, Pitts M. Trends in HIV testing among homosexual and bisexual men in eastern Australian states. *Sexual Health* 2008;**5**:119-123.
14. **Flowers P,** Knussen C, Church S. Psychosocial factors associated with HIV testing amongst Scottish gay men. *Psychol Health* 2003;**18**:739-752.

15. **Hart GJ**, Williamson LM. Increase in HIV sexual risk behaviour in gay men in Scotland, 1996-2002: Prevention failure? *Sex Transm Infect* 2005;**81**:367-372.
16. **O'Donnell S**. HIV testing in Edinburgh and the Lothians. Final Report. Edinburgh: Testing Barriers Project, Gay Men's Health, 2004.
17. **Rogstad KE**, Palfreeman A, Rooney G, Hart G, Lowbury R, Mortimer P, et al. United Kingdom National Guidelines on HIV Testing 2006: Clinical Effectiveness Group, British Association for Sexual Health and HIV. Available at <http://www.bashh.org/documents/63/63.pdf> [Accessed 12 August 2008].
18. **Williamson LM**, Dodds J, Mercey DE, Hart GJ, Johnson AM. Sexual risk behaviour and knowledge of HIV status among community samples of gay men in the UK. *AIDS* 2008;**22**:1063-1070.
19. **Huebner DM**, Binson D, Woods WJ, Dilworth SE, Neilands TB, Grinstead O. Bathhouse-based voluntary counseling and testing is feasible and shows preliminary evidence of effectiveness. *J Acquir Immune Defic Syndr* 2006;**43**:239-246.
20. **Prost A**, Chopin M, McOwan A, Elam G, Dodds J, Macdonald N, et al. "There is such a thing as asking for trouble": taking rapid HIV testing to gay venues is fraught with challenges. *Sex Transm Infect* 2007;**83**:185-188.
21. **Flowers P**, Duncan B, Frankis JS. Community, responsibility and culpability: HIV risk management amongst Scottish gay men. *J Community Appl Soc Psychol* 2000;**10**:258-300.
22. **Flowers P**, Duncan B, Knussen C. Re-appraising HIV testing: an exploration of the psychosocial costs and benefits associated with learning one's HIV

- status in a purposive sample of Scottish gay men. *Br J Health Psychol* 2003;**8**:179-194.
23. **Flowers P**, Church S. To test or not? HIV antibody testing amongst gay men. *Cult Health Sex* 2002;**4**:43-65.
24. **Butler DM**, Smith DM. Serosorting can potentially increase HIV transmissions. *AIDS* 2007;**21**:1218-1220.
25. **Eaton LA**, Kalichman SC, Cain DN, Cherry C, Stearns HL, Amaral CM, et al. Serosorting sexual partners and risk for HIV among men who have sex with men. *Am J Prev Med* 2007;**33**:479-485.
26. **Elford J**, Bolding G, Sherr L, Hart G. No evidence of an increase in serosorting with casual partners among HIV-negative gay men in London, 1998-2005. *AIDS* 2007;**21**:243-245.
27. **Mao L**, Crawford JM, Hospers HJ, Prestage GP, Grulich AE, Kaldor JM, et al. 'Serosorting' in casual anal sex of HIV-negative gay men is noteworthy and is increasing in Sydney, Australia. *AIDS* 2006;**20**:1204-1206.
28. **Koblin BA**, Husnik MJ, Colfax G, Huang Y, Madison M, Mayer K, et al. Risk factors for HIV infection among men who have sex with men. *AIDS* 2006;**20**:731-739.
29. **Macdonald N**, Elam G, Hickson F, Imrie J, McGarrigle CA, Fenton KA, et al. Factors associated with HIV seroconversion in gay men in England at the start of the 21st century. *Sex Transm Infect* 2008;**84**:8-13.
30. **Flowers P**, Knussen C, Duncan B. Re-appraising HIV testing among Scottish gay men: the impact of new HIV treatments. *J Health Psychol* 2001;**6**:665-678.

Table 1: Survey year, source, location and numbers approached, participating and included in the analyses

Year of survey	Source	Location	Number approached	Number participating (N=9613)	Response rate	Number included in analyses (N=8305)*
1996	MRC	Edinburgh/Glasgow	2881	2276	79%	2097
1999	MRC	Edinburgh/Glasgow	3202	2498	78%	2125
2000	HGS	Edinburgh/Glasgow	1029	803	78%	652
2002 ^a	MRC	Edinburgh/Glasgow	2796	1734	62%	1468
2002 ^b	GMH	Edinburgh	456	283	62%	262
2003	GMH	Edinburgh	429	275	64%	230
2005	MRC	Edinburgh/Glasgow	2642	1744	66%	1471

MRC = Medical Research Council, HGS = Healthy Gay Scotland, GMH = Gay Men’s Health.

* Those who responded to all of the following questions: age, ever had HIV test, date of last HIV test, and numbers of unprotected anal intercourse partners in previous year.

Table 2: Age of respondents by year of survey (N=8305)

	Year of survey													
	1996		1999		2000		2002 ^a		2002 ^b		2003		2005	
	N=2097		N=2125		N=652		N=1468		N=262		N=230		N=1471	
	N	%	n	%	n	%	n	%	n	%	n	%	n	%
<i>Age group</i>														
<25 years	475	22.7	479	22.5	173	26.5	438	29.8	82	31.3	43	18.7	400	27.2
25-34 years	1010	48.2	942	44.3	263	40.3	567	38.6	96	36.6	77	33.5	488	33.2
35-44 years	455	21.7	504	23.7	161	24.7	366	24.9	63	24.0	77	33.5	407	27.7
>44 years	157	7.5	200	9.4	55	8.4	97	6.6	21	8.0	33	14.3	176	12.0

Table 3: HIV testing by year of survey and age of respondents (N=8305)

		Year of survey													
		1996		1999		2000		2002 ^a		2002 ^b		2003		2005	
		N=2097		N=2125		N=652		N=1468		N=262		N=230		N=1471	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%
<i>Age group</i>															
<25 years	Recent HIV test*	161	33.9	127	26.5	57	32.9	118	26.9	33	40.2	22	51.1	137	34.2
	Last tested >1-5 years ago	40	8.4	49	10.2	12	6.9	50	11.4	4	4.9	3	7.0	31	7.8
	Last tested 6+ years ago	4	0.8	3	0.6	0	0	0	0	0	0	0	0	6	1.5
	Never had HIV test	270	56.8	300	62.6	104	60.1	270	61.6	45	54.9	18	41.9	226	56.5
25-34 years	Recent HIV test*	300	29.7	245	26.0	71	27.0	155	27.3	37	38.5	32	41.6	202	41.4
	Last tested >1-5 years ago	179	17.7	194	20.6	66	25.1	131	23.1	20	20.8	17	22.1	78	16.0
	Last tested 6+ years ago	68	6.7	62	6.6	17	6.5	43	7.6	7	7.3	7	9.1	27	5.5

	Never had HIV test	463	45.8	441	46.8	109	41.4	238	42.0	32	33.3	21	27.3	181	37.1
35-44 years	Recent HIV test*	102	22.4	102	20.2	38	23.6	76	20.8	18	28.6	30	39.0	106	26.0
	Last tested >1-5 years ago	74	16.3	109	21.6	29	18.0	84	23.0	15	23.8	16	20.8	86	21.1
	Last tested 6+ years ago	47	10.3	60	11.9	24	14.9	58	15.8	11	17.5	16	20.8	74	18.2
	Never had HIV test	232	51.0	233	46.2	70	43.5	148	40.4	19	30.2	15	19.5	141	34.6
>44 years	Recent HIV test*	32	20.4	37	18.5	9	16.4	20	20.6	7	33.3	9	27.3	43	24.4
	Last tested >1-5 years ago	25	15.9	30	15.0	5	9.1	15	15.5	2	9.5	8	24.2	26	14.8
	Last tested 6+ years ago	10	6.4	31	15.5	13	23.6	16	16.5	4	19.0	4	12.1	34	19.3
	Never had HIV test	90	57.3	102	51.0	28	50.9	46	47.4	8	38.1	12	36.4	73	41.5
Total	Recent HIV test*	595	28.4	511	24.0	175	26.8	369	25.1	95	36.3	93	40.4	488	33.2
	Last tested >1-5 years ago	318	15.2	382	18.0	112	17.2	280	19.1	41	15.6	44	19.1	221	15.0
	Last tested 6+ years ago	129	6.2	156	7.3	54	8.3	117	8.0	22	8.4	27	11.7	141	9.6

Never had HIV test	1055	50.3	1076	50.6	311	47.7	702	47.8	104	39.7	66	28.7	621	42.2
AOR** of having recently had an	1		0.81		0.94		0.85		1.45		1.87		1.31	
HIV test (95% CI)			(0.71-0.93)		(0.77-		(0.73-		(1.11-		(1.41-		(1.13-	
					1.15)		0.99)		1.91)		2.48)		1.52)	

* Recent HIV testing includes all men tested in the calendar year of the survey and the calendar year prior to the survey (e.g. for the 2005 survey, all men tested in 2004 and 2005 are included in the recent testers category).

** Odds ratio of having had a recent HIV test by year of survey, adjusted for age.

Table 4: Recent HIV testing by reported number of unprotected anal intercourse partners in the previous year and age of respondents

		Age group									
		<25 years		25-34 years		35-44 years		45+ years		Total	
		N=2090		N=3443		N=2033		N=739		N=8305	
		n	%	n	%	n	%	n	%	n	%
<i>No of UAI partners in previous year</i>											
0/1 partner	Had recent HIV test*	539	29.8	870	28.4	407	22.1	131	19.5	1947	26.3
	Last tested >1-5 years ago	162	9.0	616	20.1	377	20.4	98	14.6	1253	17.0
	Last tested 6+ years ago	13	0.7	212	6.9	263	14.3	106	15.8	594	8.0
	Never had HIV test	1093	60.5	1369	44.6	796	43.2	337	50.1	3595	48.7
2+ partners	Had recent HIV test*	116	41.0	172	45.7	65	34.2	26	38.8	379	41.4
	Last tested >1-5 years ago	27	9.5	69	18.3	36	18.9	13	19.4	145	15.8
	Last tested 6+ years ago	0	0	19	5.1	27	14.2	6	9.0	52	5.7

Never had HIV test	140	49.5	116	30.9	62	32.6	22	32.8	340	37.1
--------------------	-----	------	-----	------	----	------	----	------	-----	------

UAI = unprotected anal intercourse.

* Recent HIV testing includes all men tested in the calendar year of the survey and the calendar year prior to the survey (e.g. for the 2005 survey, all men tested in 2004 and 2005 are included in the recent testers category).