



HAL
open science

Dynamical properties of the negative beta transformation

Lingmin Liao, Wolfgang Steiner

► **To cite this version:**

Lingmin Liao, Wolfgang Steiner. Dynamical properties of the negative beta transformation. 2011. hal-00555127v1

HAL Id: hal-00555127

<https://hal.science/hal-00555127v1>

Preprint submitted on 12 Jan 2011 (v1), last revised 28 Jun 2011 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DYNAMICAL PROPERTIES OF THE NEGATIVE BETA TRANSFORMATION

LINGMIN LIAO AND WOLFGANG STEINER

ABSTRACT. We analyse basic dynamical properties of the negative beta transformation, which has been studied recently by Ito and Sadahiro. Contrary to the classical beta transformation, the absolutely continuous invariant measure of the negative beta transformation may be zero on certain intervals. By investigating this property in detail, we prove that for any $\beta > 1$ the $(-\beta)$ -transformation is exact, confirming a conjecture of Góra, and intrinsic, which completes a study of Faller. We also show that the limit behaviour of the $(-\beta)$ -expansion of 1 when β tends to 1 is related to the Thue-Morse sequence.

A consequence of the exactness is that every $(-\beta)$ -number, which is a $\beta > 1$ such that the $(-\beta)$ -expansion of 1 is eventually periodic, is a Perron number. This generalises a well-known property of Parry numbers (which were called β -numbers by Parry). Nevertheless, the properties β -number and $(-\beta)$ -number are not equivalent.

1. INTRODUCTION

For a real number $\beta > 1$, the β -*transformation* is defined by

$$T_\beta : [0, 1) \rightarrow [0, 1), \quad x \mapsto \beta x - \lfloor \beta x \rfloor.$$

Rényi [Rén57] was the first to use it for representing real numbers in base β , generalising expansions in integer bases. The (greedy) β -*expansion* of $x \in [0, 1)$ is

$$x = \frac{\lfloor \beta x \rfloor}{\beta} + \frac{\lfloor \beta T_\beta(x) \rfloor}{\beta^2} + \frac{\lfloor \beta T_\beta^2(x) \rfloor}{\beta^3} + \dots.$$

In the present paper, we study the $(-\beta)$ -*transformation* (negative beta transformation)

$$T_{-\beta} : (0, 1] \rightarrow (0, 1], \quad x \mapsto -\beta x + \lfloor \beta x \rfloor + 1.$$

Note that $T_{-\beta}(x) = -\beta x - \lfloor -\beta x \rfloor$ except for finitely many points, hence $T_{-\beta}$ is a natural modification of the β -transformation, avoiding the discontinuity of $x \mapsto -\beta x - \lfloor -\beta x \rfloor$ at 0. For $x \in (0, 1]$, set $d_{-\beta}(x) = d_{-\beta,1}(x) d_{-\beta,2}(x) \dots$ with

$$d_{-\beta,1}(x) = \lfloor \beta x \rfloor + 1, \quad d_{-\beta,n}(x) = d_{-\beta,1}(T_{-\beta}^{n-1}(x)) \quad \text{for } n \geq 1,$$

then the $(-\beta)$ -*expansion* of x is

$$x = \sum_{k=1}^{\infty} \frac{-d_{-\beta,k}(x)}{(-\beta)^k} = \frac{d_{-\beta,1}(x)}{\beta} - \frac{d_{-\beta,2}(x)}{\beta^2} + \frac{d_{-\beta,3}(x)}{\beta^3} - \frac{d_{-\beta,4}(x)}{\beta^4} + \dots.$$

Examples of a β -transformation and a $(-\beta)$ -transformation are depicted in Figure 1.

Date: January 12, 2011.




FIGURE 1. β -transformation (left) and $(-\beta)$ -transformation (right), $\beta = \frac{1+\sqrt{5}}{2}$.

The map $T_{-\beta}$ has been studied recently by Góra [Gór07] and Faller [Fal08]. Ito and Sadahiro [IS09] considered a $(-\beta)$ -transformation on the interval $[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1})$, defined by

$$x \mapsto -\beta x - \lfloor -\beta x + \frac{\beta}{\beta+1} \rfloor.$$

We remark that their transformation is conjugate to our $T_{-\beta}$ through the conjugacy function $\phi(x) = \frac{1}{\beta+1} - x$. So, all the results in [IS09] can be translated into our case directly, in particular their $(-\beta)$ -expansion of $\phi(x) \in [\frac{-\beta}{\beta+1}, \frac{1}{\beta+1})$ is

$$\phi(x) = \sum_{k=1}^{\infty} \frac{d_{-\beta,k}(x) - 1}{(-\beta)^k}.$$

Both β -transformation and $(-\beta)$ -transformation are examples of piecewise monotonic transformations. The dynamical properties of general piecewise monotonic transformations have been investigated in the literature [Kel78, Wag79, Hof79, Hof81a, Hof81b]. However, for the specific case of the $(-\beta)$ -transformation, the detailed dynamical properties are unknown. As pointed out in [IS09], for any $\beta > 1$, $T_{-\beta}$ admits a unique absolutely continuous invariant measure (hence ergodic), with the density

$$h_{-\beta}(x) = \sum_{n \geq 1, T_{-\beta}^n(1) \geq x} \frac{1}{(-\beta)^n}.$$

Contrary to the classical β -transformation, Ito and Sadahiro [IS09] gave an example such that the density $h_{-\beta}$ is zero on some intervals.

In the present paper, we fully study this phenomenon. An interval on which the density equals zero will be called a *gap*. From a result of Keller [Kel78], one can deduce that, for fixed $\beta > 1$, the number of gaps is finite. In Theorem 2.1, we describe the set of the gaps and show that, as β decreases, the number of gaps forms an increasing sequence

$$0 < 1 < 2 < 5 < 10 < \dots < \lfloor 2^{n+1}/3 \rfloor < \dots.$$

Figures 2 and 3 show examples with 2 gaps and 5 gaps respectively. The endpoints of the gaps are determined by the orbit of 1, which is described by the sequence $d_{-\beta}(1)$.

We also show (Theorem 2.5) that

$$\lim_{\beta \rightarrow 1} d_{-\beta}(1) = 211222112112112221122 \dots,$$

which is the fixed point of the following morphism:

$$\varphi : 1 \mapsto 2, \quad 2 \mapsto 211.$$

Here, φ is a morphism on finite words on the alphabet $\{1, 2\}$, which is naturally extended to the set of infinite words $\{1, 2\}^{\mathbb{N}}$ as in [Lot97], and $\lim_{\beta \rightarrow 1} d_{-\beta}(1) = u$ means that longer and longer prefixes of $d_{-\beta}(1)$ agree with prefixes of u when $\beta \rightarrow 1$. We remark that this sequence has been studied e.g. in [AAB⁺95, Dub06, Dub07], and is known to be the smallest aperiodic sequence in $\{1, 2\}^{\mathbb{N}}$ with the property that all its proper suffixes are smaller than itself with respect to the alternate order. Moreover, by adding a 1 at the beginning, we obtain the sequence of run lengths in the Thue-Morse sequence. It is shown in [IS09] that every sequence $d_{-\beta}(1)$, $\beta > 1$, has the property that all its proper suffixes are smaller than itself with respect to the alternate order.

A dynamical system (X, T) is said to be *locally eventually onto* or topologically exact if, for any non-empty open subset $U \subset X$, there exists a positive integer n such that $f^n(U) = X$. The main theorem (Theorem 2.2) of this paper asserts that $T_{-\beta}$ is locally eventually onto on $(0, 1] \setminus G(\beta)$, where $G(\beta)$ is the (finite) union of the gaps.

The notion of topological exactness is derived from the exactness in ergodic theory introduced by Rohlin [Roh61]. A measure-preserving transformation T on a probability space (X, \mathcal{B}, μ) is *exact* if and only if, for any positive measure subset A with $T^n(A) \in \mathcal{B}$ ($n \geq 0$), one has $\lim_{n \rightarrow \infty} \mu(T^n(A)) = 1$ (see [PY98], p. 125). As a corollary of our main theorem, we confirm Góra's conjecture [Gór07] that all $(-\beta)$ -transformations are exact.

An invariant measure of a dynamical system is called a *maximal entropy measure* if it maximises entropy. If there is a unique maximal entropy measure, the dynamical system is called *intrinsic*. As another application of our main theorem, we give a complete answer about the uniqueness of maximal entropy measures discussed in the thesis of Faller [Fal08]. We prove that all $(-\beta)$ -transformations are intrinsic (Corollary 2.4).

Finally, we use the main result to show that every $(-\beta)$ -number, that is a number $\beta > 1$ with the property that the $T_{-\beta}$ -orbit of 1 is eventually periodic, is a Perron number. Numbers $\beta > 1$ with the corresponding property for T_{β} are called β -numbers [Par60] or Parry numbers, and are known to be Perron numbers as well. In Section 6, we give examples showing that the set of $(-\beta)$ -numbers is different from the set of β -numbers.

The paper is organised as follows. In Section 2, we state the main results. In Section 3, we establish properties of the sequences $d_{-\beta}(1)$, which allow describing the structure of the gaps in Section 4. In Section 5, we show that $T_{-\beta}$ is locally eventually onto on $F(\beta)$. The proofs of the main results are completed in Section 6.

2. MAIN RESULTS

For each $n \geq 1$, let γ_n be the (unique) positive real number defined by

$$\gamma_n^{g_n+1} = \gamma_n + 1, \quad \text{with } g_n = \lfloor 2^{n+1}/3 \rfloor,$$

and set $\gamma_0 = \infty$. Then

$$2 > \gamma_1 > \gamma_2 > \gamma_3 > \cdots > 1.$$

Note that γ_1 is the golden ratio and that γ_2 is the smallest Pisot number.

For each $n \geq 0$ and $1 < \beta < \gamma_n$, set

$$\mathcal{G}_n(\beta) = \left\{ G_{m,k}(\beta) \mid 0 \leq m < n, 0 \leq k < \frac{2^{m+1} + (-1)^m}{3} \right\},$$

with open intervals

$$G_{m,k}(\beta) = \begin{cases} (T_{-\beta}^{2^{m+1}+k}(1), T_{-\beta}^{(2^{m+2}-(-1)^m)/3+k}(1)) & \text{if } k \text{ is even,} \\ (T_{-\beta}^{(2^{m+2}-(-1)^m)/3+k}(1), T_{-\beta}^{2^{m+1}+k}(1)) & \text{if } k \text{ is odd,} \end{cases}$$

in particular $\mathcal{G}_0(\beta)$ is the empty set.

Theorem 2.1. *For any $\gamma_{n+1} \leq \beta < \gamma_n$, $n \geq 0$, the set of gaps of the transformation $T_{-\beta}$ is $\mathcal{G}_n(\beta)$, which consists of $g_n = \lfloor 2^{n+1}/3 \rfloor$ disjoint non-empty intervals.*

We define therefore

$$G(\beta) = \bigcup_{I \in \mathcal{G}_n(\beta)} I \quad \text{if } \gamma_{n+1} \leq \beta < \gamma_n, n \geq 0.$$

Theorem 2.2. *For any $\beta > 1$, $T_{-\beta}$ is locally eventually onto on $(0, 1] \setminus G(\beta)$,*

$$T_{-\beta}^{-1}(G(\beta)) \subset G(\beta) \quad \text{and} \quad \lim_{k \rightarrow \infty} \lambda(T_{-\beta}^{-k}(G(\beta))) = 0.$$

Góra [Gór07] proved that, for $\beta > \gamma_2$, the transformation $T_{-\beta}$ is exact, and he conjectured that this would hold for all $\beta > 1$. We confirm his conjecture.

Corollary 2.3. *For any $\beta > 1$, the transformation $T_{-\beta}$ is exact with respect to its unique absolutely continuous invariant measure.*

Proof. The unicity of the absolutely continuous invariant measure follows from [LY78], as already noted in [IS09]. Now, the exactness is a direct consequence of Theorem 2.2. \square

Faller [Fal08] proved that, for $\beta > \sqrt[3]{2}$, $T_{-\beta}$ admits a unique maximal entropy measure and left the case $\beta \leq \sqrt[3]{2}$. We give a complete answer.

Corollary 2.4. *For any $\beta > 1$, the transformation $T_{-\beta}$ has a unique maximal entropy measure, hence is intrinsic.*

Proof. By Theorem 2.2, for any $\beta > 1$, the transformation $T_{-\beta}$ is locally eventually onto on $(0, 1] \setminus G(\beta)$. A result of Walters [Wal78, Theorem 16, p. 140] completes the proof. \square

The key to understanding the $(-\beta)$ -transformation is to know the $(-\beta)$ -expansion of 1. Here, the main tool will be the morphism $\varphi : 1 \mapsto 2, 2 \mapsto 211$. The expansion of 1 under $T_{-\gamma_n}$ can be described by this morphism, and the expansion of 1 under $T_{-\beta}$ tends to the fixed point of the morphism when β tends to 1 (from above).

Theorem 2.5. *For every $n \geq 1$, we have*

$$d_{-\gamma_n}(1) = \varphi^{n-1}(2\bar{1}),$$

where $\bar{1} = 111\cdots$. If $1 < \beta \leq \gamma_n$, then $d_{-\beta}(1)$ starts with $\varphi^n(2)$, hence

$$\lim_{\beta \rightarrow 1} d_{-\beta}(1) = \lim_{n \rightarrow \infty} \varphi^n(2\bar{1}) = 211222112112221122\cdots.$$

Masáková and Pelantová [MP] showed that the $(-\beta)$ -numbers are algebraic integers with all conjugates having modulus less than 2, thus all $(-\beta)$ -numbers with $\beta \geq 2$ are Perron numbers. Again, we give a complete answer.

Theorem 2.6. *Every $(-\beta)$ -number is a Perron number.*

At the end of this section, we recall some identities. Note that

$$|\varphi^n(1)| = \frac{2^{n+1} + (-1)^n}{3} = g_n + \frac{1 + (-1)^n}{2}, \quad (2.1)$$

$$|\varphi^n(2)| = |\varphi^{n+1}(1)| = \frac{2^{n+2} - (-1)^n}{3} = g_{n+1} + \frac{1 - (-1)^n}{2}, \quad (2.2)$$

$$|\varphi^n(11)| = 2 \cdot |\varphi^n(1)| = \frac{2^{n+2} + 2 \cdot (-1)^n}{3} = g_{n+1} + \frac{1 + (-1)^n}{2}, \quad (2.3)$$

where $|w|$ denotes the length of the word $|w|$. In particular, $|\varphi^n(1)|$ and $|\varphi^n(2)|$ are odd, $|\varphi^n(21)| = 2^{n+1}$, g_n is odd if and only if n is odd, and we can write

$$\mathcal{G}_n(\beta) = \{G_{m,k}(\beta) \mid 0 \leq m < n, 0 \leq k < |\varphi^m(1)|\},$$

$$G_{m,k}(\beta) = \begin{cases} (T_{-\beta}^{|\varphi^m(21)|+k}(1), T_{-\beta}^{|\varphi^m(2)|+k}(1)) & \text{if } k \text{ is even,} \\ (T_{-\beta}^{|\varphi^m(2)|+k}(1), T_{-\beta}^{|\varphi^m(21)|+k}(1)) & \text{if } k \text{ is odd.} \end{cases}$$

3. POLYNOMIALS AND EXPANSIONS

To study the trajectories of 1 under $T_{-\beta}$, we define the maps

$$f_{\beta,a} : \mathbb{R} \rightarrow \mathbb{R}, \quad x \mapsto -\beta x + a, \quad a \in \{1, 2\}.$$

The composition of maps $f_{\beta,a_1}, \dots, f_{\beta,a_k}$, is denoted by

$$f_{\beta,a_1 \dots a_k} = f_{\beta,a_k} \circ \cdots \circ f_{\beta,a_1}.$$

Since

$$f_{\beta,a_1 \dots a_k}(1) = (-\beta)^k + \sum_{j=1}^k a_k (-\beta)^{k-j},$$

we define the polynomial

$$P_{a_1 \dots a_k} = (-X)^k + \sum_{j=1}^k a_k (-X)^{k-j} \in \mathbb{Z}[X]$$

for every word $a_1 \cdots a_k \in \{1, 2\}^k$, $k \geq 1$. Then $f_{\beta, a_1 \cdots a_k}(1) = P_{a_1 \cdots a_k}(\beta)$, in particular

$$T_{-\beta}^k(1) = f_{\beta, d_{-\beta, 1}(1) \cdots d_{-\beta, k}(1)}(1) = P_{d_{-\beta, 1}(1) \cdots d_{-\beta, k}(1)}(\beta).$$

For the proof of Theorem 2.5, we use the following polynomial identities.

Lemma 3.1. *For $1 \leq j < k$, we have*

$$P_{a_1 \cdots a_k} = (-X)^{k-j} (P_{a_1 \cdots a_j} - 1) + P_{a_{j+1} \cdots a_k}.$$

Proof. We can deduce the identity directly from the definition. \square

Lemma 3.2. *For every $n \geq 0$, we have*

$$X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(2)} + X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(11)} = X + 1 = X^{\frac{1+(-1)^n}{2}} + X^{\frac{1-(-1)^n}{2}}.$$

Proof. The second equation holds for even and odd n , thus for all $n \in \mathbb{Z}$. Since $X P_2 + P_{11} = X(-X + 2) + X^2 - X + 1$, the first equation holds for $n = 0$. For every $n \geq 0$, we have

$$X^{\frac{1+(-1)^{n+1}}{2}} P_{\varphi^{n+1}(2)} + X^{\frac{1-(-1)^{n+1}}{2}} P_{\varphi^{n+1}(11)} = X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(211)} + X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(22)}.$$

Then, using that $|\varphi^n(11)|$ is even, $|\varphi^n(2)|$ is odd and Lemma 3.1, we obtain

$$\begin{aligned} X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(211)} + X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(22)} &= X^{\frac{1-(-1)^n}{2} + |\varphi^n(11)|} (P_{\varphi^n(2)} - 1) + X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(11)} \\ &\quad - X^{\frac{1+(-1)^n}{2} + |\varphi^n(2)|} (P_{\varphi^n(2)} - 1) + X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(2)} \\ &= X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(2)} + X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(11)}. \end{aligned}$$

Therefore, the lemma follows inductively. \square

The following lemma is a consequence of Lemma 3.2.

Lemma 3.3. *For every $n \geq 0$, the words $\varphi^n(2)$ and $\varphi^n(11)$ agree on the first $g_{n+1} - 1$ letters and differ on the g_{n+1} -st letter.*

Proof. The polynomials $X^{\frac{1+(-1)^n}{2}} P_{\varphi^n(2)}$ and $X^{\frac{1-(-1)^n}{2}} P_{\varphi^n(11)}$ have degree $g_{n+1} + 1$ by (2.2) and (2.3). Therefore, the coefficient of $X^{g_{n+1}+1-j}$ in their sum is given by the difference of the j -th letter in $\varphi^n(2)$ and $\varphi^n(11)$, for $1 \leq j \leq \min(|\varphi^n(2)|, |\varphi^n(11)|) = g_{n+1}$. Since the sum of these polynomials is equal to $X + 1$, by Lemma 3.2, the first $g_{n+1} - 1$ letters in $\varphi^n(2)$ and $\varphi^n(11)$ are equal, and the g_{n+1} -st letters differ. \square

Lemma 3.4. *For every $n \geq 0$, we have*

$$1 - P_{\varphi^n(1)} = X^{\frac{1+(-1)^n}{2}} \prod_{m=0}^{n-1} (X^{|\varphi^m(1)|} - 1), \quad (3.1)$$

$$P_{\varphi^n(21)} - 1 = \left(X^{g_{n+1}+1} - X^{g_n+1} - X^{\frac{1+(-1)^n}{2}} \right) \prod_{m=0}^{n-1} (X^{|\varphi^m(1)|} - 1), \quad (3.2)$$

$$P_{\varphi^n(21)} - P_{\varphi^n(2)} = (X^{g_{n+1}+1} - X - 1) \prod_{m=0}^{n-1} (X^{|\varphi^m(1)|} - 1), \quad (3.3)$$

$$P_{\varphi^n(2)} - P_{\varphi^{n+1}(2)} = (X^{g_{n+1}+1} - X - 1) \prod_{m=0}^n (X^{|\varphi^m(1)|} - 1). \quad (3.4)$$

Proof. Since $1 - P_1 = X$, (3.1) holds for $n = 0$. By Lemmas 3.2 and 3.1, we obtain

$$\begin{aligned} 1 - P_{\varphi^{n+1}(1)} &= 1 - P_{\varphi^n(2)} \\ &= X^{-(1)^n} (P_{\varphi^n(11)} - 1) \\ &= X^{-(1)^n} (-X^{|\varphi^n(1)|} (P_{\varphi^n(1)} - 1) + P_{\varphi^n(1)} - 1) \\ &= X^{-(1)^n} (X^{|\varphi^n(1)|} - 1) (1 - P_{\varphi^n(1)}) \end{aligned}$$

for every $n \geq 0$, hence (3.1) holds inductively for all $n \geq 0$.

Applying first Lemma 3.1, then equation (3.1) for n and $n + 1$, we obtain

$$\begin{aligned} P_{\varphi^n(21)} - 1 &= -X^{|\varphi^n(1)|} (P_{\varphi^n(2)} - 1) + P_{\varphi^n(1)} - 1 \\ &= \left(X^{|\varphi^n(1)| + \frac{1-(-1)^n}{2}} (X^{|\varphi^n(1)|} - 1) - X^{\frac{1+(-1)^n}{2}} \right) \prod_{m=0}^{n-1} (X^{|\varphi^m(1)|} - 1). \end{aligned}$$

By (2.1) and (2.3), we have $|\varphi^n(1)| + \frac{1-(-1)^n}{2} = g_n + 1$ and $2 \cdot |\varphi^n(1)| + \frac{1-(-1)^n}{2} = g_{n+1} + 1$, which gives (3.2).

Combining (3.2) and (3.1) for $n + 1$, $X^{\frac{1-(-1)^n}{2}} (X^{|\varphi^n(1)|} - 1) = X^{g_n+1} - X^{\frac{1-(-1)^n}{2}}$, we obtain (3.3) since $X^{\frac{1+(-1)^n}{2}} + X^{\frac{1-(-1)^n}{2}} = X + 1$.

Finally, (3.4) follows from (3.1) for $n + 1$ and $n + 2$, with

$$X^{\frac{1+(-1)^n}{2}} (X^{|\varphi^{n+1}(1)|} - 1) - X^{\frac{1-(-1)^n}{2}} = X^{g_{n+1}+1} - X^{\frac{1+(-1)^n}{2}} - X^{\frac{1-(-1)^n}{2}}. \quad \square$$

With the help of Lemma 3.4, we will show the following proposition, where $\eta_n > 1$, $n \geq 1$, is defined by

$$\eta_n^{g_n+1} = \eta_n^{g_{n-1}+1} + \eta_n^{\frac{1-(-1)^n}{2}}.$$

It can be easily verified that

$$2 = \eta_1 > \gamma_2^2 > \gamma_1 > \eta_2 > \gamma_2 > \eta_3 > \cdots > 1.$$

Proposition 3.5. *Let $\beta > 1$ and $n \geq 1$. Then $d_{-\beta}(1)$ starts with $\varphi^n(2)$ if and only if $\beta < \gamma_2^2$ in case $n = 1$, $\beta \leq \eta_n$ in case n is even, $\beta < \eta_n$ in case $n \geq 3$ is odd.*

For the proof of the proposition, we use the following lemmas. For the sake of readability, we often omit the dependence on β in $T_{-\beta}$, $d_{-\beta}$ and $f_{\beta,a}$ in the following.

Lemma 3.6. *Let $x \in (0, 1]$ and $1 < \beta < 2$. Then*

- $d_1(x) = 2$ if and only if $f_2(x) \leq 1$,
- $d_1(x) = 1$ if and only if $f_1(x) > 0$,
- $d_1(x)d_2(x) = 11$ if and only if $0 < f_{11}(x) < 1$.

Proof. We have $d_1(x) = a$ if and only if $f_a(x) \in (0, 1]$. The inequalities $f_2(x) > -\beta + 2 > 0$ and $f_1(x) < 1$ prove the first two points. If $f_1(f_1(x)) = f_{11}(x) < 1$, then we must have $f_1(x) > 0$, thus $d_1(x) = 1$ and $d_2(x) = d_1(f_1(x)) = 1$. \square

Lemma 3.7. *Let $\beta > 1$. If $d(1)$ starts with $a_1 \cdots a_k$ and $d(f_{a_1 \cdots a_k}(1))$ starts with $b_1 \cdots b_j$, then $d(1)$ starts with $a_1 \cdots a_k b_1 \cdots b_j$.*

Proof. If $d_1(1) \cdots d_k(1) = a_1 \cdots a_k$, then $T^k(1) = f_{a_1 \cdots a_k}(1)$, and the lemma follows from $d_{k+1}(1) \cdots d_{k+j}(1) = d_1(T^k(1)) \cdots d_j(T^k(1))$. \square

Lemma 3.8. *Let $x \in (0, 1]$, $n \geq 1$, such that $d(x)$ starts with the first $g_n - 1$ letters of $\varphi^n(1)$. Then $d(x)$ starts with $\varphi^n(1)$ if and only if $f_{\varphi^n(1)}(x) \leq 1$ in case n is odd, $0 < f_{\varphi^n(1)}(x) < 1$ in case n is even.*

Proof. Let $x \in (0, 1]$, $n \geq 1$, such that $d(x)$ starts with the first $g_n - 1$ letters of $\varphi^n(1)$.

If n is odd, then $g_n = |\varphi^n(1)|$, and $\varphi^n(1)$ terminates by 2, thus $f_2 T^{|\varphi^n(1)|-1}(1) = f_{\varphi^n(1)}$. By Lemma 3.6, we have $d_{|\varphi^n(1)|}(1) = d_1(T^{|\varphi^n(1)|-1}(1)) = 2$ if and only if $f_2 T^{|\varphi^n(1)|-1}(1) \leq 1$. If n is even, then $g_n = |\varphi^n(1)| - 1$, and $\varphi^n(1)$ terminates by 11, thus $f_{11} T^{|\varphi^n(1)|-2}(1) = f_{\varphi^n(1)}$. By Lemma 3.6, $d_{|\varphi^n(1)|-1}(1) d_{|\varphi^n(1)|}(1) = 11$ is equivalent to $0 < f_{11} T^{|\varphi^n(1)|-2}(1) < 1$. \square

Lemma 3.9. *Let $\beta > 1$ and $n \geq 1$. If $d(1)$ starts with $\varphi^n(2)$, then T^{g_n-1} is continuous on $[T^{|\varphi^n(1)|}(1), 1]$, and $1/\beta$ is in the interior of $T^{g_n-1}([T^{|\varphi^n(1)|}(1), 1])$.*

Proof. Let $\beta > 1$ and $n \geq 1$ such that $d(1)$ starts with $\varphi^n(2) = \varphi^{n-1}(211)$, then $d(T^{|\varphi^n(1)|}(1))$ starts with $\varphi^{n-1}(11)$. By Lemma 3.3, the words $\varphi^{n-1}(2)$ and $\varphi^{n-1}(11)$ share the first $g_n - 1$ letters and differ on the g_n -th letter. This proves that T^{g_n-1} is continuous on $[T^{|\varphi^n(1)|}(1), 1]$, and that $1/\beta$ is in the interior or the right endpoint of $T^{g_n-1}([T^{|\varphi^n(1)|}(1), 1])$. If $T^{g_n-1}(1) = 1/\beta$ with odd n , then $T^{|\varphi^n(1)|}(1) = 1$. If $T^{|\varphi^n(1)|+g_n-1}(1) = T^{g_{n+1}-1}(1) = 1/\beta$ with even n , then $T^{|\varphi^n(2)|}(1) = 1$. Both situations are impossible since $f_{\varphi^n(1)}(1) < 1$ for all $\beta > 1$, $n \geq 0$, by (3.1). Therefore, $1/\beta$ is in the interior of $T^{g_n-1}([T^{|\varphi^n(1)|}(1), 1])$. \square

Lemma 3.10. *For any $1 < \beta \leq \eta_n$, $n \geq 1$, we have $f_{\varphi^{n-1}(1)}(1) \leq f_{\varphi^n(2)}(1) < 1$.*

Proof. By (3.1), we have $f_{\varphi^n(1)}(1) < 1$ for all $\beta > 1$, $n \geq 0$.

For all $1 < \beta \leq \eta_n$, $n \geq 1$, we have $f_{\varphi^{n-1}(21)}(1) \leq 1$ by (3.2). Since $f_{\varphi^{n-1}(1)}$ is order reversing, we get that $f_{\varphi^n(2)}(1) = f_{\varphi^{n-1}(211)}(1) \geq f_{\varphi^{n-1}(1)}(1)$. \square

Proof of Proposition 3.5. Let $\beta > 1$. Lemma 3.6 states that, if $d_1(1) = 2$, then $d(1)$ starts with $\varphi(2)$ if and only if $0 < f_{\varphi(2)}(1) < 1$. By (3.1), we have $1 - f_{\varphi(2)}(1) = \beta(\beta - 1)^2 > 0$,

and $1 - f_{\varphi(2)}(1) < 1$ if and only if $\beta - 1 < \beta^{-1/2}$, i.e., $\beta < \gamma_2^2$. Since $d_1(1) = 2$ for all $\beta < \gamma_2^2 < 2$, the case $n = 1$ of the proposition is proved.

Suppose now that $d(1)$ starts with $\varphi^{n+1}(2) = \varphi^n(211)$ for some $n \geq 1$. Then we must have $f_{\varphi^n(21)}(1) \leq 1$, which is equivalent to $\beta \leq \eta_{n+1}$ by (3.2). Suppose moreover that the proposition is already shown for n . Since $\eta_{n+1} < \min(\eta_n, \gamma_2^2)$, this implies that $d(1)$ starts with $\varphi^n(2)$. Moreover, $\eta_{n+1} < \gamma_n$ and (3.4) yield that $f_{\varphi^n(2)}(1) > f_{\varphi^n(1)}(1)$. By Lemma 3.9, the first $g_n - 1$ letters of $d(x)$ are equal to those of $d(1)$ for all $x \in [f_{\varphi^n(1)}(1), 1]$. Therefore, Lemma 3.8 implies that $d(1)$ starts with $\varphi^n(21)$ if and only if $f_{\varphi^n(21)}(1) \leq 1$ in case n is odd, $0 < f_{\varphi^n(21)}(1) < 1$ in case n is even. For even n , we must therefore have $\beta < \eta_{n+1}$. Note that $f_{\varphi^n(21)}(1) > 0$ is guaranteed by $f_{\varphi^n(21)}(1) > f_{\varphi^n(1)}(1) = T^{|\varphi^n(1)|}(1)$, which follows from $f_{\varphi^n(2)}(1) < 1$ and the order reversing property of $f_{\varphi^n(1)}$. By Lemma 3.10, we have $f_{\varphi^n(1)}(1) \leq f_{\varphi^n(21)}(1) < 1$, thus $f_{\varphi^n(21)}(1) \in [f_{\varphi^n(1)}(1), 1]$, which implies that $d(f_{\varphi^n(21)}(1))$ starts with $\varphi^n(1)$. Putting everything together, we obtain that $d(1)$ starts with $\varphi^n(211) = \varphi^{n+1}(2)$ if and only if $\beta < \eta_{n+1}$ with even n , or $\beta \leq \eta_{n+1}$ with odd n . \square

Now, we can prove Theorem 2.5.

Proof of Theorem 2.5. By Proposition 3.5, $d_{-\beta}(1)$ starts with $\varphi^n(2)$ for any $1 < \beta \leq \gamma_n$, $n \geq 1$. By (3.3), we have $f_{\gamma_n, \varphi^{n-1}(21)}(1) = f_{\gamma_n, \varphi^{n-1}(2)}(1)$, thus $T_{-\gamma_n}^{|\varphi^{n-1}(21)|}(1) = T_{-\gamma_n}^{|\varphi^{n-1}(2)|}(1)$, i.e., $T_{-\gamma_n}^{|\varphi^{n-1}(2)|}(1)$ is a fixed point of $T_{-\gamma_n}^{|\varphi^{n-1}(1)|}$. Hence, by Lemma 3.7, $d_{-\gamma_n}(1) = \varphi^{n-1}(2\bar{1})$.

Since $\lim_{n \rightarrow \infty} \gamma_n = 1$, we obtain that $d_{-\beta}(1)$ starts with $\varphi^n(2)$ for larger and larger n when $\beta \rightarrow 1$, thus $\lim_{\beta \rightarrow 1} d_{-\beta}(1) = \lim_{n \rightarrow \infty} \varphi^n(2\bar{1})$. \square

4. STRUCTURE OF THE GAPS

We will show that the support of the invariant measure is

$$F(\beta) = (0, 1] \setminus G(\beta).$$

For $\beta > 1$, $n \geq 0$, let

$$\mathcal{F}_n(\beta) = \{F_{n,k}(\beta) \mid 0 \leq k \leq g_n\}$$

with

$$F_{n,k}(\beta) = \begin{cases} [T_{-\beta}^{|\varphi^n(1)|+k}(1), T_{-\beta}^k(1)] & \text{if } k \text{ is even, } k < g_n, \\ [T_{-\beta}^k(1), T_{-\beta}^{|\varphi^n(1)|+k}(1)] & \text{if } k \text{ is odd, } k < g_n, \end{cases}$$

$$F_{n,g_n}(\beta) = \begin{cases} (0, T_{-\beta}^{|\varphi^n(1)|-1}(1)] & \text{if } n \text{ is even,} \\ (0, T_{-\beta}^{|\varphi^n(2)|-1}(1)] & \text{if } n \text{ is odd.} \end{cases}$$

(We will consider these sets only in the case when the left number is smaller than the right number.) Figures 2 and 3 show examples of the decomposition of $(0, 1]$ into these sets.

As in Section 3, we often omit the dependence on β in the following.

Lemma 4.1. *Let $n \geq 2$ and $1 < \beta \leq \eta_n$ if n is even; $1 < \beta < \eta_n$ if n is odd; or $n = 1$, $1 < \beta < \gamma_2^2$. Then*




FIGURE 2. The $(-\beta)$ -transformation for $\beta = 5/4$, with set of gaps $G_{1,0} \cup G_{2,0}$ and support of the invariant measure $F_{2,2} \cup F_{2,1} \cup F_{2,0}$. Here, $t_k = T_{-\beta}^k(1)$.

- (i) $F_{n,k} = T^k(F_{n,0})$ for all $0 \leq k < g_n$,
- (ii) $1/\beta$ is in the interior of F_{n,g_n-1} ,
- (iii) $T^{g_n}(F_{n,0}) = F_{n,g_n} \cup F_{n,0}$, $T(F_{n,g_n}) = F_{n+1,0} \setminus \{1\}$, if n is odd,
- (iv) $T^{g_n}(F_{n,0}) = F_{n,g_n} \cup F_{n+1,0}$, $T(F_{n,g_n}) = F_{n,0} \setminus \{1\}$, if n is even.

Proof. Let β and n have the properties of the statement of the lemma. Then Proposition 3.5 and Lemma 3.9 give (i) and (ii). If n is odd, then $F_{n,g_n-1} = [T^{g_{n+1}-1}(1), T^{g_n-1}(1)]$, thus

$$T(F_{n,g_n-1}) = (0, T^{g_{n+1}}(1)] \cup [T^{g_n}(1), 1] = (0, T^{|\varphi^n(2)|-1}(1)] \cup [T^{|\varphi^n(1)|}(1), 1] = F_{n,g_n} \cup F_{n,0}.$$

Since $d(1)$ starts with $\varphi^n(2)$ and $\varphi^n(2)$ ends with 1, we have

$$T(F_{n,g_n}) = T((0, T^{|\varphi^n(2)|-1}(1)]) = [T^{|\varphi^n(2)|}(1), 1] = F_{n+1,0} \setminus \{1\},$$

i.e., (iii) holds. If n is even, then $F_{n,g_n-1} = [T^{g_n-1}(1), T^{g_{n+1}-1}(1)]$ and

$$T(F_{n,g_n-1}) = (0, T^{g_n}(1)] \cup [T^{g_{n+1}}(1), 1] = (0, T^{|\varphi^n(1)|-1}(1)] \cup [T^{|\varphi^n(2)|}(1), 1] = F_{n,g_n} \cup F_{n+1,0}.$$




FIGURE 3. The $(-\beta)$ -transformation for $\beta = 9/8$ with gaps $G_{m,k}$, support of the invariant measure $F_{3,0} \cup \dots \cup F_{3,5}$, and $t_k = T_{-\beta}^k(1)$. Ratios between $G_{1,0}$, $G_{2,0}$ and the rest of the picture are not respected. In reality, $G_{1,0}$ is almost 10 times the size of $G_{2,0}$, and $G_{2,0}$ is almost 5 times the size of $F_{2,0}$.

Now, $\varphi^n(1)$ ends with 1, which gives $T(F_{n,g_n}) = F_{n,0} \setminus \{1\}$, i.e., (iv) holds. □

Proposition 4.2. *Let $1 < \beta < \gamma_n$, $n \geq 0$. Then the elements of \mathcal{F}_n and \mathcal{G}_n are intervals of positive length which form a partition of $(0, 1]$. If $n \geq 1$, then*

- (i) $F_{n-1,k} = F_{n,|\varphi^{n-1}(1)|+k} \cup G_{n-1,k} \cup F_{n,k}$ for all $0 \leq k < |\varphi^{n-1}(1)|$,
- (ii) $G_{m,k} = T^k(G_{m,0})$ for all $0 \leq m < n$, $0 \leq k < |\varphi^m(1)|$,
- (iii) $T^{|\varphi^m(1)|}(G_{m,0}) = G_{m,0} \cup F_{m,0} \setminus F_{m+1,0} \supset G_{m,0}$ for all $0 \leq m < n$.

Proof. We have $\mathcal{F}_0 = \{F_{0,0}\} = \{(0, 1]\}$ and $\mathcal{G}_0 = \emptyset$, thus the elements of \mathcal{F}_0 and \mathcal{G}_0 form the trivial partition of $(0, 1]$. Let now $1 < \beta < \gamma_n$, $n \geq 1$, and assume that $\mathcal{F}_{n-1} \cup \mathcal{G}_{n-1}$ forms a partition of $(0, 1]$. Lemma 3.4 and the order reversing property of $f_{\varphi^{n-1}(1)}$ give

$$f_{\varphi^{n-1}(1)}(1) < f_{\varphi^{n-1}(2)}(1) < f_{\varphi^{n-2}(2)}(1) < f_{\varphi^n(2)}(1) < 1.$$

For $n = 1$, we have $f_{21}(1) > 0$. Therefore, $F_{n-1,0}$ splits into $F_{n,|\varphi^{n-1}(1)|}$, $G_{n-1,0}$ and $F_{n,0}$, which are intervals of positive length. This shows (i) in case $n = 1$. By Proposition 3.5 and Lemma 3.9, $T^{g_{n-1}-1}$ is continuous on $F_{n-1,0}$, thus every $F_{n-1,k}$, $0 \leq k < g_{n-1}$, also splits into $F_{n,|\varphi^{n-1}(1)|+k}$, $G_{n-1,k}$ and $F_{n,k}$. If n is even, then this proves (i) since $g_{n-1} = |\varphi^{n-1}(1)|$. In this case, we have $F_{n-1,g_{n-1}} = F_{n,g_n}$, thus $\mathcal{F}_n \cup \mathcal{G}_n$ is a refinement of the partition $\mathcal{F}_{n-1} \cup \mathcal{G}_{n-1}$. If n is odd, then we use that $T^{|\varphi^{n-1}(1)|}$ is continuous on $G_{n-1,0} \cup F_{n-1,0}$ since both $d(T^{|\varphi^{n-1}(2)|}(1))$ and $d(1)$ start with $\varphi^{n-1}(1)$. In this case, $F_{n-1,g_{n-1}} = F_{n,|\varphi^{n-1}(1)|-1}$ splits into F_{n,g_n} , $G_{n-1,|\varphi^{n-1}(1)|-1}$ and $F_{n,|\varphi^{n-1}(1)|-1}$, thus (i) holds for odd n too, and $\mathcal{F}_n \cup \mathcal{G}_n$ is again a refinement of the partition $\mathcal{F}_{n-1} \cup \mathcal{G}_{n-1}$.

For all $0 \leq m < n$, the continuity of $T^{|\varphi^m(1)|}$ on $G_{m,0}$ gives (ii). Moreover, $T^{|\varphi^m(1)|}(G_{m,0}) = (T^{|\varphi^m(2)|}(1), T^{|\varphi^{m+1}(2)|}(1))$ and $F_{m+1,0} \subset F_{m,0}$ imply (iii). \square

Corollary 4.3. *Let $1 < \beta < \gamma_n$, $n \geq 0$. Then*

$$T_{-\beta}\left(\bigcup_{I \in \mathcal{F}_n(\beta)} I\right) = \bigcup_{I \in \mathcal{F}_n(\beta)} I \quad \text{and} \quad T_{-\beta}^{-1}\left(\bigcup_{I \in \mathcal{G}_n(\beta)} I\right) \subseteq \bigcup_{I \in \mathcal{G}_n(\beta)} I.$$

If $\gamma_{n+1} \leq \beta < \gamma_n$, then $F(\beta) = \bigcup_{I \in \mathcal{F}_n(\beta)} I$ (and $G(\beta) = \bigcup_{I \in \mathcal{G}_n(\beta)} I$ by definition).

Proposition 4.4. *Let $\beta > 1$ and μ be an invariant measure of $T_{-\beta}$ which is absolutely continuous with respect to the Lebesgue measure. Then $\mu(G(\beta)) = 0$.*

Proof. If $\beta \geq \gamma_1$, then $\mu(G(\beta)) = \mu(\emptyset) = 0$. We will show, by induction on n , that

$$\mu(G_{n,k}) = 0 \quad \text{for all } n \geq 0, 0 \leq k < |\varphi^n(1)|, 1 < \beta < \gamma_{n+1}. \quad (4.1)$$

Since $G(\beta) = \bigcup_{m=0}^{n-1} \bigcup_{k=0}^{|\varphi^m(1)|-1} G_{m,k}$ for $\gamma_{n+2} \leq \beta < \gamma_{n+1}$, this will prove the proposition.

For $1 < \beta < \gamma_1$, we have $G_{0,0} = (T^2(1), T(1)) = (f_1 T(1), T(1))$ and

$$T^{-1}(G_{0,0}) = f_1^{-1}(G_{0,0}) = (f_1^{-1} T(1), T(1)) \subset G_{0,0}.$$

Since $\mu(T^{-1}(G_{0,0})) = \mu(G_{0,0})$, we obtain $\mu((f_1 T(1), f_1^{-1} T(1))) = 0$. Iteratively, we get

$$\mu((f_1^{-2k+1} T(1), f_1^{-2k-1} T(1))) = 0, \quad \mu((f_1^{-2k-2} T(1), f_1^{-2k} T(1))) = 0$$

for all $k \geq 0$. This gives $\mu((T^2(1), \frac{1}{\beta+1})) = 0$ and $\mu((\frac{1}{\beta+1}, T(1))) = 0$, since f_1^{-1} is contracting with fixed point $\frac{1}{\beta+1}$. Then, the absolute continuity of μ implies $\mu(G_{0,0}) = 0$, hence (4.1) holds for $n = 0$.

Now consider $1 < \beta < \gamma_{n+1}$, $n \geq 1$, and assume that (4.1) holds for $n - 1$. We have

$$T(G_{m,k-1}) = \begin{cases} G_{m,k} & \text{if } 1 \leq k < |\varphi^m(1)|, 0 \leq m \leq n, \\ G_{m,0} \cup F_{m+1,|\varphi^m(1)|} \cup G_{m+1,0} & \text{if } k = |\varphi^m(1)|, 0 \leq m < n, \end{cases} \quad (4.2)$$

by Proposition 4.2 (i)–(iii). Since $T^{-1}(\bigcup_{I \in \mathcal{G}_n} I) \subset \bigcup_{I \in \mathcal{G}_n} I$ by Corollary 4.3, we obtain

$$T^{-1}(G_{n,0}) \subset G_{n,|\varphi^n(1)|-1} \cup G_{n-1,|\varphi^{n-1}(1)|-1} \quad \text{and} \quad T^{1-|\varphi^n(1)|}(G_{n,|\varphi^n(1)|-1}) = G_{n,0},$$

thus, up to a set of μ -measure zero, $T^{-|\varphi^n(1)|}(G_{n,0}) = f_{\varphi^n(1)}^{-1}(G_{n,0}) \subset G_{n,0}$. We have

$$f_{\varphi^n(1)}^{-1}(G_{n,0}) = f_{\varphi^n(1)}^{-1}((f_{\varphi^n(1)} T^{|\varphi^n(2)|}(1), T^{|\varphi^n(2)|}(1))) = (f_{\varphi^n(1)}^{-1} T^{|\varphi^n(2)|}(1), T^{|\varphi^n(2)|}(1)).$$

As in the case $n = 0$, we obtain

$$\mu((f_{\varphi^n(1)}^{-2k+1} T^{|\varphi^n(2)|}(1), f_{\varphi^n(1)}^{-2k-1} T^{|\varphi^n(2)|}(1))) = 0 = \mu([f_{\varphi^n(1)}^{-2k-2} T^{|\varphi^n(2)|}(1), f_{\varphi^n(1)}^{-2k} T^{|\varphi^n(2)|}(1)])$$

for all $k \geq 0$, thus $\mu((T^{|\varphi^n(2)|}(1), y)) = 0$ and $\mu((y, T^{|\varphi^n(2)|}(1))) = 0$, where y denotes the fixed point of $f_{\varphi^n(1)}$, i.e., $\mu(G_{n,0}) = 0$. Since $\mu(G_{n,k}) = \mu(T^{-k}(G_{n,k})) = \mu(G_{n,0})$ for $1 \leq k < |\varphi^n(1)|$, (4.1) holds for n . \square

Proposition 4.5. *Let $\beta > 1$, then $\lim_{k \rightarrow \infty} \lambda(T_{-\beta}^{-k}(G(\beta))) = 0$.*

Proof. If $\beta \geq \gamma_1$, then $G(\beta) = \emptyset$ and the statement holds. For $1 < \beta < \gamma_{n+1}$, $n \geq 0$, the preimage $T^{-k}(G_{n,0})$ can be written as a disjoint union

$$T^{-k}(G_{n,0}) = \bigcup_{a_1 \cdots a_k \in L_k} f_{a_1 \cdots a_k}^{-1}(G_{n,0})$$

with $L_k \subseteq \{1, 2\}^k$. Then we have $\mu(T^{-k}(G_{n,0})) = \#(L_k) \beta^{-k} \mu(G_{n,0})$. By (4.2) and Corollary 4.3, L_k consists of the length k suffixes of all concatenations $1^{i_0}(\varphi(1))^{i_1} \cdots (\varphi^n(1))^{i_n}$, $i_0, i_1, \dots, i_n \geq 0$. Therefore, the number of elements in L_k is bounded by $\binom{k}{n}$, which grows only polynomially (in k). This yields that $\lim_{k \rightarrow \infty} \mu(T^{-k}(G_{n,0})) = 0$. By $T^{-j}(G_{n,j}) = G_{n,0}$, $1 \leq j < |\varphi^n(1)|$, the same holds for $G_{n,j}$, thus $\lim_{k \rightarrow \infty} \mu(T^{-k}(I)) = 0$ for all $I \in \mathcal{G}_{n+1}$. \square

5. LOCALLY EVENTUALLY ONTO

Lemma 5.1. *Let $1 < \beta < \gamma_1$, I be an interval of positive length in $F(\beta)$, and $n \geq 1$ such that $\eta_{n+1} < \beta < \eta_n$ with odd $n \geq 1$, or $\eta_{n+1} \leq \beta \leq \eta_n$ with even $n \geq 2$. Then there exists an $m \geq 0$ such that $T_{-\beta}^m(I) \supseteq F_{n,0}(\beta) \setminus \{1\}$.*

Proof. We use ideas of Góra's proof of Proposition 8 in [Gór07]. We distinguish four cases.

Case 1: odd $n \geq 1$, $\eta_{n+1} < \beta < \gamma_n$. In this case, we have $\beta^{2g_n} = \beta^{g_{n+1}} > \beta^{g_n} + \beta^{-1} \geq \beta + \beta^{-1} > 2$. Therefore, the largest connected components of images of I grow until they cover two consecutive discontinuities of T^{2g_n} in the interior of $F(\beta)$. Since the only

discontinuity of T is $1/\beta \in F_{n,g_n-1}$, the images eventually also cover two consecutive discontinuities of T^{2g_n} in the interior of $F_{n,g_n-1} = [T^{|\varphi^n(2)|-2}(1), T^{|\varphi^n(1)|-1}(1)]$. We have

$$\begin{aligned} T^{2g_n}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) &= T^{g_n}([1/\beta, T^{|\varphi^n(1)|-1}(1)]), \\ T^{2g_n}([f_{11\varphi^n(1)}^{-1}(1), 1/\beta]) &= T^{g_n-1}([1/\beta, T^{|\varphi^n(1)|-1}(1)]), \\ T^{2g_n}((f_{2\varphi^n(1)}^{-1}(1), T^{|\varphi^n(1)|-1}(1))) &= T^{g_n}([T^{|\varphi^n(2)|-2}(1), 1/\beta]), \\ T^{2g_n}([T^{|\varphi^n(2)|-2}(1), f_{11\varphi^n(1)}^{-1}(1)]) &= T^{g_n-1}([T^{|\varphi^n(21)|-1}(1), 1/\beta]), \end{aligned}$$

thus T^{2g_n} is continuous on these four intervals. Note that $T^{|\varphi^n(2)|-2}(1) < f_{11\varphi^n(1)}^{-1}(1)$ since $f_{11\varphi^n(1)} T^{|\varphi^n(2)|-2}(1) = f_{\varphi^n(21)}(1) > 1$. The discontinuities of T^{2g_n} in the interior of F_{n,g_n-1} are therefore $1/\beta$, $f_{2\varphi^n(1)}^{-1}(1)$ and $f_{11\varphi^n(1)}^{-1}(1)$. With

$$T^{g_n+1}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) = F_{n,0} \quad \text{and} \quad T^{g_n+1}([1/\beta, T^{|\varphi^n(1)|-1}(1)]) = F_{n,0} \setminus \{1\},$$

we obtain that $T^m(I) \supseteq F_{n,0} \setminus \{1\}$ for some $m \geq 0$.

Case 2: odd $n \geq 1$, $\gamma_{n+1} \leq \beta \leq \eta_{n+1}$. Since $\beta^{g_{n+1}+1} \geq \beta + 1 > 2$, we look at T^{2g_n+1} on F_{n,g_n-1} . Note that $T^{2g_n}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) = F_{n,g_n-1}$, thus there is a discontinuity of T^{2g_n+1} in the interior of $[1/\beta, f_{2\varphi^n(1)}^{-1}(1)]$. Contrary to Case 1, T^{2g_n+1} has no discontinuity in F_{n,g_n-1} on the left of $1/\beta$. More precisely,

$$\begin{aligned} T^{2g_n+1}([1/\beta, f_{2\varphi^n(2)}^{-1}(1)]) &= T([T^{|\varphi^n(2)|-2}(1), 1/\beta]), \\ T^{2g_n+1}([f_{2\varphi^n(2)}^{-1}(1), f_{2\varphi^n(1)}^{-1}(1)]) &= T([1/\beta, T^{|\varphi^n(1)|-1}(1)]), \\ T^{2g_n+1}((f_{2\varphi^n(1)}^{-1}(1), T^{|\varphi^n(1)|-1}(1))) &= T^{g_n+1}([T^{|\varphi^n(2)|-2}(1), 1/\beta]), \\ T^{2g_n+1}([T^{|\varphi^n(2)|-2}(1), 1/\beta]) &= T^{g_n}([T^{|\varphi^n(21)|-1}(1), T^{|\varphi^n(1)|-1}(1)]), \end{aligned}$$

Here, $f_{\varphi^n(21)}(1) \leq 1$ implies that $T^{|\varphi^n(21)|-1}(1) = f_2^{-1} f_{\varphi^n(21)}(1) \geq 1/\beta$. Hence, the discontinuities of T^{2g_n+1} in the interior of F_{n,g_n-1} are $1/\beta$, $f_{2\varphi^n(2)}^{-1}(1)$ and $f_{2\varphi^n(1)}^{-1}(1)$, with

$$T^{2g_n+2}([1/\beta, f_{2\varphi^n(2)}^{-1}(1)]) = F_{n+1,0} \setminus \{1\} \quad \text{and} \quad T^{2g_n+1}([f_{2\varphi^n(2)}^{-1}(1), f_{2\varphi^n(1)}^{-1}(1)]) = F_{n,0}.$$

This gives $T^m(I) \supseteq F_{n+1,0} \setminus \{1\}$ for some $m \geq 0$.

Case 3: even $n \geq 2$, $\eta_{n+1} \leq \beta < \gamma_n$. In this case, we have $\beta^{2g_n} = \beta^{g_{n+1}-1} > \beta^{g_n-1} + \beta^{-1} \geq \beta + \beta^{-1} > 2$, $F_{n,g_n-1} = [T^{|\varphi^n(1)|-2}(1), T^{|\varphi^n(2)|-1}(1)]$, and

$$\begin{aligned} T^{2g_n}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) &= T^{g_n}([T^{|\varphi^n(1)|-2}(1), 1/\beta]), \\ T^{2g_n}((f_{11\varphi^n(1)}^{-1}(1), 1/\beta)) &= T^{g_n-1}((T^{|\varphi^n(1)|-2}(1), 1/\beta)), \\ T^{2g_n}([f_{2\varphi^n(1)}^{-1}(1), T^{|\varphi^n(2)|-1}(1)]) &= T^{g_n}([1/\beta, T^{|\varphi^n(21)|-2}(1)]), \\ T^{2g_n}([T^{|\varphi^n(1)|-2}(1), f_{11\varphi^n(1)}^{-1}(1)]) &= T^{g_n-1}([1/\beta, T^{|\varphi^n(2)|-1}(1)]). \end{aligned}$$

Here, $f_{2\varphi^n(1)} T^{|\varphi^n(2)|-1}(1) = f_{\varphi^n(21)}(1) \geq 1$ implies $f_{2\varphi^n(1)}^{-1}(1) \leq T^{|\varphi^n(2)|-1}(1)$. The discontinuities of T^{2g_n} in the interior of F_{n,g_n-1} are at most $1/\beta$, $f_{2\varphi^n(1)}^{-1}(1)$ and $f_{11\varphi^n(1)}^{-1}(1)$, with

$$T^{g_n+2}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) = F_{n,0} \setminus \{1\}, \quad T^{g_n+3}((f_{11\varphi^n(1)}^{-1}(1), 1/\beta)) = F_{n,0} \setminus \{1, T^{|\varphi^n(1)|}(1)\},$$

similarly to Case 1. We obtain that $T^m(I) \supseteq F_{n,0} \setminus \{1, T^{|\varphi^n(1)|}(1)\}$ for some $m \geq 0$. Since $T^{g_n+1}(F_{n,0} \setminus \{1, T^{|\varphi^n(1)|}(1)\}) \supseteq F_{n,0} \setminus \{1\}$, we also have $T^m(I) \supseteq F_{n,0} \setminus \{1\}$ for some $m \geq 0$.

Case 4: even $n \geq 2$, $\gamma_{n+1} \leq \beta < \eta_{n+1}$. Since $\beta^{g_{n+1}+1} \geq \beta + 1 > 2$, we look at T^{2g_n+2} on F_{n,g_n-1} . We have $T^{2g_n+1}([1/\beta, f_{2\varphi^n(1)}^{-1}(1)]) = F_{n,g_n-1} \setminus \{T^{|\varphi^n(1)|-2}(1)\}$ with $f_{2\varphi^n(1)}^{-1}(1) > T^{|\varphi^n(2)|-1}(1)$, thus $f_{2\varphi^n(1)}^{-1}(1)$ is outside of F_{n,g_n-1} , but T^{2g_n+2} still has a discontinuity in $(1/\beta, T^{|\varphi^n(2)|-1}(1))$. More precisely,

$$\begin{aligned} T^{2g_n+2}([1/\beta, f_{2\varphi^n(2)}^{-1}(1)]) &= T([1/\beta, T^{|\varphi^n(2)|-1}(1)]), \\ T^{2g_n+2}((f_{11\varphi^n(1)}^{-1}(1), 1/\beta)) &= T^{g_n+1}((T^{|\varphi^n(1)|-2}(1), 1/\beta)), \\ T^{2g_n+2}((f_{2\varphi^n(2)}^{-1}(1), T^{|\varphi^n(2)|-1}(1))) &= T([T^{|\varphi^n(2)|-2}(1), 1/\beta]), \\ T^{2g_n+2}([T^{|\varphi^n(1)|-2}(1), f_{11\varphi^n(1)}^{-1}(1)]) &= T([T^{|\varphi^n(1)|-2}(1), T^{|\varphi^n(21)|-2}(1)]), \end{aligned}$$

with $T^{|\varphi^n(21)|-2}(1) = f_{11}^{-1} f_{\varphi^n(21)}(1) < 1/\beta$ since $f_{\varphi^n(21)}(1) < 1$. Therefore, the discontinuities of T^{2g_n+2} in the interior of F_{n,g_n-1} are $1/\beta$, $f_{2\varphi^n(2)}^{-1}(1)$ and $f_{11\varphi^n(1)}^{-1}(1)$, with

$$T^{2g_n+3}([1/\beta, f_{2\varphi^n(2)}^{-1}(1)]) = F_{n+1,0}, \quad T^{g_n+3}((f_{11\varphi^n(1)}^{-1}(1), 1/\beta)) = F_{n,0} \setminus \{1, T^{|\varphi^n(1)|}(1)\}.$$

This gives $T^m(I) \supseteq F_{n+1,0} \setminus \{1\}$ for some $m \geq 0$. \square

Lemma 5.2. *Let $1 < \beta < \gamma_1$ and $n \geq 1$ such that $\eta_{n+1} < \beta < \eta_n$ with odd $n \geq 1$, or $\eta_{n+1} \leq \beta \leq \eta_n$ with even $n \geq 2$. Then we have*

$$T_{-\beta}^{g_n(2g_n+(-1)^n)}(F_{n,0}(\beta) \setminus \{1\}) \supseteq F(\beta) \setminus \{T_{-\beta}^k(1) \mid 0 \leq k \leq g_n(2g_n + (-1)^n)\}.$$

Proof. Let $1 < \beta < \gamma_1$ and n as in the statement of the lemma. By Corollary 4.3, we have $F(\beta) = \bigcup_{I \in \mathcal{F}_n(\beta)} I$ if $\beta < \gamma_n$, and $F(\beta) = \bigcup_{I \in \mathcal{F}_{n-1}(\beta)} I$ if $\beta \geq \gamma_n$. In the latter case, $f_{\varphi^n(1)} \leq f_{\varphi^{n-1}(1)}$ implies that $F_{n-1,0} = F_{n,|\varphi^{n-1}(1)|} \cup F_{n,0}$, and we obtain, similarly to

the proof of Proposition 4.2, that $\bigcup_{I \in \mathcal{F}_{n-1}(\beta)} I = \bigcup_{I \in \mathcal{F}_n(\beta)} I$. Therefore, we always have $F(\beta) = \bigcup_{I \in \mathcal{F}_n(\beta)} I$.

Let first $n \geq 1$ be odd. For simplicity, we omit points in $\{T^k(1) \mid k \geq 0\}$ in the following statements. We have $T^{g_n+1}(F_{n,0}) = F_{n+1,0} \cup F_{n,1}$ and

$$T^{g_n-1}(F_{n+1,0}) = [T^{|\varphi^n(21)|-1}(1), T^{|\varphi^n(1)|-1}(1)].$$

Since $\beta > \eta_{n+1}$, we have $T^{|\varphi^n(21)|-1}(1) < 1/\beta$, thus $T^{g_n}(F_{n+1,0}) \supseteq F_{n,0}$, and

$$T^{2g_n+1}(F_{n,0}) \supseteq F_{n,0} \cup F_{n,1}.$$

Inductively, we get $T^{(g_n-1)(2g_n+1)}(F_{n,0}) \supseteq \bigcup_{k=0}^{g_n-1} F_{n,k}$. Then, $T(F_{n,g_n-1}) = F_{n,g_n} \cup F_{n,0}$ yields

$$T^{g_n(2g_n-1)}(F_{n,0}) = T^{(g_n-1)(2g_n+1)+1}(F_{n,0}) = \bigcup_{k=0}^{g_n} F_{n,k} = F(\beta).$$

Here, we have only omitted points in $\{T^k(1) \mid 0 \leq k \leq g_n(2g_n-1)\}$.

Let now $n \geq 2$ be even. Omitting $\{T^k(1) \mid k \geq 0\}$, we have $T^{g_n}(F_{n,0}) = F_{n,g_n} \cup F_{n+1,0}$ and

$$T^{g_n-1}(F_{n+1,0}) = [T^{|\varphi^n(1)|-2}(1), T^{|\varphi^n(21)|-2}(1)].$$

Since $\beta \geq \eta_{n+1}$, we have $T^{|\varphi^n(21)|-2}(1) \geq 1/\beta$, thus $T^{g_n+1}(F_{n+1,0}) \supseteq F_{n,0}$, and

$$T^{2g_n+2}(F_{n,0}) \supseteq F_{n,0} \cup F_{n,1}.$$

As above, we get $T^{(g_n-1)(2g_n+2)}(F_{n,0}) \supseteq \bigcup_{k=0}^{g_n-1} F_{n,k}$. Since $T^{g_n+2}(F_{n,g_n-1}) \supseteq F_{n,g_n} \cup F_{n,0}$ and $T^{g_n+2}(F_{n,k}) \supseteq F_{n,k+1}$ for $0 \leq k < g_n$, we obtain

$$T^{g_n(2g_n+1)}(F_{n,0}) = T^{(g_n-1)(2g_n+2)+g_n+2}(F_{n,0}) = \bigcup_{k=0}^{g_n} F_{n,k} = F(\beta),$$

up to some points in $\{T^k(1) \mid 0 \leq k \leq g_n(2g_n+1)\}$. □

6. PROOFS OF THE MAIN RESULTS

Now we are ready to prove Theorems 2.1 and 2.2.

Proof of Theorem 2.2. For $1 < \beta < \gamma_1$, Lemma 5.1 and 5.2 show that $T_{-\beta}$ is locally eventually onto on $F(\beta) \setminus \{T_{-\beta}^k(1) \mid 0 \leq k \leq g_n(2g_n + (-1)^n)\}$ for some $n \geq 1$. For $\beta \geq \gamma_1$, this was shown in [Gór07, Proposition 8]. The iterated $T_{-\beta}$ -images of any interval of positive length clearly contain $1/\beta$, thus they also contain $T_{-\beta}^k(1)$ for any $k \geq 0$. Therefore, $T_{-\beta}$ is locally eventually onto on $F(\beta) = (0, 1] \setminus G(\beta)$.

The second part of the theorem is in Corollary 4.3 and Proposition 4.5. □

Proof of Theorem 2.1. By Proposition 4.4, all the intervals in $\mathcal{G}_n(\beta)$ are gaps. By Theorem 2.2, they are the only gaps. Calculating directly the numbers of the intervals of $\mathcal{G}_n(\beta)$ by the definition, we complete the proof. □

Theorem 2.5 was proved in Section 3, thus it only remains to prove Theorem 2.6. Recall that a number $\beta > 1$ is a $(-\beta)$ -number if $V_{-\beta} = \{T_{-\beta}^n(1) \mid n \geq 0\}$ is a finite set.

Proof of Theorem 2.6. Let $V'_{-\beta}$ be the set of numbers in $V_{-\beta}$ which are not right endpoints of gaps, and $J_x = (\max\{y \in V_{-\beta} \cup \{0\} \mid y < x\}, x)$ for each $x \in V'_{-\beta}$. Then $\{J_x \mid x \in V'_{-\beta}\}$ is a partition of $F(\beta)$ up to finitely many points, and $T_{-\beta}(J_x)$ is a union of intervals J_y , $y \in V'_{-\beta}$, for each $x \in V'_{-\beta}$. Define a matrix

$$M_{-\beta} = (m_{x,y})_{x,y \in V'_{-\beta}} \quad \text{with} \quad m_{x,y} = \frac{\beta \lambda(J_x \cap T_{-\beta}^{-1}(J_y))}{\lambda(J_y)},$$

i.e., $m_{x,y}$ is the number of times that J_y is contained in $T_{-\beta}(J_x)$. Since

$$\sum_{y \in V'_{-\beta}} m_{x,y} \lambda(J_y) = \beta \lambda(J_x)$$

for all $x \in V'_{-\beta}$, $(\lambda(J_x))_{x \in V'_{-\beta}}$ is a positive eigenvector of $M_{-\beta}$ to the eigenvalue β . The property that $T_{-\beta}$ is locally eventually onto on $F(\beta)$ implies that $M_{-\beta}$ is primitive, thus β is the Perron-Frobenius eigenvalue of $M_{-\beta}$, which is a Perron number. \square

We finally show that the set of β -numbers and the set of $(-\beta)$ -numbers do not include each other. For the definition of a β -number, it is convenient to extend the domain of T_β to $[0, 1]$ by setting $T_\beta(1) = \beta - \lfloor \beta \rfloor$. Then β is a β -number if and only if $(T_\beta^n(1))_{n \geq 0}$ is eventually periodic. We know that all Pisot numbers are both β -numbers [Ber77, Sch80] and $(-\beta)$ -numbers [FL09]. Therefore, the symmetric difference between the β -numbers and the $(-\beta)$ -numbers can only contain Perron numbers which are not Pisot numbers.

Proposition 6.1. *Let $\beta > 1$ with $\beta^4 = \beta + 1$, i.e., $\beta \approx 1.2207$. Then $T_{-\beta}^{10}(1) = T_{-\beta}^5(1)$, and $(T_\beta^n(1))_{n \geq 0}$ is aperiodic.*

Proof. It is easy to check that $T_{-\beta}^{10}(1) = T_{-\beta}^5(1)$.

On the other hand, let α be an algebraic conjugate of β satisfying $|\alpha| > 1$, i.e., $\alpha \approx -0.2481 \pm 1.034i$, and let $\sigma : \mathbb{Q}(\beta) \rightarrow \mathbb{Q}(\alpha)$ be the field homomorphism defined by $\sigma(\beta) = \alpha$. Since $T_\beta(x) \in \{\beta x, \beta x - 1\}$, we have $|\sigma(T_\beta(x))| \geq |\alpha| |\sigma(x)| - 1$ for all $x \in [0, 1] \cap \mathbb{Q}(\beta)$, thus $|\sigma(T_\beta(x))| > |\sigma(x)|$ if $|\sigma(x)| > 1/(|\alpha| - 1)$. We have $T_\beta^{35}(1) = \beta^{35} - \beta^{34} - \beta^{26} - \beta^{13} - \beta^4$, and one can check that $|\sigma(T_\beta^{35}(1))| > 1/(|\alpha| - 1)$. This implies that $(|\sigma(T_\beta^n(1))|)_{n \geq 35}$ is a strictly increasing sequence, hence $(T_\beta^n(1))_{n \geq 0}$ is aperiodic. \square

Proposition 6.2. *Let $\beta > 1$ with $\beta^7 = \beta^6 + 1$, i.e., $\beta \approx 1.2254$. Then $T_\beta^7(1) = 0$, and $(T_{-\beta}^n(1))_{n \geq 0}$ is aperiodic.*

Proof. We have $T_\beta^7(1) = \beta^7 - \beta^6 - 1 = 0$.

Let α be an algebraic conjugate of β satisfying $|\alpha| > 1$, i.e., $\alpha \approx 0.7802 \pm 0.7053i$, and let $\sigma : \mathbb{Q}(\beta) \rightarrow \mathbb{Q}(\alpha)$ be the field homomorphism defined by $\sigma(\beta) = \alpha$. We have $T_{-\beta}(x) \in \{-\beta x + 1, -\beta x + 2\}$, thus $|\sigma(T_{-\beta}(x))| \geq |\alpha| |\sigma(x)| - 2$ for all $x \in (0, 1] \cap \mathbb{Q}(\beta)$. It is more convenient to consider $\phi(x)$ with ϕ as in the Introduction, since $\phi(T_{-\beta}(x)) \in \{-\beta \phi(x), -\beta \phi(x) - 1\}$ and $|\sigma(\phi(T_{-\beta}(x)))| \geq |\alpha| |\sigma(\phi(x))| - 1$ for all $x \in (0, 1] \cap \mathbb{Q}(\beta)$. Now one can check that $|\sigma(\phi(T_{-\beta}^{53}(1)))| > 1/(|\alpha| - 1)$, which implies that $(|\sigma(\phi(T_{-\beta}^n(1)))|)_{n \geq 53}$ is a strictly increasing sequence, hence $(T_{-\beta}^n(1))_{n \geq 0}$ is aperiodic. \square

REFERENCES

- [AAB⁺95] J.-P. Allouche, A. Arnold, J. Berstel, S. Brlek, W. Jockusch, S. Plouffe, and B. E. Sagan. A relative of the Thue-Morse sequence. *Discrete Math.*, 139(1-3):455–461, 1995. Formal power series and algebraic combinatorics (Montreal, PQ, 1992).
- [Ber77] A. Bertrand. Développements en base de Pisot et répartition modulo 1. *C. R. Acad. Sci. Paris Sér. A*, 285(6):419–421, 1977.
- [Dub06] A. Dubickas. On the distance from a rational power to the nearest integer. *J. Number Theory*, 117(1):222–239, 2006.
- [Dub07] A. Dubickas. On a sequence related to that of Thue-Morse and its applications. *Discrete Math.*, 307(9-10):1082–1093, 2007.
- [Fal08] B. Faller. *Contribution to the Ergodic Theory of Piecewise Monotone Continuous Maps*. PhD thesis, École Polytechnique Fédérale de Lausanne, 2008.
- [FL09] C. Frougny and A. C. Lai. On negative bases. In *Proceedings of DLT 09*, volume 5583 of *Lecture Notes in Comput. Sci.*, pages 252–263. Springer, Berlin, 2009.
- [Gór07] P. Góra. Invariant densities for generalized β -maps. *Ergodic Theory Dynam. Systems*, 27(5):1583–1598, 2007.
- [Hof79] F. Hofbauer. On intrinsic ergodicity of piecewise monotonic transformations with positive entropy. *Israel J. Math.*, 34(3):213–237, 1979.
- [Hof81a] F. Hofbauer. On intrinsic ergodicity of piecewise monotonic transformations with positive entropy. II. *Israel J. Math.*, 38(1-2):107–115, 1981.
- [Hof81b] F. Hofbauer. The structure of piecewise monotonic transformations. *Ergodic Theory Dynamical Systems*, 1(2):159–178, 1981.
- [IS09] S. Ito and T. Sadahiro. Beta-expansions with negative bases. *Integers*, 9:A22, 239–259, 2009.
- [Kel78] G. Keller. Piecewise monotonic transformations and exactness. In *Seminar on Probability, Rennes 1978 (French)*, pages Exp. No. 6, 32. Univ. Rennes, Rennes, 1978.
- [Lot97] M. Lothaire. *Combinatorics on words*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1997.
- [LY78] T.-Y. Li and J. A. Yorke. Ergodic transformations from an interval into itself. *Trans. Amer. Math. Soc.*, 235:183–192, 1978.
- [MP] Z. Masáková and E. Pelantová. Ito-Sadahiro numbers vs. Parry numbers. Preprint, arXiv:1010.6181v1.
- [Par60] W. Parry. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [PY98] M. Pollicott and M. Yuri. *Dynamical systems and ergodic theory*, volume 40 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1998.
- [Rén57] A. Rényi. Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.*, 8:477–493, 1957.
- [Roh61] V. A. Rohlin. Exact endomorphisms of a Lebesgue space. *Izv. Akad. Nauk SSSR Ser. Mat.*, 25:499–530, 1961.
- [Sch80] K. Schmidt. On periodic expansions of Pisot numbers and Salem numbers. *Bull. London Math. Soc.*, 12(4):269–278, 1980.
- [Wag79] G. Wagner. The ergodic behaviour of piecewise monotonic transformations. *Z. Wahrsch. Verw. Gebiete*, 46(3):317–324, 1979.
- [Wal78] P. Walters. Invariant measures and equilibrium states for some mappings which expand distances. *Trans. Amer. Math. Soc.*, 236:121–153, 1978.

LAMA CNRS UMR 8050, UNIVERSITÉ PARIS-EST - CRÉTEIL - VAL-DE-MARNE, UFR SCIENCES
ET TECHNOLOGIE, 61 AVENUE DU GÉNÉRAL DE GAULLE, 94010 CRÉTEIL CEDEX, FRANCE

E-mail address: `lingmin.liao@u-pec.fr`

LIAFA, CNRS, UNIVERSITÉ PARIS DIDEROT – PARIS 7, CASE 7014, 75205 PARIS CEDEX 13,
FRANCE

E-mail address: `steiner@liafa.jussieu.fr`