



Improved diagnosis and treatment of soft tissue sarcoma patients after implementation of national guidelines: A population based study

M.L.E.A. Jansen-Landheer, P. Krijnen, M.J. Oostindiër, W.M. Kloosterman-Boele, E.M. Noordijk, M.A. Nooij, W.H. Steup, A.H.M. Taminiau, R. Vree, P.C.W. Hogendoorn, et al.

► To cite this version:

M.L.E.A. Jansen-Landheer, P. Krijnen, M.J. Oostindiër, W.M. Kloosterman-Boele, E.M. Noordijk, et al.. Improved diagnosis and treatment of soft tissue sarcoma patients after implementation of national guidelines: A population based study. *EJSO - European Journal of Surgical Oncology*, 2009, 35 (12), pp.1326. 10.1016/j.ejso.2009.05.002 . hal-00556305

HAL Id: hal-00556305

<https://hal.science/hal-00556305>

Submitted on 16 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Improved diagnosis and treatment of soft tissue sarcoma patients after implementation of national guidelines: A population based study

Authors: M.L.E.A. Jansen-Landheer, P. Krijnen, M.J. Oostindiër, W.M. Kloosterman-Boele, E.M. Noordijk, M.A. Nooij, W.H. Steup, A.H.M. Taminiau, R. Vree, P.C.W. Hogendoorn, R.A.E.M. Tollenaar, H. Gelderblom

PII: S0748-7983(09)00161-9

DOI: [10.1016/j.ejso.2009.05.002](https://doi.org/10.1016/j.ejso.2009.05.002)


Reference: YEJSO 2834

To appear in: *European Journal of Surgical Oncology*

Received Date: 10 March 2008

Revised Date: 4 May 2009

Accepted Date: 6 May 2009



Please cite this article as: Jansen-Landheer MLEA, Krijnen P, Oostindiër MJ, Kloosterman-Boele WM, Noordijk EM, Nooij MA, Steup WH, Taminiau AHM, Vree R, Hogendoorn PCW, Tollenaar RAEM, Gelderblom H. Improved diagnosis and treatment of soft tissue sarcoma patients after implementation of national guidelines: A population based study, *European Journal of Surgical Oncology* (2009), doi: 10.1016/j.ejso.2009.05.002

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Improved diagnosis and treatment of soft tissue sarcoma patients after implementation of national guidelines: a population based study

M.L.E.A. Jansen-Landheer^a, P. Krijnen^a, M.J. Oostindier^a, W.M. Kloosterman-Boele^a, E.M. Noordijk^b, M.A. Nooij^b, W.H. Steup^c, A.H.M. Taminiau^b, R. Vree^d, P.C.W. Hogendoorn^b, R.A.E.M. Tollenaar^b, H. Gelderblom^b

^aComprehensive Cancer Centre West, Schipholweg 5a, 2316 XB Leiden, the Netherlands

^bLeiden University Medical Center, PO Box 9600, 2300 RC Leiden, the Netherlands

^cHaga Hospital, PO Box 40551, 2504 LN The Hague, the Netherlands

^dDiaconessenhuis Leiden, PO Box 9650, 2300 RD Leiden, the Netherlands

Correspondence to

Mrs. M.L.E.A. Jansen – Landheer MD PhD

Leiden Cancer Registry

Comprehensive Cancer Centre West

Schipholweg 5 A

2316 XB Leiden

the Netherlands

Tel: +31 71 52 59 759

Fax: +31 71 52 59 700

m.jansen@ikca.nl

Keywords: sarcoma, practice guidelines, multidisciplinary treatment, pathology, surgery

ABSTRACT

Aim

The majority of clinicians, radiologists and pathologists have limited experience with soft tissue sarcomas. In 2004, national guidelines were established in the Netherlands to improve the quality of diagnosis and treatment of these rare tumours. This study evaluates the compliance with the guidelines over time.

Patients

Population based series of 119 operated patients with a soft tissue sarcoma (STS) diagnosed in 1998-1999 (79 before implementation of new guidelines) and in 2006 (40 after implementation)

Methods

Coded information regarding patient and tumour characteristics as well as (the results of) pathology review was collected from the medical patient file by two experienced data-managers.

Results

Diagnostic imaging of the tumour was performed according to the guidelines in 75-100% depending on the site of the tumour (abdominal versus non-abdominal) as well as the time of diagnosis.

Adherence to the guidelines with respect to invasive diagnostic procedures in patients with non-abdominal STS improved over time. A pre-operative histological diagnosis was obtained in 42% of the patients in 1998-1999 and in 72% of the patients in 2006 ($p<0.001$). The guidelines for reporting on pathology were increasingly adhered to. In 2006, (nearly) all pathology reports mentioned tumour size, morphology, tumour grade, resection margins and radicality. This represents a major improvement compared to the pathology reports in 1998-1999, where these aspects were not mentioned in 14% to 40% of the cases. The proportion of prospective pathology reviews by (a member of) the expert panel increased from 60% in 1998-1999 to 90% in 2006 ($p=0.001$).

Discussion

The compliance with the guidelines has been optimised by the increased attention to this group of patients. Most important factors have been the reporting of the results of the first evaluation and the (discussions about) the centralisation of treatment. Further improvements could be reached by the prospective web based registry monitoring logistic aspects as well as parameters useful for the evaluation of the quality of care.

INTRODUCTION

Soft tissue sarcomas (STS) are a rare and heterogeneous group of tumours with a variable biology and clinical behaviour.¹⁻³ Subsequently, the majority of clinicians and pathologists have limited experience with the extensive variability in clinical and histopathological presentation of these tumours and their complex multimodality treatment. In the Netherlands a national guideline has been available since 1993.^{4-6*} In 1997 the guideline was regionally updated for the western part of the country.^{7*}

Pre-operative multidisciplinary discussion of the results of diagnosis and staging is considered as an essential element of the treatment strategy since the chosen surgical approach depends on the size of the tumour and its relation to surrounding structures, as well as the histological type of the tumour, the histological grade and the presence of metastases at the time of diagnosis. It is the aim of the multidisciplinary meeting to define the optimal therapeutic approach, which usually consists of resection of the tumour followed by radiotherapy if indicated.

In 2004 a new, national, evidence-based guideline was published.^{8,9} This guideline recommends the review of all available histological material by a regional expert panel. In the region of the Comprehensive Cancer Centre West (CCCW) such a panel has been available for a much longer time which enables the evaluation of the impact of pathologic review.

Compliance with guidelines is important for various reasons. Experience from clinical trials indicates that diagnosis and treatment according to strict rules affects clinical outcome and patient survival.¹⁰

Similarly, explicit guidelines improve clinical practice because appropriate preoperative investigations and accurate staging are essential for planning of appropriate treatment.¹¹

This study intends to evaluate the adherence to guidelines in the CCCW region over time (between 1998 and 2006) with the ultimate goal to further improve the quality of oncological care for patients with a soft tissue sarcoma.

* Of the 1993 and 1997 guidelines an English version is not available. Since the aspects of the guidelines studied were roughly unchanged during the years an appendix was added to the manuscript consisting of a flowchart concerning the diagnosis and treatment of soft tissue tumours.

PATIENTS

Patients with a STS diagnosed in the period 1998-1999 (before implementation of new guidelines) and in 2006 (after implementation) were included in the study. Patients were selected from the population-based Leiden Cancer Registry of the CCCW. Selected STS were located in the head and neck area, (retro)peritoneum, extremities, thorax and trunk. Patients younger than 16 years at diagnosis and patients with STS from gynaecological or urological origin were excluded. Inclusion was restricted to operated patients. Non-operated patients were not selected, because in these cases the choice of treatment was individualised which hampered the evaluation of the compliance to the guidelines.

METHODS

Datacollection and definitions

Information regarding patient and tumour characteristics as well as (the results of) pathology review was collected from the medical patient file. Items on the registration form were based on studies of the Swedish Sarcoma Group Register.¹² The data were collected and coded by two data-managers of the CCCW.

A pre-operative diagnosis STS was assumed to be *considered* if this was explicitly recorded in the medical files or if histological material had been evaluated preoperatively and/or adequate imaging had been performed.

Content of the guidelines

The guidelines^{8,9} for diagnostic work-up require physical examination and imaging techniques indicating the size of the tumour and its extension in surrounding tissues (see flowchart in the appendix). In general magnetic resonance imaging (MRI) is the imaging method of choice, followed by computed tomography (CT) and conventional X-ray. In case of intra abdominal or intra thoracic tumours a CT is preferred to MRI. Distant metastases were excluded by pulmonary X-ray and an additional CT-scan in specific cases (patients with a T2/N1 tumour with a negative pulmonary X-ray). For definitive histological diagnosis, a true-cut biopsy is advocated. Only in patients with small (≤ 3 cm), unsuspecting and superficial (above the superficial fascia) tumours a radical excisional biopsy could be performed. In patients with large (> 3 cm), suspicious or deep tumours (beneath the superficial fascia), incisional biopsy was required before surgical treatment if true-cut biopsy provided

insufficient diagnostic evidence. Referral to a centre was recommended for patients with large and complex tumours.

Surgery and pathology

Surgical treatment consisted of a wide resection of the tumour, which is defined as 'en bloc' resection of the tumour (including the biopsy tract) with at least 2 cms of normal tissue in its surroundings.

State of the art diagnostic pathology is regarded to include statements about the radicality of resection¹³⁻¹⁵, grading¹⁶ and if appropriate expression of specific markers and assessment of chromosomal translocations and/or mutations in tumour specific genes. These elements have all been taken care of in our approach although not all findings are reported explicitly.

Pathology review (prospectively or retrospectively) was performed by the regional expert panel.

The regional expert panel is made up of at least five pathologists from the regional hospitals with a special interest in soft tissue tumours. The panel is chaired by a pathologist with qualified expertise in this area and who is also a referent pathologist of the EORTC soft tissue and bone sarcoma group.

RESULTS

In total, 119 patients with an operated STS were included in the study. Of these patients, 79 were diagnosed in the years 1998 and 1999 (before the implementation of the evidence-based national guidelines), and 40 were diagnosed in 2006 (after implementation). The patients in the two periods did not differ with respect to patient and tumour characteristics (Table 1). About half of the patients were male and the majority of the patients was younger than 70 years. Eighteen percent of the STS were intra-abdominal tumours, and 82% of the STS were located in the extremities, trunk or head and neck region. In 1998-1999, more small and superficial STS were included than in 2006: 18 versus 4. More than 60% of the STS were myogenic sarcomas, myxofibrosarcomas and liposarcomas, but various other types of STS within the WHO-categories were also found. The patients received their primary treatment in 11 district hospitals and/or one university hospital in the CCCW. The annual number of patients per hospital varied from 1 to 10 in the years 1998-1999 and from 1 to 15 in 2006.

Diagnostic procedures (Table 2)

For patients with a tumour in the extremities, trunk or head and neck region, the diagnosis STS was considered pre-operatively in 46 of the 66 patients in 1998-1999 and in 25 of the 32 patients in 2006 ($p=0.47$). For the remaining patients with a non-abdominal tumour, the diagnosis STS was made only after surgery (whoops surgery). These cases concerned a tumour smaller than 3 cm and located above the superficial fascia in 11 of 20 patients in 1998-1999 and in 2 of 7 patients in 2006.

In both periods the guidelines for diagnostic imaging for non-abdominal tumours (i.e. MRI) were adhered to in 75% of the patients in whom STS was considered pre-operatively. Of the non-abdominal STS without MRI, 4 of the 12 tumours in 1998-1999 and 2 of the 6 tumours in 2006 were smaller than 3 cm and located above the superficial fascia.

The requested CT in case of an abdominal STS was performed in 10 of 13 patients in 1998-1999 and in all 8 patients in 2006 ($p=0.26$).

The use of diagnostic imaging for distant metastases did not change over time.

Adherence to the guidelines with respect to invasive diagnostic procedures in patients with non-abdominal STS improved over time. A pre-operative histological diagnosis was obtained in 42% of the patients in 1998-1999 and in 72% of the patients in 2006 ($p<0.001$). An excisional biopsy was performed in 25 of the 66 patients with non-abdominal STS in 1998-1999 compared to 8 of 32 patients

in 2006. In 11 and 1 of these patients, respectively, the excisional biopsy concerned a tumour that was smaller than 3 cm and located above the superficial fascia.

Most patients in 1998-1999 (72%) and in 2006 (75%) were discussed multidisciplinary ($p=0.83$). For the remaining patients, it was not documented whether or not a multidisciplinary discussion had taken place.

Treatment (Table 3)

Less than half of the (primary) resections were performed in a university hospital: 37% in 1998-1999 and 50% in 2006 ($p=0.17$). The proportion of patients who underwent a re-resection was constant over time: 24% of the patients diagnosed in 1998-1999 and 18% of the patients diagnosed in 2006 ($p=0.41$). Re-resection was performed more often if STS was not suspected pre-operatively (67% versus 11%, $p<0.001$).

The outcome of surgery (including re-resection) for patients with a non-abdominal STS in 2006 was less favourable than in 1998-1999: the proportion of patients whose tumour was not radically removed was 19% in 2006 compared to 6% in 1998-1999 ($p=0.02$). However, in 1998-1999 there were proportionally more small and superficial tumours than in 2006. Most of these patients received additional radiotherapy. This was not the case for 9 out of 66 patients in 1998-1999 and for 1 out of 32 patients in 2006, for whom radiotherapy may not have been feasible.

Pathology reports (Table 4)

The guidelines for reporting on pathology were increasingly adhered to. In 2006, (nearly) all pathology reports mentioned tumour size, morphology, tumour grade, resection margins and radicality. This represents a major improvement compared to the pathology reports in 1998-1999, where these aspects were not mentioned in 14% to 40% of the cases. The proportion of prospective pathology reviews by (a member of) the expert panel increased from 60% in 1998-1999 to 90% in 2006 ($p=0.001$).

DISCUSSION

Guidelines on diagnosis and treatment of soft tissue sarcomas have been available in the region for many years and updates were distributed to all involved clinicians by the regional multidisciplinary working group Bone and Soft Tissue Sarcomas. It could therefore be assumed that all clinicians were informed about the most recent version of the guideline during the study period. Through this forum continuous awareness is propagated and clinicians are in some way accounted for their decision making in patients with soft tissue sarcomas.

Increasing awareness among medical specialists

Since the majority of patients are admitted through the out-patient ward of the surgical departments (not necessarily specialised in oncology) the primary awareness at this first intake is crucial. Apart from the introduction and implementation of the revised soft tissue sarcoma guideline a shift in approach amongst the surgical oncologists in the CCW region was evolving. This shift concerned the joint regional approach in the diagnosis and treatment of so called low-volume complex solid cancers, i.e. esophageal, pancreatic, locally advanced rectal cancer and liver metastases. Through centralised data collection and frequent reporting and discussions on the volumes and outcomes of these tumours clinicians were confronted with their own results, compared to others. By this approach an increasing awareness arose with respect to the benefits of centralised treatment. This shift undoubtedly also had an effect on the diagnosis and treatment of soft tissue sarcomas, perhaps not that much on centralisation but clearly on adherence to the guidelines.

Improving the implementation of guidelines

The current evaluation gave insight in the way soft tissue sarcomas were diagnosed and treated within the region of the CCCW over time, although the number of patients is small. In many aspects the guidelines were increasingly adhered to. The most remarkable changes were found in the percentage of pre-operative histological diagnosis in patients with a non-abdominal tumour and in the percentage of patients in whom pathology review was performed prospectively. The continuous and repetitive referral to the national guidelines can be accounted for this effect since more awareness for the standardised work-up is created amongst the first clinicians in line, not always the most experienced ones, therefore having highly valuable educational impact. A secondary but important effect of this

guideline-based and multidisciplinary approach is the phenomenon that doctors better adhere to guidelines because of public confrontation and accountability.

Relation to comparable studies

In the Netherlands a similar study was performed and published earlier.²⁹ This study was performed in the Northern part of the Netherlands between 1989 and 1996. Guidelines were significantly better adhered to in specialised centres²⁹. The benefit of centralised treatment of soft tissue sarcomas was also described in other publications.³⁰⁻³³ The additional value was expected to be due to an increased familiarity with alarm signals and with the existing guidelines as well as more experience with the complex treatment which results in appropriate preoperative investigations and accurate staging, essential for planning of appropriate treatment.¹¹ This association was not found in the CCCW region possibly because the study population was too small.

How a better adherence to guidelines was and can be achieved

The increased adherence to the guidelines, however, can be explained by a few factors in addition to the process that was already described. First, the results of the first study have been communicated with all involved clinicians within the region in order to encourage the use of the available guidelines. Second, pathologic material was increasingly reviewed by a regional expert panel. These factors combined with the strong regional intention to further improve the quality of surgical oncological care have resulted in the intended centralisation of treatment in which patients with a tumour > 3 cm, located subfascially or tumours with an aberrant course in a specialised centre. The second evaluation was performed during the process of centralisation. This explains why patients were still operated on in several district hospitals. Moreover, in 2004, the national evidence-based guideline⁸ was published. Both have contributed to a higher awareness and attention for the diagnosis and treatment of patients with soft tissue sarcomas. Further improvements could be reached by the prospective webbased registry which was started in 2005 and which is used for monitoring logistic aspects as well as indicators of the quality of care.

Table 1: Patient and tumour characteristics by period of diagnosis

	1998-1999 N	2006 N	P value
Total no. of patients	79	40	
Gender			0.85
Male	37	20	
Female	42	20	
Age at diagnosis			0.41
< 50	29	12	
50-69	21	15	
70-79	19	6	
80 +	10	7	
Tumour site			0.20
Extremities	45	16	
Trunk / head / neck	21	16	
Abdomen	13	8	
Tumour size and location *			0.24
Small / superficial	18	4	
Large / deep	48	28	
Abdominal	13	8	
Morphology			0.16
Myogenic sarcoma	17	10	
Myxofibrosarcoma	18	10	
Liposarcoma	10	8	
Synoviosarcoma	4	2	
Nerve sheath sarcoma	2	4	
Sarcoma other / NOS	28	6	

* Small/superficial tumours are located in extremities/trunk/head/neck, less than 3 cm in greatest dimension, and located above the superficial fascia.

Large/deep tumours are located in extremities/trunk/head/neck, and more than 3 cm in greatest dimension or located beneath the superficial fascia.

Table 2: Adherence to the diagnostic guidelines by period of diagnosis

	1998-1999 N	2006 N	P value
Non-abdominal STS	66	32	
STS considered pre-operatively			0.47
Yes	46	25	
No	20	7	
MRI *			1.00
Yes	34/46	19/25	
No	12/46	6/25	
Invasive diagnostic procedures			<0.001
Cytology	13	1	
True cut biopsy	6	17	
Incisional biopsy	22	6	
Not performed (excisional biopsy)	25	8	
Abdominal STS	13	8	
CT scan			0.26
Yes	10	8	
No	3	0	
All STS	79	40	
Diagnostic imaging for distant metastasis			0.32
Yes	61	28	
No	16	12	
Unknown	2	0	
Discussed in multidisciplinary meeting			0.83
Yes	57	30	
No/Not recorded	22	10	

* for patients in whom STS was considered pre-operatively.

Table 3: Treatment by period of diagnosis

	1998-1999 N	2006 N	P value
All STS	79	40	
Primary resection in university hospital			0.17
Yes	29	20	
No	50	20	
Re-resection performed			0.41
Yes	19	7	
No	60	33	
Non-abdominal STS	66	32	
Resection margins			0.10
Wide	41	14	
Marginal	9	10	
Intra-lesional	4	4	
Unknown	12	4	
Optimal treatment *			0.23
Yes	49	28	
No	9	1	
Unknown	8	3	


* Treatment was considered optimal if resection margins were free or if the patient received post-operative radiotherapy after irradical surgery.

Table 4: Adherence to the guidelines for reporting on pathology by period of diagnosis

	1998-1999 N	2006 N	P value
All STS	79	40	
Pathological reports mentioning:			
Tumour size			<0.001
Yes	55	39	
No	24	1	
Morphology			0.02
Yes	68	40	
No	11	0	
Grade			0.10
Yes	69	39	
No	10	1	
Resection margins			0.004
Yes	61	39	
No	18	1	
Radicality			<0.001
Yes	47	40	
No	32	0	
Pathology reviewed by expert panel			0.001
Yes	47	36	
No	32	4	

Appendix to the guideline on soft tissue tumours, diagnosis chapter

Version: 1.0, Responsible party: Dutch working group on soft tissue tumours, Date: 01 January 2004

FLOWCHART

Acknowledgement

This study was enabled by a grant of the Stichting Samenwerking Oncologie Haaglanden.

The authors would like to thank all clinicians and pathologists concerned for their co-operation and their permission for the review of all patient data and patient material.

All hospitals of the CCCW-region participated, which were the following:

Bronovo Hospital, The Hague

Diaconessenhuis Leiden, Leiden

Groene Hart Hospital, Gouda

Haga Hospital, The Hague

't Lange Land Hospital, Zoetermeer

Leiden University Medical Center

Medical Center Haaglanden, The Hague

Reinier de Graaf Group, Delft

Rijnland Hospital, Leiderdorp

Reference List

1. Nijhuis PH, Schaapveld M, Otter R, Molenaar WM, van der Graaf WT, Hoekstra HJ. Epidemiological aspects of soft tissue sarcomas (STS)--consequences for the design of clinical STS trials. *Eur J Cancer* 1999; 35:1705-10.
2. Visser, O., Coebergh, J. W. W., van Dijck, J. A. A. M., and Siesling, S. Incidence of cancer in the Netherlands 2001. 2004.
3. Graadt van Roggen JF, Hogendoorn PC. New insights in the classification of soft tissue tumors. *Ned Tijdschr Geneesk* 2002; 146:2022-6.
4. Nederlandse werkgroep weke delen tumoren. Richtlijn diagnostiek en behandeling weke-delentumoren. Resultaat van een consensusbijeenkomst, gehouden in 1993 te Rotterdam. Utrecht: Vereniging van Integrale Kankercentra, 1993.*
5. Van Geel AN, Van Unnik JA, Keus RB. Diagnosis and treatment of soft tissue tumours: the Dutch nationwide-accepted consensus. *Sarcoma* 1998; 2:183-91.
6. Van Geel AN, Van Unnik JA, Keus RB. Consensus soft tissue tumors. Dutch Workgroup Soft-Tissue Tumours. *Ned Tijdschr Geneesk* 1995; 139:833-7.
7. Anonymous Diagnostiek en behandeling weke delen tumoren. Leiden: Integraal Kankercentrum West, 1995.*
8. Dutch working group on soft tissue tumours. National guideline on soft tissue tumours (revision) 2004. www.oncoline.nl 2007.
9. van Geel AN, Wyrdeman HK, Seynaeve C, Hogendoorn PC, Bongaerts AH, Molenaar WM. Practice guideline 'Diagnostic techniques for soft tissue tumours and treatment of soft tissue sarcomas (revision)'. *Ned Tijdschr Geneesk* 2005; 149:924-8.
10. Stiller CA. Centralised treatment, entry to trials and survival. *Br J Cancer* 1994; 70:352-362.
11. Grimshaw JM, Russell IT. Effect of clinical guidelines on medical practice: a systematic review of rigorous evaluations. *Lancet* 1993; 342:1317-22.
12. Alvegard TA, Bauer H, Blomqvist C, Rydholm A, Smeland S. The Scandinavian Sarcoma Group--background, organization and the SSG Register--the first 25 years. *Acta Orthop Scand Suppl* 2004; 75:1-7.
13. Enneking WF. A system of staging musculoskeletal neoplasms. *Clin Orthop Relat Res* 1986; 204:9-24.
14. Enneking WF. The staging system for benign and malignant tumors of the musculoskeletal system. In: Enneking WF, editor. *Clinical musculoskeletal pathology*. Gainesville, Florida: University Press of Florida, 1990:451-466.
15. Enneking WF, Spanier SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. 1980. *Clin Orthop Relat Res* 2003; 415:4-18.
16. Trojani M, Contesso G, Coindre JM, Rouesse J, Bui NB, de Mascarel A, et al. Soft tissue sarcomas of adults; study of pathological prognostic variables and definition of a histopathological grading system. *International Journal of Cancer* 1984; 33:37-42.
17. Wiklund T, Huuhtanen R, Blomqvist C, Tukiainen E, Virolainen M, Virkkunen P, et al. The importance of a multidisciplinary group in the treatment of soft tissue sarcomas. *Eur J Cancer* 1996; 32A:269-73.
18. Baker LH, Benjamin RS. Histologic frequency of disseminated soft tissue sarcomas in adults. *Proc Am Soc Clin Oncol* 1978; 19:324
19. Presant CA, Russell WO, Alexander RW, Fu YS. Soft-tissue and bone sarcoma histopathology peer review: the frequency of disagreement in diagnosis and the need for second pathology opinions. The Southeastern Cancer Study Group experience. *J Clin Oncol* 1986; 4:1658-61.
20. Alvegard TA, Berg NO. Histopathology peer review of high-grade soft tissue sarcoma: the Scandinavian Sarcoma Group experience. *J Clin Oncol* 1989; 7:1845-51.
21. Shiraki M, Enterline HT, Brooks JJ, Cooper NS, Hirschl S, Roth JA, et al. Pathologic analysis of advanced adult soft tissue sarcomas, bone sarcomas, and mesotheliomas. The Eastern Cooperative Oncology Group (ECOG) experience. *Cancer* 1989; 64:484-90.
22. Harris M, Hartley AL, Blair V, Birch JM, Banerjee SS, Freemont AJ, et al. Sarcomas in north west England: I. Histopathological peer review. *Br J Cancer* 1991; 64:315-20.

23. Harris M, Hartley AL. Value of peer review of pathology in soft tissue sarcomas. *Cancer Treat Res* 1997; 91:1-8.
24. Meis-Kindblom JM, Bjerkehage B, Bohling T, Domanski H, Halvorsen TB, Larsson O, et al. Morphologic review of 1000 soft tissue sarcomas from the Scandinavian Sarcoma Group (SSG) Register. The peer-review committee experience. *Acta Orthop Scand Suppl* 1999; 285:18-26.
25. Eilber FR, Huth JF, Mirra J, Rosen G. Progress in the recognition and treatment of soft tissue sarcomas. *Cancer* 1990; 65:660-6.
26. Fletcher CDM, Kempson RL, Weiss SW. Recommendations for the reporting of soft tissue sarcoma. Association of Directors of Anatomic and Surgical Pathology. *Virchows Arch* 1999; 434:187-91.
27. Arbiser ZK, Folpe AL, Weiss SW. Consultative (expert) second opinions in soft tissue pathology. Analysis of problem-prone diagnostic situations. *Am J Clin Pathol* 2001; 116:473-6.
28. Van Glabbeke M, van Oosterom AT, Oosterhuis JW, Mouridsen H, Crowther D, Somers R, et al. Prognostic factors for the outcome of chemotherapy in advanced soft tissue sarcoma: an analysis of 2,185 patients treated with anthracycline-containing first-line regimens--a European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group Study. *J Clin Oncol* 1999; 17:150-7.
29. Nijhuis PH, Schaapveld M, Otter R, Hoekstra HJ. Soft tissue sarcoma--compliance with guidelines. *Cancer* 2001; 91:2186-95.
30. Gustafson P, Dreinhofer KE, Rydholm A. Soft tissue sarcoma should be treated at a tumor center. A comparison of quality of surgery in 375 patients. *Acta Orthop Scand* 1994; 65:47-50.
31. Trovik CS, Gustafson P, Bauer HC, Saeter G, Klepp R, Berlin O, et al. Consequences of local recurrence of soft tissue sarcoma: 205 patients from the Scandinavian Sarcoma Group Register. *Acta Orthop Scand* 2000; 71:488-95.
32. Clasby R, Tilling K, Smith MA, Fletcher CD. Variable management of soft tissue sarcoma: regional audit with implications for specialist care. *Br J Surg* 1997; 84:1692-6.
33. Rydholm A. Centralization of soft tissue sarcoma. The southern Sweden experience. *Acta Orthop Scand Suppl* 1997; 273:4-8.

* Of the 1993 and 1997 guidelines an english version is not available. Since the aspects of the guidelines studied were roughly unchanged during the years an appendix was added to the manuscript consisting of a flowchart concerning the diagnosis and treatment of soft tissue tumours.