



## **Circumcision among men who have sex with men in Scotland: Limited potential for HIV prevention**

Lisa M Mcdaid, Helen A Weiss, Graham J Hart

### **► To cite this version:**

Lisa M Mcdaid, Helen A Weiss, Graham J Hart. Circumcision among men who have sex with men in Scotland: Limited potential for HIV prevention. Sexually Transmitted Infections, 2010, 86 (5), pp.404. 10.1136/sti.2010.042895 . hal-00557474

**HAL Id: hal-00557474**

**<https://hal.science/hal-00557474>**

Submitted on 19 Jan 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Circumcision among men who have sex with men in Scotland:**

## **Limited potential for HIV prevention**

**Lisa M McDaid (nee Williamson)<sup>a</sup>**

**Helen A Weiss<sup>b</sup>**

**Graham J Hart<sup>c</sup>**

<sup>a</sup> MRC/CSO Social and Public Health Sciences Unit, Glasgow, UK

<sup>b</sup> MRC Tropical Epidemiology Group, London School of Hygiene and Tropical  
Medicine, London, UK

<sup>c</sup> Centre for Sexual Health and HIV Research, Research Department of Infection and  
Population Health, University College London, London, UK

Correspondence to: Lisa M McDaid, MRC/CSO Social and Public Health Sciences Unit,  
4 Lilybank Gardens, Glasgow, G12 8RZ, UK.

Tel: +44 (0) 141 357 3949. Fax: +44 (0) 141 337 2389.

Email: [l.mcdaid@sphsu.mrc.ac.uk](mailto:l.mcdaid@sphsu.mrc.ac.uk)

Word Count: 960

Keywords: circumcision, HIV, men who have sex with men, unprotected anal intercourse

## **ABSTRACT**

**Objective:** Male circumcision has been shown to reduce risk of HIV acquisition among heterosexual men but the impact among men who have sex with men (MSM) is not known. In this paper, we explore the feasibility of research into circumcision for HIV prevention among MSM in Scotland.

**Methods:** Anonymous, self-complete questionnaires and Orasure™ oral fluid collection kits were distributed to men visiting the commercial gay scenes in Glasgow and Edinburgh.

**Results:** 1508 men completed questionnaires (70.5% response rate) and 1277 provided oral fluid samples (59.7% response rate). Overall, 1405 men were eligible for inclusion in the analyses. 16.6% reported having been circumcised. HIV prevalence was similar among circumcised and uncircumcised men (4.2% and 4.6%, respectively). Although biologically, circumcision is most likely to protect against HIV for men practicing unprotected insertive anal intercourse (UIAI), only 7.8% (91/1172) of uncircumcised men reported exclusive UIAI in the past 12 months. Relatively few men reported being willing to participate in a research study on circumcision and HIV prevention (13.9%), and only 11.3% of uncircumcised men did so.

**Conclusion:** The lack of association between circumcision and HIV status, low levels of exclusive IUIAI, and low levels of willingness to take part in circumcision research studies suggest circumcision is unlikely to be a feasible HIV prevention strategy for MSM in the UK. Behaviour change should continue to be the focus of HIV prevention in this population.

Three randomised controlled trials (RCT) have shown that male circumcision reduces the risk of HIV acquisition among heterosexual men by about 60%,[1] and male circumcision is now recommended as an additional HIV prevention strategy for this population.[2] However, the evidence for such an association among men who have sex with men (MSM) is weak and inconsistent.[3-5] Biologically, circumcision may provide partial protection against HIV acquisition among MSM practicing unprotected insertive anal intercourse (UIAI), in a similar way as it protects against vaginal-penile transmission. Possible mechanisms include the fact the thin inner surface of the foreskin is susceptible to microtears and abrasions, and contains a high density of superficial Langerhan's cells; and the preputial space provides an environment thought to favour pathogen survival and replication.[4, 6-9]

In the UK, MSM are the group most at risk of acquiring HIV, with estimated prevalence of 5%,[10] and new HIV prevention strategies are needed. Circumcision is not common in the UK,[11] and attitudes towards circumcision are largely unknown. One recent survey of London MSM found only one in ten uncircumcised men were willing to participate in future research on circumcision for HIV prevention.[12] The aim of our study was to describe sexual practices by circumcision status, and to explore the feasibility of conducting research on male circumcision for HIV prevention among MSM in Scotland.

## **METHODS**

Circumcision questions were included in the 2008 MRC Gay Men's Survey, which collected anonymous, self-complete questionnaires and (Orasure™) oral fluid specimens (screened for anti-HIV using an enzyme immunoassay; positives re-screened, and repeat reactives confirmed using Western Blot). Time and location sampling was used to recruit representative samples in the commercial gay scenes of Glasgow and Edinburgh.[13] Data were analysed with SPSS 15.0. Logistic regression was used to estimate odds ratios and 95% confidence intervals (CI). Odds ratios were adjusted for age and nationality, which were significantly associated with circumcision status. Ethical approval was granted by the University of Glasgow, Faculty of Medicine Ethics Committee.

## **RESULTS**

2138 men were approached, 1508 completed questionnaires (70.5% response rate) and 1277 provided oral fluid samples (59.7% response rate). In multivariate analysis comparing men who did and did not provide samples, only age was significant (reduced odds among men aged 26+ years, AOR=0.63, 95% CI 0.43-0.92). 54 men who reported they were heterosexual and had no sexual contact with men in the previous 12 months and 49 men who did not provide their circumcision status are excluded (N=1405).

233 men (16.6%) reported having been circumcised; 32.3% had been circumcised before they were one year old and, overall, 73.1% by age 18. Compared with the 16-25 years age group, men aged 36-45 and 46+ were more likely to be circumcised (21.4% and 18.2% respectively, compared with 11.8%;  $p=0.007$ ). Nationals from non-European

countries (mainly America, Canada, Australia and New Zealand) were more likely to be circumcised than Scottish men (50.0% and 13.1% respectively,  $p < 0.001$ ).

HIV prevalence was comparable among uncircumcised and circumcised men (4.6% and 4.2% respectively; Table 1). Similarly, there was little difference in prevalence of other, self-reported, STIs in the past 12 months (though it should be noted that STI transmission may also occur by means other than anal intercourse).

Overall, 39.9% of men reported any UAI in the past 12 months and this was similar by circumcision status (Table 1). Of these, 22.4% reported exclusive UIAI, and circumcised men were more likely to report this than uncircumcised men (Table 1). Among men who reported always being the insertive partner, none of the circumcised men tested HIV-positive (0/27), compared with 2.5% (2/79) of uncircumcised men.

Few men (13.9%) reported being willing to take part in an HIV prevention research study on circumcision, while 20.1% reported that they did not know if they would be willing to do so (Table 1). Only 11.3% of uncircumcised men reported being willing to take part.

## **DISCUSSION**

This is the first study to assess the association between circumcision and HIV among MSM in Scotland, though some study limitations should be noted. The data are cross-sectional and causality cannot be inferred. The oral fluid sample response rate was relatively low, and may not be representative. However, the overall survey response rate

was higher (71%), and only age differed between the men who did and did not provide oral fluid specimens; suggesting that the men who provided these were representative of the larger venue-based sample. Finally, circumcision status was self-reported, though this has been shown to be a valid measure.[14]

We found no evidence of an association between circumcision and HIV or other self-reported STIs among MSM in Scotland, similar to findings elsewhere.[3-5] Among men reporting UAI, less than one quarter exclusively favoured the insertive role, which might plausibly provide partial protection against HIV infection. In contrast to a recent US study,[15] this behaviour was more common among circumcised than uncircumcised men; and none of the circumcised men who were exclusively the insertive partner tested HIV-positive. Although findings are inconsistent across studies, a possibly protective effect among this group has been reported,[4] and merits further investigation.

There are few studies of the willingness of uncircumcised MSM to be circumcised.[4] One, from the US, found that 53% were willing to be circumcised.[16] However, our findings are similar to those of the London study, in which only one in ten uncircumcised men were willing to participate in circumcision research.[12] Together with the low levels of exclusive UIAI in these populations, these findings suggest that a RCT of male circumcision for HIV prevention is unlikely to be feasible in the UK. Although circumcision may be partially protective against HIV infection for those men who are exclusively insertive partners, our results suggest that comprehensive HIV prevention

strategies, including the promotion of consistent condom use, should continue to be the focus of prevention efforts among MSM.



## **KEY MESSAGES**

- There was no evidence of an association between circumcision and HIV or other self-reported STIs among MSM in Scotland.
- Among men reporting unprotected anal intercourse, less than one quarter exclusively favoured the insertive role.
- With only 11% of uncircumcised men reporting willingness to take part in a HIV prevention research study on male circumcision, a randomised controlled trial of male circumcision for HIV prevention is unlikely to be feasible in this population.

## **ACKNOWLEDGEMENTS**

We thank the survey staff and fieldworkers in each city, the venue managers, their staff, and the men who agreed to participate in the survey.

## **COMPETING INTERESTS**

None.

## **FUNDING**

The UK Medical Research Council funds LMCD and the Gay Men's Survey as part of the Sexual and Reproductive Health Programme (WBS U.1300.00.005) at the Social and Public Health Sciences Unit, and funds HW as part of the MRC Tropical Epidemiology Group.

## **AUTHOR CONTRIBUTIONS**

LMCD, HW and GH devised the paper. LMCD conducted the analyses and wrote the first draft. All authors contributed to subsequent drafts and approved the final version of the manuscript.

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in STI and any other BMJPGGL products to

exploit all subsidiary rights, as set out in our licence  
<http://sextrans.bmj.com/ifora/licence.pdf>."

## REFERENCES

1. **Weiss HA**, Halperin D, Bailey RC, et al. Male circumcision for HIV prevention: from evidence to action? *AIDS* 2008;**22**:567-574.
2. **WHO/UNAIDS**. *New data on male circumcision and HIV prevention: policy and programme implications*. Geneva: WHO, 2007. Available at [http://www.who.int/hiv/pub/malecircumcision/research\\_implications/en/index.html](http://www.who.int/hiv/pub/malecircumcision/research_implications/en/index.html). [Accessed 21 December 2009]
3. **Millett GA**, Flores SA, Marks G, et al. Circumcision status and risk of HIV and sexually transmitted infections among men who have sex with men. *JAMA* 2008;**300**:1674-1684.
4. **Templeton DJ**, Millett GA, Grulich AE. Male circumcision to reduce the risk of HIV and sexually transmitted infections among men who have sex with men. *Curr Opin Infect Dis* 2010;**23**:45-52.
5. **Gust DA**, Wiegand RE, Kretsinger K, et al. Circumcision status and HIV infection among MSM: reanalysis of a Phase III HIV vaccine clinical trial. *AIDS* 2010; Epub ahead of print, doi:10.1097/QAD.0b013e328337b8bd.
6. **Donoval BA**, Landay AL, Moses S, et al. HIV-1 target cells in foreskins of African men with varying histories of sexually transmitted infections. *Am J Clin Pathol* 2006;**125**:386-391.
7. **McCoombe SG**, Short RV. Potential HIV-1 target cells in the human penis. *AIDS* 2006;**20**:1491-1495.

8. **Patterson BK**, Landay A, Siegel JN, et al. Susceptibility to human immunodeficiency virus-1 infection of human foreskin and cervical tissue grown in explant culture. *Am J Pathol* 2002;**161**:867-873.
9. **Szabo R**, Short RV. How does male circumcision protect against HIV infection? *BMJ* 2000;**320**:1592-1594.
10. **Health Protection Agency**. *Sexually transmitted infections and men who have sex with men in the UK: 2008 report*. London: Health Protection Agency, 2008.
11. **Dave SS**, Johnson AM, Fenton KA, et al. Male circumcision in Britain: findings from a national probability sample survey. *Sex Transm Infect* 2003;**79**:499-500.
12. **Thornton A**, Lattimore S, Delpech V, et al. Circumcision among men who have sex with men in London [Abstract No. P4.62]. *18th Biennial Meeting of the International Society for STD Research*. London, UK, 2009.
13. **Williamson LM**, Hart GJ. HIV prevalence and undiagnosed infection among a community sample of gay men in Scotland. *J Acquir Immune Defic Syndr* 2007;**45**:224-230.
14. **Templeton DJ**, Mao L, Prestage GP, et al. Self-report is a valid measure of circumcision status in homosexual men. *Sex Transm Infect* 2008;**84**:187-188.
15. **Jameson DR**, Celum CL, Manhart L, et al. The association between lack of circumcision and HIV, HSV-2, and other sexually transmitted infections among men who have sex with men. *Sex Transm Dis* Epub ahead of print, doi: 10.1097/OLQ.1090b1013e3181bd1090ff1090.

16. **Begley EB**, Jafa K, Voetsch AC, et al. Willingness of men who have sex with men (MSM) in the United States to be circumcised as adults to reduce the risk of HIV infection. *PLoS ONE* 2008;**3**:e2731.

**Table 1: HIV status, sexual behaviour, and willingness to take part in a research study on circumcision and HIV prevention: comparing circumcised and uncircumcised men**

	Circumcision Status						Adjusted OR (95% CI) among circumcised men*
	Uncircumcised (N=1172)		Circumcised (N=233)		Total (N=1405)		
	n	%	n	%	n	%	
<b><u>HIV and other sexually transmitted infections (STIs)</u></b>							
<b>HIV status (oral fluid specimen)†</b>							
HIV-negative	951	95.4	184	95.8	1135	95.5	1
HIV- positive	46	4.6	8	4.2	54	4.5	0.78 (0.35-1.73)
<b>Self-reported STI in the past 12 months</b>							
No	1052	90.8	211	91.7	1263	91.0	1
Yes	106	9.2	19	8.3	125	9.0	0.95 (0.56-1.61)
<b><u>Sexual Behaviour</u></b>							
<b>Any unprotected anal intercourse (UAI) partners in the past 12 months</b>							
No	700	60.8	130	56.3	830	60.1	1
Yes	451	39.2	101	43.7	552	39.9	1.25 (0.93-1.69)
<b>UAI with more than one partner in the past 12 months</b>							
0/1 partner	1006	87.4	202	87.4	1208	87.4	1
2 or more partners	145	12.6	29	12.6	174	12.6	1.07 (0.68-1.66)
<b>UAI with partners of unknown or discordant HIV status in the past 12 months</b>							
No	861	74.8	174	75.3	1035	74.9	1
Yes	290	25.2	57	24.7	347	25.1	1.06 (0.75-1.49)

### Sexual position during UAI in the past

#### 12 months†

Always insertive	91	20.5	31	31.0	122	22.4	1.67 (1.01-2.76)¶
Mostly insertive	38	8.6	11	11.0	49	9.0	
Equally both	203	45.7	36	36.0	239	43.9	1
Mostly receptive	73	16.4	19	19.0	92	16.9	
Always receptive	39	8.8	3	3.0	42	7.7	

#### Research on circumcision

#### Willingness to take part in a research study on circumcision and HIV prevention§

No	758	68.8	94	50.0	852	66.0	1
Don't know	220	20.0	39	20.7	259	20.1	
Yes	124	11.3	55	29.3	179	13.9	3.36 (2.29-4.92)

\* Adjusted for age and nationality.

† Among men who provided oral fluid specimens (N=1189).

‡ Among men reporting any UAI (N=552).

¶ Adjusted odds ratio for men reporting always being the insertive UAI partner in the past 12 months; for men reporting always or mostly being the insertive UAI partner in the past 12 months AOR=1.82, 95%CI 1.14-2.92.

§ The willingness question used was 'We are looking for new ways to prevent HIV. Should the following research studies take place, which would you be willing to take part in?', with participants asked to select 'yes', 'no' or 'don't know' for circumcision (and also for behaviour change, rectal microbicides and HIV vaccines).