



HAL
open science

Asymptomatic immunoglobulin light chain amyloidosis (AL) at the time of diagnostic bone marrow biopsy in newly diagnosed patients with multiple myeloma and smoldering myeloma. A series of 144 cases and a review of the literature

Sergio Siragusa, William Morice, Morie A. Gertz, Robert A. Kyle, Philip R. Greipp, John A. Lust, Thomas E. Witzig, Martha Q. Lacy, Steven R. Zeldenrust, S. Vincent Rajkumar, et al.

► **To cite this version:**

Sergio Siragusa, William Morice, Morie A. Gertz, Robert A. Kyle, Philip R. Greipp, et al.. Asymptomatic immunoglobulin light chain amyloidosis (AL) at the time of diagnostic bone marrow biopsy in newly diagnosed patients with multiple myeloma and smoldering myeloma. A series of 144 cases and a review of the literature. *Annals of Hematology*, 2010, 90 (1), pp.101-106. 10.1007/s00277-010-1028-8 . hal-00557757

HAL Id: hal-00557757

<https://hal.science/hal-00557757>

Submitted on 20 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Editorial Manager(tm) for Annals of Hematology
Manuscript Draft

Manuscript Number: AOHE-D-10-00270

Title: Asymptomatic immunoglobulin light chain amyloidosis (AL) at the time of diagnostic bone marrow biopsy in newly diagnosed patients with multiple myeloma and smoldering myeloma. A series of 144 cases and a review of the literature

Article Type: Original Article

Keywords: Amyloid, Multiple Myeloma, Smoldering Myeloma, Asymptomatic, Bone Marrow Biopsy

Corresponding Author: Prof. Sergio Siragusa, MD

Corresponding Author's Institution: University Hospital of Palermo

First Author: Sergio Siragusa, MD

Order of Authors: Sergio Siragusa, MD; William Morice; Morie A Gertz; Robert A Kyle; Philipp R Greipp; John Lust; Thomas E Witzig; Martha Q Lacy; Steven R Zeldenrust; Vincent S Rajkumar; Stephen J Russel; Suzanne R Hayman; Francis Baudi; Shaji K Kumar; David Dingli; Angela Dispenzieri

Abstract: The rate of asymptomatic amyloidosis (AL) among patients with newly diagnosed multiple myeloma (MM) or smoldering multiple myeloma (SMM) is unknown. We evaluated number and clinical significance of asymptomatic AL in consecutive MM and SMM patients, not having recognition of symptomatic AL at the time of their diagnostic bone marrow biopsy. Bone marrow biopsies were stained with Congo Red and considered diagnostic for AL in case of positive Congo Red staining with apple green birefringence. Biopsies from 144 patients were evaluated: 77 had a diagnosis of MM and 67 of SMM. The median age was 59 (range 26-84) years; the median follow-up was of 76 months (range 0-216). Immunoglobulin isotypes were: 96/144 (67%), IgG; 23/144 (16%), IgA; 12/144 (8%), light chain only; 1/77 (1%) IgD; and biclonal or indeterminate, 12/144 (8%). Fifty-eight percent (84/144) were κ restricted. The presence of amyloid was found in 2 cases (1%, 95% CI - 0.6 to 3.2), 1 in MM and 1 in SMM group and none had or developed signs or symptoms suggestive of organ involvement by amyloid. Among the 142 other patients without amyloid deposition in their index bone marrow, 1 (0.7%, 95% CI -0.6 to 2.0) developed symptomatic AL after 119 months.

1
2
3
4 **Asymptomatic immunoglobulin light chain amyloidosis (AL) at the time of**
5
6
7 **diagnostic bone marrow biopsy in newly diagnosed patients with multiple**
8
9
10 **myeloma and smoldering myeloma. A series of 144 cases and a review of the**
11
12 **literature**
13
14
15
16
17

18 Running title: Asymptomatic Amyloidosis in Multiple Myeloma
19
20
21
22

23 ¹Sergio Siragusa M.D., ²William Morice, M.D., ³Morie A Gertz, M.D., ³Robert A. Kyle, ³Philip
24 R. Greipp, M.D., ³John A. Lust, M.D., Ph.D., ³Thomas E. Witzig, M.D., ³Martha Q. Lacy, M.D.,
25
26 ³Steven R. Zeldenrust, M.D., Ph.D., ³S. Vincent Rajkumar, M.D, ³Stephen J. Russell, M.D.,
27
28 Ph.D., ³Suzanne R. Hayman, M.D., ³Francis Buadi, M.D., ³Shaji K. Kumar, M.D., M.D., ³David
29
30 Dingli, M.D., ³Angela Dispenzieri
31
32
33
34
35
36
37

38 ¹Cattedra ed U.O. di Ematologia con trapianto, Azienda Ospedaliera Universitaria Policlinico di
39
40 Palermo, Palermo, Italy
41

42 ²Division of Pathology, Mayo Clinic Rochester, Rochester, MN, USA
43
44

45 ³Division of Hematology, Mayo Clinic Rochester, Rochester, MN, USA
46
47
48
49

50 Correspondence to:

51 Sergio Siragusa, MD
52

53 Cattedra ed U.O. di Ematologia con trapianto,
54
55

56 Azienda Ospedaliera Universitaria Policlinico,
57
58

59 via del Vespro 127
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

90127 Palermo, Italy

Ph: + 39 091 655 4419

Fax: + 39 091 655 4402

e-mail: sergio.siragusa@unipa.it

Abstract words: 200

Text words: 2537

1
2
3
4 **SUMMARY**
5
6
7
8

9 The rate of asymptomatic amyloidosis (AL) among patients with newly diagnosed multiple
10 myeloma (MM) or smoldering multiple myeloma (SMM) is unknown. We evaluated number and
11 clinical significance of asymptomatic AL in consecutive MM and SMM patients, not having
12 recognition of symptomatic AL at the time of their diagnostic bone marrow biopsy. Bone
13 marrow biopsies were stained with Congo Red and considered diagnostic for AL in case of
14 positive Congo Red staining with apple green birefringence. Biopsies from 144 patients were
15 evaluated: 77 had a diagnosis of MM and 67 of SMM. The median age was 59 (range 26-84)
16 years; the median follow-up was of 76 months (range 0-216). Immunoglobulin isotypes were:
17 96/144 (67%), IgG; 23/144 (16%), IgA; 12/144 (8%), light chain only; 1/77 (1%) IgD; and
18 biconal or indeterminate, 12/144 (8%). Fifty-eight percent (84/144) were κ restricted. The
19 presence of amyloid was found in 2 cases (1%, 95% CI – 0.6 to 3.2), 1 in MM and 1 in SMM
20 group and none had or developed signs or symptoms suggestive of organ involvement by
21 amyloid. Among the 142 other patients without amyloid deposition in their index bone marrow,
22 1 (0.7%, 95% CI -0.6 to 2.0) developed symptomatic AL after 119 months.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Key words:** Amyloid, Multiple Myeloma, Smoldering Myeloma, Asymptomatic, Bone Marrow
46 Biopsy
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Multiple myeloma (MM), smoldering multiple myeloma (SMM), immunoglobulin light chain amyloidosis (AL) and monoclonal gammopathy of undetermined significance (MGUS) represent a spectrum of plasma cells disorders (PCDs). AL is a rare and serious disorder characterized by the deposition of amyloid fibrils in different tissues with more than 80% of patients dead within 2 years of diagnosis (1, 2)

Immunoglobulin light chain amyloidosis may coexist with any of the other PCDs (2, 3). It has been reported that approximately 10% of patients with AL may have MM at the time of diagnosis (2), while only a minority will developed delayed MM (4). Conversely, it has been reported that up to 30% of MM patients may have subclinical amyloid deposits (in subcutaneous fat aspirate, bone marrow biopsy or biopsies of other vital organs) (5, 6, 7, 8) and in up to 15% of all MM patients a clinically overt AL appear during the course of index MM (3, 5)).

The rate of asymptomatic amyloidosis (asympt-amyloid) in patients with MM or SMM as well as its clinical impact, is uncertain. This topic is significant because unrecognized AL can be associated with significant mortality and morbidity. In the case of patients with SMM, no treatment will be directed against the plasma cell clone, potentially allowing for unbridled progression of the AL. Unrecognized AL in patients diagnosed with MM is also significant because of the potential for higher rates of treatment related morbidity and mortality in patients with occult visceral involvement with AL. However, at the present we deal with discordant information. Some authors found out MM-associated asympt- amyloid in up to 26% of patients at the time of diagnosis (8); this was also an independent high-risk prognostic factor for MM patients, thus suggesting need for searching amyloid and need for aggressive treatment. Others

1
2
3
4 plasmacytosis, consistent with co-existing multiple myeloma. Of these, only 18 (46.1% of
5
6 patients with > 10% of BM plasma cells) or 2.6% of the entire AL population had symptoms
7
8 consistent with myeloma, i.e. bone lesions, anemia (Hb <11 g/dL), calcium greater than 10.5
9
10 mg/dL or myeloma related kidney disease, which was defined as a creatinine greater than 1.5
11
12 mg/dL and a 24-hour urinary M-protein to 24-hour urinary total protein ratio of greater than 0.5.
13
14
15
16
17
18

19 **Diagnosis of asymptomatic AL**

20
21 Bone marrow biopsies were reviewed and examined for Congo red staining. A patient
22
23 was considered to have amyloid if the stain was positive and apple green birefringence was
24
25 found on polarized microscopy. Posterior superior iliac crest was the preferred biopsy site and
26
27 the specimens were obtained with a modified Jamshidi needle as an outpatient procedure.
28
29 Following decalcification the bone marrows were paraffin embedded and H&E stained and
30
31 Congo red stained sections were reviewed by a single author (WB). Amyloid deposits were
32
33 detected by examination at 200x under polarized light. Only data from patients with an adequate
34
35 follow-up period were considered in the final analysis. In addition, serum samples were obtained
36
37 at diagnosis to determine LDH, β 2 microglobulin, albumin, calcium, phosphorus, creatinine, and
38
39 liver function tests. Serum and urine electrophoresis and immunofixation, as well as bone
40
41 marrow aspiration and bone X-ray studies were taken at diagnosis.
42
43
44
45
46
47
48
49

50 **Statistical analysis**

51
52
53 Baseline differences between groups were assessed by the chi-square test (Fisher's Exact
54
55 Test) for categorical variables and univariate analysis of variance (ANOVA test) or Mann-
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Whitney U statistic test for parametric and nonparametric analyses. Survival was estimated using
5
6 the method of Kaplan-and Meier.
7
8
9

10 11 **RESULTS** 12 13 14

15 16 **Overall characteristics of MM and SMM patients** 17

18
19 Of the 1180 patients with a diagnosis of MM (959) or SMM (221) seen at Mayo Clinic
20
21 within 30 days of their diagnosis, biopsies were available from 144 (F 59, M 84). Among
22
23 patients evaluated 77 had a diagnosis of MM and 67 of SM. Characteristics of MM and SM
24
25 patients at time of diagnosis are reported in Table 1. One hundred-thirty-six (95.8%) patients had
26
27 a positive serum M-spike before diagnostic BM. Median serum M-spike at the time of diagnostic
28
29 BM biopsy was 2.93 (range 0.6-9.5). At a median follow-up of living patients of 76.2 months
30
31 (range 0-215.9), 46 (31.9%) patients were alive, 93 (64.5%) dead and 5 (3.4%) lost to follow-up.
32
33
34
35
36 The overall median survival was 70.4 months (figure 1).
37

38
39 At presentation, SMM patients were older, less anemic, had higher creatinine, urine
40
41 protein, but lower urine M-spikes levels than MM patients. These last patients were more likely
42
43 to have fatigue and pain, while SMM patients had more frequently paresthesia (Table 1).
44
45
46
47
48
49
50
51
52

53 **Patients with asymptomatic amyloid deposit at presentation** 54

55
56 Amyloid was retrospectively detected in the index bone marrow biopsy of 2 of the 144
57
58 cases (1.38%, 95% CI – 0.6 to 3.2), 1 in MM and 1 in SMM group, respectively. The first (OE,
59
60
61
62
63
64
65

1
2
3
4 female, 66 y.o.) was diagnosed having MM; at presentation, she had moderate anemia (Hg 9.4
5
6 g/l), pain, multiple fractures but not hypercalcemia, renal or liver tests alteration. No measurable
7
8 M-spike was identified on electrophoresis, but there was a monoclonal kappa light chain
9
10 detectable in the serum by immunofixation. Urine protein electrophoresis showed a total urine
11
12 protein of 280 mg with an M-spike of 249 mg/24h that was confirmed to be kappa light-chain.
13
14 Quantitative immunoglobulins were not done. The bone marrow biopsy revealed plasma-cells
15
16 comprising 30% of the total cellularity. Beta-2-microglobulin was elevated at 2.28 mcg/mL.
17
18 Cytogenetic studies by FISH did not show IGH translocation, monosomy or deletion 13, deletion
19
20 of p53 or trisomy 3, 7, 9, 11, 13, 15 or 17. Cytogenetics studies of BM showed a normal
21
22 karyotype (46, XX). The second patient (RW, male, 77 y.o.) was diagnosed with SMM (IgG-κ).
23
24 He had a history of weight loss and severe chronic obstructive pulmonary disease. At
25
26 presentation, physical exam did not reveal any abnormalities. The BM biopsy revealed plasma-
27
28 cells up to 25% of the total cellularity; β2-microglobulin was 1.52 mcg /mL. Bone survey was
29
30 negative for metastases or myeloma. Clinical and laboratory characteristics are listed in Table 2.
31
32 After more than seven years of follow-up, neither of these patients had or developed during their
33
34 disease signs or symptoms suggestive for organ involvement by amyloid.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **Patients without bone marrow amyloid deposition at presentation but subsequent systemic** 51 **amyloidosis** 52

53
54
55 There was one patient (R.J., male, 51 y.o.) who was diagnosed having MGUS (IgG-λ) in
56
57 1992 followed by SMM in 1993. Clinical and laboratory characteristics at presentation are listed
58
59
60
61
62
63
64
65

1
2
3
4 in Table 2. At the presentation physical examination was normal. He did not have weakness,
5
6 fatigue, bleeding or increased bruising, not bone pain, change in his voice or tongue, steatorrhea,
7
8 paresthesia, claudication or weight loss. He developed symptomatic AL after 72.6 months
9
10 (February 1999) from initial diagnosis of SMM. He was treated with melphalan/prednisone. At
11
12 the time of last contact in 2002, he considered himself to be in good health.
13
14
15
16
17
18

19 **DISCUSSION**

20
21
22
23 Rate and clinical impact of asymp-amyloid in patients with MM or SMM is uncertain.
24
25 While asymptomatic amyloid deposits have been reported to coexist at the time of MM diagnosis
26
27 in up to 30% of patients (1, 5), less information is available in patients with established MM or
28
29 SMM who are seen at a facility at which an extensive amyloid directed review of systems and
30
31 signs is the norm. This issue is topical since unrecognized AL can be associated with significant
32
33 mortality and morbidity and its early detection could change the management of MM and SMM
34
35 patients.
36
37
38
39

40
41 We found that only 1.4% of patients with retrospective Congo Red staining of their bone
42
43 marrow had amyloid present on their presenting bone marrow. This may be a limitation of the
44
45 present investigation, since it is well known that the sensitivity of bone marrow biopsy is only
46
47 about 50% in comparison to that of abdominal fat aspirate with a sensitivity of about 80% of
48
49 confirmed amyloidosis (1, 14). It is possible that our approach yielded a low estimate. For
50
51 example, the one patient, whose initial bone marrow was negative for amyloidosis but who was
52
53 diagnosed with AL 6 years later, might have had a positive fat aspirate at the time of his earlier
54
55 bone marrow; if so, he would have had very indolent amyloidosis.
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Our data, which may appear to conflict with prior reports, would suggest that a careful
5 amyloid specific review at the time of diagnosis of SMM or MM misses only 1-2% of cases even
6
7 with a 6.4 person-years of follow-up and therefore abrogates the need to do routine amyloid
8
9 staining on all cases of SMM or MM. It also supports the Mayo Clinic practice of having a low
10
11 threshold for amyloidosis screening if non-specific symptoms are present in a patient with a
12
13 monoclonal gammopathy.
14
15
16
17

18
19 Table 3 puts our findings into the context of prior reports. One of the first reports
20
21 addressing this question was by Desikan *et al* (6), who prospectively evaluated fat pad aspirate
22
23 and BM biopsy for amyloid deposit in a cohort of 84 MM patients. Amyloid was detected in
24
25 38% of patients; fat pad aspirate was positive in 25% and BM in 10% of cases and other organ
26
27 sites in 7 patients. MM or SMM patients did not differ in disease characteristics and, particularly,
28
29 no λ predominance was found. The median overall and event free survival of the entire cohort is
30
31 66+ and 52 months respectively; the median overall (59+ months vs 66+ months, $p=0.9$) and
32
33 event free survival (59+ months vs 52+ months, $p=0.9$) were similar in cohort with and without
34
35 amyloid respectively. These authors concluded therefore that, although the incidence of light-
36
37 chain asymptomatic AL in MM patients is high, it does not influence the outcome of these
38
39 patients treated with autologous transplantation.
40
41
42
43
44

45
46 Vela-Ojeda *et al* evaluated 201 consecutive patients with *de novo* MM in whom a fat-pad
47
48 biopsy needle aspiration (stained with Congo red) was performed (8). Overall, 68 patients (34%)
49
50 were diagnosed having concomitant AL; 16 (8%) had symptoms consistent with amyloidosis at
51
52 the time of MM diagnosis while in 52 (25.8%) fat pad aspiration was useful to detect incidental
53
54 asymptomatic AL. The median follow-up was 36 months (range 12-199). Among patients
55
56 without AL at the time of diagnostic BM for MM, 69% of them became symptomatic for AL
57
58
59
60
61
62
63
64
65

1
2
3
4 after a median follow-up of 14 months (range 6 to 30). Even excluding 16 patients with obvious
5
6 symptoms of AL at diagnosis, overall survival was worse in patients who developed late AL (13
7
8 vs 64 months of those without AL). Cox regression model for overall survival detected three
9
10 variables having independent prognostic significance: the presence of AL (RR 3.4, P<0.004),
11
12 serum albumin levels <3.5 g/dl (RR 3.2, p<0.005) and patients not achieving complete remission
13
14 or very good partial remission (RR 2.9, p<0.02). The authors then concluded that MM-associated
15
16 AL represents a poorer prognosis disease even in the absence of symptoms at diagnosis and this
17
18 specific association may be considered as an independent high-risk prognostic factor. Thus,
19
20 Authors concluded that routine study of peri-umbilical fat-pad tissue should be mandatory in all
21
22 patients with MM.
23
24
25
26
27

28
29 Petruzziello *et al* (2009) evaluated retrospectively the bone marrow smears of 166
30
31 unselected patients affected by MM (126 at diagnosis and 40 after treatment); no data were
32
33 reported about the presence of signs or symptoms suggestive for concomitant systemic AL (12).
34
35 All samples were stained with Congo red and studied by transmission and birefringence
36
37 microscopy. Both focal and diffuse involvement were considered positive. Overall, 67 patients
38
39 were positive for AL and 99 were negative to Congo red and apple-green birefringence. In
40
41 particular, 51 of the 126 patients studied at diagnosis and 16 of the 40 patients with advanced
42
43 disease were positive. Authors failed to find correlations between bone marrow amyloid deposits
44
45 and immunoglobulin type, disease stage, plasma cells percentage, hemoglobin, calcium,
46
47 creatinine, albumin, or β 2-microglobulin.
48
49
50
51
52

53
54 Although the main objective of our investigation was that of evaluate the rate of asym-
55
56 amyloid at the time of first BM, we were able to look at the clinical outcome of our cohort who
57
58 had a median follow up of 6 years. Neither of the patients who were retrospectively found to
59
60
61
62
63
64
65

1
2
3
4 have amyloid in their index bone marrows developed symptomatic AL, and their survival did not
5
6 differ from the other patients. Among Congo Red negative patients, we detected one (SMM
7
8 patient) who developed symptomatic AL during a follow up of 72.6 months. He had serum λ
9
10 light-chain, an isotype frequently reported in patients with AL (13).
11
12

13
14 One potential caveat in studies looking at the detection of asymp-amyloid relies on
15
16 patients' selection. In fact, AL requires a high index of suspicion and most of AL cases are often
17
18 misdiagnosed with other PCDs, mainly with MM (The UK Myeloma Forum AL Amyloidosis
19
20 Guidelines Working Group, 2004). It is not unlikely that the true occurrence of asymptomatic
21
22 AL has been previously overestimated. Because of the high level of suspicion for amyloid at our
23
24 Amyloid Center, we are confident that all patients evaluated did not have (even sub-clinical) AL
25
26 at the time of diagnostic BM for MM or SM. Therefore, evaluation of adequate specimens in
27
28 experienced laboratories is necessary to maintain high diagnostic sensitivity and specificity.
29
30
31

32
33 In conclusion, we evaluated rates and clinical significance of amyloid deposit at the time
34
35 of first diagnostic BM biopsy in newly diagnosed MM and SMM. Our findings show that fewer
36
37 than 2% of patients in both groups have asymp-amyloid thus discouraging any specific search for
38
39 amyloid deposit when careful evaluation of clinical presentation and patient's history are not
40
41 suggestive for systemic AL.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **REFERENCES**
5
6
7
8

- 9 1. Gertz MA, Lacy MQ, Dispenzieri A. (1999) Amyloidosis. *Hematol Oncol Clin North Am*
10 13:1211–1233
11
12
13
14
15
16 2. Kyle RA, Gertz MA. (1995) Primary systemic amyloidosis: clinical and laboratory features in
17 474 cases. *Semin Hematol* 32:45–59.
18
19
20
21
22
23 3. Rajkumar SV, Dispenzieri A, Kyle RA. (2006) Monoclonal gammopathy of undetermined
24 significance, Waldenström macroglobulinemia, AL amyloidosis, and related plasma cell
25 disorders: diagnosis and treatment. *Mayo Clin Proc* 81(5):693-703
26
27
28
29
30
31
32
33 4. Rajkumar SV, Gertz MA, Kyle RA. (1998) Primary systemic amyloidosis with delayed
34 progression to multiple myeloma, *Cancer* 82:1501–1505
35
36
37
38
39
40 5. Kyle RA. Multiple Myeloma and other plasma cell disorders. (1995) In: Hoffman R, Benz EJ,
41 Shattil SJ, Furie B Cohen HJ, Silberstein LE, editors. *Hematology, basic principles and practice.*
42 2nd edition. New York: Churchill Livingstone 1354-74
43
44
45
46
47
48
49
50 6. Desikan KR, Dhodapkar MV, Hough A, et al. (1997) Incidence and impact of light chain
51 associated (AL) amyloidosis on the prognosis of patients with multiple myeloma treated with
52 autologous transplantation. *Leuk Lymphoma* 27: 315–319
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 7. Bahlis NJ, Lazarus HM. (2006) Multiple myeloma-associated AL amyloidosis: is a distinctive
5
6 therapeutic approach warranted? Bone Marrow Transplant 38(1):7-15
7
8
9
- 10
11 8. Vela-Ojeda J, García-Ruiz Esparza MA, Padilla-González Y, et al. (2009) Multiple myeloma-
12
13 associated amyloidosis is an independent high-risk prognostic factor. Ann Hematol 88(1):59-66.
14
15
16
17
18
- 19 9. The UK Myeloma Forum AL Amyloidosis Guidelines Working Group. Guidelines on the
20
21 diagnosis and management of AL amyloidosis. (2004) Br J Haematol 125:681–700
22
23
24
25
- 26 10. Kyle RA, Remstein ED, Therneau TM, et al. (2007) Clinical course and prognosis of
27
28 smoldering (asymptomatic) multiple myeloma. N Engl J Med 356(25):2582-90
29
30
31
32
- 33 11. Duston, M.A., Skinner, M., Shraham, T., et al (1987) Diagnosis of amyloidosis by abdominal
34
35 fat aspiration: analysis of 4 years experience. Am J Med 82:412–414
36
37
38
39
- 40 12. Petruzzello F, Zeppa P, Catalano L, et al. (2010) Amyloid in bone marrow smears of
41
42 patients affected by multiple myeloma. Ann Hematol 89(5):469-74
43
44
45
46
47
- 48 13. Solomon A, Weiss DT, Kattine AA. Et al. (1991) Nephrotoxic potential of Bence Jones
49
50 proteins. N Engl J Med 324(26):1845-51
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Baseline patients' characteristics

Characteristics	MM population (n=77)	SM population (n=67)
Median months from biopsy to last follow up (range)	65 (10.4-169.2)	64.6 (0-210)
Such Fatigue, %	13 (16.8)	5 (7.4)
Pain, %	42 (54.5)	1 (1.5)
Paresthesia, %	1 (1.3)	3 (4.4)
Bleeding, %	0	1 (1.5)
Edema, %	0	1 (1.5)
Dyspnea, %	6 (7.8)	1 (1.5)
Median age, years (range)	55.5 (33-70)	63 (26-84)
Hemoglobin (g/dL), median (range)	11.0 (7.6-15.3)	12.5 (7.6-14.8)
Platelets (x 10 ⁹ /L), median (range)	222.0 (8-399)	224.0 (65-786)
Serum protein (g/dL), median (range)	8.0 (6-16.1)	8.4 (6.3-11.8)
Urine protein (grams/24h)	0.13 (0.01-10.7)	0.1 (0.01-2.7)
Serum Albumin, median (range)	3.64 (2.3-5.7)	3.60 (2.3-4.7)
Alkaline Phosphate (U/L), median (range)	175.5 (53-1199)	155.0 (62-480)
Creatinine (mg/dL), median (range)	1.1 (0.7-2.2)	1.1 (0.9-6.6)

BM: Bone Marrow; FU: Follow-up

Table 2. Characteristics of two patients with detection of amyloid deposit at presentation and one patient without amyloid deposit but subsequent systemic amyloidosis

Clinical/laboratory characteristics at diagnosis	Patients with amyloid deposit at presentation		Patient without amyloid deposit at presentation but subsequent AL
	MM patient	SM patient	
Months from biopsy to last follow-up	86.4	87.6	72.6
Age (y)	66	77	51
Hg (g/dl)	9.4	13	12.3
Platelets (n. x 10 ⁹ /l)	328	201	446
Pro-thrombin time (sec.)	10.1	NA	NA
Serum protein (g/dl)	NA	7.4	6.9
Urine protein (mg/dl)	280	8	13
Serum Albumin (g/dl)	2.9	3.6	3.5
Alkaline Phosphate (U/l)	NA	145	152
Creatinine (mg/100 ml)	0.8	1.1	0.9
B2-microglobulin (µg/ml)	2.28	1.56	1.5
C-reactive protein level (mg/dl)	NA	NA	0.59
Serum light-chain isotype	kappa	kappa	lambda

NA: Not Available

Table 3. Studies considering relationship between asymptomatic AL in myeloma patients

MM cases with incidental amyloidosis				
Author^{ref}	MM cases, n	AL cases, n (%)	Tissues stained for amyloid	Prognostic factor?
Desikan ⁶	84	32 (30.8)	Subcutaneous abdominal fat aspirate (SAFA) in 25, bone marrow in 8 and other organ sites in 7 patients	No
Vela-Ojeda ⁸	201	68 (34)*	SAFA	Yes
Petruziello ¹²	166	67 (40.3)	Cytological techniques in bone marrow smears	No
Present study MM	77	1 (1.3)	Bone marrow	No
Present study SMM	67	1 (1.5)	Bone marrow	No
AL cases with asymptomatic MM				
	AL cases,	MM case,		Comments

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

	n	n (%)		
Kyle ²	474			
AL cases with symptomatic MM				
	AL cases, n	MM case, n (%)		Comments
Rajkumar ⁴	1596	6 (0.3)	Biopsy-proven AL	No laboratory or clinical evidence of MM at the diagnosis of AL. MM developed 10-81 months thereafter

*16 were symptomatic for AL

***Conflict of interest**

[Click here to download Conflict of interest: Conflict of Interest Disclosure Form.doc](#)