

Christophe Chesneau

# ► To cite this version:

Christophe Chesneau. Adaptive wavelet regression in random design and general errors with weakly dependent data. 2011. hal-00562228

# HAL Id: hal-00562228 https://hal.science/hal-00562228

Preprint submitted on 2 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Christophe Chesneau

University of Caen (LMNO), Caen, France. Email: chesneau@math.unicaen.fr

#### Abstract

We investigate function estimation in a nonparametric regression model having the following particularities: the design is random (with a known distribution), the errors admit finite moments of order 2 and the data are weakly dependent; the exponentially strongly mixing case is considered. In this general framework, we construct a new adaptive estimator. It is based on wavelets and the combination of two hard thresholding rules. We determine an upper bound of the associated mean integrated squared error and prove that it is sharp for a wide class of regression functions.

**Keywords:** Nonparametric regression, weakly dependent data, minimax estimation, Besov balls, wavelets, hard thresholding.

2000 Mathematics Subject Classification: 62G07, 62G20.

### **1** Introduction

Let  $(X_i, Y_i)_{i \in \mathbb{Z}}$  be a bivariate stationary random process where, for any  $i \in \mathbb{Z}$ ,

$$Y_i = f(X_i) + \xi_i, \tag{1.1}$$

 $f:[0,1] \to \mathbb{R}$  is an unknown function,  $(X_i)_{i\in\mathbb{Z}}$  is a sequence of identically distributed random variables having the common known density  $g:[0,1] \to [0,\infty)$  and  $(\xi_i)_{i\in\mathbb{Z}}$  is a sequence of identically distributed variables independent of  $(X_i)_{i\in\mathbb{Z}}$  satisfying  $\mathbb{E}(\xi_1) =$ 0 and  $\mathbb{E}(\xi_1^2) < \infty$ . We suppose that  $(X_i, Y_i)_{i\in\mathbb{Z}}$  is strongly mixing (to be defined in Section 2). Given *n* observations  $(X_1, Y_1), \ldots, (X_n, Y_n)$  drawn from  $(X_i, Y_i)_{i\in\mathbb{Z}}$ , we aim to estimate *f* globally on [0, 1].

To measure the performance of an estimator  $\hat{f}$  of f, we use the Mean Integrated Squared Error (MISE) defined by:

$$\mathbf{R}(\widehat{f},f) = \mathbb{E}\left(\int_0^1 (\widehat{f}(x) - f(x))^2 dx\right).$$

Our goal is to construct  $\hat{f}$  such that the associated MISE is as small as possible. Many methods can be considered (kernel, spline, wavelets,...) (see e.g. [31]). In this study, we

This work is supported by ANR grant NatImages, ANR-08-EMER-009

#### Christophe Chesneau

focus our attention on the wavelet methods. They are attractive for nonparametric function estimation because of their spatial adaptivity, computational efficiency and asymptotic optimality properties. Further details can be found in [1] and [22].

In the litterature, when  $(X_1, Y_1), \ldots, (X_n, Y_n)$  are *i.i.d.*, various wavelet methods have been developed. See, e.g., [14–16], [17], [21], [2, 3], [4], [30], [5, 6] [7, 8], [10], [29], [13], [33], [23] and [9]. When  $\xi_1, \ldots, \xi_n$  have some kind of dependence (long memory,  $\rho$ -mixing,...), see e.g. [18], [19], [25], [26] and [24]. To the best of our knowledge, the wavelet estimation of f when  $(X_1, Y_1), \ldots, (X_n, Y_n)$  are weakly dependent has only been investigated by [26]. More precisely, [26] constructs a linear nonadaptive wavelet estimator of f which attained a sharp rate of convergence under the uniform risk over Besov balls. However, the adaptive wavelet estimation of f, more realistic, has never been addressed earlier and motivates this study. In addition to this new challenge, we relax some classical assumptions on the errors: the common distribution of  $\xi_1, \ldots, \xi_n$  can be unknown; only  $\mathbb{E}(\xi_1) = 0$  and  $\mathbb{E}(\xi_1^2) < \infty$  are required.

We construct a new adaptive wavelet estimator based on the following steps: we estimate the unknown wavelet coefficients of f by a new thresholded versions of the empirical ones, we operate a term-by-term selection of these estimators via a hard thresholding rule, then we reconstruct the selected estimators by taking the initial wavelet basis and choosing appropriate levels. Naturally, the definitions of both thresholds take into account the dependence of the data and are chosen to minimize the associated MISE. Assuming that fbelongs to a Besov balls  $B_{p,q}^s(M)$  (to be defined in Section 3), we prove that our estimator  $\hat{f}$  satisfies

$$\mathbf{R}(\widehat{f}, f) \le C\left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}$$

where C > 0 is a constant (independent of n),  $n_{\theta} = n^{\theta/(\theta+1)}$  and  $\theta$  refers to the exponentially strong mixing case.

The paper is organized as follows. Section 2 clarifies the assumptions on the model and introduces some notations. Section 3 describes the wavelet basis on [0, 1] and the Besov balls  $B_{p,q}^s(M)$ . Our wavelet hard thresholding estimator is presented in Section 4. The results are set in Section 5. Section 6 is devoted to the proofs.

## **2** Assumptions and notations

**Dependence assumption on**  $(X_i, Y_i)_{i \in \mathbb{Z}}$ . Set, for any  $i \in \mathbb{Z}$ ,  $Z_i = (X_i, Y_i)$ . We suppose that  $Z = (Z_i)_{i \in \mathbb{Z}}$  is stationary and exponentially strongly mixing. Let us now clarify this kind of dependence.

For any  $m \in \mathbb{Z}$ , we define the *m*-th strongly mixing coefficient of Z by

$$a_m = \sup_{(A,B)\in \mathcal{F}^Z_{-\infty,0}\times \mathcal{F}^Z_{m,\infty}} \left| \mathbb{P}(A\cap B) - \mathbb{P}(A)\mathbb{P}(B) \right|,$$

where, for any  $i \in \mathbb{Z}$ ,  $\mathcal{F}_{-\infty,i}^{Z}$  is the  $\sigma$ -algebra generated by  $\ldots, Z_{i-1}, Z_i$  and  $\mathcal{F}_{i,\infty}^{Z}$  is the  $\sigma$ -algebra generated by  $Z_i, Z_{i+1}, \ldots$ . The bivariate random process Z is said to be strongly mixing if  $\lim_{m\to\infty} a_m = 0$ . The exponentially strongly mixing condition is characterized by the following inequality: there exists three known constants,  $\gamma > 0, c > 0$  and  $\theta > 0$  such that, for any  $m \in \mathbb{Z}$ ,

$$a_m \le \gamma exp\left(-c|m|^{\theta}\right). \tag{2.1}$$

This assumption is satisfied by a large class of processes (ARMA processes, ARCH process, ...). See e.g. [32], [20] and [28]. Remark that, when  $\theta \to \infty$ ,  $Z = (Z_i)_{i \in \mathbb{Z}}$  becomes a bivariate sequence of *i.i.d.* random variables.

**Boundedness assumption on** f. We suppose that there exists a known constant  $C_* > 0$  such that

$$\sup_{x \in [0,1]} |f(x)| \le C_*.$$
(2.2)

**Boundedness ssumption on** g. We suppose that there exists a known constant  $c_* > 0$  such that

$$\inf_{x \in [0,1]} g(x) \ge c_*.$$
(2.3)

### **3** Wavelet bases and Besov balls

#### 3.1 Wavelet basis

Let  $N \in \mathbb{N}^*$  (the set of positive integers), and  $\phi$  and  $\psi$  be the initial wavelet functions of the Daubechies wavelets dbN. Set

$$\phi_{j,k}(x) = 2^{j/2}\phi(2^jx - k), \qquad \psi_{j,k}(x) = 2^{j/2}\psi(2^jx - k).$$

Then, with an appropriate treatments at the boundaries, there exists an integer  $\tau$  satisfying  $2^{\tau} \geq 2N$  such that, for any integer  $\ell \geq \tau$ ,

$$\mathcal{B} = \{\phi_{\ell,k}(.), k \in \{0, \dots, 2^{\ell} - 1\}; \psi_{j,k}(.); j \in \mathbb{N} - \{0, \dots, \ell - 1\}, k \in \{0, \dots, 2^{j} - 1\}\},\$$

is an orthonormal basis of  $\mathbb{L}^2([0,1]) = \{h : [0,1] \to \mathbb{R}; \int_0^1 h^2(x) dx < \infty\}$ . See [11].

For any integer  $\ell \geq \tau$ , any  $h \in \mathbb{L}^2([0,1])$  can be expanded on  $\mathcal B$  as

$$h(x) = \sum_{k=0}^{2^{\ell}-1} \alpha_{\ell,k} \phi_{\ell,k}(x) + \sum_{j=\ell}^{\infty} \sum_{k=0}^{2^{j}-1} \beta_{j,k} \psi_{j,k}(x),$$

where  $\alpha_{j,k}$  and  $\beta_{j,k}$  are the wavelet coefficients of h defined by

$$\alpha_{j,k} = \int_0^1 h(x)\phi_{j,k}(x)dx, \qquad \beta_{j,k} = \int_0^1 h(x)\psi_{j,k}(x)dx.$$
(3.1)

Christophe Chesneau

#### 3.2 Besov balls

Let M > 0, s > 0,  $p \ge 1$  and  $r \ge 1$ . A function h belongs to  $B^s_{p,r}(M)$  if, and only if, there exists a constant  $M^* > 0$  (depending on M) such that the associated wavelet coefficients (3.1) satisfy

$$2^{\tau(1/2-1/p)} \left(\sum_{k=0}^{2^{\tau}-1} |\alpha_{\tau,k}|^p\right)^{1/p} + \left(\sum_{j=\tau}^{\infty} \left(2^{j(s+1/2-1/p)} \left(\sum_{k=0}^{2^j-1} |\beta_{j,k}|^p\right)^{1/p}\right)^r\right)^{1/r} \le M^*.$$

In this expression, s is a smoothness parameter and p and r are norm parameters. For a particular choice of s, p and r, the Besov balls contain the standard Hölder and Sobolev balls. See [27].

## 4 Estimators

#### 4.1 Wavelet coefficient estimators and some of their statistical properties

The first step to estimate f in (1.1) consists in expanding f on the wavelet basis  $\mathcal{B}$ . Then we aim to estimate the unknown wavelet coefficients:  $\alpha_{j,k} = \int_0^1 f(x)\phi_{j,k}(x)dx$  and  $\beta_{j,k} = \int_0^1 f(x)\psi_{j,k}(x)dx$ . The considered estimators are described below. Set

$$\gamma_n = \mu \sqrt{\frac{n_\theta}{\ln n_\theta}}$$

where  $n_{\theta} = n^{\theta/(\theta+1)}$ ,  $\theta$  is the one in (2.1) and  $\mu = \sqrt{(C_*^2 + \mathbb{E}(\xi_1^2))/c_*}$ . For any integer  $j \ge \tau$  and any  $k \in \{0, \ldots, 2^j - 1\}$ ,

• we estimate  $\alpha_{j,k}$  by

$$\widehat{\alpha}_{j,k} = \frac{1}{n} \sum_{i=1}^{n} \frac{Y_i}{g(X_i)} \phi_{j,k}(X_i) \mathbb{I}_{\left\{ \left| \frac{Y_i}{g(X_i)} \phi_{j,k}(X_i) \right| \le \gamma_n \right\}},\tag{4.1}$$

where, for any random event  $\mathcal{A}$ ,  $\mathbb{I}_{\mathcal{A}}$  is the indicator function on  $\mathcal{A}$ .

• we estimate  $\beta_{j,k}$  by

$$\widehat{\beta}_{j,k} = \frac{1}{n} \sum_{i=1}^{n} \frac{Y_i}{g(X_i)} \psi_{j,k}(X_i) \mathbb{I}_{\left\{ \left| \frac{Y_i}{g(X_i)} \psi_{j,k}(X_i) \right| \le \gamma_n \right\}}.$$
(4.2)

Remark that  $\widehat{\alpha}_{j,k}$  and  $\widehat{\beta}_{j,k}$  are thresholded versions of the standard empirical wavelet estimators for (1.1) (see, e.g., [14–16]). Such an observation thresholding has been introduced by [13] for (1.1) when  $(X_1, Y_1), \ldots, (X_n, Y_n)$  are *i.i.d.*. In our study, it allows

4

us to have non restrictive assumptions on  $\xi_1, \ldots, \xi_n$  and treat the weak dependence of  $(X_1, Y_1), \ldots, (X_n, Y_n)$ .

Propositions 4.1 and 4.2 below show moments properties for (4.1) and (4.2).

**Proposition 4.1.** Consider (1.1) under the assumptions of Section 2. For any integer  $j \ge \tau$  and any  $k \in \{0, ..., 2^j - 1\}$ , let  $\beta_{j,k} = \int_0^1 f(x)\psi_{j,k}(x)dx$  and  $\hat{\beta}_{j,k}$  be (4.2). Then there exists a constant C > 0 such that

$$\mathbb{E}\left((\widehat{\beta}_{j,k} - \beta_{j,k})^2\right) \le C \frac{\ln n_\theta}{n_\theta}.$$

This inequality holds with  $\alpha_{j,k} = \int_0^1 f(x)\phi_{j,k}(x)dx$  instead of  $\beta_{j,k}$  and  $\widehat{\alpha}_{j,k}$  defined by (4.1) instead of  $\widehat{\beta}_{j,k}$ .

**Proposition 4.2.** Consider (1.1) under the assumptions of Section 2. For any integer  $j \ge \tau$  and any  $k \in \{0, ..., 2^j - 1\}$ , let  $\beta_{j,k} = \int_0^1 f(x)\psi_{j,k}(x)dx$  and  $\hat{\beta}_{j,k}$  be (4.2). Then there exists a constant C > 0 such that

$$\mathbb{E}\left((\widehat{\beta}_{j,k}-\beta_{j,k})^4\right) \le C.$$

Proposition 4.3 below determines a concentration inequality for (4.2).

**Proposition 4.3.** Consider (1.1) under the assumptions of Section 2. For any integer  $j \ge \tau$  and any  $k \in \{0, \ldots, 2^j - 1\}$ , let  $\beta_{j,k} = \int_0^1 f(x)\psi_{j,k}(x)dx$ ,  $\widehat{\beta}_{j,k}$  be (4.2) and

$$\lambda_n = \mu \sqrt{\frac{\ln n_\theta}{n_\theta}}.$$

Then, for any  $\kappa \ge 2 + 16/(3u) + 4\sqrt{(1/u)(16/9u^2 + 2)}$  with  $u = (1/2)(c/8)^{1/(\theta+1)}$ , we have

$$\mathbb{P}\left(|\widehat{\beta}_{j,k} - \beta_{j,k}| \ge \kappa \lambda_n/2\right) \le 2(1 + 4e^{-2}\gamma)\frac{1}{n_{\theta}^4}.$$

#### 4.2 Hard thresholding estimator

We define the hard thresholding estimator  $\hat{f}$  by

$$\widehat{f}(x) = \sum_{k=0}^{2^{\tau}-1} \widehat{\alpha}_{\tau,k} \phi_{\tau,k}(x) + \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^{j}-1} \widehat{\beta}_{j,k} \mathbb{I}_{\left\{|\widehat{\beta}_{j,k}| \ge \kappa \lambda_n\right\}} \psi_{j,k}(x),$$
(4.3)

where  $\widehat{\alpha}_{\tau,k}$  is defined by (4.1) with  $j = \tau$ ,  $\widehat{\beta}_{j,k}$  by (4.2),  $j_1$  is the integer satisfying

$$\frac{1}{2}n_{\theta} < 2^{j_1} \le n_{\theta},$$
  
 $\kappa \ge 2 + 16/(3u) + 4\sqrt{(1/u)(16/9u^2 + 2)}$  with  $u = (1/2)(c/8)^{1/(\theta+1)}$  and  
 $\lambda_n = \mu \sqrt{\frac{\ln n_{\theta}}{n_{\theta}}}.$ 

The feature of  $\hat{f}$  is to only estimate the "large" unknown wavelet coefficients of f because they are those which contain his main characteristics.

# **5** Results

**Theorem 5.1.** Consider (1.1) under the assumptions of Section 2. Let  $\hat{f}$  be (4.3). Suppose that  $f \in B_{p,r}^s(M)$  with  $r \ge 1$ ,  $\{p \ge 2 \text{ and } s > 0\}$  or  $\{p \in [1, 2) \text{ and } s > 1/p\}$ . Then there exists a constant C > 0 such that

$$\mathbf{R}(\widehat{f}, f) \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)},$$

where  $n_{\theta} = n^{\theta/(\theta+1)}$ .

The proof of Theorem 5.1 uses a suitable decomposition of the MISE with the results in Propositions 4.1, 4.2 and 4.3.

If we restrict our study to independent  $(X_1, Y_1), \ldots, (X_n, Y_n)$  i.e.  $\theta \to \infty$ , our rate of convergence becomes  $(\ln n/n)^{2s/(2s+1)}$  which is the standard "near optimal" one in the minimax sense. See e.g. [22] and [13].

Note that the rate of convergence  $(\ln n_{\theta}/n_{\theta})^{2s/(2s+1)}$  is also achieved by the abstract minimum complexity regression estimator in [28, Theorem 2.1] but for a slightly different regression problem with more restrictions on  $(X_1, Y_1), \ldots, (X_n, Y_n), \xi_1, \ldots, \xi_n$  and f (see [28, Section II. B.]).

#### 6 **Proofs**

In this section, C represents a positive constant which may differ from one term to another.

**Proof of Proposition 4.1.** For any  $i \in \{1, \ldots, n\}$ , set

$$W_{i,j,k} = \frac{Y_i}{g(X_i)}\psi_{j,k}(X_i).$$

Since  $X_1$  and  $\xi_1$  are independent and  $\mathbb{E}(\xi_1) = 0$ , we have

/ --

$$\begin{split} \mathbb{E}(W_{1,j,k}) &= \mathbb{E}\left(\frac{Y_1}{g(X_1)}\psi_{j,k}(X_1)\right) \\ &= \mathbb{E}\left(\frac{f(X_1)}{g(X_1)}\psi_{j,k}(X_1)\right) + \mathbb{E}(\xi_1)\mathbb{E}(\frac{1}{g(X_1)}\psi_{j,k}(X_1)) \\ &= \mathbb{E}\left(\frac{f(X_1)}{g(X_1)}\psi_{j,k}(X_1)\right) = \int_0^1 \frac{f(x)}{g(x)}\psi_{j,k}(x)g(x)dx \\ &= \int_0^1 f(x)\psi_{j,k}(x)dx = \beta_{j,k}. \end{split}$$

Hence, since  $W_{1,j,k}, \ldots, W_{n,j,k}$  are identically distributed,

$$\beta_{j,k} = \mathbb{E}(W_{1,j,k}) = \mathbb{E}(W_{1,j,k} \mathbb{I}_{\{|W_{1,j,k}| \le \gamma_n\}}) + \mathbb{E}(W_{1,j,k} \mathbb{I}_{\{|W_{1,j,k}| > \gamma_n\}})$$
$$= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n W_{i,j,k} \mathbb{I}_{\{|W_{i,j,k}| \le \gamma_n\}}\right) + \mathbb{E}(W_{1,j,k} \mathbb{I}_{\{|W_{1,j,k}| > \gamma_n\}}).$$
(6.1)

This with the elementary inequality  $(x+y)^2 \leq 2(x^2+y^2), (x,y) \in \mathbb{R}^2$ , imply that

$$\mathbb{E}\left(\left(\widehat{\beta}_{j,k} - \beta_{j,k}\right)^2\right) \le 2(A+B),\tag{6.2}$$

where

$$A = \mathbb{E}\left(\left(\frac{1}{n}\sum_{i=1}^{n} \left(W_{i,j,k}\mathbb{1}_{\{|W_{i,j,k}| \le \gamma_n\}} - \mathbb{E}(W_{i,j,k}\mathbb{1}_{\{|W_{i,j,k}| \le \gamma_n\}})\right)\right)^2\right)$$

and

$$B = \left( \mathbb{E} \left( |W_{1,j,k}| \mathbb{1}_{\{|W_{1,j,k}| > \gamma_n\}} \right) \right)^2.$$

Let us bound B and A.

Upper bound for B. Since  $\mathrm{I\!I}_{\{|W_{1,j,k}|>\gamma_n\}} \leq |W_{1,j,k}|/\gamma_n,$  we have

$$\mathbb{E}\left(|W_{1,j,k}|\mathbf{I}_{\{|W_{1,j,k}|>\gamma_n\}}\right) \leq \frac{\mathbb{E}(W_{1,j,k}^2)}{\gamma_n}.$$

Let us now bound  $\mathbb{E}(W^2_{1,j,k})$ . Since  $X_1$  and  $\xi_1$  are independent and  $\mathbb{E}(\xi_1) = 0$ , we have

$$\begin{split} \mathbb{E}(W_{1,j,k}^2) &= \mathbb{E}\left(\frac{Y_1^2}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) = \mathbb{E}\left(\frac{(f(X_1) + \xi_1)^2}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) \\ &= \mathbb{E}\left(\frac{f^2(X_1)}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) + 2\mathbb{E}(\xi_1)\mathbb{E}\left(\frac{f(X_1)}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) \\ &+ \mathbb{E}\left(\xi_1^2\right)\mathbb{E}\left(\frac{1}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) \\ &= \mathbb{E}\left(\frac{f^2(X_1)}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) + \mathbb{E}(\xi_1^2)\mathbb{E}\left(\frac{1}{g^2(X_1)}\psi_{j,k}^2(X_1)\right). \end{split}$$

Using (2.2), (2.3) and  $\int_0^1 \psi_{j,k}^2(x) dx = 1$ , we obtain

$$\begin{split} \mathbb{E}\left(\frac{f^2(X_1)}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) &\leq C_*^2 \mathbb{E}\left(\frac{1}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) \\ &= C_*^2 \int_0^1 \frac{1}{g^2(x)}\psi_{j,k}^2(x)g(x)dx = C_*^2 \int_0^1 \frac{1}{g(x)}\psi_{j,k}^2(x)dx \\ &\leq \frac{C_*^2}{c_*} \int_0^1 \psi_{j,k}^2(x)dx = \frac{C_*^2}{c_*}. \end{split}$$

And, in a similar way,

$$\begin{split} \mathbb{E}\left(\frac{1}{g^2(X_1)}\psi_{j,k}^2(X_1)\right) &= \int_0^1 \frac{1}{g^2(x)}\psi_{j,k}^2(x)g(x)dx = \int_0^1 \frac{1}{g(x)}\psi_{j,k}^2(x)dx \\ &\leq \frac{1}{c_*}\int_0^1 \psi_{j,k}^2(x)dx = \frac{1}{c_*}. \end{split}$$

So

$$\mathbb{E}(W_{1,j,k}^2) \le \frac{1}{c_*} (C_*^2 + \mathbb{E}(\xi_1^2)) = \mu^2 = C.$$
(6.3)

Therefore

$$B = \left(\mathbb{E}\left(|W_{1,j,k}|\mathbf{1}_{\{|W_{1,j,k}| > \gamma_n\}}\right)\right)^2 \le \left(\frac{\mathbb{E}(W_{1,j,k}^2)}{\gamma_n}\right)^2 \le C\frac{1}{\gamma_n^2} \le C\frac{\ln n_\theta}{n_\theta}.$$
 (6.4)

Upper bound for A. We have

$$\begin{aligned}
A &= \mathbb{V}\left(\frac{1}{n}\sum_{i=1}^{n}W_{i,j,k}\mathbb{I}_{\{|W_{i,j,k}|\leq\gamma_{n}\}}\right) \\
&= \frac{1}{n^{2}}\sum_{v=1}^{n}\sum_{\ell=1}^{n}\mathbb{C}\left(W_{v,j,k}\mathbb{I}_{\{|W_{v,j,k}|\leq\gamma_{n}\}}, W_{\ell,j,k}\mathbb{I}_{\{|W_{\ell,j,k}|\leq\gamma_{n}\}}\right) \\
&= \frac{1}{n}\mathbb{V}(W_{1,j,k}\mathbb{I}_{\{|W_{1,j,k}|\leq\gamma_{n}\}}) + \frac{2}{n^{2}}\sum_{v=2}^{n}\sum_{\ell=1}^{v-1}\mathbb{C}\left(W_{v,j,k}\mathbb{I}_{\{|W_{v,j,k}|\leq\gamma_{n}\}}, W_{\ell,j,k}\mathbb{I}_{\{|W_{\ell,j,k}|\leq\gamma_{n}\}}\right) \\
&\leq \frac{1}{n}\mathbb{V}(W_{1,j,k}\mathbb{I}_{\{|W_{1,j,k}|\leq\gamma_{n}\}}) + \frac{2}{n^{2}}\left|\sum_{v=2}^{n}\sum_{\ell=1}^{v-1}\mathbb{C}\left(W_{v,j,k}\mathbb{I}_{\{|W_{v,j,k}|\leq\gamma_{n}\}}, W_{\ell,j,k}\mathbb{I}_{\{|W_{\ell,j,k}|\leq\gamma_{n}\}}\right)\right|.
\end{aligned}$$
(6.5)

Using (6.3), we obtain

$$\mathbb{V}(W_{1,j,k}\mathbb{I}_{\{|W_{1,j,k}| \le \gamma_n\}}) \le \mathbb{E}(W_{1,j,k}^2\mathbb{I}_{\{|W_{1,j,k}| \le \gamma_n\}}) \le \mathbb{E}(W_{1,j,k}^2) \le \mu^2 = C.$$
(6.6)

It follows from the stationarity of  $(X_i,Y_i)_{i\in\mathbb{Z}}$  that

$$\begin{split} &\sum_{v=2}^{n} \sum_{\ell=1}^{v-1} \mathbb{C} \left( W_{v,j,k} 1\!\!\mathrm{I}_{\{|W_{v,j,k}| \le \gamma_n\}}, W_{\ell,j,k} 1\!\!\mathrm{I}_{\{|W_{\ell,j,k}| \le \gamma_n\}} \right) \\ &= \left| \left| \sum_{m=1}^{n} (n-m) \mathbb{C} \left( W_{0,j,k} 1\!\!\mathrm{I}_{\{|W_{0,j,k}| \le \gamma_n\}}, W_{m,j,k} 1\!\!\mathrm{I}_{\{|W_{m,j,k}| \le \gamma_n\}} \right) \right| \\ &\leq n \sum_{m=1}^{n} \left| \mathbb{C} \left( W_{0,j,k} 1\!\!\mathrm{I}_{\{|W_{0,j,k}| \le \gamma_n\}}, W_{m,j,k} 1\!\!\mathrm{I}_{\{|W_{m,j,k}| \le \gamma_n\}} \right) \right|. \end{split}$$

By the Davydov inequality for strongly mixing processes (see [12]), the inequality  $|W_{0,j,k}| \mathbb{1}_{\{|W_{0,j,k}| \le \gamma_n\}} \le \max(\gamma_n, |W_{0,j,k}|)$  and (6.3), for any  $q \in (0, 1)$ , we have

$$\begin{aligned} & \left\| \mathbb{C} \left( W_{0,j,k} \, \mathbf{I}_{\{|W_{0,j,k}| \leq \gamma_n\}}, W_{m,j,k} \, \mathbf{I}_{\{|W_{m,j,k}| \leq \gamma_n\}} \right) \right\| \\ & \leq 10 a_m^q \left( \mathbb{E} \left( |W_{0,j,k}|^{2/(1-q)} \, \mathbf{I}_{\{|W_{0,j,k}| \leq \gamma_n\}} \right) \right)^{1-q} \\ & = 10 a_m^q \left( \mathbb{E} \left( |W_{0,j,k}|^{2q/(1-q)} \, \mathbf{I}_{\{|W_{0,j,k}| \leq \gamma_n\}} W_{0,j,k}^2 \, \mathbf{I}_{\{|W_{0,j,k}| \leq \gamma_n\}} \right) \right)^{1-q} \\ & \leq C a_m^q \left( \gamma_n^{2q/(1-q)} \right)^{1-q} \left( \mathbb{E} \left( W_{0,j,k}^2 \right) \right)^{1-q} \leq C a_m^q \left( \frac{n_\theta}{\ln n_\theta} \right)^q. \end{aligned}$$

Due to (2.1), we have  $\sum_{m=1}^{n} a_m^q \leq \gamma \sum_{m=1}^{\infty} exp\left(-cq|m|^{\theta}\right) = C$ . So

$$\left|\sum_{v=2}^{n}\sum_{\ell=1}^{v-1}\mathbb{C}\left(W_{v,j,k}\mathbb{I}_{\{|W_{v,j,k}|\leq\gamma_{n}\}},W_{\ell,j,k}\mathbb{I}_{\{|W_{\ell,j,k}|\leq\gamma_{n}\}}\right)\right| \leq Cn\left(\frac{n_{\theta}}{\ln n_{\theta}}\right)^{q}.$$
 (6.7)

Taking  $q = 1/\theta$ , we have  $n_{\theta}^q/n = 1/n_{\theta}$ . It follows from (6.5), (6.6) and (6.7) that

$$A \le C\left(\frac{1}{n} + \frac{1}{n^2}n\left(\frac{n_\theta}{\ln n_\theta}\right)^q\right) \le C\frac{n_\theta^q}{n} = C\frac{1}{n_\theta} \le C\frac{\ln n_\theta}{n_\theta}.$$
(6.8)

It follows from (6.2), (6.4) and (6.8) that

$$\mathbb{E}\left((\widehat{\beta}_{j,k} - \beta_{j,k})^2\right) \le C \frac{\ln n_\theta}{n_\theta}.$$
(6.9)

Replacing  $\phi$  instead of  $\psi$  in the previous proof, one can show that (6.9) holds with  $\alpha_{j,k} = \int_0^1 f(x)\phi_{j,k}(x)dx$  instead of  $\beta_{j,k}$  and  $\widehat{\alpha}_{j,k}$  defined by (4.1) instead of  $\widehat{\beta}_{j,k}$ .

The proof of Proposition 4.1 is complete.

# **Proof of Proposition 4.2.** We have

$$|\widehat{\beta}_{j,k} - \beta_{j,k}| \le |\widehat{\beta}_{j,k}| + |\beta_{j,k}|.$$

We have

$$\left|\widehat{\beta}_{j,k}\right| \leq \frac{1}{n} \sum_{i=1}^{n} \left| \frac{Y_i}{g(X_i)} \psi_{j,k}(X_i) \right| \mathbf{I}_{\left\{ \left| \frac{Y_i}{g(X_i)} \psi_{j,k}(X_i) \right| \leq \gamma_n \right\}} \leq \gamma_n = \mu \sqrt{\frac{n_\theta}{\ln n_\theta}}.$$

It follows from (2.2), the Cauchy-Schwarz inequality and  $\int_0^1 \psi_{j,k}^2(x) dx = 1$  that

$$\begin{aligned} |\beta_{j,k}| &\leq \int_0^1 |f(x)| |\psi_{j,k}(x)| dx \leq C_* \int_0^1 |\psi_{j,k}(x)| dx \\ &\leq C_* \left( \int_0^1 \psi_{j,k}^2(x) dx \right)^{1/2} = C_* \leq C \sqrt{\frac{n_\theta}{\ln n_\theta}}. \end{aligned}$$

Therefore

$$|\widehat{\beta}_{j,k} - \beta_{j,k}| \le C \sqrt{\frac{n_{\theta}}{\ln n_{\theta}}}.$$
(6.10)

By (6.10) and Proposition 4.1, we have

$$\mathbb{E}\left(\left(\widehat{\beta}_{j,k}-\beta_{j,k}\right)^{4}\right) \leq C\frac{n_{\theta}}{\ln n_{\theta}}\mathbb{E}\left(\left(\widehat{\beta}_{j,k}-\beta_{j,k}\right)^{2}\right) \leq C\frac{n_{\theta}}{\ln n_{\theta}}\frac{\ln n_{\theta}}{n_{\theta}} = C.$$

L		
L		
ь.		

**Proof of Proposition 4.3.** For any  $i \in \{1, \ldots, n\}$ , set

$$W_{i,j,k} = \frac{Y_i}{g(X_i)}\psi_{j,k}(X_i).$$

By (6.1), we have

$$\begin{aligned} |\widehat{\beta}_{j,k} - \beta_{j,k}| \\ &= \left| \frac{1}{n} \sum_{i=1}^{n} \left( W_{i,j,k} \mathbf{I}_{\{|W_{i,j,k}| \le \gamma_n\}} - \mathbb{E} \left( W_{i,j,k} \mathbf{I}_{\{|W_{i,j,k}| \le \gamma_n\}} \right) \right) - \mathbb{E} \left( W_{1,j,k} \mathbf{I}_{\{|W_{1,j,k}| > \gamma_n\}} \right) \right| \\ &\leq \left| \frac{1}{n} \sum_{i=1}^{n} \left( W_{i,j,k} \mathbf{I}_{\{|W_{i,j,k}| \le \gamma_n\}} - \mathbb{E} \left( W_{i,j,k} \mathbf{I}_{\{|W_{i,j,k}| \le \gamma_n\}} \right) \right) \right| + \mathbb{E} \left( |W_{1,j,k}| \mathbf{I}_{\{|W_{1,j,k}| > \gamma_n\}} \right). \end{aligned}$$

Using (6.3), we obtain

$$\mathbb{E}\left(|W_{1,j,k}|\mathbf{1}_{\{|W_{1,j,k}|>\gamma_n\}}\right) \leq \frac{\mathbb{E}(W_{1,j,k}^2)}{\gamma_n} \leq \mu^2 \frac{1}{\mu} \sqrt{\frac{\ln n_\theta}{n_\theta}} = \lambda_n.$$

Hence

$$\mathbb{P}\left(|\widehat{\beta}_{j,k} - \beta_{j,k}| \ge \kappa \lambda_n/2\right) \\
\le \mathbb{P}\left(\left|\frac{1}{w_n} \sum_{i=1}^n \left(W_{i,j,k} \mathbb{I}_{\{|W_{i,j,k}| \le \gamma_n\}} - \mathbb{E}\left(W_{i,j,k} \mathbb{I}_{\{|W_{i,j,k}| \le \gamma_n\}}\right)\right)\right| \ge (\kappa/2 - 1)\lambda_n\right). \tag{6.11}$$

Let us now present a Bernstein inequality for exponentially strongly mixing process. This is a slightly modified version of [28, Theorem 4.2].

**Lemma 6.1** ([28]). Let  $\gamma > 0$ , c > 0,  $\theta > 1$  and  $(S_i)_{i \in \mathbb{Z}}$  be a stationary process such that, for any  $m \in \mathbb{Z}$ , the associated m-th strongly mixing coefficient satisfies

$$a_m \leq \gamma exp(-c|m|^{\theta}).$$

Let  $n \in \mathbb{N}^*$ ,  $h : \mathbb{R} \to \mathbb{R}$  be a measurable function and, for any  $i \in \mathbb{Z}$ ,  $U_i = h(S_i)$ . We assume that  $\mathbb{E}(U_1) = 0$  and there exists a constant M > 0 satisfying  $|U_1| \le M < \infty$ . Then, for any  $\lambda > 0$ , we have

$$\mathbb{P}\left(\left|\frac{1}{n}\sum_{i=1}^{n}U_{i}\right| \geq \lambda\right) \leq 2(1+4e^{-2}\gamma)\exp\left(-\frac{u\lambda^{2}n^{\theta/(\theta+1)}}{2\left(\mathbb{E}\left(U_{1}^{2}\right)+\lambda M/3\right)}\right),$$

where  $u = (1/2)(c/8)^{1/(\theta+1)}$ .

Set, for any  $i \in \{1, \ldots, n\}$ ,

$$U_{i,j,k} = W_{i,j,k} \, \mathrm{I}_{\{|W_{i,j,k}| \le \gamma_n\}} - \mathbb{E} \left( W_{i,j,k} \, \mathrm{I}_{\{|W_{i,j,k}| \le \gamma_n\}} \right).$$

10

Then  $U_{1,j,k}, \ldots, U_{n,j,k}$  are identically distributed, depend on the stationary strongly mixing process  $(X_i, Y_i)_{i \in \mathbb{Z}}$  which satisfies (2.1),  $\mathbb{E}(U_{1,j,k}) = 0$ ,

$$|U_{1,j,k}| \le |W_{1,j,k}| \, \mathrm{I}\!\mathrm{I}_{\{|W_{1,j,k}| \le \gamma_n\}} + \mathbb{E}\left(|W_{1,j,k}| \, \mathrm{I}\!\mathrm{I}_{\{|W_{1,j,k}| \le \gamma_n\}}\right) \le 2\gamma_n$$

and, by (6.3),

$$\mathbb{E}(U_{1,j,k}^2) = \mathbb{V}\left(W_{1,j,k} \mathrm{I}_{\{|W_{1,j,k}| \le \gamma_n\}}\right) \le \mathbb{E}(W_{1,j,k}^2) \le \mu^2.$$

It follows from Lemma 6.1 that

$$\mathbb{P}\left(\left|\frac{1}{n}\sum_{i=1}^{n}U_{i,j,k}\right| \ge (\kappa/2-1)\lambda_n\right) \le 2(1+4e^{-2}\gamma)\exp\left(-\frac{u(\kappa/2-1)^2\lambda_n^2n_\theta}{2(\mu^2+2(\kappa/2-1)\lambda_n\gamma_n/3)}\right).$$
(6.12)

We have

$$\lambda_n \gamma_n = \mu \sqrt{\frac{\ln n_\theta}{n_\theta}} \mu \sqrt{\frac{n_\theta}{\ln n_\theta}} = \mu^2, \qquad \lambda_n^2 = \mu^2 \frac{\ln n_\theta}{n_\theta}.$$

Combining (6.11) and (6.12), for any  $\kappa \ge 2 + 16/(3u) + 4\sqrt{(1/u)(16/9u^2 + 2)}$ , we have

$$\begin{split} \mathbb{P}\left(|\widehat{\beta}_{j,k} - \beta_{j,k}| \geq \kappa \lambda_n/2\right) \\ &\leq 2(1 + 4e^{-2}\gamma) \exp\left(-\frac{u(\kappa/2 - 1)^2 \ln n_{\theta}}{2\left(1 + 2(\kappa/2 - 1)/3\right)}\right) \\ &= 2(1 + 4e^{-2}\gamma) n_{\theta}^{-\frac{u(\kappa/2 - 1)^2}{2(1 + 2(\kappa/2 - 1)/3)}} \leq 2(1 + 4e^{-2}\gamma) \frac{1}{n_{\theta}^4}. \end{split}$$

This ends the proof of Proposition 4.3.

**Proof of Theorem 5.1.** We expand the function f on  $\mathcal{B}$  as

$$f(x) = \sum_{k=0}^{2^{\tau}-1} \alpha_{\tau,k} \phi_{\tau,k}(x) + \sum_{j=\tau}^{\infty} \sum_{k=0}^{2^{j}-1} \beta_{j,k} \psi_{j,k}(x), \quad x \in [0,1],$$

where

 $\overline{}$ 

$$\alpha_{\tau,k} = \int_0^1 f(x)\phi_{\tau,k}(x)dx, \qquad \beta_{j,k} = \int_0^1 f(x)\psi_{j,k}(x)dx.$$

We have, for any  $x \in [0, 1]$ ,

$$f(x) - f(x) = \sum_{k=0}^{2^{\tau}-1} (\widehat{\alpha}_{\tau,k} - \alpha_{\tau,k}) \phi_{\tau,k}(x) + \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^{j}-1} \left( \widehat{\beta}_{j,k} \mathbb{I}_{\{|\widehat{\beta}_{j,k}| \ge \kappa \lambda_n\}} - \beta_{j,k} \right) \psi_{j,k}(x) \\ - \sum_{j=j_1+1}^{\infty} \sum_{k=0}^{2^{j}-1} \beta_{j,k} \psi_{j,k}(x).$$

Since  $\mathcal{B}$  is an orthonormal basis of  $\mathbb{L}^2([0,1])$ , we can write

$$\mathbf{R}(\widehat{f},f) = \mathbb{E}\left(\int_0^1 (\widehat{f}(x) - f(x))^2 dx\right) = R + S + T,\tag{6.13}$$

where

$$R = \sum_{k=0}^{2^{\tau}-1} \mathbb{E}\left( (\widehat{\alpha}_{\tau,k} - \alpha_{\tau,k})^2 \right), \qquad S = \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \mathbb{E}\left( (\widehat{\beta}_{j,k} \mathbb{1}_{\left\{ |\widehat{\beta}_{j,k}| \ge \kappa \lambda_n \right\}} - \beta_{j,k})^2 \right)$$

and

$$T = \sum_{j=j_1+1}^{\infty} \sum_{k=0}^{2^j - 1} \beta_{j,k}^2.$$

Let us bound R, T and S.

Upper bound for R. Using Proposition 4.1 and 2s/(2s+1) < 1, we obtain

$$R \le C 2^{\tau} \frac{\ln n_{\theta}}{n_{\theta}} \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$
(6.14)

•

Upper bound for T. For  $r \ge 1$  and  $p \ge 2$ , we have  $B^s_{p,r}(M) \subseteq B^s_{2,\infty}(M)$ . Since 2s/(2s+1) < 2s, we have

$$T \le C \sum_{j=j_1+1}^{\infty} 2^{-2j_s} \le C 2^{-2j_1s} \le C n_{\theta}^{-2s} \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s} \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}$$

For  $r \ge 1$  and  $p \in [1,2)$ , we have  $B_{p,r}^s(M) \subseteq B_{2,\infty}^{s+1/2-1/p}(M)$ . Since s > 1/p, we have s + 1/2 - 1/p > s/(2s+1). So

$$T \leq C \sum_{j=j_1+1}^{\infty} 2^{-2j(s+1/2-1/p)} \leq C 2^{-2j_1(s+1/2-1/p)}$$
$$\leq C n_{\theta}^{-2(s+1/2-1/p)} \leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2(s+1/2-1/p)} \leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$

Hence, for  $r \ge 1$ ,  $\{p \ge 2 \text{ and } s > 0\}$  or  $\{p \in [1, 2) \text{ and } s > 1/p\}$ , we have

$$T \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$
(6.15)

Upper bound for S. We have

$$S = S_1 + S_2 + S_3 + S_4, (6.16)$$

where

$$S_1 = \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \mathbb{E}\left( (\widehat{\beta}_{j,k} - \beta_{j,k})^2 \mathbf{I}_{\left\{ |\widehat{\beta}_{j,k}| \ge \kappa \lambda_n \right\}} \mathbf{I}_{\left\{ |\beta_{j,k}| < \kappa \lambda_n / 2 \right\}} \right),$$

$$S_{2} = \sum_{j=\tau}^{j_{1}} \sum_{k=0}^{2^{j}-1} \mathbb{E}\left( (\widehat{\beta}_{j,k} - \beta_{j,k})^{2} \mathbb{I}_{\left\{ |\widehat{\beta}_{j,k}| \ge \kappa\lambda_{n} \right\}} \mathbb{I}_{\left\{ |\beta_{j,k}| \ge \kappa\lambda_{n}/2 \right\}} \right),$$
$$S_{3} = \sum_{j=\tau}^{j_{1}} \sum_{k=0}^{2^{j}-1} \mathbb{E}\left( \beta_{j,k}^{2} \mathbb{I}_{\left\{ |\widehat{\beta}_{j,k}| < \kappa\lambda_{n} \right\}} \mathbb{I}_{\left\{ |\beta_{j,k}| \ge 2\kappa\lambda_{n} \right\}} \right)$$

and

$$S_4 = \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j - 1} \mathbb{E}\left(\beta_{j,k}^2 \mathrm{I}\!\mathrm{I}_{\left\{|\widehat{\beta}_{j,k}| < \kappa\lambda_n\right\}} \mathrm{I}\!\mathrm{I}_{\left\{|\beta_{j,k}| < 2\kappa\lambda_n\right\}}\right).$$

Let us investigate the bounds of  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_4$ . Upper bounds for  $S_1$  and  $S_3$ . We have

$$\left\{ |\widehat{\beta}_{j,k}| < \kappa \lambda_n, \ |\beta_{j,k}| \ge 2\kappa \lambda_n \right\} \subseteq \left\{ |\widehat{\beta}_{j,k} - \beta_{j,k}| > \kappa \lambda_n/2 \right\},$$
$$\left\{ |\widehat{\beta}_{j,k}| \ge \kappa \lambda_n, \ |\beta_{j,k}| < \kappa \lambda_n/2 \right\} \subseteq \left\{ |\widehat{\beta}_{j,k} - \beta_{j,k}| > \kappa \lambda_n/2 \right\}$$

and

$$\left\{ |\widehat{\beta}_{j,k}| < \kappa \lambda_n, \ |\beta_{j,k}| \ge 2\kappa \lambda_n \right\} \subseteq \left\{ |\beta_{j,k}| \le 2|\widehat{\beta}_{j,k} - \beta_{j,k}| \right\}.$$

So

$$\max(S_1, S_3) \le C \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \mathbb{E}\left( (\widehat{\beta}_{j,k} - \beta_{j,k})^2 \mathbf{I}_{\left\{ |\widehat{\beta}_{j,k} - \beta_{j,k}| > \kappa \lambda_n/2 \right\}} \right).$$

It follows from the Cauchy-Schwarz inequality, Propositions 4.2 and 4.3 that

$$\mathbb{E}\left((\widehat{\beta}_{j,k}-\beta_{j,k})^{2}\mathbb{I}_{\left\{|\widehat{\beta}_{j,k}-\beta_{j,k}|>\kappa\lambda_{n}/2\right\}}\right) \\
\leq \left(\mathbb{E}\left((\widehat{\beta}_{j,k}-\beta_{j,k})^{4}\right)\right)^{1/2} \left(\mathbb{P}\left(|\widehat{\beta}_{j,k}-\beta_{j,k}|>\kappa\lambda_{n}/2\right)\right)^{1/2} \\
\leq C\left(\frac{1}{n_{\theta}^{4}}\right)^{1/2} = C\frac{1}{n_{\theta}^{2}}.$$

Since 2s/(2s+1) < 1, we have

$$\max(S_1, S_3) \le C \frac{1}{n_{\theta}^2} \sum_{j=\tau}^{j_1} 2^j \le C \frac{1}{n_{\theta}^2} 2^{j_1} \le C \frac{1}{n_{\theta}} \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$
 (6.17)

Upper bound for  $S_2$ . Using again Proposition 4.1, we obtain

$$\mathbb{E}\left((\widehat{\beta}_{j,k} - \beta_{j,k})^2\right) \le C \frac{\ln n_\theta}{n_\theta}.$$

Hence

$$S_2 \leq C \frac{\ln n_\theta}{n_\theta} \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \mathrm{I}_{\{|\beta_{j,k}| > \kappa \lambda_n/2\}}.$$

Let  $j_2$  be the integer defined by

$$\frac{1}{2} \left(\frac{n_{\theta}}{\ln n_{\theta}}\right)^{1/(2s+1)} < 2^{j_2} \le \left(\frac{n_{\theta}}{\ln n_{\theta}}\right)^{1/(2s+1)}.$$
(6.18)

We have

$$S_2 \le S_{2,1} + S_{2,2},$$

where

$$S_{2,1} = C \frac{\ln n_{\theta}}{n_{\theta}} \sum_{j=\tau}^{j_2} \sum_{k=0}^{2^j - 1} \mathrm{I}_{\{|\beta_{j,k}| > \kappa \lambda_n/2\}}$$

and

$$S_{2,2} = C \frac{\ln n_{\theta}}{n_{\theta}} \sum_{j=j_{2}+1}^{j_{1}} \sum_{k=0}^{2^{j}-1} \mathrm{I}_{\{|\beta_{j,k}| > \kappa \lambda_{n}/2\}}.$$

We have

$$S_{2,1} \le C \frac{\ln n_{\theta}}{n_{\theta}} \sum_{j=\tau}^{j_2} 2^j \le C \frac{\ln n_{\theta}}{n_{\theta}} 2^{j_2} \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$

For  $r\geq 1$  and  $p\geq 2,$  since  $B^s_{p,r}(M)\subseteq B^s_{2,\infty}(M),$  we have

$$\begin{split} S_{2,2} &\leq C \frac{\ln n_{\theta}}{n_{\theta} \lambda_n^2} \sum_{j=j_2+1}^{j_1} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \leq C \sum_{j=j_2+1}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \leq C 2^{-2j_2s} \\ &\leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}. \end{split}$$

For  $r \ge 1$ ,  $p \in [1,2)$  and s > 1/p, using  $I\!\!I_{\{|\beta_{j,k}| > \kappa \lambda_n/2\}} \le C |\beta_{j,k}|^p / \lambda_n^p$ ,  $B_{p,r}^s(M) \subseteq B_{2,\infty}^{s+1/2-1/p}(M)$  and (2s+1)(2-p)/2 + (s+1/2-1/p)p = 2s, we have

$$S_{2,2} \leq C \frac{\ln n_{\theta}}{n_{\theta} \lambda_{n}^{p}} \sum_{j=j_{2}+1}^{j_{1}} \sum_{k=0}^{2^{j}-1} |\beta_{j,k}|^{p} \leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{(2-p)/2} \sum_{j=j_{2}+1}^{\infty} 2^{-j(s+1/2-1/p)p}$$
$$\leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{(2-p)/2} 2^{-j_{2}(s+1/2-1/p)p} \leq C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$

So, for  $r\geq 1,$   $\{p\geq 2 \text{ and } s>0\}$  or  $\{p\in [1,2) \text{ and } s>1/p\},$  we have

$$S_2 \le C \left(\frac{\ln n_\theta}{n_\theta}\right)^{2s/(2s+1)}.$$
(6.19)

*Upper bound for*  $S_4$ . We have

$$S_4 \leq \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \mathbb{I}_{\{|\beta_{j,k}| < 2\kappa\lambda_n\}}.$$

Let  $j_2$  be the integer (6.18). Then

$$S_4 \le S_{4,1} + S_{4,2},$$

where

$$S_{4,1} = \sum_{j=\tau}^{j_2} \sum_{k=0}^{2^j - 1} \beta_{j,k}^2 \mathbb{I}_{\{|\beta_{j,k}| < 2\kappa\lambda_n\}}, \qquad S_{4,2} = \sum_{j=j_2+1}^{j_1} \sum_{k=0}^{2^j - 1} \beta_{j,k}^2 \mathbb{I}_{\{|\beta_{j,k}| < 2\kappa\lambda_n\}}.$$

We have

$$S_{4,1} \le C \sum_{j=\tau}^{j_2} 2^j \lambda_n^2 = C \frac{\ln n_\theta}{n_\theta} \sum_{j=\tau}^{j_2} 2^j \le C \frac{\ln n_\theta}{n_\theta} 2^{j_2} \le C \left(\frac{\ln n_\theta}{n_\theta}\right)^{2s/(2s+1)}$$

For  $r \geq 1$  and  $p \geq 2$ , since  $B^s_{p,r}(M) \subseteq B^s_{2,\infty}(M)$ , we have

$$S_{4,2} \le \sum_{j=j_2+1}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k}^2 \le C 2^{-2j_2s} \le C \left(\frac{\ln n_\theta}{n_\theta}\right)^{2s/(2s+1)}.$$

For  $r \ge 1, p \in [1, 2)$  and s > 1/p, using  $\beta_{j,k}^2 \mathbb{I}_{\{|\beta_{j,k}| < 2\kappa\lambda_n\}} \le C\lambda_n^{2-p} |\beta_{j,k}|^p$ ,  $B_{p,r}^s(M) \subseteq B_{2,\infty}^{s+1/2-1/p}(M)$  and (2s+1)(2-p)/2 + (s+1/2-1/p)p = 2s, we have

$$S_{4,2} \leq C\lambda_n^{2-p} \sum_{j=j_2+1}^{j_1} \sum_{k=0}^{2^j-1} |\beta_{j,k}|^p = C\left(\frac{\ln n_\theta}{n_\theta}\right)^{(2-p)/2} \sum_{j=j_2+1}^{j_1} \sum_{k=0}^{2^j-1} |\beta_{j,k}|^p$$
  
$$\leq C\left(\frac{\ln n_\theta}{n_\theta}\right)^{(2-p)/2} \sum_{j=j_2+1}^{\infty} 2^{-j(s+1/2-1/p)p} \leq C\left(\frac{\ln n_\theta}{n_\theta}\right)^{(2-p)/2} 2^{-j_2(s+1/2-1/p)p}$$
  
$$\leq C\left(\frac{\ln n_\theta}{n_\theta}\right)^{2s/(2s+1)}.$$

So, for  $r \ge 1$ ,  $\{p \ge 2 \text{ and } s > 0\}$  or  $\{p \in [1, 2) \text{ and } s > 1/p\}$ , we have

$$S_4 \le C \left(\frac{\ln n_\theta}{n_\theta}\right)^{2s/(2s+1)}.$$
(6.20)

It follows from (6.16), (6.17), (6.19) and (6.20) that

$$S \le C \left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$
(6.21)

Combining (6.13), (6.14), (6.15) and (6.21), we have, for  $r \ge 1$ ,  $\{p \ge 2 \text{ and } s > 0\}$  or  $\{p \in [1, 2) \text{ and } s > 1/p\}$ ,

$$\mathbf{R}(\widehat{f}, f) \le C\left(\frac{\ln n_{\theta}}{n_{\theta}}\right)^{2s/(2s+1)}.$$

The proof of Theorem 5.1 is complete.

#### Christophe Chesneau

# References

- [1] A. Antoniadis, (1997). Wavelets in statistics: a review (with discussion), *Journal of the Italian Statistical Society* Series B, 6, 97-144.
- [2] A. Antoniadis, J. Bigot and T. Sapatinas, (2001). Wavelet estimators in nonparametric regression: a comparative simulation study, *Journal of Statistical Software*, **6**, i06.
- [3] A. Antoniadis, D. Leporini and J.-C. Pequet, (2002). Wavelet thresholding for some classes of non-Gaussian noise, *Statistica Neerlandica*, **56**, 4, 434-453.
- [4] R. Averkamp and C. Houdré, (2005). Wavelet thresholding for non-necessarily Gaussian noise: functionality, *The Annals of Statistics*, **33**, 2164-2193.
- [5] T. Cai, (1999). Adaptive wavelet estimation: a block thresholding and oracle inequality approach, *The Annals of Statistics*, **27**, 898-924.
- [6] T. Cai, (2002). On block thresholding in wavelet regression: adaptivity, block size and threshold level, *Statistica Sinica*, **12**, 1241-1273.
- [7] T. Cai and L. D. Brown, (1998). Wavelet shrinkage for nonequispaced samples, *The Annals of Statistics*, 26, 1783-1799.
- [8] T. Cai and L. D. Brown, (1999). Wavelet estimation for samples with random uniform design, *Statistics and Probability Letters*, **42**, 313-321.
- [9] C. Chesneau, (2007). Wavelet block thresholding for samples with random design: a minimax approach under the  $\mathbb{L}^p$  risk, *Electronic Journal of Statistics*, 1, 331-346.
- [10] E. Chicken, (2003). Block thresholding and wavelet estimation for nonequispaced samples, *Journal of Statistical Planning and Inference*, **116**, 113-129.
- [11] A. Cohen, I. Daubechies, B. Jawerth and P. Vial, (1993). Wavelets on the interval and fast wavelet transforms, *Applied and Computational Harmonic Analysis*, **24**, 1, 54-81.
- [12] Y. Davydov, (1970). The invariance principle for stationary processes. *Theor. Probab. Appl.*, 15, **3**, 498-509.
- [13] B. Delyon and A. Juditsky, (1996). On minimax wavelet estimators, *Applied Compu*tational Harmonic Analysis, 3, 215–228.
- [14] D. L. Donoho and I. M. Johnstone, (1994). Ideal spatial adaptation by wavelet shrinkage, *Biometrika* 81, 425-455.
- [15] D. L. Donoho and I. M. Johnstone, (1995). Adapting to unknown smoothness via wavelet shrinkage, *Journal of the American Statistical Association*, **90**, 432, 1200-1224.
- [16] D. L. Donoho and I. M. Johnstone, (1998). Minimax estimation via wavelet shrinkage, *The Annals of Statistics*, 26, 879-921.
- [17] D. L. Donoho, I. M. Johnstone, G. Kerkyacharian and D. Picard, (1995). Wavelet shrinkage: asymptopia (with discussion), *Journal of the Royal Statistical Society* Serie B, 57, 301-369.
- [18] H. Doosti, M. Afshari and H.A. Niroumand, (2008). Wavelets for nonparametric stochastic regression with mixing stochastic process. *Commun. Stat., Theory Methods*, 37, 3, 373-385.

- [19] H. Doosti and H.A. Niroumand, (2009). Multivariate Stochastic Regression Estimation by Wavelets for Stationary Time Series, *Pakistan journal of Statistics*, 27, 1, 37-46.
- [20] P. Doukhan, (1994). Mixing. Properties and Examples. Lecture Notes in Statistics 85. Springer Verlag, New York.
- [21] P. Hall and B. A. Turlach, (1997). Interpolation methods for nonlinear wavelet regression with irregularly spaced design, *The Annals of Statistics*, 25, 1912-1925.
- [22] W. Härdle, G. Kerkyacharian, D. Picard and A. Tsybakov, (1998). Wavelet, Approximation and Statistical Applications, Lectures Notes in Statistics New York 129, Springer Verlag.
- [23] G. Kerkyacharian and D. Picard, (2004). Regression in random design and warped wavelets, *Bernoulli*, **10**, (6), 1053-1105.
- [24] R. Kulik and M. Raimondo, (2009). L<sup>p</sup> wavelet regression with correlated errors and inverse problems, Statistica Sinica, 19, 1479-1489.
- [25] L. Li, J. Liu and Y. Xiao, (2009). On wavelet regression with long memory infinite moving average errors, *Journal of Applied Probability and Statistics*, 4, 183-211.
- [26] E. Masry, (2000). Wavelet-Based estimation of multivariate regression functions in besov spaces, *Journal of Nonparametric Statistics*, 12, 2, 283-308.
- [27] Y. Meyer, (1990). Ondelettes et Opérateurs, Hermann, Paris.
- [28] D. Modha and E. Masry, (1996). Minimum complexity regression estimation with weakly dependent observations. *IEEE Trans. Inform. Theory*, 42, 2133-2145.
- [29] M. Neumann and V. Spokoiny, (1995). On the efficiency of wavelet estimators under arbitrary error distributions, *Mathematical Methods of Statistics*, 4, 137-166.
- [30] M. Pensky and T. Sapatinas, (2007). Frequentist optimality of Bayes factor estimators in wavelet regression models, *Statistica Sinica*, 17, 599-633.
- [31] A. Tsybakov, (2004). Introduction à l'estimation nonparamétrique. Springer Verlag, Berlin.
- [32] C.S. Withers, (1981). Conditions for linear processes to be strong-mixing. Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete, 57, 477-480.
- [33] S. Zhang and Z. Zheng, (1999). Nonlinear wavelet estimation of regression function with random design. *Science in China Series A: Mathematics*, **42**, 8, 825-833.