



Cell cycle machinery and stroke

J. Rashidian, G.O. Iyirhiaro, D.S. Park

► To cite this version:

J. Rashidian, G.O. Iyirhiaro, D.S. Park. Cell cycle machinery and stroke. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, 2007, 1772 (4), pp.484. 10.1016/j.bbadis.2006.11.009 . hal-00562743

HAL Id: hal-00562743

<https://hal.science/hal-00562743>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript


Cell cycle machinery and stroke

J. Rashidian, G.O. Iyirhiaro, D.S. Park

PII: S0925-4439(06)00266-3
DOI: doi: [10.1016/j.bbadis.2006.11.009](https://doi.org/10.1016/j.bbadis.2006.11.009)
Reference: BBADIS 62670

To appear in: *BBA - Molecular Basis of Disease*

Received date: 30 August 2006
Revised date: 22 November 2006
Accepted date: 29 November 2006



Please cite this article as: J. Rashidian, G.O. Iyirhiaro, D.S. Park, Cell cycle machinery and stroke, *BBA - Molecular Basis of Disease* (2006), doi: [10.1016/j.bbadis.2006.11.009](https://doi.org/10.1016/j.bbadis.2006.11.009)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Cell Cycle Machinery and Stroke

J. Rashidian[†], G.O. Iyirhiaro[†] and D.S. Park^{*}

*Ottawa Health Research Institute, Neuroscience Group, Centre for Stroke Recovery, University of
Ottawa, K1H 8M5 Canada*

[†] J.R and G.O contributed equally to this review

Keywords: Stroke; CDKs; cyclins; cell cycle; apoptosis

***Correspondence author:**

David S. Park, Ph.D
Ottawa Health Research Institute – Neuroscience
University of Ottawa, 451 Smyth Road
Ottawa, ON, Canada
K1H 8M5
Telephone: (613) 562-5800 ext. 8269
Fax: (613) 562-5403
E-mail: dpark@uottawa.ca

Abstract

Stroke results from a transient or permanent reduction in blood flow to the brain. The mechanisms involving neuronal death following ischemic insult are complex and not fully understood. One signal which may control ischemic neuronal death is the inappropriate activation of cell cycle regulators including cyclins, cyclin dependent kinases (CDKs) and endogenous cyclin dependent kinase inhibitors (CDKIs). In dividing cells, activation of cell cycle machinery induces cell proliferation. In the context of terminally differentiated-neurons, however, aberrant activation of these elements triggers neuronal death. Indeed, there are several lines of correlative and functional evidence supporting this “cell cycle/neuronal death hypothesis”. The objective of this review is to summarize the findings implicating cell cycle machinery in ischemic neuronal death from in vitro and in vivo studies. Importantly, determining and blocking the signaling pathway(s) by which these molecules act to mediate ischemic neuronal death, in conjunction with other targets may provide a viable therapeutic strategy for stroke damage.

Introduction

Stroke occurs primarily as a result of a transient or permanent interruption of blood supply to the brain. This condition can stem from an occluded or ruptured blood vessels and in some cases cardiac arrest. Consequently, neurons in the affected brain region, or the whole brain in the case of a cardiac arrest, are deprived of oxygen and glucose. This sets in motion a cascade of cellular activities that ultimately culminate in neuronal cell death [1-3] .

Presently, stroke is a leading cause of death and permanent disability in industrialized nations. Stroke occurs on average every 45 seconds in the USA (www.strokecenter.org). The American Heart Association (AHA) estimates the cost of treating stroke related injury and disability at \$58 billion for 2006 alone (www.americanheart.org). Currently the treatment of stroke is mainly reliant on the use of thrombolytics such as tissue plasminogen activator (TPA), which themselves can pose an inherent risk of intracerebral hemorrhage. This limits the use of TPA to only certain cases of stroke. Furthermore, the efficacy of TPA depends on timely presentation of <3 hours which excludes 95% of stroke patients[4, 5] . Thus there is a need to develop new and efficacious neuroprotective strategies for the treatment of stroke.

The development of new strategies for stroke hinges on better understanding of the complex cellular and molecular interplay that ensue following stroke. A maelstrom of dysregulated molecules and potential perpetrators of ischemic neuronal death have been recently suggested. One of these exciting new developments involves the role of cell cycle molecules such as the cyclin dependent kinases. The notion that cell cycle machinery may mediate ischemic neuronal death is not unique to stroke. Indeed research evidence from numerous labs have demonstrated correlative relationship between the dysregulation of cell cycle machinery and neuronal death models of neurodegenerative diseases such as Parkinson's disease (PD) [6, 7] , Alzheimer's disease (AD) [8-10], Amyotrophic

Lateral Sclerosis (ALS) [11, 12], and Niemann-Pick type C disease [13, 14] . Whether or not cell cycle mechanism is similar in neurodegenerative diseases and acute conditions such stroke is unknown. This review will focus only on evidence implicating cell cycle molecules in ischemic neuronal death. We will also briefly discuss the potential involvement of other members of the CDK family not directly involved in cell cycle control, but only as potential activators of cell cycle machinery in stroke.

Ischemic Neuronal Death

The mechanism(s) of ischemic neuronal death is complex and may be determined by factors such as the location, severity and duration of insult. For example, neuronal death in the ischemic core, the region most severely affected by the lack of blood flow, occurs within minutes to a few hours and is marked predominantly by necrotic and excitotoxic cell death . Neuronal death in the ischemic penumbra, the region less severely affected by ischemia, is marked mainly by delayed apoptotic-like death that can progress over a period of days.

The precise sequence of events leading to neuronal death following stroke is at present not fully defined. However, a core picture is developing. Following the disruption of blood flow, the affected brain region(s) undergo a period of hypoxia resulting in a decrease in cellular ATP, due primarily to impaired mitochondrial oxidative respiration, a process that generates majority of the cellular energy required to maintain proper cell functions. The reduction of cellular energy results in the impairment of vital cellular functions such as the maintenance of the Na⁺/K⁺ pump, massive depolarization and excessive release of glutamate. Over activation of the glutamate channel particularly the N-methyl-D-aspartate receptor (NMDAR)-type channels mediates massive influx of Ca²⁺ resulting in cellular excitotoxicity and neuronal death. Increase in intracellular free calcium results in the activation of Ca²⁺ sensitive enzymes such as calpains proteases which then act to activate other molecules and

cleave cellular structures. In addition to the events described above other stressors such as oxidative stress, and DNA damage have been demonstrated to mediate ischemic neuronal death. Furthermore, extrinsic stressors such as the activation of glial cells and inflammation may impinge on ischemic neurons to mediate their demise. These later findings are particularly relevant when it comes to therapeutic interventions. In fact, cell cycle regulation may also play a critical role with these non neuronal cell types in brain injury [15, 16]. However, we will not mention this further in this review, but will instead focus on how the cell cycle machinery may impact neurons more directly.

Cell cycle regulation

Cell cycle is a highly regulated process. Timing progression of cell cycle through different phases, G₀, G₁, S, G₂, and M requires an orchestrated functions of several elements, including cyclins, cyclin-dependent kinases (CDKs), retinoblastoma protein (Rb; pocket proteins) and E2F complex proteins [17]. Different complexes of cyclin-CDK drive each phase of cell cycle [18]. In this regard, the current model is that cyclin D-cdk4/6 and cyclin E-cdk2 complexes regulate G₁/S progression, cyclin A-cdk2 complexes mediate S/G₂ transitions, and cyclin B-cdc2 complexes mediate M-phase progression [19, 20]. In addition, a recent report has suggested that cyclin C-cdk3 complexes also regulate G₀/G₁ transition [21] (Figure1). CDKs activity is regulated by binding to activating cyclin partners [22] as well as endogenous CDK inhibitors such as members of the INK4 (p15, p16, p18, p19) and Cip/Kip (p21, p27, p57) families [23, 24]. Finally, phosphorylation also plays a critical role in CDK regulation. For example, cyclin H-cdk7-Mat1 (Cdk Activating Kinase; CAK complex) by phosphorylation of a central threonine [25] and cdc25 by dephosphorylation of thr14 (on cdc2) or tyr15 (on cdk2) modulate CDKs conformation to a fully active structure so they recognize their substrates more efficiently [23]. CDKs mediate cell cycle progression by phosphorylating downstream targets [26]. For example, an important target of G₁ CDKs is the

tumor suppressor, Rb [27]. In mid- to late-G1, Rb is sequentially phosphorylated by cyclin D1-cdk4/6 and cyclin E-cdk2 complexes. Once hyperphosphorylated, Rb repression of E2F containing complexes is alleviated. E2F, along with a required binding partner DP activate genes required for S phase progression [28-30].

A growing body of exciting evidence indicates that CDKs may have a function beyond its canonical role in cell cycle regulation. Indeed, it has been suggested that neuronal death may be controlled by the inappropriate activation of some of cell cycle elements [31]. The hypothesis, compatible with the general cell cycle/neuronal death notion, is that activation of these regulatory elements, in the context of post-mitotic neurons, signals for death rather than cell cycle progression. Several lines of in vitro and in vivo evidence support this hypothesis that aberrant activation of cell cycle involves in different pathological conditions such as stroke [32-34], AD [35], PD [6, 7], ALS [11].

In this review, we will first start our discussion with deregulation/upregulation of cell cycle regulators in in vitro paradigms of ischemic neuronal death and then will provide in vivo evidence suggesting that cell cycle might control death following ischemic neuronal injury in the adult animal. Finally, we will touch upon the role of other CDKs, in particular the neuronal cdk5 which has important roles in neuronal development and death and which might also link to cell cycle signals.

In vitro evidence for involvement of cell cycle elements in stroke

Generally, terminally differentiated cells such as neuron do not divide. Indeed as neurons undertake the process of terminal differentiation they irreversibly withdraw from the cell cycle. The levels and activity of key cell cycle regulators are downregulated in differentiated neurons [36, 37]. For example, Cdk4, a regulator of G1/S transition becomes increasingly associated with the CDK inhibitor p27 and its activity declines in parts due to loss of phosphorylation by its activator CAK

[36]. Similarly, the levels and activity of cdk2 another important regulator of the G1/S is downregulated in differentiated neurons. Consequently, the levels of hypophosphorylated Rb increases resulting in greater E2F1 sequestration [36]. Silencing E2F1 in neurons is particularly important since its overexpression promotes aberrant S-phase entry in neurons [38]. Furthermore, overexpression of E2F1 alone in the context of a post-mitotic neuron and in the absence of insult induces neuronal death [38, 39]. Downregulation of cell cycle machinery and inhibition of E2F1 activity through its sequestration by Rb is thus an important feature of terminally differentiated neurons. This is underscored by observations that overexpression of Rb or the CDK inhibitor p27 is sufficient to induce neuronal differentiation [36]. In addition to downregulation of CDK levels and activity, observations by Sumrejkanchanakij et al. [40] suggests that the cell cycle may be held in check in post-mitotic neurons through cellular redistribution of cell cycle components. For example, in undifferentiated neurons cyclin D1 appears both cytoplasmic and nuclear but becomes predominantly cytoplasmic in differentiated neuroblastoma and cortical neurons [37, 40]. Ectopic expression of cyclin D1 in differentiated neurons leads to cytoplasmic sequestration whereas forced nuclear expression by fusing to a nuclear localization signal induces apoptosis [37]. This suggests that cellular redistribution of cell cycle machinery is an important aspect in the maintenance of quiescent state in adult neurons. While the above evidence indicates what happens under normal developmental conditions, accumulating research evidence also suggests that cell cycle machinery are activated in adult neurons in response to numerous stressors such as DNA damage, NGF deprivation, proteasome inhibition and β -amyloid toxicity and models of ischemic neuronal death [39, 41-47].

In regards to the latter type of injury, multiple lines of evidence in vitro suggest a role for cell cycle regulators as mediators of ischemic injury. Firstly, Katchanov and colleagues [42] reported loss of the CDK inhibitor p27 following oxygen glucose deprivation (OGD) of neocortical neurons. This

group also reported increases in cyclin D1 protein levels and activation of cdk2 following OGD [42]. Similar increase in cyclin D1, cyclin E, and cdk2 is observed following kainic acid (KA) induced death in cerebellar granule neurons (CGNs) [48, 49]. Secondly, Rb, a downstream target of cyclinD/Cdk4 is increasingly phosphorylated following hypoxia/reoxygenation and KA induced death [34, 50] suggesting inactivation of Rb and activation of the cell cycle in these neurons. Additionally, increase in E2F1 (downstream effector of Cdk4) protein levels and proliferating cell nuclear antigen (PCNA) another S-phase marker (required for completion of DNA synthesis and repair) is observed following KA induced death [48, 51]. Finally, increase in E2F1 mRNA transcript is also observed following OGD [52]. These observations discussed above seem to implicate the reactivation of cell cycle machinery in ischemic neuronal death and have lead to the hypothesis that inappropriate activation of the cell cycle in the context of differentiated neurons may underlie ischemic neuronal death. However, this evidence only showed the activation of cell cycle components and does not address the issue of whether this signal is required for death. In other words, does the activation of the cell cycle in adult neurons underlie neuronal death in stroke or are these observations simply an artifact of the cell death process? Studies using pharmacological inhibitors of cell cycle regulators seem to lend support to the former. For example, treatment of neurons with the CDK inhibitors olomoucine, roscovitine, 3-amino thioacridone (3-ATA) and flavopiridol has been shown to protect cortical neurons and CGNs from OGD and KA induced cell death [42, 47, 51, 53]. While these pharmacological studies supports the involvement of cell cycle components in ischemic neuronal death, the caveat to these studies is the relatively non-specific nature of inhibitors such as flavopiridol which in addition to the cell cycle CDKs can potently inhibit non-mitotic CDKs such as cdk5 [54], cdk7 [25] as well as non-CDK-related kinases such as GSK-3 β [55]. To address this, the functional relevance of cell cycle machinery in ischemic neuronal death was demonstrated by utilizing genetic manipulations of components of the cell cycle. Firstly,

Rashidian et al. [34] showed that adenoviral delivered kinase dead cdk4 protects CGNs from hypoxia mediated delayed death. Secondly, they showed that neurons derived from mice expressing kinase dead cdk4 or null for its regulator cyclin D1 are resistant to hypoxia mediated ischemic death [34] . Further support for the functional role of cell cycle pathway in ischemic neuronal death is provided by observations that cortical neurons and CGNs derived from E2F1 null mice are less susceptible to death mediated by OGD and KA induced death respectively [50, 52] . Furthermore, E2F1 deficiency improves the recovering of CA1 neurons from loss of synaptic transmission following anoxic insult of hippocampal slices in vitro [52] . The above in vitro evidence is summarized in tables 1 and 2.

Taken together these in vitro evidence strongly implicate the reactivation of cell cycle components in ischemic neuronal death. What evidence is there to implicate the involvement of cell cycle machinery in vivo? Are these observations relevant in intact physiological systems?

In vivo evidence for involvement of cell cycle elements in stroke

The above evidence indicates that neurons grown in culture can re-activate cell cycle signals upon an exogenous stress leading to death. This suggests that terminally differentiated, post mitotic neurons from embryonic or early post natal sources have retained their capacity to reactivate cell cycle CDK mediated signals. However, these neurons have only recently become post-mitotic and might represent a unique situation where the cell cycle machinery is still present. Indeed, reports have shown that while downregulated, multiple CDKs, cyclins as well as there regulators exist in cultured neurons [36, 37] . Questions remain about the more relevant situation where ischemic injury occurs in fully matured adult neurons where there is likely very little if any cell cycle machinery present under basal conditions. Here too, there is accumulating evidence that upregulation/activation of cell cycle machinery occurs and this leads to death in neurons following stroke insult.

The first body of work linking cell cycle to ischemic injury in the adult context is the correlative data demonstrating that CDKs are changed following stroke injury *in vivo*. In these situations, a central player may again be the cdk4-cyclin D complex. For example, Cdk4/6 kinase activity/levels along with its activator cyclin D1 are upregulated in different models of ischemia *in vivo*. It has been shown that cyclin D1 is expressed in infarct region following MCAO model of focal ischemia [32, 42, 56-58] and in vulnerable regions of the hippocampus following transient global ischemia [59-63]. Consistent with these observations, increased levels of cyclin D1 has been demonstrated in response to an excitotoxic death induced by systemic injection of KA [61, 64-66] and transient spinal cord ischemia [67]. Cdk4 itself is also changed during ischemic insult. Aberrant expression of cdk4 has been shown in MCAO [68] and clip models of focal ischemia [32]. Furthermore, other groups have reported induction of cdk4 expression in other models such as transient global ischemia [63], KA-mediated cortical death [66] and transient spinal cord ischemia [67].

Although the above evidence implicates cyclinD1-cdk4 complexes in ischemic injury, it must be kept in mind that other components of G1 phase might have potential roles *in vivo* as well. For instance, cdk6 is also activated by cyclin D [17]. In addition, there are some lines of data suggesting deregulation of cdk2 after MCAO and a hypoxic cerebral ischemia [42, 69]. Consistent with this, Li et al. [70] provided evidence that cdk2, as well as its partner, cyclin A, are deregulated following focal ischemia.

In line with the upregulation/activation of cell cycle CDK complexes, evidence also point to a modulation of endogenous regulators of CDKs, the CDKIs. For example, Katchanov et al. [42] demonstrated that p16 is downregulated very early in dying neurons, followed by upregulation of cyclin D1 after a mild cerebral ischemia. Additional intriguing observations suggesting that CDKIs might have a functional role in ischemic death were provided by Van Lookeren et al. [71, 72]. They

indicated that p21 expression is enhanced in surviving neurons surrounding dead area in both focal and transient forebrain ischemia. They suggested that elevation of p21 is actually a response to the ischemic insult to protect the neurons by arresting them in first steps of the cell cycle. In line with these findings, p21 mRNA was elevated in perifocal ischemia regions following focal ischemia, while no significant change was observed in ischemic area [73]. More importantly, Kuan et al. [69] demonstrated that p16 and p27 are depleted in dying hippocampal neurons and this may promote G1/S phase transition following hypoxia-ischemia insult.

The above mentioned in vivo evidence does not indicate whether these signals actually play a role in ischemic damage. To address this, several labs, including ours, have provided more direct data by carrying out additional functional studies. In support of a require role, inhibition of the cell cycle regulating proteins is protective in some of the ischemic models. For instance, administration of a synthetic cdk4 inhibitor, or alternatively, cyclin D1 and cdk4 antisense oligonucleotides has been shown to be neuroprotective in MCAO and KA-inducing excitotoxicity models, respectively [58, 66]. We have recently shown that flavopiridol, a CDK inhibitor, administrated before or after the insult, blocks the delayed death evoked by focal or global ischemia [32, 33]. Importantly, flavopiridol blocked activation of the cdk-Rb pathway, reduced E2F1 levels and improved behavioral performance [32, 33]. More critically, we also investigated involvement of specific CDKs by expression of kinase dead dominant negative forms of several cell cycle CDKs, as well as cdk5, in delayed and excitotoxic types of ischemia. In this research for the first time we provided functional evidence that in adult system, cyclin D1-cdk4 complex has a key role in inducing delayed neuronal death [34]. We also showed that phosphorylation of Rb, mediated by this complex, occurs very early following global ischemia and that this activation is inhibited by DN cdk4 expression [34]. Expression of DN cdk4 also improves behavioral performance in rats suggesting that brains are not

only neuroprotected, but that this leads to improved function [34] . This final point is particularly relevant to the goal of human therapy.

How do cyclin-CDK complexes mediate a potential ischemic death signal, *in vivo*? In other words, what is (are) the downstream target(s) for these complexes in adult systems? Recent findings suggest that inactivation of Rb leads to neuronal apoptosis, *in vivo*. Supporting this, overexpression of large T antigen in cerebellar Purkinje cells initially resulted in DNA synthesis but eventually caused neuronal degeneration in transgenic mice [74]. Rb deficiency is also embryonically lethal because of widespread apoptosis in CNS and PNS [75, 76]. However, in this case, much of the death may be due to non neuron autonomous events and not directly caused by Rb deficiency [77]. Observations in different *in vivo* models of ischemia have suggested that CDKs activity induced by apoptotic insults lead to phosphorylation and inactivation of Rb. For example, we have shown that Rb phosphorylation (on ser-795, a consensus site for CDKs) is an early that event happens a few hours after focal ischemia [32], global ischemia [33, 34] and KA injection [65]. Moreover, other groups have also shown that Rb is progressively phosphorylated at CDKs consensus sites after focal ischemia induced by MCAO and hypoxic cerebral ischemia and that this might trigger apoptotic neuronal death pathways [58, 69, 78].

To investigate the mechanism by which Rb inactivation may promote death, E2F family proteins, as a group of proteins regulated by Rb, have been studied *in vivo*. In this regard, it has been suggested that E2F may play an important role in neuronal death in select conditions such as stroke. For example, E2F1 deficient mice has been shown to exhibit smaller brain injury and improved behavior after focal ischemia [79, 80] . In agreement with this, upregulation of E2F1 has been implicated in animal models of ischemia. For instance, we have shown that focal ischemia induces E2F1 expression which occurs following Rb phosphorylation and administration of CDK inhibitor diminishes this effect and protects the neurons [32]. Microarray analysis also has also revealed that

E2F1 is among the upregulated genes after global ischemia [81]. Finally, a very recent study demonstrated that E2F1 and its target gene, c-myc, is upregulated in hippocampus following global ischemia [82]. What factor(s) determine which responsive genes, whether cell cycle or apoptotic, be regulated? It is not clear yet, but there is an idea suggesting that there is a pool of transcriptional activity for E2F. When this activity gets to a specific threshold the genes regulating cell cycle progression are induced. But when this activity reaches a second higher threshold, apoptotic genes are induced and death mechanisms start [83]. Almost nothing is known about the targets of E2F1 in ischemic death, in vivo, and this area needs to be intensively investigated.

Although what has been discussed so far suggests that cell cycle is aborted mostly at G1/S checkpoint in ischemic death, some reports have shown S phase entry. Indeed, neurons start to replicate DNA before they die in some systems. To investigate whether DNA synthesis is induced in dying neurons, induction of PCNA and incorporation of Brdu have been used as markers for S phase. In this regard, induction of PCNA has been demonstrated in several models of focal and global ischemia [58, 63, 70, 84]. In other reports, incorporation of Brdu in apoptotic neurons is indicative of tendency for DNA synthesis in these cells [58, 69]. Interestingly, these injured, “S phase positive” neurons contained elevated G1 cyclin-CDK complexes and co-localized with apoptotic markers (TUNEL positive); suggesting that cell cycle is already triggered and they have transited to S phase to resume DNA synthesis but for some reasons they undergo apoptosis and do not replicate. Therefore, there is no in vivo evidence of G2 entry in ischemic neurons, to our knowledge. Nevertheless, expression of cdc2, as E2F target and a G2 marker has been investigated in other neuronal death systems [85]. This might be an important area of future research in the stroke field.

In spite of the accumulating in vivo experimental studies demonstrating aberrant deregulation of cell cycle in stroke, there is little known about how/whether these processes occur in humans. Recently, a group investigated cell cycle elements in brains of postmortem patients who died after

cardiac arrest or focal brain infarction. They reported elevated levels of cyclin D1, cdk2 and cdk4 [86]. This study shows that results obtained from experimental studies in animals may be relevant to the human condition.

The in vivo evidence discussed above is summarized in tables 1 and 2.

Role of other CDK members including cdk5 in stroke

In addition to the core cell cycle components discussed above, other cyclins and CDK members which are not exclusively or fully linked with cell cycle regulation may also participate in ischemic response. For example, cyclin H, a component of the CAK complex is increased in CA3, dentate gyrus, and cortex after global ischemia [87]. Since these areas are resistant to this type of ischemia, it has been suggested that cyclin H might have a role in survival through its DNA repair function in ischemic injury. Cyclin G is another member of the cyclins family but its role in cell cycle is controversial. While some evidence shows its stimulating role in growth [88, 89], others suggesting it inhibits growth [90, 91] in proliferating cells. The function of cyclin G in neurons is also unclear. Nevertheless, it has been shown that cyclin G1 is increased following MCAO and global ischemia [71, 72], as well as spinal cord injury [92]. Furthermore, Maeda et al. [93] demonstrated that cyclin G1 is translocated into the nucleus of degenerating neurons following transient common carotid artery occlusion (CCAO) and MCAO. Therefore, they suggested that the cytoplasmic cyclin G1 might be associated with cell survival while nuclear cyclin G1 might signal death.

Perhaps the strongest evidence for the involvement of a CDK member not classically associated with the cell cycle involves cdk5. Cdk5 is a unique member of cyclin-dependent kinases family. Unlike the other CDKs, it does not require a cyclin to be activated. P35 and p39, instead, are activators of cdk5 and are present mainly in neurons [94, 95]. For this reason, although is ubiquitously expressed, cdk5 activity is mainly restricted to nervous system [96]. Even though there

is no functional evidence for involvement of cdk5 directly in the core cell cycle machinery, some recent observations have led to the idea that it might be associated with cell cycle components and, therefore, affects its transition. Initially it was shown that cdk5 was able to bind to cyclins D1, D3 and E [97-99]. However, this binding did not lead to increased CDK activity. Further studies showed that cdk5 could phosphorylate Rb in vitro [100]. In this study, overexpression of p25 in inducible- neuroblastoma cells led to deregulation of cell cycle elements such as elevation of cyclins A and B1 and cdc2, downregulation of p27 and finally apoptosis. Interestingly, Rb phosphorylation on ser807/811 sites was an early event in this study and was mediated by cdk5 activity since it was abolished by a cdk5 inhibitor [100]. Finally, a very recent report revealed that cdk5 is necessary for neuronal cell cycle arrest[101] . This group suggested that cdk5 connects and inhibits cell cycle possibly by competing with cdk4/6 for binding to cyclins or through phosphorylation and inhibition of Rb or, alternatively, by phosphorylation of RNA polymerase.

Cdk5 has been implicated in different neurodegenerative disorders including brain ischemia. The generally accepted concept about cdk5 is that cdk5/p35 complexes are regulators of normal biological functions of neuron. But under some neuropathological conditions p35 is cleaved to a more potent activator, p25 [102]. Production of p25 unmasks pathological face of cdk5 since it has prolonged half life, over activates, dislocates and might change cdk5 substrate specificity. There is increasing evidence, though, that argue this model. Several studies have debated role of p25 in pathogenicity of cdk5 in some circumstances [103-107] . While an exhaustive review of this issue is not possible here, we will suggest that, depending upon different factors; both p35 and p25 are able to drive “good-normal” cdk5 to “bad-pathogenic” cdk5.

Contribution of cdk5 in stroke has not been intensively studied and our knowledge about desregulated cdk5 in stroke comes from very recent research. One of the initial reports came from Hayashi et al. study [108] . They showed that immunoreactivity of cdk5 and p35/p25 is enhanced in

MCAO model. Likewise, we [34] as well as Wang et al. [109], have recently provided evidence that cdk5 is a mediator of excitotoxic neuronal death in global and focal ischemia.

Similar to classic cell cycle CDKs, the important question of the mechanism by which aberrant cdk5 causes neuronal damage remains. Several possibilities may exist. For example, it has been reported that cdk5 can phosphorylate the NMDA receptors regulating calcium influx [109]. Interestingly tau, a microtubule-associated protein which is connected to Alzheimer's disease pathology, has also been recently shown to be as a substrate for cdk5 following ischemia [110]. What this means is unknown, however. There might also be other candidates such as MEF2 (Myocyte Enhance Factor-2) or p53 transcription factors. Both have been shown to be regulated by cdk5 and both have been implicated in ischemic/excitotoxic death [111-113]. However, the link between cdk5 and these factors have not been yet addressed in in vivo stroke studies.

Conclusion:

As summarized above, there is increasing evidence of the involvement of multiple cell cycle regulatory signals in ischemic injury. Several important questions remain however, and include: 1) What are the mechanism(s) by which cell cycle signals are actually activated following stroke injury; 2) Are there other mechanisms by which cell cycle CDKs may impact death other than through modulation of Rb; 3) Does cdk5 regulate cell cycle signals in models of stroke 4) Are cell cycle targets in and of themselves a viable therapeutic option for stroke.

The latter question is of particular importance since most pharmacotherapeutic strategies tested in the clinic for treatment of stroke have failed. One reason for this is likely that preclinical testing in animal models has not been sufficiently stringent prior to clinical trials. In this regard, vigorous benchmarks of a) efficacy of any strategy in multiple animal models, b) behavioral and functional improvements beyond neuroprotection and c) long lasting efficacy have been established. Cell cycle

inhibitors appear to at least meet some of these criteria. For example, the CDK inhibition has been shown to be neuroprotective in multiple models and lead to improved behavior/function. However, the CDK inhibitor flavopiridol, at least, seems to lack long term efficacy. This may be due to the non selective nature of flavopiridol or and indication that CDK inhibition by itself can not completely normalize neuron function. These issues will have to be carefully addressed before CDK inhibitors can be contemplated for future clinical trials.

Acknowledgements:

This review was supported by funds from Heart and Stroke of Ontario, Canadian Institute for Health Research, The Centre for Brain Recovery, and Canadian Stroke Network.

References

- [1] U. Dirnagl, C. Iadecola and M. A. Moskowitz, Pathobiology of ischaemic stroke: An integrated view, *Trends Neurosci* 22 (1999) 391-7.
- [2] M. Endres and U. Dirnagl, Ischemia and stroke, *Adv Exp Med Biol* 513 (2002) 455-73.
- [3] M. Nedergaard and U. Dirnagl, Role of glial cells in cerebral ischemia, *Glia* 50 (2005) 281-6.
- [4] P. A. Barber, J. Zhang, A. M. Demchuk, M. D. Hill and A. M. Buchan, Why are stroke patients excluded from tpa therapy? An analysis of patient eligibility, *Neurology* 56 (2001) 1015-20.
- [5] M. O'Hare, F. Wang and D. S. Park, Cyclin-dependent kinases as potential targets to improve stroke outcome, *Pharmacol Ther* 93 (2002) 135-43.
- [6] B. F. El-Khodir, T. F. Oo, N. Kholodilov and R. E. Burke, Ectopic expression of cell cycle markers in models of induced programmed cell death in dopamine neurons of the rat substantia nigra pars compacta, *Exp Neurol* 179 (2003) 17-27.
- [7] K. L. Jordan-Sciutto, R. Dorsey, E. M. Chalovich, R. R. Hammond and C. L. Achim, Expression patterns of retinoblastoma protein in parkinson disease, *J Neuropathol Exp Neurol* 62 (2003) 68-74.
- [8] N. G. Milton, The amyloid-beta peptide binds to cyclin b1 and increases human cyclin-dependent kinase-1 activity, *Neurosci Lett* 322 (2002) 131-3.
- [9] Z. Nagy, M. M. Esiri and A. D. Smith, Expression of cell division markers in the hippocampus in alzheimer's disease and other neurodegenerative conditions, *Acta Neuropathol (Berl)* 93 (1997) 294-300.
- [10] A. McShea, P. L. Harris, K. R. Webster, A. F. Wahl and M. A. Smith, Abnormal expression of the cell cycle regulators p16 and cdk4 in alzheimer's disease, *Am J Pathol* 150 (1997) 1933-9.
- [11] M. D. Nguyen, M. Boudreau, J. Kriz, S. Couillard-Despres, D. R. Kaplan and J. P. Julien, Cell cycle regulators in the neuronal death pathway of amyotrophic lateral sclerosis caused by mutant superoxide dismutase 1, *J Neurosci* 23 (2003) 2131-40.
- [12] S. Ranganathan, S. Scudiere and R. Bowser, Hyperphosphorylation of the retinoblastoma gene product and altered subcellular distribution of e2f-1 during alzheimer's disease and amyotrophic lateral sclerosis, *J Alzheimers Dis* 3 (2001) 377-385.
- [13] J. W. Husseman, D. Nochlin and I. Vincent, Mitotic activation: A convergent mechanism for a cohort of neurodegenerative diseases, *Neurobiol Aging* 21 (2000) 815-28.
- [14] M. Zhang, J. Li, P. Chakrabarty, B. Bu and I. Vincent, Cyclin-dependent kinase inhibitors attenuate protein hyperphosphorylation, cytoskeletal lesion formation, and motor defects in niemann-pick type c mice, *Am J Pathol* 165 (2004) 843-53.
- [15] I. Cernak, B. Stoica, K. R. Byrnes, S. Di Giovanni and A. I. Faden, Role of the cell cycle in the pathobiology of central nervous system trauma, *Cell Cycle* 4 (2005) 1286-93.
- [16] S. Di Giovanni, V. Movsesyan, F. Ahmed, I. Cernak, S. Schinelli, B. Stoica and A. I. Faden, Cell cycle inhibition provides neuroprotection and reduces glial proliferation and scar formation after traumatic brain injury, *Proc Natl Acad Sci U S A* 102 (2005) 8333-8.
- [17] J. Pines, Cyclins and their associated cyclin-dependent kinases in the human cell cycle, *Biochem Soc Trans* 21 (1993) 921-5.
- [18] P. Nurse, A long twentieth century of the cell cycle and beyond, *Cell* 100 (2000) 71-8.

- [19] S. V. Ekholm and S. I. Reed, Regulation of g(1) cyclin-dependent kinases in the mammalian cell cycle, *Curr Opin Cell Biol* 12 (2000) 676-84.
- [20] J. Pines, Cyclins and cyclin-dependent kinases: A biochemical view, *Biochem J* 308 (Pt 3) (1995) 697-711.
- [21] S. Ren and B. J. Rollins, Cyclin c/cdk3 promotes rb-dependent g0 exit, *Cell* 117 (2004) 239-51.
- [22] E. Lees, Cyclin dependent kinase regulation, *Curr Opin Cell Biol* 7 (1995) 773-80.
- [23] D. O. Morgan, Principles of cdk regulation, *Nature* 374 (1995) 131-4.
- [24] C. J. Sherr and J. M. Roberts, Cdk inhibitors: Positive and negative regulators of g1-phase progression, *Genes Dev* 13 (1999) 1501-12.
- [25] R. P. Fisher and D. O. Morgan, A novel cyclin associates with mo15/cdk7 to form the cdk-activating kinase, *Cell* 78 (1994) 713-24.
- [26] D. O. Morgan, Cyclin-dependent kinases: Engines, clocks, and microprocessors, *Annu Rev Cell Dev Biol* 13 (1997) 261-91.
- [27] R. A. Weinberg, The retinoblastoma protein and cell cycle control, *Cell* 81 (1995) 323-30.
- [28] S. P. Chellappan, S. Hiebert, M. Mudryj, J. M. Horowitz and J. R. Nevins, The e2f transcription factor is a cellular target for the rb protein, *Cell* 65 (1991) 1053-61.
- [29] N. Dyson, The regulation of e2f by prb-family proteins, *Genes Dev* 12 (1998) 2245-62.
- [30] J. W. Harbour, R. X. Luo, A. Dei Santi, A. A. Postigo and D. C. Dean, Cdk phosphorylation triggers sequential intramolecular interactions that progressively block rb functions as cells move through g1, *Cell* 98 (1999) 859-69.
- [31] E. B. Becker and A. Bonni, Beyond proliferation--cell cycle control of neuronal survival and differentiation in the developing mammalian brain, *Semin Cell Dev Biol* 16 (2005) 439-48.
- [32] H. Osuga, S. Osuga, F. Wang, R. Fetni, M. J. Hogan, R. S. Slack, A. M. Hakim, J. E. Ikeda and D. S. Park, Cyclin-dependent kinases as a therapeutic target for stroke, *Proc Natl Acad Sci U S A* 97 (2000) 10254-9.
- [33] F. Wang, D. Corbett, H. Osuga, S. Osuga, J. E. Ikeda, R. S. Slack, M. J. Hogan, A. M. Hakim and D. S. Park, Inhibition of cyclin-dependent kinases improves ca1 neuronal survival and behavioral performance after global ischemia in the rat, *J Cereb Blood Flow Metab* 22 (2002) 171-82.
- [34] J. Rashidian, G. Iyirhiaro, H. Aleyasin, M. Rios, I. Vincent, S. Callaghan, R. J. Bland, R. S. Slack, M. J. During and D. S. Park, Multiple cyclin-dependent kinases signals are critical mediators of ischemia/hypoxic neuronal death in vitro and in vivo, *Proc Natl Acad Sci U S A* 102 (2005) 14080-5.
- [35] K. Herrup and T. Arendt, Re-expression of cell cycle proteins induces neuronal cell death during alzheimer's disease, *J Alzheimers Dis* 4 (2002) 243-7.
- [36] O. Kranenburg, V. Scharnhorst, A. J. Van der Eb and A. Zantema, Inhibition of cyclin-dependent kinase activity triggers neuronal differentiation of mouse neuroblastoma cells, *J Cell Biol* 131 (1995) 227-34.
- [37] P. Sumrejkanchanakij, M. Tamamori-Adachi, Y. Matsunaga, K. Eto and M. A. Ikeda, Role of cyclin d1 cytoplasmic sequestration in the survival of postmitotic neurons, *Oncogene* 22 (2003) 8723-30.
- [38] M. Azuma-Hara, H. Taniura, T. Uetsuki, M. Niinobe and K. Yoshikawa, Regulation and deregulation of e2f1 in postmitotic neurons differentiated from embryonal carcinoma p19 cells, *Exp Cell Res* 251 (1999) 442-51.

- [39] A. Giovanni, E. Keramaris, E. J. Morris, S. T. Hou, M. O'Hare, N. Dyson, G. S. Robertson, R. S. Slack and D. S. Park, E2f1 mediates death of b-amyloid-treated cortical neurons in a manner independent of p53 and dependent on bax and caspase 3, *J Biol Chem* 275 (2000) 11553-60.
- [40] P. Sumrejkanchanakij, K. Eto and M. A. Ikeda, Cytoplasmic sequestration of cyclin d1 associated with cell cycle withdrawal of neuroblastoma cells, *Biochem Biophys Res Commun* 340 (2006) 302-8.
- [41] A. Giovanni, F. Wirtz-Brugger, E. Keramaris, R. Slack and D. S. Park, Involvement of cell cycle elements, cyclin-dependent kinases, prb, and e2f x dp, in b-amyloid-induced neuronal death, *J Biol Chem* 274 (1999) 19011-6.
- [42] J. Katchanov, C. Harms, K. Gertz, L. Hauck, C. Waeber, L. Hirt, J. Priller, R. von Harsdorf, W. Bruck, H. Hortnagl, U. Dirnagl, P. G. Bhidé and M. Endres, Mild cerebral ischemia induces loss of cyclin-dependent kinase inhibitors and activation of cell cycle machinery before delayed neuronal cell death, *J Neurosci* 21 (2001) 5045-53.
- [43] D. S. Park, S. E. Farinelli and L. A. Greene, Inhibitors of cyclin-dependent kinases promote survival of post-mitotic neuronally differentiated pc12 cells and sympathetic neurons, *J Biol Chem* 271 (1996) 8161-9.
- [44] D. S. Park, B. Levine, G. Ferrari and L. A. Greene, Cyclin dependent kinase inhibitors and dominant negative cyclin dependent kinase 4 and 6 promote survival of ngf-deprived sympathetic neurons, *J Neurosci* 17 (1997) 8975-83.
- [45] D. S. Park, E. J. Morris, L. A. Greene and H. M. Geller, G1/s cell cycle blockers and inhibitors of cyclin-dependent kinases suppress camptothecin-induced neuronal apoptosis, *J Neurosci* 17 (1997) 1256-70.
- [46] H. J. Rideout, Q. Wang, D. S. Park and L. Stefanis, Cyclin-dependent kinase activity is required for apoptotic death but not inclusion formation in cortical neurons after proteasomal inhibition, *J Neurosci* 23 (2003) 1237-45.
- [47] E. Verdaguer, E. G. Jorda, A. Stranges, A. M. Canudas, A. Jimenez, F. X. Sureda, M. Pallas and A. Camins, Inhibition of cdks: A strategy for preventing kainic acid-induced apoptosis in neurons, *Ann N Y Acad Sci* 1010 (2003) 671-4.
- [48] E. Verdaguer, E. Garcia-Jorda, A. M. Canudas, E. Dominguez, A. Jimenez, D. Pubill, E. Escubedo, J. C. Pallas and A. Camins, Kainic acid-induced apoptosis in cerebellar granule neurons: An attempt at cell cycle re-entry, *Neuroreport* 13 (2002) 413-6.
- [49] S. F. Giardina, N. S. Cheung, M. T. Reid and P. M. Beart, Kainate-induced apoptosis in cultured murine cerebellar granule cells elevates expression of the cell cycle gene cyclin d1, *J Neurochem* 71 (1998) 1325-8.
- [50] R. A. Smith, T. Walker, X. Xie and S. T. Hou, Involvement of the transcription factor e2f1/rb in kainic acid-induced death of murine cerebellar granule cells, *Brain Res Mol Brain Res* 116 (2003) 70-9.
- [51] E. Verdaguer, A. Jimenez, A. M. Canudas, E. G. Jorda, F. X. Sureda, M. Pallas and A. Camins, Inhibition of cell cycle pathway by flavopiridol promotes survival of cerebellar granule cells after an excitotoxic treatment, *J Pharmacol Exp Ther* 308 (2004) 609-16.
- [52] T. F. Gendron, G. A. Mealing, J. Paris, A. Lou, A. Edwards, S. T. Hou, J. P. MacManus, A. M. Hakim and P. Morley, Attenuation of neurotoxicity in cortical cultures and hippocampal slices from e2f1 knockout mice, *J Neurochem* 78 (2001) 316-24.
- [53] S. F. Giardina and P. M. Beart, Kainate receptor-mediated apoptosis in primary cultures of cerebellar granule cells is attenuated by mitogen-activated protein and cyclin-dependent kinase inhibitors, *Br J Pharmacol* 135 (2002) 1733-42.

- [54] W. F. De Azevedo, Jr., H. J. Mueller-Dieckmann, U. Schulze-Gahmen, P. J. Worland, E. Sausville and S. H. Kim, Structural basis for specificity and potency of a flavonoid inhibitor of human cdk2, a cell cycle kinase, *Proc Natl Acad Sci U S A* 93 (1996) 2735-40.
- [55] S. Leclerc, M. Garnier, R. Hoessel, D. Marko, J. A. Bibb, G. L. Snyder, P. Greengard, J. Biernat, Y. Z. Wu, E. M. Mandelkow, G. Eisenbrand and L. Meijer, Indirubins inhibit glycogen synthase kinase-3 beta and cdk5/p25, two protein kinases involved in abnormal tau phosphorylation in alzheimer's disease. A property common to most cyclin-dependent kinase inhibitors? *J Biol Chem* 276 (2001) 251-60.
- [56] C. Guegan, V. Levy, J. P. David, F. Ajchenbaum-Cymbalista and B. Sola, C-jun and cyclin d1 proteins as mediators of neuronal death after a focal ischaemic insult, *Neuroreport* 8 (1997) 1003-7.
- [57] Y. Li, N. Jiang, C. Powers and M. Chopp, Neuronal damage and plasticity identified by microtubule-associated protein 2, growth-associated protein 43, and cyclin d1 immunoreactivity after focal cerebral ischemia in rats, *Stroke* 29 (1998) 1972-80; discussion 1980-1.
- [58] Y. Wen, S. Yang, R. Liu and J. W. Simpkins, Cell-cycle regulators are involved in transient cerebral ischemia induced neuronal apoptosis in female rats, *FEBS Lett* 579 (2005) 4591-9.
- [59] C. Wiessner, I. Brink, P. Lorenz, T. Neumann-Haefelin, P. Vogel and K. Yamashita, Cyclin d1 messenger rna is induced in microglia rather than neurons following transient forebrain ischaemia, *Neuroscience* 72 (1996) 947-58.
- [60] Y. Li, M. Chopp and C. Powers, Granule cell apoptosis and protein expression in hippocampal dentate gyrus after forebrain ischemia in the rat, *J Neurol Sci* 150 (1997) 93-102.
- [61] S. Timsit, S. Rivera, P. Ouaghi, F. Guischard, E. Tremblay, Y. Ben-Ari and M. Khrestchatisky, Increased cyclin d1 in vulnerable neurons in the hippocampus after ischaemia and epilepsy: A modulator of in vivo programmed cell death? *Eur J Neurosci* 11 (1999) 263-78.
- [62] D. L. Small, R. Monette, M. C. Fournier, B. Zurakowski, H. Fiander and P. Morley, Characterization of cyclin d1 expression in a rat global model of cerebral ischemia, *Brain Res* 900 (2001) 26-37.
- [63] H. Kato, A. Takahashi and Y. Itoyama, Cell cycle protein expression in proliferating microglia and astrocytes following transient global cerebral ischemia in the rat, *Brain Res Bull* 60 (2003) 215-21.
- [64] W. Liu, X. Bi, G. Tocco, M. Baudry and S. S. Schreiber, Increased expression of cyclin d1 in the adult rat brain following kainic acid treatment, *Neuroreport* 7 (1996) 2785-9.
- [65] D. S. Park, A. Obeidat, A. Giovanni and L. A. Greene, Cell cycle regulators in neuronal death evoked by excitotoxic stress: Implications for neurodegeneration and its treatment, *Neurobiol Aging* 21 (2000) 771-81.
- [66] H. Ino and T. Chiba, Cyclin-dependent kinase 4 and cyclin d1 are required for excitotoxin-induced neuronal cell death in vivo, *J Neurosci* 21 (2001) 6086-94.
- [67] M. Sakurai, T. Hayashi, K. Abe, Y. Itoyama, K. Tabayashi and W. I. Rosenblum, Cyclin d1 and cdk4 protein induction in motor neurons after transient spinal cord ischemia in rabbits, *Stroke* 31 (2000) 200-7.
- [68] T. Hayashi, M. Sakurai, K. Abe and Y. Itoyama, DNA fragmentation precedes aberrant expression of cell cycle-related protein in rat brain after mca occlusion, *Neurol Res* 21 (1999) 695-8.

- [69] C. Y. Kuan, A. J. Schloemer, A. Lu, K. A. Burns, W. L. Weng, M. T. Williams, K. I. Strauss, C. V. Vorhees, R. A. Flavell, R. J. Davis, F. R. Sharp and P. Rakic, Hypoxia-ischemia induces DNA synthesis without cell proliferation in dying neurons in adult rodent brain, *J Neurosci* 24 (2004) 10763-72.
- [70] Y. Li, M. Chopp, C. Powers and N. Jiang, Apoptosis and protein expression after focal cerebral ischemia in rat, *Brain Res* 765 (1997) 301-12.
- [71] M. van Lookeren Campagne and R. Gill, Cell cycle-related gene expression in the adult rat brain: Selective induction of cyclin g1 and p21waf1/cip1 in neurons following focal cerebral ischemia, *Neuroscience* 84 (1998) 1097-112.
- [72] M. van Lookeren Campagne and R. Gill, Increased expression of cyclin g1 and p21waf1/cip1 in neurons following transient forebrain ischemia: Comparison with early DNA damage, *J Neurosci Res* 53 (1998) 279-96.
- [73] R. Schmidt-Kastner, J. Truettner, W. Zhao, L. Belayev, C. Krieger, R. Busto and M. D. Ginsberg, Differential changes of bax, caspase-3 and p21 mrna expression after transient focal brain ischemia in the rat, *Brain Res Mol Brain Res* 79 (2000) 88-101.
- [74] R. M. Feddersen, R. Ehlenfeldt, W. S. Yunis, H. B. Clark and H. T. Orr, Disrupted cerebellar cortical development and progressive degeneration of purkinje cells in sv40 t antigen transgenic mice, *Neuron* 9 (1992) 955-66.
- [75] T. Jacks, A. Fazeli, E. M. Schmitt, R. T. Bronson, M. A. Goodell and R. A. Weinberg, Effects of an rb mutation in the mouse, *Nature* 359 (1992) 295-300.
- [76] E. Y. Lee, C. Y. Chang, N. Hu, Y. C. Wang, C. C. Lai, K. Herrup, W. H. Lee and A. Bradley, Mice deficient for rb are nonviable and show defects in neurogenesis and haematopoiesis, *Nature* 359 (1992) 288-94.
- [77] K. L. Ferguson, J. L. Vanderluit, J. M. Hebert, W. C. McIntosh, E. Tibbo, J. G. MacLaurin, D. S. Park, V. A. Wallace, M. Vooijs, S. K. McConnell and R. S. Slack, Telencephalon-specific rb knockouts reveal enhanced neurogenesis, survival and abnormal cortical development, *Embo J* 21 (2002) 3337-46.
- [78] T. Hayashi, K. Sakai, C. Sasaki, W. R. Zhang and K. Abe, Phosphorylation of retinoblastoma protein in rat brain after transient middle cerebral artery occlusion, *Neuropathol Appl Neurobiol* 26 (2000) 390-7.
- [79] J. P. MacManus, C. J. Koch, M. Jian, T. Walker and B. Zurakowski, Decreased brain infarct following focal ischemia in mice lacking the transcription factor e2f1, *Neuroreport* 10 (1999) 2711-4.
- [80] J. P. MacManus, M. Jian, E. Preston, I. Rasquinha, J. Webster and B. Zurakowski, Absence of the transcription factor e2f1 attenuates brain injury and improves behavior after focal ischemia in mice, *J Cereb Blood Flow Metab* 23 (2003) 1020-8.
- [81] K. Jin, X. O. Mao, M. W. Eshoo, T. Nagayama, M. Minami, R. P. Simon and D. A. Greenberg, Microarray analysis of hippocampal gene expression in global cerebral ischemia, *Ann Neurol* 50 (2001) 93-103.
- [82] I. K. Hwang, K. Y. Yoo, B. M. Cho, H. S. Hwang, S. M. Kim, S. M. Oh, S. K. Choi, Y. Hwang do, M. H. Won and S. M. Moon, The pattern of e2f1 and c-myb immunoreactivities in the ca1 region is different from those in the ca2/3 region of the gerbil hippocampus induced by transient ischemia, *J Neurol Sci* 247 (2006) 192-201.
- [83] J. M. Trimarchi and J. A. Lees, Sibling rivalry in the e2f family, *Nat Rev Mol Cell Biol* 3 (2002) 11-20.
- [84] G. Tomasevic, F. Kamme and T. Wieloch, Changes in proliferating cell nuclear antigen, a protein involved in DNA repair, in vulnerable hippocampal neurons following global cerebral ischemia, *Brain Res Mol Brain Res* 60 (1998) 168-76.

- [85] Y. Konishi and A. Bonni, The e2f-cdc2 cell-cycle pathway specifically mediates activity deprivation-induced apoptosis of postmitotic neurons, *J Neurosci* 23 (2003) 1649-58.
- [86] S. Love, Neuronal expression of cell cycle-related proteins after brain ischaemia in man, *Neurosci Lett* 353 (2003) 29-32.
- [87] K. Jin, T. Nagayama, J. Chen, A. R. Stetler, K. Kawaguchi, R. P. Simon and S. H. Graham, Molecular cloning of a cell cycle regulation gene cyclin h from ischemic rat brain: Expression in neurons after global cerebral ischemia, *J Neurochem* 73 (1999) 1598-608.
- [88] M. L. Smith, H. U. Kontny, R. Bortnick and A. J. Fornace, Jr., The p53-regulated cyclin g gene promotes cell growth: P53 downstream effectors cyclin g and gadd45 exert different effects on cisplatin chemosensitivity, *Exp Cell Res* 230 (1997) 61-8.
- [89] M. Skotzko, L. Wu, W. F. Anderson, E. M. Gordon and F. L. Hall, Retroviral vector-mediated gene transfer of antisense cyclin g1 (cycg1) inhibits proliferation of human osteogenic sarcoma cells, *Cancer Res* 55 (1995) 5493-8.
- [90] S. Bates, S. Rowan and K. H. Vousden, Characterisation of human cyclin g1 and g2: DNA damage inducible genes, *Oncogene* 13 (1996) 1103-9.
- [91] K. Okamoto and C. Prives, A role of cyclin g in the process of apoptosis, *Oncogene* 18 (1999) 4606-15.
- [92] S. Di Giovanni, S. M. Knoblach, C. Brandoli, S. A. Aden, E. P. Hoffman and A. I. Faden, Gene profiling in spinal cord injury shows role of cell cycle in neuronal death, *Ann Neurol* 53 (2003) 454-68.
- [93] M. Maeda, K. Ampo, S. Kiryu-Seo, H. Konishi, N. Ohba, C. Kadono and H. Kiyama, The p53-independent nuclear translocation of cyclin g1 in degenerating neurons by ischemic and traumatic insults, *Exp Neurol* 193 (2005) 350-60.
- [94] L. H. Tsai, I. Delalle, V. S. Caviness, Jr., T. Chae and E. Harlow, P35 is a neural-specific regulatory subunit of cyclin-dependent kinase 5, *Nature* 371 (1994) 419-23.
- [95] D. Tang, J. Yeung, K. Y. Lee, M. Matsushita, H. Matsui, K. Tomizawa, O. Hatase and J. H. Wang, An isoform of the neuronal cyclin-dependent kinase 5 (cdk5) activator, *J Biol Chem* 270 (1995) 26897-903.
- [96] L. H. Tsai, T. Takahashi, V. S. Caviness, Jr. and E. Harlow, Activity and expression pattern of cyclin-dependent kinase 5 in the embryonic mouse nervous system, *Development* 119 (1993) 1029-40.
- [97] Y. Xiong, H. Zhang and D. Beach, D type cyclins associate with multiple protein kinases and the DNA replication and repair factor pcna, *Cell* 71 (1992) 505-14.
- [98] M. Miyajima, H. O. Nornes and T. Neuman, Cyclin e is expressed in neurons and forms complexes with cdk5, *Neuroreport* 6 (1995) 1130-2.
- [99] H. Zhang, Y. Xiong and D. Beach, Proliferating cell nuclear antigen and p21 are components of multiple cell cycle kinase complexes, *Mol Biol Cell* 4 (1993) 897-906.
- [100] M. Hamdane, A. Bretteville, A. V. Sambo, K. Schindowski, S. Begard, A. Delacourte, P. Bertrand and L. Buee, P25/cdk5-mediated retinoblastoma phosphorylation is an early event in neuronal cell death, *J Cell Sci* 118 (2005) 1291-8.
- [101] S. Cicero and K. Herrup, Cyclin-dependent kinase 5 is essential for neuronal cell cycle arrest and differentiation, *J Neurosci* 25 (2005) 9658-68.
- [102] G. N. Patrick, L. Zukerberg, M. Nikolic, S. de la Monte, P. Dikkes and L. H. Tsai, Conversion of p35 to p25 deregulates cdk5 activity and promotes neurodegeneration, *Nature* 402 (1999) 615-22.
- [103] A. Tandon, H. Yu, L. Wang, E. Rogaeva, C. Sato, M. A. Chishti, T. Kawai, H. Hasegawa, F. Chen, P. Davies, P. E. Fraser, D. Westaway and P. H. St George-Hyslop,

- Brain levels of cdk5 activator p25 are not increased in alzheimer's or other neurodegenerative diseases with neurofibrillary tangles, *J Neurochem* 86 (2003) 572-81.
- [104] B. C. Yoo and G. Lubec, P25 protein in neurodegeneration, *Nature* 411 (2001) 763-4; discussion 764-5.
 - [105] A. Takashima, M. Murayama, K. Yasutake, H. Takahashi, M. Yokoyama and K. Ishiguro, Involvement of cyclin dependent kinase5 activator p25 on tau phosphorylation in mouse brain, *Neurosci Lett* 306 (2001) 37-40.
 - [106] S. Takahashi and A. B. Kulkarni, Mutant superoxide dismutase 1 causes motor neuron degeneration independent of cyclin-dependent kinase 5 activation by p35 or p25, *J Neurochem* 88 (2004) 1295-304.
 - [107] J. L. Hallows, R. E. Iosif, R. D. Biasell and I. Vincent, P35/p25 is not essential for tau and cytoskeletal pathology or neuronal loss in niemann-pick type c disease, *J Neurosci* 26 (2006) 2738-44.
 - [108] T. Hayashi, H. Warita, K. Abe and Y. Itoyama, Expression of cyclin-dependent kinase 5 and its activator p35 in rat brain after middle cerebral artery occlusion, *Neurosci Lett* 265 (1999) 37-40.
 - [109] J. Wang, S. Liu, Y. Fu, J. H. Wang and Y. Lu, Cdk5 activation induces hippocampal cell death by directly phosphorylating nmda receptors, *Nat Neurosci* 6 (2003) 1039-47.
 - [110] M. Morioka, T. Kawano, S. Yano, Y. Kai, H. Tsuiki, Y. Yoshinaga, J. Matsumoto, T. Maeda, J. Hamada, H. Yamamoto, K. Fukunaga and J. Kuratsu, Hyperphosphorylation at serine 199/202 of tau factor in the gerbil hippocampus after transient forebrain ischemia, *Biochem Biophys Res Commun* 347 (2006) 273-8.
 - [111] X. Gong, X. Tang, M. Wiedmann, X. Wang, J. Peng, D. Zheng, L. A. Blair, J. Marshall and Z. Mao, Cdk5-mediated inhibition of the protective effects of transcription factor mef2 in neurotoxicity-induced apoptosis, *Neuron* 38 (2003) 33-46.
 - [112] P. D. Smith, M. P. Mount, R. Shree, S. Callaghan, R. S. Slack, H. Anisman, I. Vincent, X. Wang, Z. Mao and D. S. Park, Calpain-regulated p35/cdk5 plays a central role in dopaminergic neuron death through modulation of the transcription factor myocyte enhancer factor 2, *J Neurosci* 26 (2006) 440-7.
 - [113] J. Zhang, P. K. Krishnamurthy and G. V. Johnson, Cdk5 phosphorylates p53 and regulates its activity, *J Neurochem* 81 (2002) 307-13.

Table 1. Correlative evidence for the involvement of cell cycle components in stroke models.

IN VITRO				IN VIVO		
Component	Model	Effect	References	Model	Effect	References
Cyclin D1	OGD	Increased level	42	MCAO	Increased level	42,56-58
				Focal cerebral ischemia	Increased level	32
				Global Ischemia	Increased level	59-63
				KA injection	Increased level	61,64-66
				Spinal cord ischemia	Increased level	67
				Cardiac arrest	Increased level	86
Cyclin E	KA	Increased level	48			
Cyclin A				MCAO	Increased level & nuclear localization	70
Cyclin H				Global Ischemia	Increase level in resistant neurons	87
Cyclin G1				MCAO	Increased level	71
				Global Ischemia	Increased level	72
				Spinal cord injury	Increased level	92
				CCAO and MCAO	Increased level & nuclear translocation	93
Cdk4				MCAO	Increased level	68
				Focal cerebral ischemia	Increased level	32
				Global Ischemia	Increased level	63
				KA injection	Increased level	66
				Spinal cord ischemia	Increased level	67
				Cardiac arrest	Increased level	86
Cdk2	OGD	Activated	42	MCAO	Increased activity	42
	KA	Increased level	48	Hypoxia-ischemia	Increased activity	69
					Increased level & nuclear translocation	70
				Cardiac arrest	Increased level	86
p16				MCAO	Decreased level	42
				Hypoxia-ischemia	Decreased level	69
p21				MCAO & global ischemia	Increase level in surviving neurons	71-73
p27	OGD	Loss	42	Hypoxia-ischemia	Decreased level	69
Rb	Hypoxia	Phosphorylation	34	MCAO	Phosphorylation	32,58,78
	KA	Phosphorylation	50	Global Ischemia	Phosphorylation	33,34

				Hypoxia-ischemia KA injection	Phosphorylation Phosphorylation	69 65
E2F1	OGD KA	Increased mRNA Increased level	52 48,51	Focal cerebral ischemia Global Ischemia	Increased level Increased level	32 81,82
PCNA	KA	Increased level	51	MCAO Global Ischemia	Increased level Increased level	58,70 63,84
BrdU				MCAO Hypoxia-ischemia	Incorporation in neurons Incorporation in neurons	58 69
Cdk5				MCAO	Increased level	108

Table 2. Functional evidence for the involvement of cell cycle components in stroke models.

IN VITRO				IN VIVO		
Component	Model	Method; effect	References	Model	Method; effect	References
<u>Genetic Studies</u>						
Cyclin D1	Hypoxia	Null mutant neurons protective	34	KA injection	Antisense; protective	66
Cdk4	Hypoxia	Dominant negative expression; protective	34	MCAO	Synthetic inhibitor; protective	58
				KA injection Global ischemia	Antisense; protective Dominant negative expression; protective	66 34
E2F1	OGD KA	Null neurons; protective Null neurons;	52 50	Focal cerebral ischemia	Null mice; resistant	79,80

	Anoxia	protective Null mutant hippocampal slices; protective	52			
Cdk5				Global ischemia	Dominant negative expression; protective	34,109
<u>Pharmacological Studies</u>						
Flavopiridol	KA	Protective	51	Focal cerebral ischemia	Protective	32
	KA	Protective	47	Global ischemia	Protective	33
Olomucine	KA	Protective	53			
	OGD	Protective	42			
Roscovitine	KA	Partial protection	53			
3-ATA	KA	Protective	47			

Figure 1: Schematic representation of our current understanding of the mammalian cell cycle. The cell cycle is broadly divided into four phases culminating in cell duplication. Each phase of the cell cycle is regulated by different complement of cyclin dependent kinases together with a cognate cyclin. During the G1/S-phase, the downstream target, the retinoblastoma gene product, pRb is sequentially phosphorylated by the G1 CDKs resulting in the release of E2F transcription factor and DP. This results in the transcription of E2F responsive genes and those required for the progression thorough S-phase.

