



HAL
open science

Semipermeable membrane devices link site-specific contaminants to effects: Part 1 - induction of cyp1a in rainbow trout from contaminants in prince william sound, Alaska

Kathrine R. Springman, Jeffrey W. Short, Mandy R. Lindeberg, Jacek M. Maselko, Colin Khan, Peter V. Hodson, Stanley D. Rice

► To cite this version:

Kathrine R. Springman, Jeffrey W. Short, Mandy R. Lindeberg, Jacek M. Maselko, Colin Khan, et al.. Semipermeable membrane devices link site-specific contaminants to effects: Part 1 - induction of cyp1a in rainbow trout from contaminants in prince william sound, Alaska. *Marine Environmental Research*, 2008, 66 (5), pp.477. 10.1016/j.marenvres.2008.07.001 . hal-00563041

HAL Id: hal-00563041

<https://hal.science/hal-00563041>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Semipermeable membrane devices link site-specific contaminants to effects:
Part 1 - induction of cyp1a in rainbow trout from contaminants in prince william
sound, Alaska

Kathrine R. Springman, Jeffrey W. Short, Mandy R. Lindeberg, Jacek M.
Maselko, Colin Khan, Peter V. Hodson, Stanley D. Rice

PII: S0141-1136(08)00178-5
DOI: [10.1016/j.marenvres.2008.07.001](https://doi.org/10.1016/j.marenvres.2008.07.001)
Reference: MERE 3268

To appear in: *Marine Environmental Research*

Received Date: 18 March 2008
Revised Date: 26 June 2008
Accepted Date: 1 July 2008

Please cite this article as: Springman, K.R., Short, J.W., Lindeberg, M.R., Maselko, J.M., Khan, C., Hodson, P.V., Rice, S.D., Semipermeable membrane devices link site-specific contaminants to effects: Part 1 - induction of cyp1a in rainbow trout from contaminants in prince william sound, Alaska, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.07.001](https://doi.org/10.1016/j.marenvres.2008.07.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **SEMIPERMEABLE MEMBRANE DEVICES LINK SITE-SPECIFIC CONTAMINANTS**
2 **TO EFFECTS: PART 1 - INDUCTION OF CYP1A IN RAINBOW TROUT FROM**
3 **CONTAMINANTS IN PRINCE WILLIAM SOUND, ALASKA**

4

5

6 KATHRINE R. SPRINGMAN*, JEFFREY W. SHORT¹, MANDY R. LINDEBERG¹, JACEK
7 M. MASELKO¹, COLIN KHAN², PETER V. HODSON², STANLEY D. RICE¹

8

9

10

11

12 ¹Auke Bay Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service,
13 NOAA, 11305 Glacier Highway, Juneau, Alaska 99801-8626 USA

14

15 ²Queen's University, Kingston, Ontario, K7L 3N6 Canada

16

17

18 *Corresponding author contact information:

19 Telephone: 530-400-4141; fax: 707-937-6212; e-mail: krspringman@gmail.com

20 Current address: P.O. Box 315, Littleriver, CA 95456

21

22

23

24

March 2008

25

26 Abstract

27

28 Extracts from semi-permeable membrane devices (SPMDs) deployed on beaches in
29 Prince William Sound (PWS), Alaska, were used to evaluate if complex contaminant mixtures
30 from different sources can be distinguished by the resulting cytochrome P450 1A (CYP1A)
31 activity in exposed test animals. Deployment sites included canneries, salmon hatcheries, and
32 beaches where lingering oil remains from discharges during the 1964 earthquake or the 1989
33 *Exxon Valdez* oil spill. Other sites were selected at random to evaluate region-wide contaminant
34 inputs or were located in salmon streams to evaluate contaminants carried and released by
35 migrating salmon carcasses following reproduction. Following standard deployments of
36 approximately 28 d, an aliquot of the accumulated contaminants was intraperitoneally injected
37 without cleanup into juvenile rainbow trout (*Oncorhynchus mykiss*). After 2d and 7d, the
38 activity of CYP1A was measured by the ethoxyresorufin-*o*-deethylase (EROD) assay. Exposure
39 to extracts from the oiled sites and one hatchery site with numerous creosote pilings elicited
40 strong EROD responses, whereas fish exposed to salmon stream extracts elicited weak but
41 significant responses during late summer compared to late spring. Responses from the other
42 sites were not significant, indicating contaminants from these sources are unlikely to cause
43 CYP1A induction in resident biota. Rather than simply assessing extant contaminants, this
44 method evaluates the potency of the different sites for bringing about aryl hydrocarbon receptor
45 responses in resident biota.

46

47 1. Introduction

48

49 Biochemical responses can be sensitive indicators of pollution exposure. Such responses,
50 termed biomarkers, are indicators in biological fluids, cells or tissues signaling modifications in
51 the biological system due to contaminants (NRC 1987). Biomarker responses have been
52 extensively investigated and provide a convenient and toxicologically relevant means of
53 detecting exposure to low concentrations of contaminants (Dickerson et al., 1994). While
54 biomarkers may indicate exposure to specific classes of contaminants, these responses may
55 provide little information regarding the contaminant source, beyond that inferred from the
56 geographic and temporal patterns of biomarker expression (Short and Springman 2006). Even
57 the location of contaminant sources may be ambiguous when biomarkers are assessed in mobile
58 biota. Moreover, biomarkers usually do not account for chemical speciation or availability
59 (Bucheli and Fent 1995).

60 Biomarker expression can be linked more directly to contaminant sources via an
61 intermediary contaminant sampler such as semipermeable membrane devices (SPMDs). These
62 passive samplers may be deployed to concentrate lipophilic organic contaminants from
63 acknowledged or probable sources. At equilibrium with the exposure medium, SPMDs
64 accumulate contaminants according to their octanol-water partition coefficient (K_{ow}) values
65 (Huckins et al., 2006). The contaminants retained in the SPMD are capable of membrane
66 passage, and as such are potentially a greater risk to biota. Once retrieved by dialysis, these
67 contaminants may be injected into test organisms such as fish, which can then be examined for
68 resulting changes. This relates the response expression to bioavailable contaminants present at
69 the deployment site. Previous studies have assessed the potency of cytochrome P450 1A
70 (CYP1A) in fish liver cells (PLHC-1) to inducers absorbed by environmentally exposed SPMDs
71 (Parrott et al., 1999; Villeneuve et al., 1997). Sundberg et al., (2005) injected trout eggs with

72 fractionated SPMD extracts to evaluate the induction potency of different contaminant classes
73 found in a region where a single point source was dominant. In all these studies, the
74 contaminants recovered from the SPMDs were purified by silica gel chromatography or other
75 means to facilitate chemical analysis prior to their use for the biological assays. Because
76 purification may remove contaminants that affect the selected biomarker, none of these methods
77 assess the induction potential of the environment *per se*, but only the target contaminants of the
78 study. Our aim in this work is to extend these earlier methods to evaluate the CYP1A-induction
79 potential of *all* nonpolar, lipophilic contaminants absorbed by SPMDs deployed in the
80 environment under study by injecting recovered contaminants into juvenile rainbow trout
81 without clean-up.

82 In situations where multiple contaminant sources are plausible, the biomarker response of
83 injected contaminants extracted from SPMDs provides a method of evaluating the relative
84 potency of different sources. The biomarker response of animals injected with the SPMD extract
85 provides an indication of the induction potency of contaminants from the source(s) sampled by
86 the deployed SPMD. This approach quantifies the biochemical effect triggered by the mixture
87 of nonpolar compounds as a group, and does not require identification of the particular
88 contaminants causing induction. This may be a considerable advantage in cases where a positive
89 biomarker response is evident in resident biota, multiple potential sources are present, and the
90 identity of the contaminants that are driving the biomarker response is not apparent.

91 Prince William Sound (PWS), Alaska provided an excellent setting to test the utility of
92 this method. It is a sparsely populated region where relatively few candidate sources are
93 present, one of which is oil from the 1989 *Exxon Valdez* oil spill that is still found on beaches in
94 western PWS. This oil is a CYP1A inducer (Woodin et al., 1997). Induction of CYP1A has

95 been reported in resident sea otters (*Enhydra lutris*) and harlequin ducks (*Histrionicus*
96 *histrionicus*) that frequent beaches where oil persists (Short et al., 2006, Bodkin et al., 2002,
97 Trust et al., 2000). However, confounding potential sources of CYP1A inducers are possible.
98 These include: (1) asphalt and heavy bunker fuels released from storage tanks damaged by the
99 1964 Alaska earthquake (Kvenvolden et al 1995); (2) fuel spills from commercial fishing,
100 marinas, salmon hatchery docks and other marine traffic (Page et al., 1999); (3) broad-scale
101 background pollutants from atmospheric deposition of persistent organic pollutants (Stern et al.,
102 1997); (4) polycyclic aromatic hydrocarbons (PAH) associated with oil seeps and erosion of
103 hydrocarbon-rich source rocks and coal outcrops (Short et al., 2004; Short et al., 1999; Page et
104 al., 1995); (5) PAH from forest fires (Page et al., 1999); (6) contaminants from marine vessels
105 (Page et al., 1995); and (7) persistent organic pollutants associated with spawning migrations of
106 adult salmon returning to their natal streams (Krümmel et al 2003; Ewald et al., 1998).
107 Concerns regarding these potential sources of CYP1A inducers in the region continue to be
108 raised (Boehm et al., 2007, Huggett et al., 2006, Neff et al., 2006, Lee and Anderson, 2005, Page
109 et al., 2004, Huggett et al., 2003, Jewett et al., 2002). Injecting fish with extracts from SPMDs
110 and measuring their CYP1A activity provides a way of identifying which sources are capable of
111 stimulating a response. Those that do not, can be eliminated as causes of the CYP1A induction
112 observed in wildlife in that region.

113 Here, we extended the approach used by Sundberg et al. (2005) to evaluate which
114 pollution sources in PWS were capable of stimulating CYP1A induction. In fish, the activity of
115 this enzyme, as determined with the ethoxyresorufin-*o*-deethylase (EROD) assay, is an
116 established biomarker of exposure to PAH and structurally related compounds. The activity of
117 this enzyme is particularly suited to monitor discharges of petroleum or other PAH sources, co-

118 planar polychlorinated biphenyls (PCB), and dioxins (Whyte et al., 2000; Stegeman and Hahn
119 1994). Our overall objectives were to identify which of these sources contained compounds that
120 induced CYP1A in exposed biota, and to determine which of the bioavailable hydrocarbons were
121 responsible. We report our results in two parts. Here (Part I), our objectives are to describe and
122 validate the method, which includes an assessment of the precision, sensitivity, and dynamic
123 range. We present the results of our field deployments of SPMDs in PWS to evaluate the
124 CYP1A induction potential of prospective contaminant sources. In Part II, (Short et al., 2008),
125 we relate the CYP1A induction results to the chemical composition of the contaminants
126 accumulated by the SPMDs.

127

128 **2. Methods**

129

130 *2.1 Sampling Design*

131

132 Prince William Sound is a complex, fjord-type ecosystem with a sea surface area of about
133 8,800 km² (Schmidt 1977). Most of its 7,000 inhabitants reside in Cordova, Valdez or Whittier
134 (Fig. 1), while fewer than 200 live in the vicinity of Chenega or the regional salmon hatcheries.
135 The latter group is near the path followed by oil released by the T/V Exxon Valdez during the oil
136 spill in 1989. Past industrial activities within the spill path were sporadic. These include 2 large
137 mine sites at Latouche, 6 fish processing plants, and a number of smallholdings devoted to
138 mineral prospecting, artisanal sawmills and fur-farming. All of these were abandoned during the
139 mid-20th century (Wooley 2000). Current industrial activities include the Alyeska oil terminal in

140 Port Valdez, commercial fishing and oyster farming, 5 large salmon hatcheries, and seasonal
141 tourism.

142 We deployed SPMDs to evaluate the CYP1A induction potential of 6 classes of
143 contaminant sources: (1) lingering, subsurface, Alaska North Slope (ANS) oil from the *Exxon*
144 *Valdez* oil spill ($N = 5$ sites, 3 of which were replicated); (2) human activity ($N = 6$) sites,
145 including an abandoned industrial site with Monterey Formation bunker oil from storage tanks
146 damaged by the 1964 Alaska earthquake; (3) salmon hatcheries ($N = 5$); (4) salmon streams (N
147 $= 5$), because salmon carcasses can release accumulated CYP1A-inducing pollutants as they
148 decompose in their natal streams following reproduction; (5) random sites ($N = 9$) to evaluate
149 area-wide pollution sources including atmospherically transported contaminants; and (6) a site at
150 Constantine Harbor where PAH associated with organic-rich rocks eroded from geologic sources
151 east of PWS are incorporated into the intertidal sediments (Short et al., 1996). At the ANS sites,
152 we deployed additional SPMDs at adjacent paired control sites to account for local inducing
153 agents other than ANS. We also deployed SPMDs at a remote site at Graves Harbor in southeast
154 Alaska ~ 300 km east of PWS and upcurrent of the geological hydrocarbon sources to serve as a
155 negative regional field control site, and at Cordova Harbor to serve as a positive field control.

156 The SPMDs were deployed at most sites in spring from mid-May to mid-June 2004, prior
157 to the annual spawning migration of salmon. At the salmon stream sites, they were again
158 deployed in summer from mid-August to mid-September 2004, after adult salmon had returned
159 to spawn. Five of the random site deployments were during spring, and the other 4 during
160 summer to assess region-wide seasonal differences. The random selections for these sites were
161 made within each of 5 sectors partitioning PWS (see Fig. 1 in Short et al., 2008). Each sector

162 was allocated 1 or 2 sites to ensure dispersion, and each site was located on a randomly-selected
163 shore segment within a sector as described in Part II (Short et al., 2008).

164

165 *2.2 Semipermeable membrane devices*

166

167 The SPMDs were obtained from Environmental Sampling Technologies (St. Joseph, MO),
168 and deployed in 30 cm long cylindrical cages, each loaded with 5 SPMDs woven on stainless
169 steel carriers stacked within the cage. Each SPMD had dimensions of 91.4 × 2.5 cm and
170 contained 1 ml of ultra-high purity (UHP) triolein. The cages were half buried in pits excavated
171 at the mid-tide level of deployment site shorelines. After placement in a pit, the excavated
172 material was replaced to the level of the beach surface, surrounding, but not covering, the cage.
173 This arrangement allowed the SPMDs to sample the atmosphere at low tide (Lohmann et al.,
174 2001, Ockenden et al., 1998), surface seawater at high tide, and interstitial water flow within the
175 upper 15 cm of beach while minimizing biofouling. Cages were deployed for the standard
176 deployment period of 27 – 28 days, then retrieved for dialysis. At the ANS and human activity
177 sites where contamination was greater, the cages were placed in patches of subsurface oil to
178 ensure uptake of organic contaminants. A field blank cage with 5 UHP SPMDs was briefly
179 exposed to the atmosphere during two of the deployments and then stored for dialysis.

180

181 *2.3 SPMD processing*

182

183 Contaminants were recovered by dialysis into hexane at the laboratory of Environmental
184 Sampling Technologies, following physical removal of external sediment and biofouling as per

185 Huckins et al. (2000). The dialysates from all 5 SPMDs within a cage were pooled. Following
186 concentration of the combined dialysates to ~1 ml, samples were sent to Auke Bay Laboratory in
187 sealed ampules. The contents were partitioned into subsamples for chemical analysis (3 aliquots
188 each containing 5% of the dialysate) and for injection into juvenile trout for CYP1A induction (2
189 aliquots, each with 42.5% of the dialysate). One each of the aliquots for injection and for
190 chemical analysis were archived in the event of accidental losses. Results of the chemical
191 analysis of the dialysates are presented in Part II (Short et al., 2008). The aliquot used for
192 injection was exchanged into peanut oil (Spectrum Chemical Mfg., Gardena, CA) under nitrogen
193 at ~ 35 °C. Samples for induction tests were coded with un-informative sample identification
194 numbers and sent to California for blind bioassay.

195 Quality assurance samples included field and dialysis blanks, a solvent control, and a
196 positive laboratory control (β -naphthoflavone or BNF, a model CYP1A inducer). The β -
197 naphthoflavone was suspended in peanut oil and doses of 2.5 mg kg⁻¹ were prepared for three
198 fish weight classes, ≤ 7.5 g, 7.5-10.0 g, ≥ 10.0 g, to maintain a uniform injection volume of 50 μ l
199 fish⁻¹.

200

201 2.4 Animal Exposure

202

203 Juvenile rainbow trout (*Oncorhynchus mykiss*), Mount Shasta strain, average weight 9.2
204 ± 2.6 g, were obtained from California Department of Fish and Game Mount Shasta Hatchery.
205 They were maintained at 13 ± 1 °C in aerated, flow-through water systems for at least a week
206 prior to testing and fed a commercial diet (Silver Cup, #3, Nelson & Sons Inc., Murray, UT) until
207 the day prior to dosing.

208 The dialysate aliquot from each sample was partitioned further into 10 sub-aliquots, each
209 injected into a single juvenile trout, so each trout received 4.2% of the dialysate from an SPMD
210 deployment in a 50 μ l injection. For each sample, 10 fish were anesthetized in a solution of ~50
211 mg l⁻¹ tricaine methane sulfonate (MS222) and injected intraperitoneally (i.p.) with the extract.

212 Many compounds can induce CYP1A, including planar hydrocarbons that meet the
213 requisite structural criteria, but these compounds may reach peak induction at different times.
214 The positive control, β -naphthoflavone (BNF), is at its peak induction ~ 48h after i.p. injection in
215 rainbow trout, which is similar to the response from PAHs in this species (Lemaire et al., 1996).
216 Of 209 polychlorinated biphenyls (PCBs), those that can induce CYP1A strongly in rainbow
217 trout show peak induction at 6 d following i.p. injection (Huuskonen et al., 1996). We sacrificed
218 5 fish of each group in an overdose of MS222 at 2 d to examine the CYP1A responses of those
219 compounds that are metabolized more readily, such as PAH, while allowing the remaining 5 fish
220 7 d until sacrifice to evaluate effects from compounds which reach peak induction later, such as
221 PCBs. The weight and standard length of each individual were recorded, and the liver excised
222 and frozen in liquid nitrogen within two minutes of death. This was repeated with the remaining
223 5 fish from each group at 7 d post-injection. The livers were shipped frozen to Queen's
224 University, Kingston, Ontario, Canada for evaluation of CYP1A induction.

225
226 2.5 *EROD* bioassay

227
228 CYP1A induction was estimated by measuring the activity of ethoxyresorufin-*o*-
229 deethylase (EROD), based on an original method by Pohl and Fouts (1980) modified for a
230 microplate spectrofluorometer (Hodson et al., 1996). Liver samples stored at -84°C were

231 thawed on ice and a sub-sample of 15-60 mg was homogenized in 500 – 750 μL of ice-cold 20
232 mM Na-HEPES buffer in 150 mM KCl (pH7.5). Homogenates were centrifuged for 20 minutes
233 at 2°C and 9000 x G, and the supernatant and microsome layer (S-9 fraction) were recovered and
234 stored at -84°C. Each S-9 fraction was thawed on ice and triplicate aliquots of 50 μL were
235 incubated with 50 μL of 2.2 μM 7-ethoxyresorufin in 100 mM HEPES buffer (pH 7.8) in a 96-
236 well microplate held in the dark for 10 minutes at room temperature. The reaction was initiated
237 with 10 μL of 1.1 μM of NADPH. Fluorescence of resorufin, the reaction product, was
238 measured every minute for 12 minutes at excitation and emission wavelengths of 530 and 586
239 nm with a SpectraMax Gemini spectrofluorometer (Molecular Devices, Sunnyvale CA, USA).
240 Crude activity was estimated as the rate of increase in fluorescence and converted to a molar
241 basis by reference to a resorufin standard curve (0-20 pM) included with each plate. Molar
242 specific activity (picomole resorufin per mg protein per minute; $\text{pmol mg protein}^{-1} \text{ min}^{-1}$) was
243 calculated by normalizing crude activity to S-9 protein concentrations. Protein was determined
244 against a standard curve of bovine serum albumin (0-2.5 mg l^{-1}) by mixing 10 μL of each S-9
245 fraction with 200 μL of Biorad Reagent (Biorad, Hercules, CA, USA) and measuring absorption
246 at 600 nm with a SpectraMax Plus spectrophotometer (Molecular Devices, Sunnyvale CA,
247 USA). Data were acquired and analyzed with Molecular Devices SOFTMAX PRO software.
248 Samples were analyzed blind, and the analytical batches were composed of samples selected at
249 random.

250

251 *2.6 Data analysis*

252

253 The CYP1A induction caused by contaminants recovered from the 5 SPMDs in a
254 deployment device is presented as the antilog of the logarithmic mean EROD value of the 5 fish
255 injected for each time group (2 d and 7 d). The logarithmic data transformation is used as the
256 EROD activities are log-normally distributed (Hodson et al. 1996), justifying parametric
257 statistical comparisons. Consequently, following logarithmic transformation, we used analysis
258 of variance (ANOVA) to assess the significance of differences between EROD values. Groups
259 of SPMD results within contaminant source classes are presented as the mean for individual
260 SPMD deployments. Variation is based on the logarithmic standard error, presented as the
261 antilog, which gives an asymmetric interval.

262 To clarify notation in these statistical tests, we use " N " to represent the number of sites in
263 a class, as described in Section 2.1. Because the number of individuals compared is the number
264 of fish injected, we use " n " to denote the number of fish within a group, and " m " to denote the
265 number of SPMDs that were used to prepare the extract for that group. Although 5 fish were
266 usually injected with the equivalent of an extract aliquot from one SPMD, the number of
267 corresponding EROD analyses was occasionally fewer because of processing errors, and greater
268 in some cases because all the extract was used to evaluate the response at 2 d.

269

270 **3. Results and Discussion**

271

272 *3.1 Analytical Performance*

273

274 The ability of the method to detect and distinguish contaminant burdens injected into the
275 juvenile trout depends on the sensitivity, precision, and dynamic range of the EROD response,

276 which are demonstrated by the laboratory and field control treatments. The basal EROD activity
277 of unhandled fish, $0.932 \text{ pmol mg protein}^{-1} \text{ min}^{-1}$ (Table 1), defines the minimum activity
278 expected in the absence of stress. Injection of the peanut oil solvent caused a significant
279 doubling of activity after 2 d ($P = 0.002$), subsiding by 7 d to a level not significantly different
280 than the basal activity ($P = 0.56$; Table 1). The lower responses at 7 d compared with 2 d were
281 the result of recovery from the stress of the initial injection and handling (Lemaire et al., 1996,
282 Blom and Förlin, 1997). Consequently, increases that more than double the basal activity
283 following injection were probably the result of factors other than handling stress.

284 Injection of some of the field- and dialysis-blank SPMD extracts caused greater EROD
285 activity than expected from handling stress, suggesting low-level contamination introduced
286 during manufacture. The mean EROD activity of these blanks was twice that of the
287 corresponding activities caused by injection of peanut oil at 2 d and at 7d (Table 1), both
288 significant increases ($P < 0.05$). Analysis of variance showed marginally significant differences
289 among the field blanks (FB; $m = 4$) at 2 d ($P = 0.063$) and clearly significant differences at 7 d
290 ($P < 0.001$). The 7 d activities of one of the field blanks and one of the dialysis blanks were not
291 significantly different than their corresponding 2 d activities ($P > 0.20$), the other field blank and
292 dialysis blank are significantly lower ($P < 0.01$) at 7 d. These results suggest that all the field
293 and dialysis blanks contained traces of contaminants capable of stimulating small increases in
294 EROD activity 2 d following extract injection, extending in some cases to 7 d. Such low-level
295 traces of contaminants in un-deployed SPMDs have been noted previously by Boehm et al.,
296 (2005). However, the alkylated PAH recovered from SPMDs by Boehm et al. (2005) may have
297 been introduced as laboratory artifacts during their non-standard dialysis of the samplers
298 (Schwartz, 2006).

299 Oleic acid, an impurity in the triolein used for SPMDs, may contribute to some of the
300 observed responses. The fraction containing these compounds has produced positive responses
301 in the Microtox assay (Sabaliûnas et al., 1999). However, Huckins et al. (2006) noted that while
302 deployed, any remaining oleic acid or methyl oleate impurities diffuse to the exterior surface of
303 the membrane where they are removed. This environmental release does not occur in quality
304 control SPMDs (field blanks and dialysis blanks), creating the potential for differential
305 responses. Purified (UHP) triolein substantially decreases these impurities (Lebo et al, 2003;
306 also, summary of SPMD screening in Supplementary Data), thus decreasing but not precluding
307 this risk, which could be magnified without sample cleanup. These results indicate that
308 additional purification of SPMDs may be necessary to achieve maximum sensitivity when the
309 contaminants are extracted, injected into fish and the EROD response measured. In our study,
310 this additional purification was provided by environmental exposure when SPMDs were
311 deployed at sites distant from obvious contaminant sources within PWS.

312 The low EROD responses produced by SPMDs deployed at random sites, Constantine
313 Harbor, regional field control and the local ANS control sites provide a robust estimate of the
314 precision of the method in the absence of contaminants. The mean EROD response for these
315 sites was not significantly different from the responses from the peanut oil injections at either 2 d
316 or 7 d ($P > 0.47$, $m = 16$; Table 1), but were significantly below the corresponding field and
317 dialysis blank responses ($P < 0.001$). This indicates that impurities initially present in the
318 SPMDs had diffused into the environment during deployment, an effect that is also evident in
319 PAH data presented for reference sites in PWS presented by Boehm et al. (2005). As with the
320 peanut oil injections, the 2 d responses were significantly greater than the 7 d responses by a
321 factor of 2 ($P < 0.001$), confirming the effect of handling stress. Analysis of variance did not

322 shows significant differences among the individual SPMDs at 2 d following injection ($P =$
323 0.318), but did at 7 d ($P < 0.001$), which we attribute to slight variation in fish responses to
324 handling stress.

325 Comparison of the positive and negative control samples illustrates the analytical range
326 of the method. The positive laboratory control dose of 2.5 2.5 mg kg⁻¹ BNF is near the threshold
327 for immune system impairment in juvenile trout similar to those we used (Springman et al.,
328 2005), providing a biologically meaningful laboratory benchmark. The corresponding 2 d and 7
329 d EROD responses near 70 pmol mg protein⁻¹ min⁻¹ (Table 1) were about 40-fold greater than
330 the response from the peanut oil injection. The contaminants accumulated by the SPMDs
331 deployed at the Cordova Harbor field control site elicited about half the response of the BNF
332 dose (Table 1), confirming the ability of the method to detect CYP1A-inducing contaminants
333 when present. Cordova Harbor, a confined harbor with limited seawater exchange, is one of the
334 most polluted in the United States based on petroleum-derived contaminant concentrations in
335 mussels within the harbor (Miles et al., 2001).

336 The EROD responses of the rainbow trout we injected are comparable with responses
337 expected for feral fish and other biota occupying the environment of the deployed SPMD. A
338 valid comparison of induction from bioavailable CYP1A inducers depends on a comparable
339 absorption capacity in both fish and SPMD. Direct experimental comparison of uptake rates and
340 accumulated burdens for SPMDs vs. fish shows that SPMDs accumulate polychlorinated
341 biphenyls (PCBs) about twice as fast per gram SPMD as per gram wet fish tissue, resulting in
342 body burdens about twice as great (Echols et al., 2000; Meadows et al., 1998). With our method,
343 42.5% of the contaminants accumulated by 22.5 g of SPMDs was injected into 10 rainbow trout
344 of 9.2 g mean wet weight, so the effective ratio of SPMD to wet fish tissue is 1:10, which is

345 comparable to that in Meadows et al. (1998). If lipophilic contaminants are accumulated twice
346 as fast by SPMDs than fish, then our fish received about one-fifth the body burden they would
347 have accumulated, had they been exposed to the deployment conditions of the SPMD that
348 produced the extract we injected into them. However, the entire complement of contaminants
349 accumulated by the SPMD during the 28 d SPMD deployment period was directly administered
350 to our naïve test fish. Assuming fish absorb, biotransform and eliminate xenobiotics according
351 to first-order kinetics, the time required to attain 20% of the equilibrium body burden when
352 exposed to a constant aqueous contaminant concentration is $(0.22 * t_{1/2}/0.693)$, where $t_{1/2}$ is the
353 half-life of the contaminant in the fish (Meador et al., 1995). Half-lives of 3- and 4-ring PAH in
354 rainbow trout are 2 to 9 days (Niimi and Dookhran 1989; Niimi and Palazzo 1986), implying
355 20% accumulation within 0.6 – 2.9 days. Even longer times would be required for more
356 persistent contaminants such as PCBs due to their longer half-lives. Our intraperitoneal injection
357 lead to complete absorption of contaminants within a few hours, and would therefore be about as
358 effective for stimulating CYP1A induction as placing the fish in the environment sampled by the
359 deployed SPMD. Consequently, it is unlikely that biota exposed to the same environment as our
360 deployed SPMDs would have CYP1A activities above their basal level if an injection of
361 contaminants from the SPMD fails to stimulate induction.

362 Having described the analytical performance and characteristics of our SPMD-extract
363 injection method, we now assess results from SPMD deployments at sites within or near obvious
364 local contaminant sources within PWS.

365

366 *3.2 ANS Sites*

367

368 The SPMDs deployed at the ANS sites consistently accumulated inducers responsible for
369 the strongest CYP1A responses of our study (Fig. 2). The overall mean response was 39 pmol
370 mg protein⁻¹ min⁻¹ for 2 d, significantly exceeding the ANS control site responses ($P < 0.001$).
371 Mean responses for these sites ranged from 23 – 69 pmol mg protein⁻¹ min⁻¹. At one site where
372 3 SPMDs were deployed within 2 m of each other, the mean responses ranged from 24 – 36
373 pmol mg protein⁻¹ min⁻¹, providing an indication of field reproducibility of the method. The
374 consistency of the responses from the ANS sites shows that the lingering ANS oil remains a
375 potent source of CYP1A-inducing compounds, and that those compounds are in a form that can
376 be absorbed by SPMDs and presumably by biota. The subsurface oil patches where the SPMDs
377 were deployed consisted of semi-liquid oil that readily produced surface sheens on contact with
378 water, so the CYP1A inductive potential of the extracts derived from these deployments is not
379 surprising, given the demonstrated CYP1A induction potency of ANS oil (Woodin et al., 1997).

380 Induction of EROD activity persisted for 7 d following injection of SPMD extracts from
381 most of the ANS sites (Fig. 2). The 7 d response was greater than the 2 d response at one of the
382 sites on Disk Island, but otherwise were lower by factors of 2 – 3 at 3 other sites, and by factors
383 of 5 -10 at the remaining 2 sites (KN114A and KN109A). The results from the ANS sites
384 suggest more complex interactions which incorporate site-specific characteristics, such as those
385 found at Disk Island and areas of Knight Island (KN136A). Exposure to some forms of peat
386 may induce CYP1A (Kopponen et al., 1993), and co-exposure to humic acids can modulate
387 induction (Matsuo et al. 2006). These sites are located in areas where there is peat beneath the
388 oil, and one of the Disk Island deployments was within a stream. The 2 d responses from all
389 three sites (Fig. 2) are similar to each other and to other oiled sites. However, the 7 d responses
390 corroborate the elevated CYP1A activity that results from co-exposure to humic substances and

391 petroleum (Matsuo et al., 2006), in that the 7d response is more persistent at the Disk Island
392 location where the peat is not flushed by the surface stream. ANS sites received the same oil in
393 1989, yet the response elicited by bioavailable hydrocarbons in 2004 varies in persistence and
394 intensity from site to site. Such location-specific factors may sometimes affect the potential
395 toxicity of bioavailable PAH.

396

397 *3.3 Human Activity and Salmon Hatchery Sites*

398

399 The EROD responses from the human activity and salmon hatchery sites were usually
400 similar or only slightly greater than responses at the random and other uncontaminated sites (Fig.
401 3), with two exceptions: McClure Bay and Armin F. Koernig (AFK) hatchery. Excluding these
402 two sites, the mean 2 d and 7 d EROD responses of and salmon hatchery sources ranged from 1.9
403 – 5.7 pmol mg protein⁻¹ min⁻¹ ($m = 9, n = 45$) and from 1.3 – 2.9 pmol mg protein⁻¹ min⁻¹ ($n =$
404 44), respectively. The responses that are significantly greater than those of the uncontaminated
405 sites are indicated in Fig. 3. These sites include surface patches of Monterey Formation asphalt
406 at Latouche, Port Ashton, Port Audrey and Thumb Bay, and unidentified contaminants near the
407 hatchery at Main Bay. At Latouche, the EROD response remained significantly elevated
408 through 7 d.

409 These small but significant EROD responses indicate that the CYP1A induction
410 potencies of the contaminants at most human activity and salmon hatchery sites are minor when
411 compared with induction potential from oiled ANS sites. This may be because either these
412 human activity and salmon hatchery contaminants are less bioavailable, or are not strong
413 inducers, or both. Page et al. (2006) reported substantial concentrations of PAH associated with

414 asphaltic remnants of Monterey Formation bunker oil contaminating a total of ~ 2.1 ha of
415 surficial sediments at Latouche, Port Ashton, Port Audrey and Thumb Bay. Our results indicate
416 the CYP1A induction potency of these sediments is weak in contrast to lingering ANS oil
417 (compare Figs. 2 and 3). Latouche, Port Ashton, Port Audrey and Thumb Bay account for 95%
418 of the total area contaminated by Monterey Formation bunker oil that also lies within the path of
419 the *Exxon Valdez* oil spill (Page et al., 2006). This combined area amounts to ~ 20 % of the area
420 contaminated by lingering *Exxon Valdez* oil (Short et al., 2004, 2008). The combination of
421 weaker induction and the smaller size of the combined Monterey Formation bunker oil spills
422 demonstrates these sources of EROD-inducing contaminants are minor in comparison with
423 lingering *Exxon Valdez* oil.

424 In contrast, SPMD extracts from the AFK hatchery and McClure Bay sites both produced
425 substantial EROD responses (Fig. 3), indicating the presence of stronger and more bioavailable
426 CYP1A inducing contaminants. The 2 d mean EROD responses are greater than 31 pmol mg
427 protein⁻¹ min⁻¹ and remained elevated after 7 d, especially at McClure Bay, where the 7 d
428 response was nearly half the 2 d response. The deployment site at AFK is near a large pier
429 supported by several hundred creosote pilings, and creosote pilings support numerous other piers
430 or their remnants within ~ 1- 2 km of the deployment site, unlike the other hatcheries where steel
431 or concrete pilings are in use. These creosote pilings are the likely source of the CYP1A-
432 inducing contaminants at AFK. Creosote contains numerous PAH that are potent CYP1A
433 inducers (Hyötyläinen and Oikari 1999) and the large surface area of the pilings near the AFK
434 deployment site would promote dissolution of PAH into surrounding waters. Similarly, the
435 SPMD deployment at McClure Bay was within a patch of semi-liquid, subsurface Monterey
436 Formation bunker fuel oil that was released from a nearby storage tank ruptured during the 1964

437 Alaska earthquake. In contrast with surficial deposits of Monterey Formation asphalt at the
438 other human activity sites, the EROD response from the McClure Bay deployment shows that
439 burial can impede weathering such that the associated contaminants remain readily bioavailable
440 after 40 years. This corroborates the continuing bioavailability of contaminants associated with
441 the ANS oil, as well as the extraordinary persistence of remaining ANS oil (Short et al., 2004,
442 2008). Also, the elevated 7 d EROD responses confirms our results from the oiled ANS sites, in
443 that prolonged EROD responses are associated with this hydrocarbon source as well.

444 The prolonged, increased EROD responses from SPMD extracts deployed at some sites
445 suggests possible interactive effects, since elevated 7 d EROD activity occurs only at those sites
446 with a history of hydrocarbon exposure. For example, PAH oxidation may contribute to CYP1A
447 induction at favorable sites for microbial growth. This prolonged EROD response may increase
448 the duration that significant EROD responses are detectable in monitored, mobile biota exposed
449 to lingering oil in PWS, such as sea otters (Bodkin et al., 2002), and harlequin ducks (Trust et al.
450 2000).

451

452 3.4 Salmon Streams

453

454 The EROD responses from the summer SPMD deployments in the 5 salmon streams
455 consistently exceeded those of the spring deployments, implying slight seasonal variability at
456 these sites. The 2 d and 7 d responses from salmon stream spring deployments are not
457 significantly different ($P > 0.105$) than the respective responses from the random sites sampled
458 in spring (Fig. 4). In contrast, the 2 d and 7 d responses from summer deployments, averaged
459 over the 5 salmon stream sites, are 3.6 and 2.0 pmol mg protein⁻¹ min⁻¹ ($n = 25$), which are 41%

460 and 44% greater than respective responses from the random sites sampled during summer ($P <$
461 0.036). The increases of the 2 d responses are consistent for all 5 salmon streams, and for 4 of
462 the streams for the 7 d responses. The increase during late summer but not during late spring
463 indicates a seasonally transient input that subsides from late summer to the following spring.
464 This summer increase is localized to the salmon streams. Differences between random site
465 deployments during spring compared with summer are not significant ($P > 0.15$; Fig. 4) and are
466 within the range of responses from peanut oil injections. Thus the seasonal increase observed at
467 the salmon stream sites is probably not the result of a region-wide contaminant input. The
468 timing of this increase is in agreement with contaminants released from decomposing salmon
469 carcasses, following their migration to their natal stream to reproduce.

470

471 *3.5 Constraints on EROD Induction by Background Contaminants*

472

473 The consistent absence of CYP1A induction following injection of contaminants
474 extracted from our random and control site SPMDs into juvenile trout shows that the
475 contaminant background is a negligible source of dissolved, lipophilic CYP1A inducers in PWS.
476 Background contaminants are by definition broadly distributed, so they should routinely cause
477 induction if they include CYP1A inducers. But our results show that none of the $m = 16$ SPMD
478 deployments at random and control sites elicited CYP1A induction above the stress caused by
479 peanut oil injection, and that deployment at these sites actually removed CYP1A inducing
480 compounds that were initially present in the SPMDs at deployment. We therefore conclude that,
481 apart from small, localized pollution sources that are usually readily identifiable (e.g. creosote
482 pilings, remnant oil deposits from the 1964 earthquake or the 1989 *Exxon Valdez* oil spill, etc.),

483 the marine waters of PWS are among the least contaminated coastal waters with respect to
484 dissolved, lipophilic CYP1A inducers.

485 The low levels of dissolved CYP1A-inducing contaminants in the marine waters of PWS
486 implied by our random and control site results are consistent with most other studies of
487 contaminants in the region. As reviewed recently by Boehm et al. (2007), measured background
488 concentrations of total PAH in the marine waters of PWS are usually less than 20 ng L^{-1} (often
489 near or below the detection limits of the methods used), nearly all of which are 2- and 3-ring
490 PAH that are not potent CYP1A inducers (Barron et al., 2004). The SPMDs deployed by Boehm
491 et al. (2005) on the beach surface at their reference site contained a mean total PAH amount of
492 39 ng SPMD^{-1} , implying aqueous concentrations in the neighborhood of 1 ng L^{-1} , again with
493 compounds capable of inducing CYP1A likely to be a very small proportion. In a survey of
494 PAH bioavailability at PWS sites oiled by the 1989 *Exxon Valdez* oil spill, Neff et al. (2006)
495 measured PAH concentrations of sea lettuce (*Ulva fenestrata*), reasoning that the mucilaginous
496 surface coating of this macroalgae would efficiently collect dissolved and particle-bound PAH
497 from seawater at higher tides and from the atmosphere at lower tides, much as our SPMDs
498 absorbed dissolved PAH from these media. They found mean concentrations of 11.3 ng g^{-1} dry
499 wt., again implying ambient total PAH concentrations below 1 ng L^{-1} (assuming a wet:dry wt
500 ratio of 10:1, and an effective K_{ow} of 10^4). In a study of PAH in lipid-rich copepods in central
501 PWS, Carls et al. (2006) found total PAH of 409 ng g^{-1} , entirely of 2- and 3-ring PAH, and
502 equivalent to aqueous total PAH concentrations below about 20 ng l^{-1} . Finally, the PAH
503 concentrations detected in the SPMDs deployed at the random and control sites of this study also
504 imply ambient dissolved total PAH concentrations in the low ng L^{-1} range, with the more potent
505 CYP1A inducers such as PCBs and chrysenes usually below their quantitation limits in the

506 SPMDs (Short et al., 2008). These studies all indicate ambient concentrations of the more potent
507 CYP1A inducers such as chrysene that are well below 1 ng L^{-1} in the marine waters of PWS, and
508 hence well below concentrations that might plausibly stimulate a background elevation of
509 CYP1A in PWS biota.

510 Neff et al. (2003) concluded that background PAH in deep-water sediments of PWS are
511 bioavailable based on differences in concentration patterns detected in polychaete worms (*Nereis*
512 *virens*) following prolonged exposure, but the supporting evidence presented is equivocal. The
513 PAH burden of the polychaete worms declined during exposure to the PWS sediments, were near
514 typical detection limits of the method used, and no evidence of statistical significance was
515 reported, suggesting the PAH pattern differences may simply reflect analytical variability.

516 Huggett et al. (2003) and Page et al. (2004) have argued that marine fish in the northern
517 Gulf of Alaska are chronically exposed to CYP1A-inducing hydrocarbons derived from oil seeps
518 and eroding hydrocarbon source rocks, but this conclusion seems premature. Their argument is
519 based on detecting measurable liver EROD responses in 4 species of fish, and fluorescent
520 aromatic compounds (FAC) in their bile. However, these authors failed to consider the
521 possibility that the low values they reported are merely the basal responses for the fish species
522 they tested. A thorough familiarity of this endogenous activity is necessary for accurate
523 interpretation of EROD induction. Of several biomarkers examined in 15 species of fish, EROD
524 induction showed the most variable basal activity between species (Förlin et al., 1995). An
525 accurate determination of the factors that can modulate it is a central concern when using this
526 biomarker to gauge contaminant impacts (Flammarion and Garric, 1997). Basal activity levels
527 were not reported for any of the fish species tested by Huggett et al. (2003), so there is no basis
528 for evaluating the significance of the EROD activities they presented. Similarly, the bile FAC

529 responses for a single fish species considered by Huggett et al. (2003) show a marginally
530 significant increase above three samples from the lower of two reference sites that were collected
531 a decade earlier by different researchers and analyzed by a different laboratory. Moreover, the
532 identity of the compounds that caused the bile FAC response, presumably peaks within the same
533 HPLC retention time interval as 3-ring PAH conjugates, is not clear. Unfortunately, Huggett et
534 al. (2003) failed to verify the fluorescent compounds they detected in their bile samples as
535 conjugates of PAH by gas chromatography-mass spectrometry (Krahn et al. 1992). These peaks
536 may have been interferences from naturally occurring compounds (Krahn et al. 1984), including
537 naturally occurring PAH such as retene (Lipiatou and Saliot 1992, 1991), or the metabolites of
538 organic pigments ingested during feeding. Also, the fish sampled were not controlled for sex,
539 age, reproductive status or differences in recently ingested prey, which may affect bile FAC
540 results (Collier and Varanasi 1992). Indeed, it appears that the basal response for bile
541 fluorescent compounds has not been established unequivocally for any species, making
542 interpretation of the evidence presented by Huggett et al. (2003) and Page et al. (2004)
543 problematic.

544 Woodin et al. (1997) did detect a slight but apparently significant increase in liver EROD
545 responses in an intertidal fish (gunnel, *Anoplarchus purpurescens*) collected from two reference
546 areas in PWS compared with basal responses, but it is not clear whether this CYP1A-inducing
547 agent was from a widespread background source or from undetected localized sources.

548 Overall, it seems unlikely that marine sediments contaminated by PAH from natural
549 petrogenic sources are capable of causing significant CYP1A induction in biota, when the extract
550 from the SPMD deployed at the Constantine Harbor failed to do so following injection into
551 juvenile trout. This deployment site is a mudflat containing ~ 560 ng PAH g^{-1} sediment derived

552 entirely from the eroded geologic sources (Short et al., 1996), and this concentration is within a
553 factor of 3 of the greatest background concentration found anywhere within PWS (Page et al.,
554 1996). The failure to accumulate sufficient PAH to stimulate a detectable EROD response when
555 injected, after deployment of the SPMD in intimate contact with the Constantine Harbor
556 sediments for nearly a month with the Constantine Harbor sediments, is strong evidence that the
557 PAH associated with these sediments is not sufficiently bioavailable to induce CYP1A in
558 exposed biota. Also, the general absence of EROD responses from SPMDs deployed at random
559 and control sites is consistent with the low intensity of human impacts (apart from the *Exxon*
560 *Valdez* oil spill) in PWS. Although human activities have affected scores of sites within PWS
561 (Page et al., 2006; Wooley, 2000), most of these sites were either concentrated around Sawmill
562 Bay or else were smallholdings that usually impacted less than 100 m of shoreline each. These
563 impacts were distributed over more than a century and their cumulative total affects less than
564 0.2% of the present day shoreline in PWS (Boehm et al., 2004). These small scale impacts,
565 coupled with a low population density, results in fewer and more localized contaminant sources.
566 The absence of CYP1A-inducing contaminants in our SPMDs from random and control sites is
567 therefore not surprising.

568

569 **4. Conclusions**

570

571 The results of this study demonstrate the value of SPMD dialysate injection for
572 elucidating contaminant sources and exposure effects. Method sensitivity is illustrated by the
573 small but significant differences among EROD responses, such as those between spring and
574 summer salmon stream EROD activities. Significant EROD induction occurs following injection

575 of dialysates from SPMDs exposed to part per trillion aqueous concentrations of potent inducers,
576 especially chrysenes (Short et al., 2008). This shows that the SPMD/EROD assay as described
577 in this study to be a cost-effective method for detecting these contaminants in comparison with
578 direct chemical analysis of the dialysate. Another advantage over direct chemical analysis is that
579 the identity of the EROD-inducing agents need not be known. Instead, this method assesses the
580 inductive potential of the bioavailable complex mixture, with all its inherent interactions, at a
581 given site. The dynamic range is sufficiently large to indicate contamination gradients that may
582 suggest sources, and to quantify the effects of the bioavailable contaminant fraction following
583 direct exposure. Method reproducibility is demonstrated by the biochemical responses from the
584 oiled sites, particularly EL56C where the replicates were 2 meters apart, and the apparent
585 absence of false negatives. Thus, the method provides a reliable means of evaluating exposure
586 to the bioavailable fraction of hydrophobic contaminants, as the response elicited is based on the
587 absorption of contaminants into the SPMD, which serves as a standardized surrogate for local
588 biota.

589 This method overcomes potential drawbacks associated with biomarkers when compared
590 to chemical analysis, as presented by Forbes et al. (2006), including cost, relevance, sensitivity,
591 reproducibility, and consistency. Technological advances, such as enzyme-linked
592 immunosorbent assay (ELISA) columns and assay kits, have led to substantial increases in ease
593 of use for EROD and other biomarkers. The EROD assay is relatively simple in both method
594 and required equipment; by comparison, determination of chemical residues in tissue or
595 environmental samples is much more complex and equipment-intensive, leading to higher per
596 sample costs than those spent on biomarkers (Dickerson, 1994; EPA, 1986). For each site in this
597 study, the cost of EROD assays was about 16% of that for chemical analyses. Finally, this

598 method allows an assessment of the total contribution of the complex mixture that can be taken
599 up by the targets, whether these compounds are detected analytically or not.

600 The SPMD extract injection method complements field testing of biota from
601 contaminated areas. This method measures the response that the bioavailable contaminant
602 fraction can produce with few confounding variables such as gender, age, reproductive status or
603 condition. CYP1A induction was selected here as it is a well-established biomarker of PAH
604 exposure. For breadth of applicability, the test organism we used is an appropriate choice as
605 rainbow trout P450 systems may be as complex as their mammalian counterparts (Buhler and
606 Wang-Buhler 1998). Ultimately, the suspected non-polar contaminants and the potential targets
607 determine which response(s) to measure and which test organism to use, but the method
608 presented here is flexible and can accommodate other types of analysis and different test
609 animals.

610

611

612 **Acknowledgements**

613

614 This study was supported in part by the *Exxon Valdez* Oil Spill Trustee Council, but the findings
615 and conclusions are those of the authors and do not necessarily reflect the views or positions of
616 the Council. Mention of products or services herein does not imply endorsement by the US
617 government.

618

619 **References**

620

- 621 Barron, M.G., Heintz, R., & Rice, S.D. (2004). Relative potency of PAHs and heterocycles as
622 aryl hydrocarbon receptor agonists in fish. *Marine Environmental Research*, 58, 95-100.
- 623 Blom, S. & Förlin, L. (1997). Effects of PCB on xenobiotic biotransformation enzyme activities
624 in the liver and 21-hydroxylation in the head kidney of juvenile rainbow trout. *Aquatic
625 Toxicology*, 39, 215-230.
- 626 Bodkin, J.L., Ballachey, B.E., Dean, T.A., Fukuyama, A.K., Jewett, S.C., McDonald, L.,
627 Monson, D.H., O'Clair, C.E., & VanBlaricom, G.R. (2002). Sea otter population status and
628 the process of recovery from the 1989 'Exxon Valdez' oil spill. *Marine Ecology Progress
629 Series*, 241, 237-253.
- 630 Boehm, P.D., Page, D.S., Brown, J.S., Neff, J.M., & Burns, W.A. (2004). Polycyclic aromatic
631 hydrocarbon levels in mussels from Prince William Sound, Alaska, USA, document the
632 return to baseline conditions. *Environmental Toxicology and Chemistry*, 23, 2916-2929.
- 633 Boehm, P.D., Page, D.S., Brown, J.S., Neff, J.M., & Bence, A.E. (2005). Comparison of
634 mussels and semi-permeable membrane devices as intertidal monitors of polycyclic
635 aromatic hydrocarbons at oil spill sites. *Marine Pollution Bulletin*, 50, 740-750.
- 636 Boehm, P.D., Neff, J.M., & Page, D.S. (2007). Assessment of polycyclic aromatic hydrocarbon
637 exposure in the waters of Prince William Sound after the Exxon Valdez oil spill: 1989-
638 2005. *Marine Pollution Bulletin*, 54, 339-367.
- 639 Bucheli, T.D., & Fent, K. (1995). Induction of cytochrome P450 as a biomarker for
640 environmental contamination in aquatic ecosystems. *Critical Reviews in Environmental
641 Science & Technology*, 25, 201-268.

- 642 Buhler, D.R., & Wang-Buhler, J.-L. (1998). Rainbow trout cytochrome P450s: purification,
643 molecular aspects, metabolic activity, induction and role in environmental monitoring.
644 *Comparative Biochemistry and Physiology Part C*, 121, 107-137.
- 645 Carls, M.G., Short, J.W., & Payne, J. (2006). Accumulation of polycyclic aromatic
646 hydrocarbons by *Neocalanus* copepods in Port Valdez, Alaska. *Marine Pollution Bulletin*,
647 52, 1480-1489.
- 648 Collier, T.K., & Varanasi, U. (1992). Hepatic activities of xenobiotic metabolizing enzymes and
649 biliary levels of xenobiotics in English Sole (*Parophrys vetulus*) exposed to environmental
650 contaminants. *Archives of Environmental Contamination and Toxicology*, 20, 462-473.
- 651 Dickerson, R.L., Hooper, M.J., Gard, N.W., Cobb, G.P., & Kendall, R.J. (1994). Toxicological
652 foundations of ecological risk assessment: biomarker development and interpretation
653 based on laboratory and wildlife species. *Environmental Health Perspectives*, 102, 65-69.
- 654 Echols, K.R., Gale, R.W., Schwartz, T.R., Huckins, J.N., Williams, L.L., Meadows, J.C., Morse,
655 D., Petty, J.D., Orazio, C.E., & Tillitt, D.E. (2000). Comparing polychlorinated biphenyl
656 concentrations and patterns in the Saginaw River using sediment, caged fish, and
657 semipermeable membrane devices. *Environmental Science & Technology*, 34, 4095-4102.
- 658 Ewald, G., Larsson, P., Linge, H., Okla, L., & Szarzi, N. (1998). Biotransport of organic
659 pollutants to an inland Alaska lake by migrating sockeye salmon (*Oncorhynchus nerka*).
660 *Arctic*, 51, 40-47.

- 661 Flammarion, P., & Garric, J. (1997). Cyprinids EROD activities in low contaminated rivers: a
662 relevant statistical approach to estimate reference levels for EROD biomarker?
663 *Chemosphere*, 35, 2375-2388.
- 664 Förlin, L., Andersson, T., Balk, L., Larsson, Å. (1995). Biochemical and physiological effects in
665 fish exposed to bleached kraft mill effluents. *Ecotoxicology and Environmental Safety*, 30,
666 164-170.
- 667 Forbes, V.E., Palmqvist, A., & Bach, L. (2006). The use and misuse of biomarkers in
668 ecotoxicology. *Environmental Toxicology and Chemistry*, 25, 272-280.
- 669 Hodson, P.V., Efler, S., Wilson, J.Y., El-Shaarawi, A., Maj, M., & Williams, T.G. (1996).
670 Measuring the potency of pulp mill effluents for induction of hepatic mixed function
671 oxygenase activity in fish. *Journal of Toxicology and Environmental Health*, 49, 83-110.
- 672 Huckins, J.N., Petty, J.D., Prest, H.F., Clark, R.C., Alvarez, D.A., Orazio, C.E., Lebo, J.A.,
673 Cranor, W.L., & Johnson, B.T. (2000). *A Guide for the Use of Semipermeable Membrane*
674 *Devices (SPMDs) as Samplers of Waterborne Hydrophobic Contaminants*. Columbia,
675 MO, USA: Columbia Environmental Research Center, U.S. Geological Survey.
- 676 Huckins, J.N., Petty, J.D., & Booij, K. (2006). *Monitors of Organic Chemicals in the*
677 *Environment: Semipermeable Membrane Devices*. New York, NY, USA: Springer.
- 678 Huggett, R.J., Neff, J.M., Stegeman, J.J., Woodin, B., Parker, K.R., & Brown, J.S. (2006).
679 Biomarkers of PAH exposure in an intertidal fish species from Prince William Sound,
680 Alaska: 2004-2005. *Environmental Science & Technology*, 40, 6513-6517.

- 681 Huggett, R.J., Stegeman, J.J., Page, D.S., Parker, K.R., & Brown, J.S. (2003). Biomarkers in fish
682 from Prince William Sound and the Gulf of Alaska: 1999-2000. *Environmental Science &*
683 *Technology*, 37, 4043-4051.
- 684 Huuskonen, S., Lindström-Seppä, P., Koponen, K., & Roy, S. (1996). Effects of non-ortho-
685 substituted polychlorinated biphenyls (congeners 77 and 126) on cytochrome P4501A and
686 conjugation activities in rainbow trout (*Oncorhynchus mykiss*). *Comparative Biochemistry*
687 *and Physiology*, 113C, 205-213.
- 688 Hyötyläinen, T., & Oikari, A. (1999). Assessment of the bioactivity of creosote-contaminated
689 sediment by liver biotransformation system of rainbow trout. *Ecotoxicology and*
690 *Environmental Safety*, 44, 253-258.
- 691 Jewett, S.C., Dean, T.A., Woodin, B.R., Hoberg, M.K., & Stegeman, J.J. (2002). Exposure to
692 hydrocarbons 10 years after the Exxon Valdez oil spill: evidence from cytochrome P4501A
693 and biliary FACs in nearshore demersal fishes. *Marine Environmental Research*, 54, 21-
694 48.
- 695 Kopponen, P., Tarhainen, J., Ruuskanen, J., Törrönen, R., & Kärenlampi, S. (1993). Peat
696 induces cytochrome P450IA1 in HEPA-1 cell line. Comparison with fly ashes from
697 combustion of peat, coal, heavy fuel oil and hazardous waste. *Chemosphere*, 26, 1409-
698 1506.
- 699 Krahn, M.M., Myers, M.S., Burrows, D.G., & Malins, D.C. (1984). Determination of
700 metabolites of xenobiotics in the bile of fish from polluted waterways. *Xenobiotica*, 14,
701 633-646.

- 702 Krahn, M.M., Burrows, D.G., Ylitalo, G.M., Brown, D.W., Wigren, C.A., Collier, T.K., Chan,
703 S.-L., & Varanasi, U. (1992). Mass spectrometric analysis for aromatic compounds in bile
704 of fish sampled after the Exxon Valdez oil spill. *Environmental Science & Technology*, 26,
705 116-126.
- 706 Krümmel, E.M., Macdonald, R.W., Kimpe, L.E., Gregory-Eaves, I., Demers, M.J., Smol, J.P.,
707 Finney, B., & Blais, J.M. (2003). Delivery of pollutants by spawning salmon. *Nature*, 425,
708 255-256.
- 709 Kvenvolden, K.A., Hostettler, F.D., Carlson, P.R., & Rapp, J.B. (1995). Ubiquitous tar balls with
710 a California-source signature on the shorelines of Prince William Sound, Alaska.
711 *Environmental Science & Technology*, 29, 2684-2694.
- 712 Lebo, J.A., Almeida, F.V., Cranor, W.L., Petty, J.D., Huckins, J.N., Rastall, A.C., Alvarez, D.A.,
713 Mogensen, B.B., & Johnson, B.T. Purification of the triolein to be used in SPMDs. SETAC
714 North America, November 12, 2003, at Austin, TX (USA). SETAC North America
715 Abstracts 2003.
- 716 Lee, R.F., & Anderson, J.W. (2005). Significance of cytochrome P450 system responses and
717 levels of bile fluorescent aromatic compounds in marine wildlife following oil spills.
718 *Marine Pollution Bulletin*, 50, 705-723.
- 719 Lemaire, P., Förlin, L., & Livingstone, D.R. (1996). Responses of hepatic biotransformation and
720 antioxidant enzymes to CYP1A-inducers (3-methylcholanthrene, β -naphthoflavone) in sea
721 bass (*Dicentrarchus labrax*), dab (*Limanda limanda*), and rainbow trout (*Oncorhynchus*
722 *mykiss*). *Aquatic Toxicology*, 36, 141-160.

- 723 Lipiatou, E., & Saliot, A. (1991). Fluxes and transport of anthropogenic and natural polycyclic
724 aromatic hydrocarbons in the western Mediterranean Sea. *Marine Chemistry*, 32, 51-71.
- 725 Lipiatou, E., & Saliot, A. (1992). Biogenic aromatic hydrocarbon geochemistry in the Rhone
726 River delta and in surface sediments from the open North-Western Mediterranean Sea.
727 *Estuarine, Gulf and Shelf Science*, 34, 515-531.
- 728 Lohmann, R., Corrigan, B.P., Howsam, M., Jones, K.C., & Ockenden, W.A. (2001). Further
729 developments in the use of semipermeable membrane devices (SPMDs) as passive air
730 samplers for persistent organic pollutants: field application in a spatial survey of PCDD/Fs
731 and PAHs. *Environmental Science & Technology*, 35, 2576-2582.
- 732 Matsuo, A.Y.O., Woodin, B.R., Reddy, C.M., Val, A.L., & Stegeman, J.J. (2006). Humic
733 substances and crude oil induce cytochrome P450 1A expression in the Amazonian fish
734 species. *Environmental Science & Technology*, 40, 2851-2858.
- 735 Meador, J.P., Stein, J.E., Reichert, W.L., & Varanasi, U. (1995). Bioaccumulation of polycyclic
736 aromatic hydrocarbons by marine organisms. In G. W. Ware (Ed.). *Reviews of*
737 *Environmental Contamination and Toxicology* (pp.79-165). New York, NY, USA:
738 Springer-Verlag,
- 739 Meadows, J.C., Echols, K.R., Huckins, J.N., Borsuk, F.A., Carline, R.F., & Tillitt, D.E. (1998).
740 Estimation of uptake rate constants for PCB congeners accumulated by semipermeable
741 membrane devices and brown trout (*Salmo trutta*). *Environmental Science & Technology*,
742 32, 1847-1852.

- 743 Miles, M.S., Ashton, B.M., East, R., Bourke, S., Griswold, M., & Overton, E.B. (2001).
744 Summary of chemistry results: shoreline monitoring study in Prince William Sound,
745 Alaska during the summer of 1999. IES/RCAT 01-01. Hazardous Materials Response and
746 Assessment Division, NOAA, Seattle, WA.
- 747 Neff, J.M., Bence, A.E., Parker, K.R., Page, D.S., Brown, J.S., & Boehm, P.D. (2006).
748 Bioavailability of polycyclic aromatic hydrocarbons from buried shoreline oil residues
749 thirteen years after the Exxon Valdez oil spill: A multispecies assessment. *Environmental*
750 *Toxicology and Chemistry*, 25, 947-961.
- 751 Neff, J.M., Boehm, P.D., Stubblefield, W.A., Kropp, R., & Page, D.S. (2003). Monitoring
752 recovery of Prince William Sound, Alaska, following the *Exxon Valdez* oil spill:
753 bioavailability of PAH in offshore sediments. In *Proceedings of the International Oil Spill*
754 *Conference, American Petroleum Institute, Washington, DC*, pp. 1-7.
- 755 Niimi, A.J., & Palazzo, V. (1986). Biological half-lives of eight polycyclic aromatic
756 hydrocarbons (PAHs) in rainbow trout (*Salmo gairdneri*). *Water Research*, 20, 503-507.
- 757 Niimi, A.J., & Dookhran, G.P. (1989). Dietary absorption efficiencies and elimination rates of
758 polycyclic aromatic hydrocarbons (PAHs) in rainbow trout (*Salmo gairdneri*).
759 *Environmental Toxicology and Chemistry*, 8, 719-722.
- 760 NRC: Committee on Biological Markers of the National Research Council. (1987). Biological
761 markers in environmental health research. *Environmental Health Perspectives*, 74, 3-9.
- 762 Ockenden, W.A., Sweetman, A.J., Prest, H.F., Steinnes, E., Jones, K.C. (1998). Toward an
763 understanding of the global atmospheric distribution of persistent organic pollutants: the

- 764 use of semipermeable membrane devices as time-integrated passive samplers.
765 *Environmental Science & Technology*, 32, 2795-2803.
- 766 Page, D.S., Boehm, P.D., Douglas, G.S., Bence, A.E., Burns, W.A., & Mankiewicz, P.J. (1999).
767 Pyrogenic polycyclic aromatic hydrocarbons in sediments record past human activity: a
768 case study in Prince William Sound, Alaska. *Marine Pollution Bulletin*, 38, 247-260.
- 769 Page, D.S., Boehm, P.D., Douglas, G.S., Bence, A.E., & Mankiewicz, P.J., (1996). The natural
770 petroleum hydrocarbon background in subtidal sediments of Prince William Sound,
771 Alaska, USA. *Environmental Toxicology and Chemistry*, 15, 1266-1281.
- 772 Page, D.S., Huggett, R.J., Stegeman, J.J., Parker, K.R., Woodin, B., Brown, J.S., & Bence, A.E.
773 (2004). Polycyclic aromatic hydrocarbon sources related to biomarker levels in fish from
774 Prince William Sound and the Gulf of Alaska. *Environmental Science & Technology*, 38,
775 4928-4936.
- 776 Page, D.S., Brown, J.S., Boehm, P.D., Bence, A.E., & Neff, J.M. (2006). A hierarchical
777 approach measures the aerial extent and concentration levels of PAH-contaminated
778 shoreline sediments at historic industrial sites in Prince William Sound, Alaska. *Marine
779 Pollution Bulletin*, 52, 367-379.
- 780 Parrott, J.L., Backus, S.M., Borgmann, A.I., & Swyripa, M. (1999). The use of semipermeable
781 membrane devices to concentrate chemicals in oil refinery effluent on the Mackenzie
782 River. *Arctic*, 52, 125-138.

- 783 Pohl, R.J., & Fouts, J.R. (1980). A rapid method for assaying the metabolism of 7-
784 ethoxyresorufin by microsomal subcellular fractions. *Analytical Biochemistry*, 107, 150-
785 155.
- 786 Sabaliūnas, D., Ellington, J., & Sabliūnienė, I. (1999). Screening bioavailable hydrophobic
787 toxicants in surface waters with semipermeable membrane devices: role of inherent oleic
788 acid in toxicity evaluations. *Ecotoxicology and Environmental Safety*, 44, 160-167.
- 789 Schmidt, G.M. (1977). The exchange of water between Prince William Sound and the Gulf of
790 Alaska. MSc thesis, University of Alaska, Fairbanks, AK
- 791 Schwartz, R.M. (2006). Comparison of mussels and semi-permeable membrane devices as inter-
792 tidal monitors of polycyclic aromatic hydrocarbons at oil spill sites (Boehm et al., 2005).
793 *Marine Pollution Bulletin*, 52, 121.
- 794 Short, J.W., & Babcock, M.M. (1996). Prespill and postspill concentrations of hydrocarbons in
795 mussels and sediments in Prince William Sound. In S. D. Rice, R. B. Spies, D. A. Wolfe &
796 B. A. Wright (Eds.). *Proceedings of the Exxon Valdez Oil Spill Symposium*. American
797 Fisheries Society Symposium 18, Bethesda, Maryland, 149-166.
- 798 Short, J.W., Irvine, G.V., Mann, D.H., Maselko, J.M., Pella, J.J., Lindeberg, M.R., Payne, J.R.,
799 Driskell, W.B., & Rice, S.D. (2007). Slightly weathered *Exxon Valdez* oil persists on Gulf
800 of Alaska beach sediments after 16 years. *Environmental Science & Technology*, 41, 1245-
801 1250.

- 802 Short, J.W., Kvenvolden, K.A., Carlson, P.R., Hostettler, F.D., Rosenbauer, R.J., & Wright, B.A.
803 (1999). Natural hydrocarbon background in benthic sediments of Prince William Sound,
804 Alaska: oil vs. coal. *Environmental Science & Technology*, *33*, 34-42.
- 805 Short, J.W., Lindeberg, M.R., Harris, P.M., Maselko, J.M., Pella, J.J., & Rice, S.D. (2004).
806 Estimate of oil persisting on the beaches of Prince William Sound 12 years after the *Exxon*
807 *Valdez* oil spill. *Environmental Science & Technology*, *38*, 1564-1572.
- 808 Short, J.W., Maselko, J.M., Lindeberg, M.R., Harris, P.M., & Rice, S.D. (2006). Vertical
809 distribution and probability of encountering intertidal Exxon Valdez oil on shorelines of
810 three embayments within Prince William Sound, Alaska. *Environmental Science &*
811 *Technology*, *40*, 3723-3729.
- 812 Short, J.W., & Springman, K.R. (2006). Identification of hydrocarbons in biological samples for
813 source determination. In: Z. Wang and S. A. Stout (Eds.). *Oil Spill Environmental*
814 *Forensics - Fingerprinting and Source Identification*. London: Elsevier Science.
- 815 Short, J.W., Springman, K.R., Lindeberg, M.R., Holland, L.G., Larsen, M.L., Sloan, C.A., Khan,
816 C., Hodson, P.V., & Rice, S.D. (2008). Semipermeable membrane devices link site-
817 specific contaminants to effects: Part II – a comparison of lingering Exxon Valdez oil with
818 other potential sources of CYP1A inducers in Prince William Sound, Alaska. *Marine*
819 *Environmental Research* (submitted).
- 820 Springman, K.R., Kurath, G., Anderson, J.J., & Emlen, J.M. (2005). Contaminants as viral
821 cofactors: assessing indirect population effects. *Aquatic Toxicology*, *71*, 13-23.

- 822 Stegeman, J.J., & Hahn, M.E. (1994). Biochemistry and molecular biology of monooxygenases:
823 current perspectives on forms, functions, and regulation of cytochrome P450 in aquatic
824 species. In: D. C. Malins and G. K. Ostrander (Eds.). *Aquatic Toxicology: Molecular,*
825 *Biochemical and Cellular Perspectives*. Lewis Publishers, Boca Raton, Florida, pp. 87-206.
- 826 Stern, G.A., Halsall, C.J., Barrie, L.A., Muir, D.C.G., Fellin, P., Rosenberg, B., Rovinsky, F.Y.,
827 Kononov, E.Y., & Pastuhov, B. (1997). Polychlorinated biphenyls in Arctic air .1.
828 Temporal and spatial trends: 1992-1994. *Environmental Science & Technology*, 31, 3619-
829 3628.
- 830 Sundberg, H., Tjärnlund, U., Åkerman, G., Blomberg, M., Ishaq, R., Grunder, K., Hammar, T.,
831 Broman, D., & Balk, L. (2005). The distribution and relative toxic potential of organic
832 chemicals in a PCB contaminated bay. *Marine Pollution Bulletin*, 50, 195-207.
- 833 Trust, K.A., Esler, D., Woodin, B.R., & Stegeman, J.J. (2000). Cytochrome P450 1A induction
834 in sea ducks inhabiting nearshore areas of Prince William Sound, Alaska. *Marine*
835 *Pollution Bulletin*, 40, 397-403.
- 836 U.S. EPA. (1986). Laboratory Manual Physical/Chemical Methods, SW 846. Washington, D.C.:
837 U.S. EPA.
- 838 Villeneuve, D.L., Crunkilton, R.L., & DeVita, W.M. (1997). Aryl hydrocarbon receptor-
839 mediated toxic potency of dissolved lipophilic organic contaminants collected from
840 Lincoln Creek, Milwaukee, Wisconsin, USA, to PLHC-1 (*Poeciliopsis lucida*) fish
841 hepatoma cells. *Environmental Toxicology and Chemistry*, 16, 977-984.

- 842 Whyte, J.J., Jung, R.E., Schmitt, C.J., & Tillitt, D.E. (2000). Ethoxyresorufin-*o*-deethylase
843 (EROD) activity in fish as a biomarker of chemical exposure. *Critical Reviews in*
844 *Toxicology*, 30, 347-570.
- 845 Woodin, B.R., Smolowitz, R.M., & Stegeman, J.J. (1997). Induction of cytochrome P4501A in
846 the intertidal fish *Anoplarchus purpureus* by Prudhoe Bay crude oil and environmental
847 induction in fish from Prince William Sound. *Environmental Science & Technology*, 31,
848 1198-1205.
- 849 Wooley, C. (2002). The myth of the "pristine environment": past human impacts in Prince
850 William Sound and the northern Gulf of Alaska. *Spill Science & Technology Bulletin*, 7,
851 89-104.
- 852
- 853

854 **Figure Legends**


855 Figure 1. Location of SPMD deployment sites in Prince William Sound, Alaska.

856 Figure 2. EROD response to SPMD extracts from the oiled ANS sites used in this study. Sites
857 KN114A1, KN114A2, KN136A and KN109A are located on Knight Island; EL56C1, EL56C2
858 and EL56C3 are sites on Eleanor Island; DI67A1 and DI67A2 are on Disk Island (See Short et
859 al., 2008 for a detailed map). The field blank (FB), field control (FC) and positive control (β -
860 naphthoflavone; BNF) responses are included for comparison. Mean EROD activities are derived
861 from the logarithmic mean of (n) individuals, with ± 1 SE. White bars indicate 2 d response and
862 grey bars are 7 d. No data are available where noted.

863
864 Figure 3. EROD response to SPMD extracts from the human activity and salmon hatchery sites
865 (as noted) including Armin F. Koernig (AFK) hatchery. White bars indicate 2 d response and
866 grey bars are 7 d. Asterisks indicate significant increases (t -test $P < 0.05$).


867
868 Figure 4. EROD responses to SPMD extracts from salmon streams (SS) and random sites (RS)
869 in both fall and spring. Responses from the peanut oil injections (PNUT) are included for
870 comparison. White bars indicate 2 d response, grey bars are 7 d. Asterisks indicate significant
871 increases compared to PNUT ($P < 0.036$).

872




873
 874
 875
 876
 877

Figure 1. Location of SPMD deployment sites in Prince William Sound, Alaska.




878
 879 Figure 2. EROD response to SPMD extracts from the oiled ANS sites used in this study. Sites
 880 KN114A1, KN114A2, KN136A and KN109A are located on Knight Island; EL56C1, EL56C2
 881 and EL56C3 are sites on Eleanor Island; DI67A1 and DI67A2 are on Disk Island (See Short et
 882 al., 2008 for a detailed map). The field blank (FB), field control (FC) and positive control (β -
 883 naphthoflavone; BNF) responses are included for comparison. Mean EROD activities are derived
 884 from the logarithmic mean of (n) individuals, with ± 1 SE. White bars indicate 2 d response and
 885 grey bars are 7 d. No data are available where noted.

886



887
 888 Figure 3. EROD response to SPMD extracts from the human activity and salmon hatchery sites
 889 (as noted) including Armin F. Koernig (AFK) hatchery. White bars indicate 2 d response and
 890 grey bars are 7 d. Asterisks indicate significant increases (t -test $P < 0.05$).

891



892
893
894
895
896 Figure 4. EROD responses to SPMD extracts from salmon streams (SS) and random sites (RS)
897 in both fall and spring. Responses from the peanut oil injections (PNUT) are included for
898 comparison. White bars indicate 2 d response, grey bars are 7 d. Asterisks indicate significant
899 increases compared to PNUT ($P < 0.036$).

900

901 Table 1. Basal EROD activity of juvenile rainbow trout (denoted “not injected”) compared with
 902 EROD responses to negative and positive control treatments and responses from SPMD
 903 deployments at un-contaminated sites. Mean EROD activities are given as $\text{pmol mg}^{-1} \text{ min}^{-1}$,
 904 derived from the logarithmic mean of (*n*) individuals, with ± 1 SE in parentheses below.
 905

<u>Site/Treatment Category</u>	<u>EROD Activity</u>	
Not Injected	0.932 (5) (0.849 – 1.02)	
	<u>2-Day</u>	<u>7-Day</u>
Peanut Oil	1.87 (5) (1.66 – 2.10)	1.13 (5) (0.835 – 1.53)
Field Blanks	3.78 (22) (3.41 – 4.21)	2.45 (20) (2.07 – 2.90)
Un-Contaminated Sites	2.24 (80) (2.10 – 2.38)	1.19 (78) (1.11 – 1.27)
BNF	75.3 (4) (64.6 – 87.7)	65.8 (5) (51.5 – 84.0)
Cordova Harbor	40.3 (10) (34.9 – 46.5)	no data

906
 907
 908