



HAL
open science

Polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-PCBs) in harbor seal () livers from San Francisco Bay, California and Gulf of Maine

June-Soo Park, Olga Ioanna Kalantzi, Dianne Kopec, Myrto Petreas

► **To cite this version:**

June-Soo Park, Olga Ioanna Kalantzi, Dianne Kopec, Myrto Petreas. Polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-PCBs) in harbor seal () livers from San Francisco Bay, California and Gulf of Maine. *Marine Environmental Research*, 2009, 67 (3), pp.129. 10.1016/j.marenvres.2008.12.003 . hal-00563064

HAL Id: hal-00563064

<https://hal.science/hal-00563064v1>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-PCBs) in harbor seal (*Phoca vitulina*) livers from San Francisco Bay, California and Gulf of Maine

June-Soo Park, Olga Ioanna Kalantzi, Dianne Kopec, Myrto Petreas

PII: S0141-1136(08)00251-1
DOI: [10.1016/j.marenvres.2008.12.003](https://doi.org/10.1016/j.marenvres.2008.12.003)
Reference: MERE 3308

To appear in: *Marine Environmental Research*

Received Date: 9 August 2007
Revised Date: 11 December 2008
Accepted Date: 12 December 2008

Please cite this article as: Park, J-S., Kalantzi, O.I., Kopec, D., Petreas, M., Polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-PCBs) in harbor seal (*Phoca vitulina*) livers from San Francisco Bay, California and Gulf of Maine, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.12.003](https://doi.org/10.1016/j.marenvres.2008.12.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-PCBs) in**
2 **harbor seal (*Phoca vitulina*) livers from San Francisco Bay, California and Gulf of**
3 **Maine**

4

5

6 June-Soo Park¹, Olga Ioanna Kalantzi¹, Dianne Kopec², Myrto Petreas¹

7

8 ¹Department of Toxic Substances Control, California Environmental Protection Agency,
9 Berkeley, California 94710, USA; ²Department of Biological Sciences, University of
10 Maine, Orono, Maine 04469, USA

11

12

13

14

15

16

17

18

19

20

21

22

*Corresponding author. Email address: jpark@dtsc.ca.gov (J.S. Park)

23 **Abstract**

24 Bioaccumulation of endocrine disruptors in marine mammals positioned at the top of
25 the food chain is of toxicological concern. Stranded four pup and ten adult harbor seal
26 (*Phoca vitulina*) livers were collected from San Francisco Bay and the Gulf of Maine and
27 analyzed for polychlorinated biphenyls (PCBs) and their hydroxylated metabolites (OH-
28 PCBs). We used GC-ECD and GC-NCI/MS to investigate the presence of 28 PCBs and
29 8 OH-PCB metabolites, respectively. Σ_{28} PCB concentrations (di- to octa-CBs) ranged
30 from 1.81 to 35.9 $\mu\text{g/g}$ lipid with a median of 6.53 for the seal pups and 2.31 to 249 $\mu\text{g/g}$
31 lipid with a median of 28.9 for the adult seals. Σ_8 OH-PCB concentrations (penta- to
32 hepta-OH-PCBs) ranged from 0.02 to 0.69 $\mu\text{g/g}$ lipid with a median of 0.04 for the adult
33 seals, i.e., at much lower concentrations than those for PCBs. Ratios of OH-PCBs to
34 PCBs (0.24% on average) were comparable to those in beluga whale, but were lower than
35 ratios in human livers. The OH-PCB profiles were slightly different between SFB and
36 GOM seal livers although similar PCB congener patterns were observed. Generally, 4-
37 OH-CB107 was found predominantly in seal livers and was the only OH-PCB detectable
38 in most of seal pup livers. This study provides information on OH-PCBs in seals, adding
39 to the scarce exposure data for these chemicals.

40

41

42 *Key words:* hydroxylated PCB metabolites, polychlorinated biphenyls, harbor seal liver,

43 San Francisco Bay, Gulf of Maine

44 1. Introduction

45

46 Polychlorinated biphenyls (PCBs) are known endocrine disruptors that are persistent
47 and lipophilic, and biomagnify through the food chain (Wan et al., 2005). Since marine
48 mammals such as seals are positioned at the top of the marine food web and have a
49 relatively long life span (25-30 years), they are exposed to organochlorines (Neale et al.,
50 2005a; Shaw et al., 2005) and are consequently vulnerable to endocrine disruptive effects
51 such as reproductive impairment (Reijnders, 1986; Reddy et al., 2001), cancer (Martineau
52 et al., 2002; Ylitalo, 2005), thyroid alteration (Sormo et al., 2005; Tabuchi et al., 2006),
53 and immune suppression/infectious diseases (Ross, 2002; Beckmen et al., 2003; Hall et
54 al., 2006; Hammond et al., 2005). PCBs are stored in the insulating blubber (lipid) and
55 can be subsequently remobilized to the blood through lipid metabolism during the
56 seasonal fasting (Debier et al., 2006), posing a health risk.

57 PCB metabolites; hydroxyl and methylsulfone PCBs (OH- and MeSO₂-PCBs) exert
58 similar toxicological effects to PCBs in mammals, particularly thyroid alteration in
59 laboratory rats (Meerts et al 2002), suggesting the need for their inclusion in risk
60 assessment studies. Marine mammals, including seals, are capable of producing
61 cytochrome P450 enzymes (Teramitsu et al., 2000) that biotransform PCBs to more
62 soluble/excretable forms, i.e., OH-PCBs. Accordingly, 837 mono OH-PCB congeners
63 can be theoretically formed in the body via the mechanisms of arene oxidation/1,2 shift
64 and/or direct oxygen insertion (Letcher et al., 2000). While most of the OH-PCBs tend to
65 be excreted (Letcher et al. 2000), some specific OH-PCBs have been retained
66 predominantly in the blood of humans (Fängström et al., 2002; Park et al., 2007a) and

67 marine mammals (Hoekstra et al., 2003; Houde et al., 2006) due to their preferential
68 binding to blood proteins (Letcher et al. 2000). OH-PCBs are also retained in the livers
69 of rats (Bergman et al., 1994), humans (Guvenius et al., 2002), and beluga whales
70 (McKinney et al., 2006), presumably by mechanisms similar to pentachlorophenol
71 conjugation to liver fatty acids and/or liver protein/enzymes (Leighty and Fentiman,
72 1982). A set of structurally known OH-PCB metabolites preferentially retained in the
73 body have an OH- group bound to a *para* carbon of a phenyl ring and chlorine atoms
74 bound to carbon adjacent to the OH-group (Lans et al., 1993).

75 In this pilot research study we used liver samples collected from stranded, dead
76 harbor seals from San Francisco Bay (SFB) and from an island off the Gulf of Maine
77 (GOM). Although the causes of death of the harbor seals stranded on SFB and GOM
78 shores were not known, it is possible that liver accumulation of PCBs and their
79 metabolites may have contributed directly/indirectly to some causative effects, such as
80 infectious disease and liver dysfunction. Thus, the circulation and bioaccumulation of
81 these contaminants in the liver of seals is of concern.

82 Therefore, we investigated the distribution of possible PCB precursors and OH-PCB
83 metabolites retained in liver tissues of harbor seals stranded in the SFB and the GOM
84 coast where there are the scarce exposure data for OH PCBs in seals which often carry
85 high PCB body burden (tens of ppm). The present study was also undertaken to provide
86 data to compare contaminant exposure in other marine mammals and humans.

87

88

89 **2. Materials and methods**

90

91 *2.1. Seal liver tissue sampling*

92

93 Seal liver samples analyzed in this study were collected from SFB as part of a larger
94 study that examined harbor seal population dynamics, health, contaminant residues and
95 prey selection (Kopec and Harvey, 1995). Liver samples were collected from stranded,
96 dead harbor seals (*Phoca vitulina*) between 1989 and 1998 found along the shores SFB,
97 an urban/industrialized estuary that hosts a year-round resident population of about 500
98 harbor seals. Liver tissues were also collected from stranded harbor seals between 2002
99 and 2003 from Mt. Desert Rock, an isolated island 20 miles offshore of the central Maine
100 coast. Seals hauling at Mt. Desert Rock are part of the larger regional seal population in
101 the GOM numbering over 74,000 harbor seals. As a pilot study, a subset of four pup and
102 five adult seal livers from SFB and five male seal livers from GOM were included in the
103 present study and analyzed for PCBs and OH-PCBs, as summarized in Table 1. All
104 harbor seal tissue collections were authorized by the National Marine Fisheries Service
105 (NMFS) in coordination with local stranding network facilities (College of the Atlantic,
106 Bar Harbor, Maine and Museum of Vertebrate Zoology, University of California,
107 Berkeley, California). SFB samples consisted of males and females while all GOM livers
108 were from males. All seal liver samples were collected during necropsy and were
109 immediately placed in clean vials (I-Chem) at -20 °C prior to analysis (Kopec and Harvey,
110 1995). Seals were grouped by age class based on standard length (Bigg, 1969).

111

112 *2.2. Chemicals and standards*

113

114 The following OH-PCB standards were purchased from Wellington Laboratory
115 (TerraChem Inc., USA) and used as reference standards for the identification and
116 quantification of the analytes: 2,3,3',4',5-pentachlorobiphenyl-4-ol (4-OH-CB107),
117 2,2',3,4',5,5'-hexachlorobiphenyl-4-ol (4-OH-CB146), 2,2',3',4,4',5-hexachlorobiphenyl-
118 3-ol (3'-OH-CB138), 2,2',3,3',4',5-hexachlorobiphenyl-4-ol (4'-OH-CB130),
119 2,2',3,4',5,5',6-heptachlorobiphenyl-4-ol (4-OH-CB187), 2,2',3',4,4',5,5'-
120 heptachlorobiphenyl-3-ol (3'-OH-CB180), 2,2',3,3',4',5,5'-heptachlorobiphenyl-4-ol (4'-
121 OH-CB172). 2,2',4,4',5,5'-hexachlorobiphenyl-3-ol (3-OH-CB153) was provided by
122 Professor Åke Bergman (Stockholm University, Sweden). The numbering of PCBs and
123 OH-PCBs is based on that specified by Ballschmitter and Zell (1980) and Letcher et al
124 (2000), respectively. Internal surrogate standards for PCBs and OH-PCBs were CB-14,
125 65 and 166 for PCBs and 4'-OH-CB159 for OH-PCBs. Injection standards for PCBs and
126 OH-PCB analyses were CB-30 and 204, and CB-209, respectively. Diazomethane was
127 synthesized in hexane by using N-nitroso-N-methylurea (Sigma-Aldrich, USA) as
128 described elsewhere (Sandau, 2000). Other chemicals and solvents used for the analysis
129 include dichloromethane and hexane (trace analysis, Burdick & Jackson), methanol,
130 methyl-tert butyl ether, and water (HPLC grade, Fisher Sci., USA), 2-propanol (99.9%,
131 pesticide grade, Fisher Sci., USA), hydrochloric acid, sulfuric acid (98%), potassium
132 hydroxide, potassium chloride, sodium hydroxide, and ethyl alcohol (94-96%, 200 proof)
133 (Fisher Sci., USA), silica (200-400 mesh) (Sigma-Aldrich, USA). Organics in Cod liver
134 Oil (SRM1588b, National Institute of Standards and Technology, Gaithersburg, MD) was
135 used as a standard reference material for PCB analysis.

136

137 *2.3. Analysis and clean up*

138

139 The analyses were conducted at the ultra-clean laboratory of the Department of Toxic
140 Substances Control, Berkeley, CA. The analytical method was modified from analyses
141 of seal blubber (She et al., 2002) and blood serum (Park et al., 2007a). In summary, liver
142 samples (0.4-3 g) along with a method blank (sodium sulfate), and a standard reference
143 material (NIST 1588b) were homogenized, spiked with PCB and OH-PCB surrogate
144 standards (1-5 ng and 1 ng, respectively), and extracted with a mixture of methyl-*tert*-
145 butyl ether (MTBE):hexane (1:1, v/v) (75 mL) with 2-propanol (10 mL). The samples
146 went through two more extractions by successive shaking and centrifuging with
147 MTBE:hexane mixture. Lipid content was determined gravimetrically using 7% (v/v) of
148 the extracts. The extracts were treated with 1% KCl wash and KOH phase separation.
149 Only 7% (v/v) of the organic extracts were used for the PCB analysis to avoid saturation
150 of the ECD. The phenolic compounds retained in the KOH solution were re-protonated
151 by using 2M HCl (pH<2), extracted with MTBE:hexane (1:9, v/v) (4 mL), and
152 derivatized overnight by using diazomethane. The organic extract was cleaned up by
153 using concentrated sulfuric acid (98%). Further clean up and fractionation was
154 conducted on the Pasteur pipette column packed with acidic silica gel (1:2 w/w, 0.6g) and
155 activated silica (0.4g). The columns were eluted with 5 mL hexane and then 10 mL
156 dichloromethane:hexane (1:1). The OH-PCB derivatives were cleaned up with
157 concentrated sulfuric acid (98%), followed by passing through a Pasteur pipette column

158 packed with sulfuric acidic silica gel (0.5g) and activated silica (0.1g). Injection
159 standards were added before GC analysis.

160

161 2.4. Instruments

162

163 Twenty eight PCB congeners (CB-28, 49, 52, 56, 66, 70, 74, , 99, 101, 105, 110, 113,
164 118, 137, 138, 146, 153, 156, 157, 170, 177, 180, 183, 187, 189, 190, 194, 203) were
165 analyzed on a Varian 3800 gas chromatograph (GC) with ⁶³Ni electron capture detection
166 (Varian Inc., Walnut Creek, CA) equipped with dual capillary columns; RTX-5MS (60m
167 × 0.25 mm i.d., 0.25 μm thickness, Restek, Bellefonte, PA) and DB-XLB (60m × 0.25
168 mm i.d., 0.25 μm thickness, J&W Scientific, Folsom, CA). Carrier and make up gases
169 were helium and nitrogen, respectively and set in constant flow mode. Injection (2 μL)
170 was made in splitless mode with an injector temperature of 280 °C. The initial GC
171 temperature was set to 80 °C and held for 1.6 min followed by a 15 °C/min increase to
172 135 °C, 1 °C/min to 261 °C, 3 °C/min to 295 °C and 1 °C/min to 300 °C held for 3.5 min.
173 Post run was set to 320 °C for 1 min.

174 Eight OH-PCBs were determined as methyl derivatives by using a Varian 3800 gas
175 chromatograph equipped with a 1200L mass spectrometer (MS) (Varian Inc., Walnut
176 Creek, CA). The MS was operated in negative chemical ionization (NCI) mode with
177 electron energy of 70 eV and emission current of 300 μA using selected ion monitoring
178 (SIM). The GC was equipped with a DB-5MS capillary column (60m × 0.25 mm i.d.,
179 0.25 μm thickness, J&W Scientific, Folsom, CA, USA). Injection (2 μL) was made in
180 split/splitless mode with an injector temperature of 250 °C. The initial GC temperature

181 was set to 80 °C and held for 2 min followed by a 50 °C/min increase to 200 °C, 1 °C/min
182 to 230 °C, and 30 °C/min to 300 °C and held for 4 min. Post run was set to 320 °C for 1
183 min. The temperatures for both ion source and quadrupole were set to 150 °C. We
184 monitored the most intense ions, either molecular ion or fragmentation ion; [M⁻] for 4'-
185 MeO-CB159, [(M+2-CH₃)⁻] for 4-MeO-CB187 and 4'-MeO-CB172, and [(M-HCl)⁻] or
186 [(M+2-HCl)⁻] for the rest of the congeners. Carrier and reagent gases were helium and
187 methane, respectively.

188

189 2.5. Quantification and QA/QC

190

191 All glassware were washed, rinsed with acetone and hexane, and baked at 550 °C for
192 8 hours. Each batch consisted of one procedural blank, one SRM, and six samples. The
193 five points of external calibration curves were used for the quantifications of PCBs and
194 OH-PCBs. CB-28, 49, 52,66, 70, 74, 101, 105, 110, 118, 138, 146, 153, 156, 157, 177,
195 180, 183, 187 and 194 were quantitated on the RTX-5MS column while CB-99, 170, 190,
196 199, and 203 were quantitated on the DB-XLB column. We tested precision for the PCB
197 and OH-PCB analytical methods as well as the gravimetric lipid determination by using
198 duplicate liver samples within and between batches. We tested accuracy by using the
199 SRM samples. Both precision and accuracy were within reasonable error ranges (±25%).
200 To monitor the performance of the experimental procedures, 2.00 ng of 4'-OH-CB159,
201 which, to our knowledge, has not been detected in any animal liver to date, were added to
202 all samples. We derivatized OH-PCB standards simultaneously with the sample extracts
203 for more accurate quantification. Any values lower than the LOQ (~0.48 ng/g fat) were

204 replaced by LOQ/2 when calculating summary statistics. We conducted non-parametric
205 tests (e.g., Spearman correlation) for PCB and OH-PCB data to assess their relationships
206 by using Minitab statistical software.

207 The average (\pm standard deviation) surrogate recoveries of CB-14, CB-65, and CB-
208 166 for PCB analysis were $92\pm6\%$, $82\pm7\%$, and $82\pm19\%$, respectively. The recoveries
209 from Standard Reference Materials (SRM 1588b Organics in Cod Liver Oil) ranged from
210 63% for CB-194 to 125% for CB-170. The average (\pm standard deviation) recovery of
211 the OH-PCB surrogate (4'-OH-CB159) was $77\pm15\%$.

212

213

214 **3. Results and discussion**

215

216 In Table 2 we report the ranges and medians of PCBs and OH-PCBs (on a lipid
217 weight basis) measured in harbor seal liver tissues for four pups and five adults from SFB
218 and five adult males from GOM.

219

220 *3.1. PCBs*

221

222 Concentrations of PCBs measured from SFB seal pup livers ranged from 1.81 to 35.9
223 $\mu\text{g/g}$ fat while adult seal livers showed a wider range (2.31-249 $\mu\text{g/g}$ fat). Because of the
224 limited sample size we did not examine the impact of gender on SFB samples and
225 geographical comparisons. In other study, females tend to have lower contaminant
226 burdens than males since reproductively active females can transfer a significant portion

227 of their PCB body burden to their offsprings through gestation or nursing (Neale et al.,
228 2005b). During the winter, GOM seals may migrate southward along the coast to
229 urban/industrialized areas (e.g., Massachusetts Bay) (Waring et al., 2006) where they
230 may be exposed to elevated PCB levels, while SFB seals reside in the Bay for most of the
231 time.

232 CB-153 was the primary congener in both SFB and GOM adult seal livers,
233 comprising 22 and 31% on average of Σ_{28} PCBs, respectively (Table 2) and was a good
234 indicator for Σ_{28} PCBs ($r=0.98$, albeit the highest point was excluded). CB-153 together
235 with CB-138, 187, and 180, comprised 67% and 66% on average of Σ_{28} PCBs in SFB and
236 GOM, respectively. The nine PCB congeners presented in Table 2 accounted for more
237 than 80% of Σ_{28} PCBs in SFB and GOM seal livers. The average concentrations of
238 Σ_{28} PCBs measured in GOM harbor seal livers (28.3 ug/g fat) were comparable to
239 Σ_{38} PCBs measured in beluga whale livers (31.9 ug/g fat) from the St. Lawrence River
240 (McKinney et al., 2006). Although the number of congeners measured was different
241 between the McKinney and our study, most PCB congeners predominantly found in the
242 liver matrices, as well as the sample collection time, overlapped. In our study, 8 out of
243 14 seal livers (1 SFB male pup, 1 SFB female adult, 3 SFB male adults and 3 GOM male
244 adults) showed PCB levels within or exceeding the estimated threshold level for adverse
245 effects on immune/reproductive function (17-77 $\mu\text{g/g}$ blubber fat) in aquatic animals as
246 summarized elsewhere (Shaw et al., 2005). Seal liver (this study) and blubber PCB
247 concentrations (Park et al., unpublished data) showed correlation ($r=0.77$), which is
248 consistent with other studies (Wolkers et al 2006). Thus, it is possible that some of the
249 stranded harbor seals, particularly one SFB seal pup with high PCB liver accumulation,

250 might have suffered PCB-related adverse effects, such as infectious diseases and/or the
251 conditions of the animals presented in Table 1.

252

253 3.2. OH-PCB metabolites

254

255 Although literature searches indicated that most of the OH-PCBs are retained in blood,
256 we attempted to measure OH-PCBs in liver because major enzyme-mediated
257 biotransformation of PCBs occurs in the liver. OH-PCBs were detected in almost all
258 liver tissues of SFB and GOM harbor seals including SFB seal pups. 4-OH-CB107 was
259 detected in almost all samples, followed by 4-OH-CB187 (67%), 4-OH-CB146 (60%), 3'-
260 OH-CB138 (60%), and 3-OH-CB153 (47%). As shown in Figure 1, the two OH-PCB
261 chromatograms analyzed as MeO-PCBs were presented for one of the calibration
262 standards (top) and a SFB adult male seal liver (bottom). The standard chromatogram
263 indicates that our GC temperature program resolved the possible co-elutions (e.g., 3-OH-
264 CB153/4-OH-CB146, 3'-OH-CB138/4'-OH-CB130, and 3'-OH-CB180/4'-OH-CB172).
265 Several unidentified OH-PCB peaks were present in the SFB seal liver chromatogram.
266 Particularly, two peaks (a hexa and a hepta OH-PCB) observed between the peaks of 4'-
267 OH-CB159 (internal standard) and 3'-OH-CB180 were fairly notable. When these
268 unidentified OH-PCBs were quantified using response factors for 4-OH-CB107 and 4-
269 OH-CB187 standards representing each hydroxyl homologue group, they comprised 16-
270 27% (22% on average) of Σ OH-PCBs (sum of identified and unidentified).

271 The harbor seal liver concentrations of OH-PCB metabolites (0.020~0.693 μ g/g fat)
272 in this study had a wider range than what has been reported for humans (0.007~0.175

273 $\mu\text{g/g}$ fat) (Guvenius et al., 2002) and beluga whales ($<0.5\text{-}0.145$ $\mu\text{g/g}$ fat) (McKinney et
274 al., 2006) due to one extreme value. Excluding that value, the range narrowed
275 ($0.020\text{-}0.064$ $\mu\text{g/g}$ fat) to previously reported levels. The $\Sigma_8\text{OH-PCB}$ levels identified in
276 the harbor seals were much lower (~ 0.59 % of $\Sigma_{28}\text{PCBs}$, 0.24% on average) than those of
277 parent PCBs. It should be noted that the sums of 28 PCBs includes congeners which are
278 not precursors of 8 OH-PCBs and that, in addition, the presence of many OH-PCB
279 congeners below the detection limits may bias their sum. These ratios were comparable
280 to data from beluga whales in St. Lawrence River, but they were lower than data in
281 human livers (usually between 1-10%). It is possible that these lower ratios in marine
282 mammals relative to humans may be due to a reduced capacity in marine mammals to
283 metabolize PCBs, or a higher Phase II conjugation process which rapidly depletes OH-
284 PCBs (McKinney et al., 2006). This OH-PCB/PCB ratio was even higher in human
285 blood (Park et al., 2007a,b) and marine mammal blood (Hoekstra et al., 2003; Houde et
286 al., 2006), indicating that OH-PCB metabolites preferentially bind to blood protein
287 relative to liver fatty acids and/or hepatic proteins and enzymes. Since we did not
288 measure lower chlorinated OH-PCBs (containing fewer than four chlorines) and also did
289 not include the ones detected but not identified, we may have underestimated the total
290 concentrations of OH-PCB metabolites in seal livers.

291 The OH-PCB profiles are presented in the bottom of Figure 2. This profile was
292 slightly different between SFB and GOM seal livers although similar PCB congener
293 patterns were observed in the two regions (Figure 2, top). This difference may result
294 from biological (e.g., nutritional/reproductive status, age, gender, polymorphism) and
295 environmental factors (e.g., temperature, migration) influencing enzyme activities and

296 thereby metabolite formation in seals. For example, Wolkers et al. (2008) reported that
297 EROD activity negatively correlated to blubber content in seals, indicating contaminants
298 get mobilized from blubber during fasting, and possibly induce the Phase I and II enzyme
299 system. Seals also go through other seasonal lipid changes via molting and reproduction.
300 However, the contributions from other various OH-PCB exposure pathways including
301 diet intake/food chain accumulation (Campbell et al., 2003), and even the abiotic aquatic
302 environments (Ueno et al., 2007) are probably minor, as observed in a recent study
303 (Verreault et al., 2008).

304 Although the profiles of OH-PCB congeners varied from sample-to-sample, 4-OH-
305 CB107 was the dominant congener, similar to the results from the St. Lawrence River
306 beluga whales (McKinney et al., 2006). This was the only congener detected in three
307 SFB seal pup livers in this study, which raises concern due to its relationship to thyroid
308 dysfunction reported for the experimental rat fetus (Meerts et al., 2002). The known
309 possible PCB precursors (CB-118 and CB-105) of 4-OH-CB107 in animals (Letcher et
310 al., 2000) were also found in our samples, albeit they were not among the most dominant
311 congeners. With 3-OH-CB153, 4-OH-CB146, 3'-OH-CB138, and 4-OH-CB187, they
312 comprised 66 to 81% of Σ OH-PCBs (identified and unidentified penta, hexa, and hepta
313 OH-PCBs). They are the OH-PCB congeners primarily found in the blood of humans
314 (Park et al., 2007a,b) and wildlife (Park et al., 2008) since their chemical structure
315 preferentially binds to the blood transthyretin (TTR) receptor protein. This binding
316 involves a hydroxyl group in either the *para*- or *meta*-position of a biphenyl ring,
317 adjacent to chlorine atoms on both sides (Lans et al., 1993; Letcher et al., 2000). This
318 also seems to apply to the seal liver matrix, indicating interaction of those OH-PCBs with

319 liver enzyme/proteins as well as fatty acid conjugation. 4'-OH-CB130 was reported at
320 considerable levels in human livers (Guvenius et al., 2002) while it was not detected or
321 detected at trace levels in our seal livers.

322 The pairs of PCB precursors and respective OH-PCB metabolites were correlated.
323 Since correlations could be driven by a single extreme value, that value was removed and
324 the resulting unbiased correlations were: CB-153/3-OH-CB146 ($r=0.67$, $p<0.05$), CB-
325 138/3-OH-CB138 ($r=0.74$, $p<0.05$), CB-187/4-OH-CB187 ($r=0.66$, $p<0.05$). However,
326 the pairs of CB-118/4-OH-CB107 and CB-105/4-OH-CB107 showed weaker correlations
327 ($r=0.24$ and 0.32 , respectively), possibly because 4-OH-CB107 is labile (Malmberg et al.
328 2004). In the body, PCBs are metabolized via the various transformation pathways
329 mediated by the cytochrome P450 enzyme series (CYP1A, CYP2B, CYP3A).
330 Preferential CYP450 enzyme mediated-metabolic capacity of marine mammals towards
331 OH-PCBs depends on the number and positioning of the chlorine atoms around the
332 biphenyl ring (Boon et al., 1997; Li et al., 2003; McKinney et al., 2004). For example,
333 CB-118 and 105, having only ortho- and meta-unsubstituted sites and one or fewer ortho-
334 Cl atoms, were subject to the CYP1A type enzyme-mediated metabolism, while CB-52,
335 49, 101 and 110, with meta- and para- vicinal H-atoms, were preferentially metabolized
336 by CYP2B or CYP3A types of enzymes

337 We here report several OH-PCBs retained in harbor seal livers in quantifiable
338 amounts. This is one of a few studies available on OH-PCB metabolites detected in liver
339 tissues; rats (Bergman et al., 1994; Haraguchi et al., 1998), humans (Guvenius et al.,
340 2002), beluga whales (McKinney et al., 2006), glaucous gulls (Verreault et al., 2007),
341 polar bears (Gebbinck et al., 2008), and predatory birds (Jaspers et al., 2008). Data on seal

342 liver OH-PCB metabolites are especially valuable because of the difficulties in sampling,
343 the rarity of the sample itself, and the complexity of liver analyses for OH-PCB
344 metabolites.

345 There are no reports available to date on the toxicological index for hepatic OH-PCB
346 metabolites. However, due to their endocrine disrupting properties, they should be
347 included in the risk assessment of marine mammals' health.

348

349

350 **4. Summary and conclusions**

351

352 PCBs and OH-PCB metabolites were detected in livers of harbor seals from both the
353 San Francisco Bay and the Gulf of Maine. OH-PCB metabolites were retained in seal
354 liver, albeit at lower OH-PCBs to PCBs ratios compared to human liver, possibly due to a
355 lower metabolic expression of CYP enzymes. The OH-PCB profiles were slightly
356 different between SFB and GOM seal livers although similar PCB congener patterns
357 were observed between the two regions. In general, 4-OH-CB107 was detected
358 predominantly in seal livers and was the only OH-PCB detectable in most seal pup livers,
359 raising toxicological concerns. These results should be treated with caution, as they are
360 based on a small sample size. They do, however, highlight the need to include OH-PCB
361 metabolites in the risk assessment of marine mammals, due to their potential endocrine
362 disrupting properties. Further investigations are needed to better understand the
363 relationship between exposure to these endocrine disruptors and possible health effects in
364 seals.

365

366

367 **Acknowledgement and disclaimer**

368

369 The authors would like to thank Joginder Dhaliwal and Dr. Reber Brown of the
370 Department of Toxic Substances Control for their valuable analytical expertise. The
371 ideas and opinions expressed herein are those of the authors and do not necessarily reflect
372 the official position of the California Department of Toxic Substances Control.

373

374

375 **References**

376

377 Ballschmitter, K., Zell, M. 1980. Analysis of polychlorinated biphenyls (PCB) by glass
378 capillary gas chromatography. Composition of technical Aroclor- and Chlophen-PCB
379 mixtures. Fresenius' journal of analytical chemistry 302, 20-31.

380 Beckmen, K. B., Blake, J. E., Ylitalo, G. M., Stott, J. L., O'Hara, T. M., 2003.

381 Organochlorine contaminant exposure and associations with hematological and
382 humoral immune functional assays with dam age as a factor in free-ranging northern
383 fur seal pups (*Callorhinus ursinus*). Marine Pollution Bulletin 46, 594-606.

384 Bergman, Å., Klassonwehler, E., Kuroki, H., 1994. Selective Retention of Hydroxylated
385 Pcb Metabolites in Blood. Environmental Health Perspectives 102, 464-469.

386 Bigg, M. A., 1969. The harbour seal in British Columbia. Fish Res Board Canada,
387 Bulletin 172. 33 pp.

- 388 Boon, J. P., vanderMeer, J., Allchin, C. R., Law, R. J., Klunsoyr, J., Leonards, P. E. G.,
389 Spliid, H., StorrHansen, E., McKenzie, C., Wells, D. E., 1997. Concentration-
390 dependent changes of PCB patterns in fish-eating mammals: Structural evidence for
391 induction of cytochrome P450. *Archives of Environmental Contamination and*
392 *Toxicology* 33, 298-311.
- 393 Campbell, L. M., Muir, D. C. G., Whittle, D. M., Backus, S., Norstrom, R. J., Fisk, A. T.,
394 2003. Hydroxylated PCBs and other chlorinated phenolic compounds in lake trout
395 (*Salvelinus namaycush*) blood plasma from the Great Lakes Region. *Environmental*
396 *Science and Technology* 37, 1720-1725.
- 397 Debier, C., Chalon, C., Le Boeuf, B. J., de Tillesse, T., Larondelle, Y., Thome, J. P., 2006.
398 Mobilization of PCBs from blubber to blood in northern elephant seals (*Mirounga*
399 *angustirostris*) during the post-weaning fast. *Aquatic Toxicology* 80, 149-157.
- 400 Fångström, B., Athanasiadou, M., Grandjean, P., Weihe, P., Bergman, Å., 2002.
401 Hydroxylated PCB metabolites and PCBs in serum from pregnant Faroese women.
402 *Environmental Health Perspectives* 110, 895-899.
- 403 Gebbink, W. A., Sonne, C., Dietz, R., Kirkegaard, M., Riget, F. F., Born, E. W., Muir, D.
404 C. G., Letcher, R. J., 2008. Tissue-specific congener composition of organohalogen
405 and metabolite contaminants in East Greenland polar bears (*Ursus maritimus*).
406 *Environmental Pollution* 152 (3), 621-629.
- 407 Guvenius, D. M., Hassanzadeh, P., Bergman, Å., Noren, K., 2002. Metabolites of
408 polychlorinated biphenyls in human liver and adipose tissue. *Environmental*
409 *Toxicology and Chemistry* 21, 2264-2269.

- 410 Hall, A. J., Hugunin, K., Deaville, R., Law, R. J., Allchin, C. R., Jepson, P. D., 2006. The
411 risk of infection from polychlorinated biphenyl exposure in the harbor porpoise
412 (*Phocoena phocoena*): A case-control approach. *Environmental Health Perspectives*
413 114, 704-711.
- 414 Hammond, J. A., Hall, A. J., Dyrinda, E. A., 2005. Comparison of polychlorinated
415 biphenyl (PCB) induced effects on innate immune functions in harbor and grey seals.
416 *Aquatic Toxicology* 74, 126-138.
- 417 Haraguchi, K., Kato, Y., Kimura, R., Masuda, Y., 1998. Hydroxylation and
418 methylthiolation of mono-ortho-substituted polychlorinated biphenyls in rats:
419 Identification of metabolites with tissue affinity. *Chemical Research in Toxicology* 11
420 (12), 1508-1515.
- 421 Hoekstra, P. F., Letcher, R. J., O'Hara, T. M., Backus, S. M., Solomon, K. R., Muir, D. C.
422 G., 2003. Hydroxylated and methylsulfone-containing metabolites of polychlorinated
423 biphenyls in the plasma and blubber of bowhead whales (*Balaena mysticetus*).
424 *Environmental Toxicology and Chemistry* 22, 2650-2658.
- 425 Houde, M., Pacepavicius, G., Wells, R. S., Fair, P. A., Letcher, R. J., Alae, M., Bossart,
426 G. D., Hohn, A. A., Sweeney, J., Solomon, K. R., Muir, D. C. G., 2006.
427 Polychlorinated biphenyls and hydroxylated polychlorinated biphenyls in plasma of
428 bottlenose dolphins (*Tursiops truncatus*) from the Western Atlantic and the Gulf of
429 Mexico. *Environmental Science and Technology* 40, 5860-5866.
- 430 Jaspers, V. L. B., Dirtu, A. C., Eens, M., Neels, H., Covaci, A. 2008. Predatory bird
431 species show different patterns of hydroxylated polychlorinated biphenyls (HO-PCBs)

- 432 and polychlorinated biphenyls (PCBs). *Environmental Science and Technology* 42,
433 3465-3471.
- 434 Kopec, A. D., Harvey, J. T., 1995. Toxic pollutants, health indices and population
435 dynamics of harbor seals in San Francisco Bay, 1989–1992. Moss Landing Marine
436 Laboratories Technical Report 96-4. Moss Landing, CA, pp. 168
- 437 Lans, M. C., Klassonwehler, E., Willemsen, M., Meussen, E., Safe, S., Brouwer, A., 1993.
438 Structure-dependent, competitive interaction of hydroxy-polychlorobiphenyls,
439 hydroxy-dibenzo-p-dioxins and hydroxy-dibenzofurans with human transthyretin.
440 *Chemico-Biological Interactions* 88, 7-21.
- 441 Leighty, E. G., Fentiman, A. F., 1982. Conjugation of pentachlorophenol to palmitic acid
442 by liver-microsomes. *Bulletin of Environmental Contamination and Toxicology* 28,
443 329-333.
- 444 Letcher, R. J., Klasson, W. E., Bergman, Å., 2000. Methylsulfon and hydroxylated
445 metabolites of polychlorinated biphenyls. In: *Handbook of Environmental Chemistry.*
446 *New Types of Persistent Halogenated Compounds, Vol. 3, Part K* (Paasivirta J, ed).
447 Berlin: Springer-Verlag.
- 448 Li, H. X., Boon, J. P., Lewis, W. E., van den Berg, M., Nyman, M., Letcher, R. J., 2003.
449 Hepatic microsomal cytochrome P450 enzyme activity in relation to in vitro
450 metabolism/inhibition of polychlorinated biphenyls and testosterone in Baltic grey
451 seal (*Halichoerus grypus*). *Environmental Toxicology and Chemistry* 22, 636-644.
- 452 Malmberg, T., Hoogstraate, J., Bergman, Å., Wehler, E. K., 2004. Pharmacokinetics of
453 two major hydroxylated polychlorinated biphenyl metabolites with specific retention
454 in rat blood. *Xenobiotica* 34, 581-589.

- 455 Martineau, D., Lemberger, K., Dallaire, A., Labelle, P., Lipscomb, T. P., Michel, P.,
456 Mikaelian, I., 2002. Cancer in wildlife, a case study: Beluga from the St. Lawrence
457 estuary, Quebec, Canada. *Environmental Health Perspectives* 110, 285-292.
- 458 McKinney, M. A., Arukwe, A., De Guise, S., Martineau, D., Beland, P., Dallaire, A., Lair,
459 S., Lebeuf, M., Letcher, R. J., 2004. Characterization and profiling of hepatic
460 cytochromes P450 and phase II xenobiotic-metabolizing enzymes in beluga whales
461 (*Delphinapterus leucas*) from the St. Lawrence River Estuary and the Canadian
462 Arctic. *Aquatic Toxicology* 69, 35-49.
- 463 McKinney, M. A., De Guise, S., Martineau, D., Beland, P., Lebeuf, M., Letcher, R. J.,
464 2006. Organohalogen contaminants and metabolites in beluga whale (*Delphinapterus*
465 *leucas*) liver from two Canadian populations. *Environmental Toxicology and*
466 *Chemistry* 25, 1246-1257.
- 467 Meerts, I., Assink, Y., Cenijs, P. H., van den Berg, J. H. J., Weijers, B. M., Bergman, Å.,
468 Koeman, J. H., Brouwer, A., 2002. Placental transfer of a hydroxylated
469 polychlorinated biphenyl and effects on fetal and maternal thyroid hormone
470 homeostasis in the rat. *Toxicological Sciences* 68, 361-371.
- 471 Neale, J. C. C., Gulland, F. M. D., Schmelzer, K. R., Harvey, J. T., Berg, E. A., Allen, S.
472 G., Greig, D. J., Grigg, E. K., Tjeerdema, R. S., 2005a. Contaminant loads and
473 hematological correlates in the harbor seal (*Phoca vitulina*) of San Francisco Bay,
474 California. *Journal of Toxicology and Environmental Health-Part a-Current Issues* 68,
475 617-633.
- 476 Neale, J. C. C., Schmelzer, K. R., Gulland, F. M. D., Berg, E. A., Tjeerdema, R. S.,
477 2005b. Rapid communication: Organohalogen levels in harbor seal (*Phoca vitulina*)

- 478 pups increase with duration of nursing. *Journal of Toxicology and Environmental*
479 *Health-Part a-Current Issues* 68, 687-691.
- 480 Park, J. S., Linderholm, L., Charles, M. J., Athanasiadou, M., Petrik, J., Kocan, A.,
481 Drobna, B., Trnovec, T., Bergman, Å., Hertz-Picciotto, I., 2007a. Polychlorinated
482 biphenyls and their hydroxylated metabolites (OH-PCBs) in pregnant women from
483 eastern Slovakia. *Environmental Health Perspectives* 115, 20-27.
- 484 Park, J. S., Petreas, M., Cohn, B. A., Factor-Litvak, P., 2007b. Measurement of
485 hydroxylated PCB metabolites (OH-PCBs) in archived serum from 1950-60's
486 California mothers. *Organohalogen Compounds* 69, 754-757.
- 487 Park, J. S., Holden, A., Wang, Y., Chang, J., Heckly, S., McKeown, K., Jewell, N.,
488 Hooper, K., 2008. Hydroxylated metabolites of polychlorinated biphenyls (PCBs) in
489 California wild birds. *Organohalogen Compounds* 70, 157-160.
- 490 Reddy, M. L., Reif, J. S., Bachand, A., Ridgway, S. H., 2001. Opportunities for using
491 Navy marine mammals to explore associations between organochlorine contaminants
492 and unfavorable effects on reproduction. *Science of the Total Environment* 274, 171-
493 182.
- 494 Reijnders, P. J. H. 1986. Reproductive failure in common seals feeding on fish from
495 polluted coastal waters. *Nature* 324: 456-457.
- 496 Ross, P. S. 2002. The role of immunotoxic environmental contaminants in facilitating the
497 emergence of infectious diseases in marine mammals. *Human and Ecological Risk*
498 *Assessment* 8: 277-292.
- 499 Sandau, C. D. 2000. Analytical chemistry of hydroxylated metabolites of PCBs and other
500 halogenated phenolic compounds in blood and their relationship to thyroid hormone

- 501 and retinol homeostasis in humans and polar bears. Carleton University, Ottawa,
502 Ontario, Canada. Ph.D. Dissertation: pp 39-44.
- 503 Shaw, S. D., Brenner, D., Bourakovsky, A., Mahaffey, C. A., Perkins, C. R., 2005.
504 Polychlorinated biphenyls and chlorinated pesticides in harbor seals (*Phoca vitulina*
505 *concolor*) from the northwestern Atlantic coast. Marine Pollution Bulletin 50, 1069-
506 1084.
- 507 She, J. W., Petreas, M., Winkler, J., Visita, P., McKinney, M. and Kopec, D., 2002.
508 PBDEs in the San Francisco Bay Area: measurements in harbor seal blubber and
509 human breast adipose tissue. Chemosphere 46, 697-707.
- 510 Sormo, E. G., Jussi, I., Jussi, M., Braathen, M., Skaare, J. U., Jenssen, B. M., 2005.
511 Thyroid hormone status in gray seal (*Halichoerus grypus*) pups from the Baltic Sea
512 and the Atlantic Ocean in relation to organochlorine pollutants. Environmental
513 Toxicology and Chemistry 24, 610-616.
- 514 Tabuchi, M., Veldhoen, A., Dangerfield, N., Jeffries, S., Helbing, C. C., Ross, P. S., 2006.
515 PCB-related alteration of thyroid hormones and thyroid hormone receptor gene
516 expression in free-ranging Harbor seals (*Phoca vitulina*). Environmental Health
517 Perspectives 114, 1024-1031.
- 518 Teramitsu, I., Yamamoto, Y., Chiba, I., Iwata, H., Tanabe, S., Fujise, Y., Kazusaka, A.,
519 Akahori, F., Fujita, S., 2000. Identification of novel cytochrome P450 1A genes from
520 five marine mammal species. Aquatic Toxicology 51, 145-153.
- 521 Ueno, D., Darling, C., Alae, M., Campbell, L., Pacepavicius, G., Teixeira, C., Muir, D.,
522 2007. Detection of hydroxylated polychlorinated biphenyls (OH-PCBs) in the abiotic

- 523 environment: Surface water and precipitation from Ontario, Canada. Environmental
524 Science and Technology 41, 1841-1848.
- 525 Verreault, J., Dietz, R., Sonne, C., Gebbink, W. A., Shahmiri, S., Letcher, R. J., 2008.
526 Comparative fate of organohalogen contaminants in two top carnivores in Greenland:
527 Captive sledge dogs and wild polar bears. Comparative Chemistry and Physiology –
528 Toxicology and Pharmacology 147 (3), 306-315.
- 529 Verreault, J., Shahmiri, S., Gabrielsen, G. W., Letcher, R.J., 2007. Organohalogen and
530 metabolically-derived contaminants and associations with whole body constituents in
531 Norwegian Arctic glaucous gulls. Environment International 33 (6), 823-830.
- 532 Wan, Y., Hu, J. Y., Yang, M., An, L. H., An, W., Jin, X. H., Hattori, T., Itoh, M., 2005.
533 Characterization of trophic transfer for polychlorinated dibenzo-p-dioxins,
534 dibenzofurans, non- and mono-ortho polychlorinated biphenyls in the marine food
535 web of Bohai Bay, north China. Environmental Science and Technology 39, 2417-
536 2425.
- 537 Waring, G.T., Gilbert, J.R., Loftin, J., Cabana, N., 2006. Short-term movements of radio-
538 tagged harbor seals in New England. Northeastern Naturalist 13, 1-14.
- 539 Wolkers, H., Hammill, M. O., van Bavel, B., 2006. Tissue-specific accumulation and
540 lactational transfer of polychlorinated biphenyls, chlorinated pesticides, and
541 brominated flame retardants in hooded seals (*Cistophora cristata*) from the Gulf of St.
542 Lawrence: Applications for monitoring. Environmental Pollution 142 (3), 476-486.
- 543 Wolkers, H., Krafft B.A., van Bavel B., Helgason L. B., Lydersen C., Kovacs K.M. 2008.
544 Biomarker responses and decreasing contaminant levels in ringed seals (*Pusa hispida*)

545 from Svalbard, Norway. *Journal of Toxicology and Environmental Health: Part A*
546 71(15), 1009-1018.

547 Ylitalo, G. M. 2005. The role of organochlorines in cancer-associated mortality in
548 California sea lions (*Zalophus californianus*). *Marine Pollution Bulletin* 50, 30-39.

549

ACCEPTED MANUSCRIPT

550 Table 1
 551 Information of seal liver samples collected from San Francisco Bay (SFB) and Gulf of Maine (GOM)

Collection ID#	Area	Age	Sex	Condition	Season Collected	Stranding Location
AP01524	SFB	Fetus	F	poor	spring	Tiburon, SFB,
AP01531	SFB	Pup	F	poor	spring	Yerba Buena Island, SFB
AP01528	SFB	Fetus	M	poor	spring	Fremont, SFB
AP01530	SFB	Pup	M	good	spring	Berkeley, SFB
AP01527	SFB	AD	F	poor	fall	East Palo Alto, SFB
AP01526	SFB	AD	F	excellent	spring	Redwood Creek, SFB
AP01529	SFB	AD	M	good	spring	Alameda, SFB
AP01523	SFB	AD	M	good	spring	Marin Co. coast (previously captured in SFB)
AP01525	SFB	AD	M	good	fall	San Francisco, SFB
AP01533	ME	AD	M	good	summer	Mount Desert Rock, Maine
AP01536	ME	AD	M	good	summer	Mount Desert Rock, Maine
AP01535	ME	AD	M	moderate	summer	Mount Desert Rock, Maine
AP01534	ME	AD	M	moderate	summer	Mount Desert Rock, Maine
AP01537	ME	AD	M	excellent	summer	Mount Desert Rock, Maine

552 *length (cm)/weight (kg) to roughly quantify blubber thickness and general health.

553 Table 2
 554 Concentrations of PCBs and OH-PCBs measured from livers of dead and stranded harbor seals along San
 555 Francisco Bay (SFB) and Gulf of Maine (GOM)

	San Francisco Bay Seals						Gulf of Maine Seals		
	Pups (N=4)			Adults (N=5)			Adults (N=5)		
	min	max	median	min	max	median	min	max	median
<i>PCBs ($\mu\text{g/g lipid}$)</i>									
PCB118	0.02	0.48	0.07	0.03	0.32	0.17	0.03	0.19	0.08
PCB146	0.03	1.52	0.24	0.05	8.88	1.36	0.24	2.84	0.89
PCB153	0.18	8.58	1.32	0.24	67.3	9.26	1.45	22.2	7.27
PCB105	0.001	0.15	0.03	0.01	0.12	0.06	0.003	0.07	0.04
PCB138	0.16	6.13	0.97	0.23	38.9	5.39	1.09	13.5	4.23
PCB187	0.09	3.45	0.68	0.62	38.6	4.53	0.37	6.73	2.25
PCB183	0.04	1.37	0.22	0.06	9.77	1.23	0.18	1.99	0.69
PCB180	0.11	4.00	0.60	0.15	42.3	3.86	0.39	4.58	2.39
PCB170	0.04	1.72	0.39	0.07	10.0	1.40	0.29	2.22	0.83
Σ_{28} PCBs	1.81	35.9	6.53	2.31	249	35.6	5.60	63.7	22.4
<i>OH-PCBs ($\mu\text{g/g lipid}$)</i>									
4-OH-CB107	<0.001	0.018	0.004	<0.001	0.123	0.026	0.008	0.033	0.017
3-OH-CB153	<0.001	<0.001	<0.001	<0.001	0.036	<0.001	<0.001	0.006	0.002
4-OH-CB146	<0.001	<0.001	<0.001	<0.001	0.218	0.002	0.003	0.018	0.008
3'-OH-CB138	<0.001	0.003	<0.001	<0.001	0.104	<0.001	0.002	0.013	0.005
4'-OH-CB130	<0.001	0.001	<0.001	<0.001	0.034	<0.001	<0.001	0.002	<0.001
4-OH-CB187	<0.001	0.011	<0.001	0.013	0.149	0.019	0.004	0.019	0.008
3'-OH-CB180	<0.001	<0.001	<0.001	<0.001	0.017	<0.001	<0.001	<0.001	<0.001
4'-OH-CB172	<0.001	<0.001	<0.001	<0.001	0.011	<0.001	<0.001	<0.001	<0.001
Σ_8 OH-PCBs	0.004	0.034	0.004	0.017	0.693	0.045	0.019	0.064	0.045

Figure captions

Fig. 1. GC chromatograms of calibration standard (top) and identified and unidentified OH-PCB metabolites from an adult harbor seal liver collected from San Francisco Bay (SFB) (bottom).

Fig. 2. Profiles of PCB precursors (top) and OH-PCB metabolites (bottom) observed in adult harbor seal livers from San Francisco Bay (SFB) and Gulf of Maine (GOM). Error bars indicate standard errors.

ACCEPTED MANUSCRIPT



