



HAL
open science

Persistent Organic Pollutants (POPs) in killer whales () from the Crozet archipelago, southern Indian Ocean

Marie Noel, Lance Barrett-Lennard, Christophe Guinet, Neil Dangerfield,
Peter S.Ross

► **To cite this version:**

Marie Noel, Lance Barrett-Lennard, Christophe Guinet, Neil Dangerfield, Peter S.Ross. Persistent Organic Pollutants (POPs) in killer whales () from the Crozet archipelago, southern Indian Ocean. *Marine Environmental Research*, 2009, 68 (4), pp.196. 10.1016/j.marenvres.2009.06.009 . hal-00563086

HAL Id: hal-00563086

<https://hal.science/hal-00563086>

Submitted on 4 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Persistent Organic Pollutants (POPs) in killer whales (*Orcinus orca*) from the Crozet archipelago, southern Indian Ocean

Marie Noel, Lance Barrett-Lennard, Christophe Guinet, Neil Dangerfield, Peter S.Ross

PII: S0141-1136(09)00070-1
DOI: [10.1016/j.marenvres.2009.06.009](https://doi.org/10.1016/j.marenvres.2009.06.009)
Reference: MERE 3347

To appear in: *Marine Environmental Research*

Received Date: 10 January 2009
Revised Date: 24 April 2009
Accepted Date: 2 June 2009

Please cite this article as: Noel, M., Barrett-Lennard, L., Guinet, C., Dangerfield, N., S.Ross, P., Persistent Organic Pollutants (POPs) in killer whales (*Orcinus orca*) from the Crozet archipelago, southern Indian Ocean, *Marine Environmental Research* (2009), doi: [10.1016/j.marenvres.2009.06.009](https://doi.org/10.1016/j.marenvres.2009.06.009)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1
2
3
4
5
6 Persistent Organic Pollutants (POPs) in killer whales (*Orcinus orca*) from the
7
8 Crozet archipelago, southern Indian Ocean
9

10 Marie Noel^{a,b}, Lance Barrett-Lennard^c, Christophe Guinet^d, Neil Dangerfield^a, and Peter
11
12 S.Ross^{a*}
13
14

15
16 ^a Institute of Ocean Sciences, Fisheries and Oceans Canada, 9860 West Saanich Road,
17 P.O.Box 6000, Sidney, British Columbia V8L 4B2, Canada
18

19 ^b School of Earth and Ocean Sciences, University of Victoria, British Columbia V8W 3P6,
20 Canada
21

22
23 ^c Vancouver Aquarium, Marine Science Center, Vancouver, British Columbia V6B 3X8,
24 Canada
25

26 ^d Centre d'Etudes Biologiques de Chizé, Centre National de la Recherche Scientifique,
27 79360 Villiers-en-Bois, France
28

29
30 *Corresponding author:
31 email: peter.s.ross@dfo-mpo.gc.ca
32 Phone: (250) 363-6806
33 Fax: (250) 363-6807
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Persistent Organic Pollutants (POPs), including polychlorinated biphenyls (PCBs), polychlorinated dibenzo-*p*-dioxins (PCDDs), and dibenzofurans (PCDFs), are ubiquitous environmental contaminants of which significant concentrations are reported in upper trophic level animals. In 1998, we collected blubber biopsy samples (n=11) from killer whales (*Orcinus orca*) inhabiting the coastal waters around Possession Island, Crozet Archipelago, southern Indian Ocean, for contaminant analyses. Despite inhabiting an isolated region far removed from industrial activities, these killer whales can presently be considered among the most PCB-contaminated cetaceans in the southern hemisphere, with concentrations ranging from 4.4 to 20.5 mg/kg lipid weight (lw). PCDD levels ranged from below the detection limit (5 ng/kg) to 77.1 ng/kg lw and PCDF levels from below the detection limit (7 ng/kg) to 36.1 ng/kg lw. Over 70 % of our study animals had PCB concentrations which exceeded a 1.3 mg/kg PCB threshold established for endocrine disruption and immunotoxicity in free-ranging harbour seals, suggesting that organic contaminants cannot be ruled out as an additional threat to this declining population.

Keywords: killer whale, Crozet Archipelago, PCBs, PCDDs, PCDFs, TEQs

1. Introduction

Polychlorinated biphenyls (PCBs) and polychlorinated dibenzo-*p*-dioxins / dibenzofurans (PCDDs/Fs) are persistent organic pollutants (POPs). They are hydrophobic, have moderate vapour pressure, and low reactivity that allow them to be readily transported in the environment. “Global distillation” was described as the process by which POPs evaporate in the warmer regions and are then transported in the atmosphere towards the poles where they condense and are deposited. Cold regions, such as the Arctic and the Antarctic, have thus been described as important sinks for these contaminants (Wania and Mackay, 2001).

PCBs were extensively used by industrialized nations until the late 70s when they were largely banned. They were detected for the first time in the Antarctic environment in the early 70s. Atmospheric transport is believed to be the dominant pathway for PCB contamination in Antarctica where local sources are thought to be restricted to electrical equipment at isolated research stations (Risebrough et al., 1990).

PCDD/Fs are formed as by-products during different industrial and thermal processes such as emissions from metallurgical industries, municipal incinerators, pulp and paper mills, and the manufacture of chlorinated chemicals. Atmospheric transport is also thought to be the dominant pathway for the delivery of dioxins and furans to the Antarctic along with minor local sources such as incinerators at research stations (Lugar et al., 1996).

As long lived, high level trophic feeders, marine mammals tend to bioaccumulate POPs in lipid tissues. Marine mammals from Europe, North America, Asia, and the Arctic have been found to be particularly contaminated with PCBs (Bergman et al., 2001);

1
2
3 Kajiwaru et al., 2006; Muir et al., 2000; Ross et al., 2000). Marine mammal-eating transient
4
5 killer whales (*Orcinus orca*) from the Northeastern Pacific Ocean are reported as among
6
7 the most PCB-contaminated marine mammals in the world (Ross et al., 2000) and killer
8
9 whales from Norway are the most PCB-contaminated animals in the Arctic (Wolkers et al.,
10
11 2007).
12
13

14 The biology of killer whales from the coastal waters of Possession Island, Crozet
15
16 Archipelago, southern Indian Ocean, has been studied since the late 1980s, but no
17
18 information exists on their exposure to POPs. Killer whales are sighted in the area year
19
20 round but some photo-identified individuals have been seen at the nearby Kerguelen Island
21
22 and close to the Antarctic continent. Killer whales in the Crozet Archipelago feed on a
23
24 variety of prey including fish, penguins (*Eudyptes sp.*), southern elephant seals (*Mirounga*
25
26 *leonina*) and, occasionally, large cetaceans (Guinet, 1991). Between 1988 and 2002, this
27
28 population declined by 50% due to the decline of an important prey species (the southern
29
30 elephant seal), interactions with the Patagonian toothfish (*Dissostichus eleginoides*)
31
32 longline fishery, and/or emigration of individuals or groups from the coastal waters of the
33
34 Crozet Archipelago (Poncelet et al., 2008).
35
36
37
38
39

40 POPs are known to cause hormone disruption, immunosuppression, and
41
42 reproductive failure in other marine mammals (Mos et al., 2007a; Reijnders, 1986; Ross et
43
44 al., 1995; Tabuchi et al., 2006). Such persistent contaminants can represent a population-
45
46 level conservation threat to killer whales (Ross, 2006), with heavy POP accumulation
47
48 largely due to their high trophic position and long lifespan (Hickie et al., 2007). In
49
50 analysing PCB, PCDD, and PCDF concentrations in blubber biopsies taken from the
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 declining Crozet Archipelago population, we aimed to provide a much-needed baseline for
4
5 these priority contaminants, and to characterize associated health risks.
6
7

8 9 10 **2. Materials and methods**

11 12 *2.1 Sampling*

13
14 The study was carried out in the Crozet Archipelago, southern Indian Ocean
15 (46°25'S; 51°59'E) (Figure 1). Lightweight pneumatic darts were used to biopsy killer
16
17 whales inhabiting the coastal waters of Possession Island. The variable-power dart projector
18
19 and its stainless-steel, 6.4 mm diameter tip, as well as a full description of the sampling
20
21 procedure, is described elsewhere (Barrett-Lennard et al., 1996). This method was adapted
22
23 to enable shore-based, rather than ship-based, collections.
24
25
26
27

28
29 Briefly, biopsy sampling was conducted from shore with a fine fishing line attached
30
31 to the dart in order to retrieve biopsies taken posterior to, and below, the dorsal fin of the
32
33 animal. Samples of epidermal, dermal, and hypodermal tissue typically weighed
34
35 approximately 0.5g. Skin from the biopsy samples was preserved in dimethyl sulphonyde
36
37 and a saturated salt solution and stored at 4°C for DNA extraction. Blubber from the
38
39 biopsies was placed in pesticide grade hexane-rinsed glass vials with aluminium foil-
40
41 covered caps and stored at -20°C for contaminant analyses. 11 blubber biopsies were
42
43 collected from nine different killer whales. Genetic results from the present samples,
44
45 combined with the photo-identification work conducted previously (Guinet, 1991), enabled
46
47 a determination of sex as well as age class for each of the nine individuals.
48
49
50
51
52
53
54

55 *2.2 Contaminant analyses*

56
57
58
59
60
61
62
63
64
65

1
2
3 Blubber samples were analyzed for PCDDs, PCDFs, and mono-ortho, di-ortho, and
4 non-ortho (planar) PCBs. Thawed blubber was ground in a porcelain mortar and pestle with
5 200 g of anhydrous sodium sulphate and spiked with a mixture of ^{13}C -labelled PCBs,
6 PCDDs, and PCDFs. The blubber-sodium sulphate mixture was transferred to an extraction
7 column and extracted with 250 mL of 1:1 dichloromethane/hexane (DCM/hex) from a glass
8 column by gravity flow. The extract was evaporated to dryness and the lipid content of the
9 samples was determined. Analyses of cleaned-up samples were conducted by high
10 resolution gas chromatography / high resolution mass spectrometry (HRGC/HRMS).
11 Procedural blanks were analyzed along with samples to assess the integrity of the analytical
12 procedures. Details on the sample clean-up, instrumental analysis condition used,
13 quantification protocols, criteria used for congener identification and the quality assurance /
14 quality control (QA/QC) measures undertaken for the HRGC/HRMS analysis of all the
15 analytes of interest are described elsewhere (Ikonou et al., 1999; Ikonou et al., 2001;
16 Rantaleinen et al., 1998).

2.3 Data Analysis

21 Two killer whales were biopsied twice allowing for replicate comparison. For one
22 of these two whales, contaminant concentrations differed between the two samples.
23 However, in the case of this individual, one of the biopsies was taken on the dorsal ridge
24 just behind the dorsal fin. This sample was low in lipid content, not representative of a
25 typical biopsy and therefore eliminated from our analysis. For the other whale, the two
26 biopsies provided similar results (2.3% and 8.6% difference between the two samples for
27 PCBs and PCDDs/Fs, respectively). Average contaminant concentrations were considered

1
2
3 for this whale. Although we were unable to age sampled individuals, we were able to divide
4
5 our study animals into three categories for POP comparisons: juvenile females (n = 3),
6
7 adult females (n = 5) and adult male (n = 1).
8
9

10 All the concentrations were expressed on a lipid weight basis (lw) and were blank-
11
12 corrected as well as recovery-corrected using the stable isotope dilution method based on
13
14 ¹³C-labelled PCB, PCDD, and PCDF spikes (Ikonomou et al., 2001) (the recovery ranges
15
16 were a mean of 60% for PCBs (range 20-95%), 82% for PCDDs (50-105%), and 81% for
17
18 PCDFs (60-100%)).
19
20

21
22 Many congeners were not detected. When congeners were undetected, substitutions
23
24 were applied according to the following rule: 1) when congeners were detected in less than
25
26 70% of the samples, concentrations of 0 was substituted; and, 2) when congeners were
27
28 detected in more than 70% of the samples, a detection limit value substitution was applied.
29
30

31 Total Toxic Equivalents (TEQ) to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD)
32
33 were calculated for all dioxin-like PCBs measured (mono-ortho PCBs 105, 114, 118, 123,
34
35 156, 157, 167, and 189; non-ortho PCBs 77, 81, 126, and 169) and 2,3,7,8-Cl substituted
36
37 PCDDs (n = 7) and PCDFs (n=10) using the international Toxic Equivalency Factors (TEF)
38
39 (Van den Berg et al., 2006).
40
41
42
43
44

45 *2.4 Statistical analysis*

46
47 Differences in contaminant levels were assessed using an unpaired t-test between
48
49 juvenile and adult females. The only adult male was considered separately due to evidence
50
51 in the literature of sex-based differences in POP concentrations (Ross et al., 2000).
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Results and discussion

PCB and PCDD/F data were available for 10 blubber biopsies taken from nine killer whales inhabiting the coastal waters of Possession Island, Crozet Archipelago. This represents the first information on POP contamination for this declining population of killer whales.

3.1 PCBs

3.1.1 Concentrations

Of the 209 PCB congeners sought, we detected 124 ± 8 (range: 119 – 134) congeners in our killer whale samples. PCB levels ranged from 2.7 to 20.5 mg/kg lw (Table 1). Concentrations in juvenile and adult females did not differ ($p > 0.05$), in contrast to reported findings for killer whales from the Northeastern Pacific Ocean, where immature whales had higher PCB concentrations than adult females (Ross et al., 2000). While the only male killer whale in the present study had the highest PCB levels of all individuals sampled in our study (Figure 2, Table 1), this could not be statistically compared with our other age/sex categories. Higher POP concentrations in males compared to females have been reported in various species of marine mammals including killer whales from the Northeastern Pacific Ocean (Ross et al., 2000) and in pilot whales (*Globicephala melaena*) (Borrell et al., 1995). In marine mammals, a continuous increase of POP concentrations with age is observed in immature individual of both sexes and in adult males. In adult females, concentrations start to decline when they reach sexual maturity because of the offload of fat-soluble contaminants to their young during gestation and nursing (Borrell et al., 1995; Ross et al., 2000; Ylitalo et al., 2001).

1
2
3 The Possession Island killer whales were approximately 10 times more
4
5 contaminated than the type C killer whales sampled recently from the southern Ross Sea
6
7 adjacent to Antarctica (Krahn et al., 2008), likely reflecting a difference in diet and possibly
8
9 the remote location of the Ross Sea. While PCB concentrations in our whales were lower
10
11 than those reported in killer whales from industrialized areas (Table 2), they were similar
12
13 to those observed in resident killer whales from Alaska (Ylitalo et al., 2001), and only
14
15 slightly lower than the levels in northern residents from the Northeastern Pacific Ocean
16
17 (Ross et al., 2000) and Norwegian killer whales (Wolkers et al., 2007). As the latter killer
18
19 whales are considered the most contaminated Arctic marine mammal, our results again
20
21 underscore the vulnerability of high trophic level marine mammals in polar and otherwise
22
23 remote regions.
24
25
26
27

28
29 However, the Norwegian killer whales and the resident killer whales of the
30
31 Northeastern Pacific Ocean are fish-eating, while the Possession Island killer whales appear
32
33 to consume a mixed diet of mostly marine mammals but also fish and penguins (Guinet,
34
35 1991). Recently, a juvenile male was found stranded on Possession Island and the stomach
36
37 content revealed remains of elephant seals, fur seals (*Arctocephalus sp*), and king penguins
38
39 but no fish. Dietary preferences are a major factor affecting contaminant concentrations in
40
41 marine mammals. In the Northeastern Pacific Ocean, transient marine mammal-eating killer
42
43 whales are far more contaminated than the resident fish-eating killer whales (Ross et al.,
44
45 2000). Based on the limited data available, the diet of the Possession Island killer whales
46
47 seems trophically comparable to transient killer whales of the Northeastern Pacific Ocean
48
49 and thus comparing contaminant burdens between these two populations provides at least a
50
51 crude evaluation of differences in regional contamination. The present levels in the
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 Possession Island killer whales were one fifth to one tenth of the levels recorded in
4
5 transients from the Northeastern Pacific Ocean (Ross et al., 2000) and Alaska (Ylitalo et al.,
6
7 2001). These differences could be the result of dietary preferences but may also be due to
8
9 the remote nature of Possession Island.
10

11
12 PCB concentrations in Crozet killer whales were higher than levels reported in
13
14 marine mammals from Antarctica, including southern elephant seals (Miranda-Filho et al.,
15
16 2007), and minke whales (*Balaenoptera acutorostrata*) (Aono et al., 1997). The Crozet
17
18 killer whales were less PCB-contaminated than only two other southern hemisphere
19
20 species, namely the estuarine dolphins (*Sotalia guianensis*) and Atlantic spotted dolphins
21
22 (*Stenella frontalis*) inhabiting the nearshore waters of Brazil (Kajiwara et al., 2004). The
23
24 Crozet whales had higher PCB concentrations than Arctic belugas (*Delphinapterus leucas*)
25
26 (Helm et al., 2002) that are at the receiving end of PCB dispersion from northern
27
28 hemispheric industries.
29
30
31
32
33
34
35

36 3.1.2 Homologue group and congener patterns

37

38 No significant differences ($p > 0.05$) in PCB homologue group and congener
39
40 patterns were observed between juvenile and adult females (Figure 3). The adult male also
41
42 revealed a similar pattern. In all the three groups, di-ortho PCBs were highly dominant
43
44 (97.5 ± 1.1 % of \sum PCBs) (Figure 2). Hexa-CBs was the dominant homologue group,
45
46 accounting for 55 ± 3 % of total PCBs, followed by hepta- (23 ± 2 %) and penta-CBs ($11 \pm$
47
48 3 %). A similar pattern was found in southern elephant seals but they had a more even
49
50 contribution of hepta and penta-CBs (Miranda-Filho et al., 2007). The six dominant
51
52 congeners in our killer whales accounted for 55 ± 1 % of \sum PCBs (Table 1).
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 Similar to our results, PCBs-153 and 138 were among the dominant congeners in
4
5 the three populations of killer whales from the Northeastern Pacific Ocean (Ross et al.,
6
7 2000), in killer whales from Alaska (Ylitalo et al., 2001), and Norway (Wolkers et al.,
8
9 2007). In addition, PCB 153 and 138 dominated PCB patterns in southern elephant seals
10
11 (Miranda-Filho et al., 2007) as well as in lower trophic levels of the Antarctic killer whale
12
13 food web (Goerke et al., 2004). This is likely due to the fact that PCB-153 was a major
14
15 component of the most heavily used PCB technical mixture, and, as it has no vicinal
16
17 hydrogen and is Cl-substituted at both para positions, is resistant to metabolism (Boon and
18
19 Eijgenraam, 1988).
20
21
22
23
24
25

26 3.2 Dioxins and furans

27 3.2.1 Concentrations

28
29 Dioxin and furan concentrations were very low with many congeners being
30
31 undetected. Out of the 75 possible dioxin congeners, only up to 3 congeners were detected.
32
33 The total PCDD concentrations ranged from below the mean detection limit (6.8 ± 2.3
34
35 ng/kg) to 77.1 ng/kg lw with a mean concentration of 17.5 ± 7.8 ng/kg lw (Table 2). Out of
36
37 the 135 possible furan congeners, only up to two congeners were detected in our whales
38
39 and the total PCDF concentrations ranged from below the mean detection limit (3.4 ± 1.4
40
41 ng/kg) to 113.1 ng/kg lw with a mean concentration of 5.1 ± 3.6 ng/kg lw.
42
43
44
45
46
47

48 There were no significant differences in PCDD/F concentrations between juvenile
49
50 and adult female killer whales from Possession Island (Table 2). The adult male was the
51
52 most PCDD/F contaminated, as observed in killer whales from the Northeastern Pacific
53
54 Ocean (Ross et al., 2000), but the small sample size precluded statistical evaluation. The
55
56
57
58
59
60
61
62
63
64
65

1
2
3 apparently higher dioxin and furan concentrations in adult females compared to juvenile
4
5 females were not supported statistically ($p > 0.05$). The high standard error reported for
6
7 adult females was due to only one highly PCDD contaminated individual. Due to our
8
9 limited sample size and therefore lack of strong statistical basis, it was not clear whether or
10
11 not the higher concentrations reported in adult females compared to the juveniles could be
12
13 an indication that dioxins and furans are not transferred as efficiently as PCBs from mother
14
15 to calf.
16
17

18
19 Data on dioxins and furans in killer whales are limited (Table 2). PCDD/F
20
21 concentrations in our study were higher than those reported for killer whales from Japan
22
23 (Kajiwara et al., 2006), but lower than levels reported for the three different communities of
24
25 killer whales living in the Northeastern Pacific Ocean (Ross et al., 2000). The higher
26
27 PCDD/F concentrations observed in the latter killer whales may reflect local contamination
28
29 by pulp and paper mills (Ross et al., 2004). Despite these observations, however, PCDD/F
30
31 concentrations in all killer whales are relatively low, something that has been attributed to
32
33 the minimal biomagnification of planar “dioxin-like” compounds in aquatic food webs as a
34
35 consequence of metabolism and excretion (Ross et al., 2000). The duality of distance from
36
37 source (remote location) and limited biomagnification in aquatic food webs likely explains
38
39 the low levels of dioxins and furans in Crozet killer whales.
40
41
42
43
44

45
46 Total PCDD/F concentrations in killer whales from Possession Island were higher
47
48 than levels previously reported for other Antarctic marine mammals such as Weddell seals
49
50 (Corsolini et al., 2002) and Antarctic fur seals (*Arctocephalus gazella*) (Oehme et al.,
51
52 1995). However, they were lower than levels reported for south polar skuas (*Catharacta*
53
54 *maccomicki*) which migrate to more contaminated areas in lower latitudes (Senthilkumar et
55
56
57
58
59
60
61
62
63
64
65

1
2
3 al., 2002). Compared to Arctic marine mammals, our observed concentrations were in the
4
5 same range as reported in ringed seals (*Pusa hispida*) (Norstrom and Simon, 1990), but
6
7 higher than levels in belugas from Baffin Island (Norstrom et al., 1992).
8
9

10 11 12 3.2.2 Homologue group and congener patterns 13

14
15 Only congeners from the hexa and octa homologue groups were detected,
16
17 accounting for 30 ± 35 % and 56 ± 37 % of \sum PCDD, respectively. 1,2,3,6,7,8 HxCDD
18
19 accounted on average for 65 ± 15 % of \sum HxCDD. This particular congener was previously
20
21 reported as being dominant in several species of marine mammals including killer whales
22
23 from Japan (Kajiwara et al., 2006), Weddell seals from Antarctica (Senthilkumar et al.,
24
25 2002), harbour seals (*Phoca vitulina*) from British Columbia (Addison et al., 2005). This
26
27 congener has, in part, been attributed to the historical use of elemental chlorine by pulp
28
29 mills, and to woodchips contaminated by pentachlorophenol (Addison et al., 2005). OCDD
30
31 was also detected in Arctic ringed seals (Norstrom and Simon, 1990). OCDD originates
32
33 from pulp mills, combustion, or high temperature industrial processes in which there is a
34
35 chlorine source (Norstrom and Simon, 1990). The remote nature of Possession Island thus
36
37 suggests that PCDD contamination is most likely due to long range atmospheric transport
38
39 but could also be in part due to small-scale combustion-related activities nearby, such as
40
41 shipping or research operations (Lugar et al., 1996).
42
43
44
45
46
47

48
49 No PCDF congeners were detected in juvenile females. 1,2,3,4,7,8 HxCDF,
50
51 1,2,3,4,6,7,8 HpCDF, 1,2,3,4,7,8,9 HpCDF, and OCDF were the only congeners detected in
52
53 the adult females. 2,3,7,8 TCDF was the only congener detected in the adult male. Even
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 though the patterns appeared different among the three groups, no clear trend can be
4
5 concluded as the sample size was small and the number of analytes detected very limited.
6

7 8 9 10 3.3 Toxic Equivalents (TEQ)

11
12 Consistent with the POP concentration patterns, juvenile females had similar TEQ
13
14 to adult females (76.5 ± 5.0 ng/kg lw and 44.2 ± 8.4 ng/kg lw, respectively) ($p > 0.05$),
15
16 while the adult male had the highest TEQ (109.1 ng/kg lw) (Table 3). Again, however, the
17
18 small sample size precluded statistical interpretation of the single male sample. PCBs
19
20 contributed up to 99.9% of the total TEQ in the killer whale samples, underscoring the
21
22 recalcitrance of this contaminant class. The adult male appeared to have a proportionately
23
24 lower contribution of PCBs to TEQ (88.3%) than did the immature ($99.8 \pm 0.1\%$) and adult
25
26 females ($97.3 \pm 1.7\%$), although the limited sample size precludes a rigorous evaluation.
27
28 Non-ortho PCB congeners contributed to most of the PCB TEQ (Table 3). This pattern
29
30 differed to those previously reported for other cetaceans species including killer whales
31
32 from the Northeastern Pacific Ocean (Ross et al., 2000), in which non-ortho and mono-
33
34 ortho PCBs contributed more equally to the \sum PCB TEQ. However, our use of new TEF
35
36 values (2005) partly explains this difference in observations; similar non-ortho and mono-
37
38 ortho PCB contributions to the \sum PCB TEQ was observed when we applied TEFs from
39
40 1998 to this dataset.
41
42

43
44 CB-118, 169, 126, and 105 were the dominant contributors to the total TEQ,
45
46
47
48 consistent with previous studies of cetaceans (Corsolini et al., 2003; Focardi et al., 1995;
49
50
51
52
53 Ylitalo et al., 2001).
54
55
56
57
58
59
60
61
62
63
64
65

3.4 Toxicological implications

Organic contaminants are a concern to the health of marine mammals, with observations of impaired reproduction, skeletal lesions, kidney damage, tumors, premature birth and skin lesions in populations inhabiting contaminated areas (Beckmen et al., 1997; Béland et al., 1993; Bergman et al., 2001; Delong et al., 1973; Olsson et al., 1992). High PCB concentrations have been linked to decreased immune function in field studies of harbour seals (Mos et al., 2006), bottlenose dolphins (*Tursiops truncatus*) (Lahvis et al., 1995) and polar bears (Lie et al., 2005), as well as captive feeding studies of harbour seals (De Swart et al., 1994; Ross et al., 1996). Contaminant-related immunotoxicity has been, in part, blamed for serious outbreaks of infectious disease outbreaks in marine mammals (Osterhaus et al., 1995). In addition, PCBs have been implicated in the disruption of vitamin A and thyroid hormone systems in harbour seals, which could lead to adverse effects on growth and development (Mos et al., 2007a; Tabuchi et al., 2006).

Different thresholds for adverse health effects have been established for PCBs in marine mammals. Immunotoxicity and endocrine disruption were observed in harbour seals at a concentration of 209 ng/kg total TEQ (De Swart et al., 1994; Ross et al., 1995). This value was adjusted to 255 ng/kg TEQ for killer whales from the Northeastern Pacific Ocean owing to a broader analytical evaluation (Ross et al., 2000). None of our Crozet killer whales surpass this threshold.

Thresholds have also been determined for PCBs in marine mammals. A threshold value of 17 mg/kg lw was established for immunotoxicity in captive harbour seals (Ross et al., 1996), while a 10 mg/kg lw threshold was associated with an increase in calf mortality in bottlenose dolphins (Hall et al., 2006). Recently, a more protective value of 1.3 mg/kg

1
2
3 was proposed for the protection of marine mammal health based on the 95% confidence
4 interval for endocrine disruption and immunotoxicity in multiple studies of free-ranging
5 harbour seals (Mos et al., 2007b). Even though the PCB concentrations in killer whales
6 from Possession Island were among the lowest reported in killer whales world-wide, 70%
7 of our study animals surpassed this latter threshold level, suggesting that contaminants
8 cannot be excluded as a possible risk factor in the decline of this population.
9
10
11
12
13
14
15
16
17
18

19 **4. Conclusion**

20
21 Despite our limited number of samples, this study provided baseline organic
22 contaminant data for the Possession Island killer whales. While these killer whales were
23 less contaminated than some of their northern hemispheric counterparts, they can be
24 counted among the most PCB-contaminated marine mammals in the southern hemisphere.
25
26 Despite a PCB ban in the industrialized world in the 1970s, the dispersion of this chemical
27 has attained the far reaches of the planet, including the killer whales that ply the remote
28 waters around Possession Island. While concentrations of POPs are decreasing in the
29 industrialized regions and in the Arctic, the global PCB distribution has not yet reached a
30 steady state and concentrations are expected to increase in the Antarctic before reaching a
31 state of equilibrium (Goerke et al., 2004). While we did not measure the organochlorine
32 pesticide dichloro diphenyl trichloroethane (DDT), other studies of marine mammals from
33 the southern hemisphere have found their concentrations to exceed that of PCBs by 2 - 9
34 times (Krahn et al., 2008; Aono et al., 1997). The results of our study provide another
35 reminder that POPs know no borders, and can significantly contaminate wildlife inhabiting
36 remote corners of the world.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

This work was conducted in part under the Program 109 of the Institut Polaire Paul Emile Victor. We would like to thank Terres Australes et Antarctiques Francaises (TAAF), the Killer Whale Adoption Program of the Vancouver Aquarium, and the staff of the regional contaminant laboratory at the Institute of Ocean Sciences, Fisheries and Oceans Canada.

Reference List

- Addison, R. F., Ikonomou, M. G., Smith, T. G., 2005. PCDD/F and PCB in harbour seals (*Phoca vitulina*) from British Columbia: response to exposure from pulp mill effluents. *Marine Environmental Research* 59, 165-176.
- Aono, S., Tanabe, S., Fujise, Y., Kato, H., Tatsukawa, R., 1997. Persistent organochlorines in minke whale (*Balaenoptera acutorostrata*) and their prey species from the Antarctic and the North Pacific. *Environmental Pollution* 98, 81-89.
- Barrett-Lennard, L. G., Smith, T. G., Ellis, G. M., 1996. A cetacean biopsy system using lightweight pneumatic darts, and its effect on the behavior of killer whales. *Marine Mammal Science* 12, 14-27.
- Beckmen, K. B., Lowenstine, L. J., Newman, J., Hill, J., Hanni, K., Gerber, J., 1997. Clinical and pathological characterization of northern elephant seal skin disease. *Journal of Wildlife Diseases* 33, 438-449.
- Béland, P., De Guise, S., Girard, C., Lagacé, A., Martineau, D., Michaud, R., Muir, D. C. G., Norstrom, R. J., Pelletier, E., Ray, S., Shugart, L. R., 1993. Toxic compounds and health and reproductive effects in St. Lawrence beluga whales. *Journal of Great Lakes Research* 19, 766-775.
- Bergman, A., Bergstrand, A., Bignert, A., 2001. Renal lesions in Baltic grey seals (*Halichoerus grypus*) and ringed seals (*Phoca hispida botnica*). *Ambio* 30, 397-409.
- Boon, J. P., Eijgenraam, F., 1988. The possible role of metabolism in determining patterns of PCB congeners in species from the Dutch Wadden Sea. *Marine Environmental Research* 24, 3-8.
- Borrell, A., Bloch, D., Desportes, G., 1995. Age trends and reproductive transfer of organochlorine compounds in long-finned pilot whales from the Faroe Islands. *Environmental Pollution* 88, 283-292.
- Corsolini, S., Ademollo, N., Romeo, T., Olmastroni, S., Focardi, S., 2003. Persistent organic pollutants in some species of a Ross Sea pelagic trophic web. *Antarctic Science* 15, 95-104.
- Corsolini, S., Kannan, K., Imagawa, T., Focardi, S., Giesy, J., 2002. Polychloronaphthalenes and other dioxin-like compounds in Arctic and Antarctic marine food webs. *Environmental Science and Technology* 36, 3490-3496.
- De Swart, R. L., Ross, P. S., Vedder, L. J., Timmerman, H. H., Heisterkamp, S. H., Van Loveren, H., Vos, J. G., Reijnders, P. J. H., Osterhaus, A. D. M. E., 1994. Impairment of immune function in harbor seals (*Phoca vitulina*) feeding on fish from polluted waters. *Ambio* 23, 155-159.

1
2 Delong, R. L., Gilmartin, W. G., Simpson, J. G., 1973. Premature births in California sea
3 lions: Association with high organochlorine pollutant residue levels. *Science* 181, 1168-
4 1170.

5
6
7 Focardi, S., Bargagli, R., Corsolini, S., 1995. Isomer-specific analysis and toxic potential
8 evaluation of polychlorinated biphenyls in Antarctic fish, seabirds and Weddell seals from
9 Terra Nova Bay (Ross Sea). *Antarctic Science* 7, 31-35.

10
11 Goerke, W., Weber, K., Bornemann, H., Ramdohr, S., Plotz, J., 2004. Increasing levels and
12 biomagnification of persistent organic pollutants (POPs) in Antarctic biota. *Marine*
13 *Pollution Bulletin* 48, 295-302.

14
15
16 Guinet, C., 1991. L'orque (*Orcinus orca*) autour de l'archipel Crozet: comparaison avec
17 d'autres localités. *Revue d'écologie (Terre Vie)* 46, 321-337.

18
19 Hall, A. J., McConnell, B. J., Rowles, T. K., Aguilar, A., Borrell, A., Schwacke, L.,
20 Reijnders, P. J. H., Wells, R. S., 2006. Individual-based model framework to assess
21 population consequences of polychlorinated biphenyl exposure in bottlenose dolphins.
22 *Environmental Health Perspectives* 114, 60-64.

23
24
25 Helm, P. A., Bidleman, T. F., Stern, G. A., Koczanski, K., 2002. Polychlorinated
26 naphthalenes and coplanar polychlorinated biphenyls in beluga whale (*Delphinapterus*
27 *leucas*) and ringed seal (*Phoca hispida*) from the eastern Canadian Arctic. *Environmental*
28 *Pollution* 119, 69-78.

29
30
31 Hickie, B. E., Ross, P. S., Macdonald, R. W., Ford, J. K. B., 2007. Killer whales (*Orcinus*
32 *orca*) face protracted health risks associated with lifetime exposure to PCBs. *Environmental*
33 *Science and Technology* 41, 6613-6619.

34
35
36 Ikonomou, M. G., Fraser, T., Crewe, N., Fischer, M. B., Rogers, I. H., He, T., Sather, P.,
37 Lamb, R., 2001. A comprehensive multiresidue ultra-trace analytical method, based on
38 HRGC/HRMS, for the determination of PCDDs, PCDFs, PCBs, PBDEs, PCDEs, and
39 organochlorine pesticides in six different environmental matrices. *Canadian Technical*
40 *Report Of Fisheries and Aquatic Sciences* 2389, 1-95.

41
42
43 Ikonomou, M. G., Sather, P., Crewe, N., Fraser, T., He, T., 1999. Analytical considerations
44 when carbon-fibre fractionation is used in conjunction with HRGC/HRMS for the full
45 congener PCBs determination in environmental samples. submitted

46
47 Kajiwara, N., Kunisue, T., Kamikawa, S., Ochi, Y., Yano, S., Tanabe, S., 2006.
48 Organohalogen and organotin compounds in killer whales mass-stranded in the Shiretoko
49 peninsula, Hokkaido, Japan. *Marine Pollution Bulletin* 52, 1066-1076.

50
51
52 Kajiwara, N., Matsuoka, M., Iwata, H., Tanabe, S., Rosas, F. C. W., Fillmann, G.,
53 Readman, J. W., 2004. Contamination by persistent organochlorines in cetaceans
54 incidentally caught along Brazilian coastal waters. *Archives of Environmental*
55 *Contamination and Toxicology* 46, 124-134.

1
2 Krahn, M., Pitman, R. L., Burrows, D. G., Herman, D. P., Pearce, R. W., 2008. Use of
3 chemical tracers to assess diet and persistent organic pollutants in Antarctic type C killer
4 whales. *Marine Mammal Science* 24, 643-663.

5
6
7 Lahvis, G. P., Wells, R. S., Kuehl, D. W., Stewart, J. L., Rhinehart, H. L., Via, C. S., 1995.
8 Decreased lymphocyte responses in free-ranging bottlenose dolphins (*Tursiops truncatus*)
9 are associated with increased concentrations of PCBs and DDT in peripheral blood.
10 *Environmental Health Perspectives Supplements* 103, 67-72.

11
12
13 Lie, E., Larsen, H. J. S., Larsen, S., Johansen, G. M., Derocher, A. E., Lunn, N. J.,
14 Norstrom, R. J., Wiig, O., Skaare, J. U., 2005. Does high organochlorine (OC) exposure
15 impair the resistance to infection in polar bears (*Ursus maritimus*)? Part 2: possible effect
16 of OCs on mitogen- and antigen-induced lymphocyte proliferation. *Journal of Toxicology*
17 *and Environmental Health, Part A* 68, 457-484.

18
19
20 Lugar, R. M., Harless, R. L., Dupuy, A. E., Daniel, D. D., 1996. Results of monitoring for
21 PCDDs and PCDFs in ambient air at McMurdo station, Antarctica. *Environmental Science*
22 *and Technology* 30, 555-561.

23
24
25 Miranda-Filho, K. C., Metcalfe, T. L., Metcalfe, C. D., Robaldo, R. B., Muelbert, M. M. C.,
26 Colares, E. P., Martinez, P. E., Bianchini, A., 2007. Residues of persistent organochlorine
27 contaminants in southern elephant seals (*Mirounga leonina*) from Elephant Island,
28 Antarctica. *Environmental Science and Technology* 41, 3829-3835.

29
30
31 Mos, L., Morsey, B., Jeffries, S. J., Yunker, M. B., Raverty, S., De Guise, S., Ross, P. S.,
32 2006. Chemical and biological pollution contribute to the immunological profiles of free-
33 ranging harbor seals. *Environmental Toxicology and Chemistry* 25, 3110-3117.

34
35
36 Mos, L., Tabuchi, M., Dangerfield, N., Jeffries, S. J., Koop, B. F., Ross, P. S., 2007a.
37 Contaminant-associated disruption of vitamin A and its receptor (RAR α) in free-ranging
38 harbour seals (*Phoca vitulina*). *Aquatic Toxicology* 81, 319-328.

39
40
41 Mos, L., Tabuchi, M., Koop, B. F., Helbing, C. C., and Ross, P. S., 2007b.
42 Toxicogenomics: a new tool for the assessment of health and toxicity in marine mammals.
43 17th Biennial Conference on the Biology of Marine Mammals

44
45
46 Norstrom, R. J., Muir, D. C. G., Ford, C. A., Simon, M., Macdonald, C. R., Beland, P.,
47 1992. Indications of P450 monooxygenase activities in Beluga (*Delphinapterus leucas*) and
48 narwhal (*Monodon monocerus*) from patterns of PCB, PCDD and PCDF accumulation.
49 *Marine Environmental Research* 34, 267-272.

50
51
52 Norstrom, R. J., Simon, M., 1990. Polychlorinated dibenzo-*p*-dioxins and dibenzofurans in
53 marine mammals in the Canadian north. *Environmental Pollution* 66, 1-19.

54
55
56 Oehme, M., Schlabach, M., Boyd, I., 1995. Polychlorinated Dibenzo-*p*-Dioxins,
57 dibenzofurans and coplanar biphenyls in Antarctic fur seal blubber. *Ambio* 24, 41-46.

58
59
60 Olsson, M., Andersson, O., Bergman, A., Blomkvist, G., Frank, A., Rappe, C., 1992.
61 Contaminants and diseases in seals from Swedish waters. *Ambio* 21, 561-562.

1
2 Osterhaus, A. D. M. E., De Swart, R. L., Vos, H. W., Ross, P. S., Kenter, M. J. H., Barrett,
3 T., 1995. Morbillivirus infections of aquatic mammals: newly identified members of the
4 genus. *Veterinary Microbiology* 44, 219-227.
5

6
7 Poncelet, E., Barbraud, C., Guinet, C., 2008. Population dynamics of killer whales in
8 Crozet Archipelago, southern Indian Ocean: exploiting opportunistic and protocol-based
9 photographs in a mark recapture study. *Journal of Cetacean Research and Management*. In
10 submission.
11

12 Rantaleinen, A. L., Ikonomou, M. G., Rogers, I. H., 1998. Lipid-containing semipermeable
13 membrane devices (SPMDs) as concentrators of toxic chemicals in the lower Fraser River,
14 Vancouver, British Columbia. *Chemosphere* 37, 1119-1138.
15

16
17 Reijnders, P. J. H., 1986. Reproductive failure in common seals feeding on fish from
18 polluted coastal waters. *Nature* 324, 456-457.
19

20
21 Risebrough, R. W., De Lappe, B. W., Younghans-haug, C., 1990. PCB and PCT
22 contamination in Winter Quarters bay, Antarctica. *Marine Pollution Bulletin* 21, 523-529.
23

24
25 Ross, P. S., 2006. Fireproof killer whales: Flame retardant chemicals and the conservation
26 imperative in the charismatic icon of British Columbia. *Canadian Journal of Fisheries and
27 Aquatic Sciences* 63, 224-234.
28

29
30 Ross, P. S., De Swart, R. L., Addison, R. F., Van Loveren, H., Vos, J. G., Osterhaus, A. D.
31 M. E., 1996. Contaminant-induced immunotoxicity in harbour seals: wildlife at risk?
32 *Toxicology* 112, 157-169.
33

34
35 Ross, P. S., De Swart, R. L., Reijnders, P. J. H., Van Loveren, H., Vos, J. G., Osterhaus, A.
36 D. M. E., 1995. Contaminant-related suppression of delayed-type hypersensitivity and
37 antibody responses in harbor seals fed herring from the Baltic Sea. *Environmental Health
38 Perspectives* 103, 162-167.
39

40
41 Ross, P. S., Ellis, G. M., Ikonomou, M. G., Barrett-Lennard, L. G., Addison, R. F., 2000.
42 High PCB concentrations in free-ranging Pacific killer whales, *Orcinus orca*: effects of age,
43 sex and dietary preference. *Marine Pollution Bulletin* 40, 504-515.
44

45
46 Ross, P. S., Jeffries, S. J., Yunker, M. B., Addison, R. F., Ikonomou, M. G., Calambokidis,
47 J., 2004. Harbour seals (*Phoca vitulina*) in British Columbia, Canada, and Washington,
48 USA, reveal a combination of local and global polychlorinated biphenyl, dioxin, and furan
49 signals. *Environmental Toxicology and Chemistry* 23, 157-165.
50

51
52 Senthilkumar, K., Kannan, K., Corsolini, S., Evans, T., Giesy, J. P., Nakanishi, J.,
53 Masunaga, S., 2002. Polychlorinated dibenzo-*p*- dioxins, dibenzofurans and
54 polychlorinated biphenyls in polar bear, penguin and south polar skua. *Environmental
55 Pollution* 119, 151-161.
56

57
58 Tabuchi, M., Veldhoen, N., Dangerfield, N., Jeffries, S. J., Helbing, C. C., Ross, P. S.,
59 2006. PCB-related alteration of thyroid hormones and thyroid hormone receptor gene
60
61
62
63
64
65

1
2 expression in free-ranging harbor seals (*Phoca vitulina*). Environmental Health
3 Perspectives 114, 1024-1031.
4

5 Van den Berg, M., Birnbaum, L. S., Denison, M., De Vito, M., Farland, W., Feeley, M.,
6 Fiedler, H., Hakansson, H., Hanberg, A., Haws, L., Rose, M., Safe, S., Schrenk, D.,
7 Tohyama, C., Tritscher, A., Tuomisto, J., Tysklind, M., Walker, N., Peterson, R. E., 2006.
8 The 2005 world health organization reevaluation of human and mammalian toxic
9 equivalency factors for dioxins and dioxin-like compounds. Toxicological Sciences 93,
10 223-241.
11
12

13 Wania, F., Mackay, D., 2001. Global fractionation and cold condensation of low volatility
14 organochlorine compounds in polar regions. Ambio 22, 10-18.
15
16

17 Wolkers, H., Corkeron, P. J., Van Parijs, S. M., Similä, T., Van Bavel, B., 2007.
18 Accumulation and transfer of contaminants in killer whales (*Orcinus orca*) from Norway:
19 indications for contaminant metabolism. Environmental Toxicology and Chemistry 26,
20 1582-1590.
21
22

23 Ylitalo, G. M., Matkin, C. O., Buzitis, J., Krahn, M., Jones, L. L., Rowles, T., Stein, J. E.,
24 2001. Influence of life-history parameters on organochlorine concentrations in free-ranging
25 killer whales (*Orcinus orca*) from Prince William Sound, AK. Science of the Total
26 Environment 281, 183-203.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FIGURE CAPTIONS

1
2 **Figure 1:** Killer whales were sampled around Possession Island, Crozet Archipelago, in the
3
4 southern Indian Ocean.
5
6

7
8 **Figure 2:** Juvenile and adult females have similar PCB (total, di-ortho, mono-ortho and
9 coplanar), PCDD, and PCDF concentrations (mean \pm standard error). The adult male shows the
10 highest PCB, PCDD and PCDF concentrations (mg/kg lw), although limited sample size
11 precluded statistical evaluation.
12
13
14
15
16

17
18 **Figure 3:** The homologue group contribution to the total PCB concentrations was similar across
19 age/sex categories. (Mean \pm standard error is available only for juvenile and adult female killer
20 whales).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1: The PCB composition was generally similar across age/sex categories as evidenced by the top six PCB congeners (concentrations in mg/kg lw). The percentage contribution to the total PCB contamination is added in brackets.

	Juvenile Females (n = 3)	Adult Females (n = 5)	Adult male (n = 1)
Σ PCBs	11.4 \pm 1.2	10.1 \pm 3.1	19.5
153 ^a	2.2 \pm 0.2 (19.3 \pm 0.4%)	153 ^a 2.0 \pm 0.6 (19.9 \pm 0.2%)	153 ^a 3.8 (19.2%)
138 ^a	1.5 \pm 0.1 (12.9 \pm 0.8%)	138 ^a 1.4 \pm 0.4 (13.1 \pm 0.2%)	138 ^a 2.7 (13.8%)
180	0.9 \pm 0.2 (7.5 \pm 1.2%)	149 0.8 \pm 0.3 (7.4 \pm 0.3%)	180 1.5 (7.8%)
149	0.8 \pm 0.2 (7.2 \pm 0.6%)	180 0.7 \pm 0.2 (7.2 \pm 0.1%)	149 1.3 (6.9%)
187 ^a	0.7 \pm 0.1 (5.8 \pm 0.6%)	187 ^a 0.6 \pm 0.2 (5.8 \pm 0.2%)	187 ^a 1.1 (5.6%)
201	0.5 \pm 0.2 (3.9 \pm 1.0%)	201 0.4 \pm 0.1 (3.4 \pm 0.1%)	118 ^a 0.6 (3.1%)
Σ top 6	6.5 \pm 0.3 (57.4 \pm 0.6%)	5.9 \pm 0.3 (54.6 \pm 0.2%)	11.0 (56.4%)

^a: dominant congener of a co-eluting group (CB-153 was co-eluting with 132; 138 with 163 and 164; 187 with 182; 118 with 106)

Table 2: PCB concentrations in the largely marine mammal-eating Crozet killer whales exceed concentrations in the fish-eating Antarctic killer whales. PCB levels approach those observed in the relatively contaminated northern resident killer whales from the Northeastern Pacific Ocean.

Location	Year	Type	Sex	Lifestage	PCBs (mg/kg lw)	PCDDs (ng/kg lw)	PCDFs (ng/kg lw)	TEQ (ng/kg lw)	Reference
Crozet Archipelago	1998		F	Juvenile	11.4 ± 1.1	3.9 ± 1.8	ND ^a	76.5 ± 5.0	Our study
			F	Adult	10.1 ± 3.0	20.6 ± 12.1	6.6 ± 5.9	44.2 ± 8.4	
			M	Adult	19.5	38.79	11.4	120.4	
Ross Sea, Antarctica	2005-2006	Type C ^b	M	Adult	1.6 ± 1.1	n.a ^c	n.a	n.a	Krahn et al.,2008
			M	Subadult	2.1 ± 2.3	n.a	n.a	n.a	
			F	Adult	0.6 ± 0.4	n.a	n.a	n.a	
Norway	2002		M	Adult/Juvenile	16-44	n.a	n.a	n.a	Wolkers et al., 2007
Japan	2005		M	Calf	47	0.62	0.77	300	Kajiwara et al., 2006
			F	Calf	51	N.D	0.96	405	
			M	Adult	57	1.50	3.90	350	
			F	Adult	30	3.50	5.80	220	
British Columbia, Canada	1993-1996	Northern Resident	M	Adult	37 ± 6			435 ± 60	Ross et al., 2000
			F	Adult	9 ± 3			111 ± 29	
		Southern Resident	M	Adult	146 ± 33	1050 ± 258	55 ± 6	845 ± 195	
			F	Adult	55 ± 19			456 ± 27	
		Transient	M	Adult	251 ± 55			699 ± 142	
			F	Adult	59 ± 21			338 ± 53	
	2004-2006	Southern Resident	M	Adult	66 ± 26	n.a	n.a	n.a	Krahn et al.,2007
			F	Adult	45	n.a	n.a	n.a	
Alaska, USA	1994-1999	Resident	M/F	Adult/Juvenile	14 ± 13	n.a	n.a	100 ± 98	Ylitalo et al., 2001
		Transient	M/F	Adult/Juvenile	230 ± 130	n.a	n.a	860 ± 640	

^a: Non detected

^b: Type C killer whales are thought to prey mainly on marine fish

^c: Non available

Table 3: In juvenile female, adult female and adult male killer whales, PCBs are the dominant contributor to the total toxic equivalents to 2,3,7,8-TCDD.

	PCB congeners	Mean TEQ \pm st error (pg/g lw)		
		Juvenile Female (n=3)	Adult female (n=6)	Adult male (n=1)
Non-ortho PCBs	77	0.01 \pm 0.00	0.01 \pm 0.00	0.02
	81	0.01 \pm 0.00	0.01 \pm 0.00	0.01
	126	10.61 \pm 2.53	7.69 \pm 1.11	16.63
	169	53.39 \pm 7.07	31.38 \pm 7.02	60.56
	Σ non-ortho PCBs	64.01 \pm 8.72	39.07 \pm 7.65	77.21
	% of Σ PCB TEQ	83.42 \pm 8.71%	87.30 \pm 2.06%	70.82%
Mono-ortho PCBs	105	1.17 \pm 0.25	0.81 \pm 0.18	4.74
	114	0.09 \pm 0.02	0.06 \pm 0.01	0.32
	118	3.77 \pm 0.74	2.71 \pm 0.66	17.89
	123	0.04 \pm 0.01	0.02 \pm 0.01	0.20
	156	0.57 \pm 0.03	0.38 \pm 0.09	2.16
	157	0.15 \pm 0.01	0.11 \pm 0.03	0.69
	167	0.51 \pm 0.04	0.39 \pm 0.12	2.68
	189	0.08 \pm 0.02	0.06 \pm 0.02	0.40
	Σ mono-ortho PCBs	6.38 \pm 0.98	4.55 \pm 1.12	29.08
	% of Σ PCB TEQ	8.28 \pm 0.97%	9.97 \pm 1.17%	26.68%
	Sum PCB TEQ	76.26 \pm 4.88	43.62 \pm 8.66	106.29
		99.77 \pm 0.15%	97.27 \pm 1.70%	88.31%
	Sum PCDD TEQ	0.19 \pm 0.13	0.45 \pm 0.34	1.59
		0.23 \pm 0.15%	1.99 \pm 1.56%	1.46%
	Sum PCDF TEQ	ND	0.17 \pm 0.14	1.14
		-	0.72 \pm 0.66%	1.04%
	TOTAL TEQ	76.45 \pm 5.00	44.24 \pm 8.37	109.02

Figure 1:




Figure 2:




Figure 3:

