



HAL
open science

The activity coefficient of high density systems with hard-sphere interactions. The application of the IGCMC method

Stanislaw Lamperski, Monika Pluciennik

► **To cite this version:**

Stanislaw Lamperski, Monika Pluciennik. The activity coefficient of high density systems with hard-sphere interactions. The application of the IGCMC method. *Molecular Simulation*, 2010, 36 (02), pp.111-117. 10.1080/08927020903124577 . hal-00564474

HAL Id: hal-00564474

<https://hal.science/hal-00564474>

Submitted on 9 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The activity coefficient of high density systems with hard-sphere interactions. The application of the IGCMC method

Journal:	<i>Molecular Simulation</i> / <i>Journal of Experimental Nanoscience</i>
Manuscript ID:	GMOS-2009-0054.R1
Journal:	Molecular Simulation
Date Submitted by the Author:	12-Jun-2009
Complete List of Authors:	Lamperski, Stanislaw; A. Mickiewicz University, Department of Chemistry Płuciennik, Monika; A. Mickiewicz University, Department of Chemistry
Keywords:	Activity coefficient, Inverse grand canonical Monte Carlo method, Hard-sphere interactions, Solvent primitive model electrolyte, GCMC simulation

SCHOLARONE™
Manuscripts

The activity coefficient of high density systems with hard-sphere interactions. The application of the IGCMC method

S. LAMPERSKI* and M. PŁUCIENNIK

*Department of Physical Chemistry, Faculty of Chemistry, A. Mickiewicz University,
Grunwaldzka 6, 60-780 Poznań, Poland*

Abstract

The inverse grand-canonical Monte Carlo technique is used to calculate the activity coefficients of the following hard-sphere systems: one-component fluid, binary mixture and solvent primitive model (SPM) electrolyte. The calculations for a one-component fluid are performed at different densities. The components of a binary mixture differ in diameters (300 and 500 pm) with the results being presented for different density and composition of a mixture. For the SPM model simulations are performed for 1:1 electrolyte at different electrolyte concentrations at the packing fraction equal to 0.3. Ions and solvent molecules of the same or different sizes are considered. The results are compared with those reported by Adams (one-component fluid), with those calculated using the Ebeling - Scherwinski equation (one-component fluid and binary mixture), and with the predictions from the symmetric Poisson-Boltzmann theory and the mean spherical approximation (SPM electrolyte).

Keywords: Activity coefficient; Inverse grand canonical Monte Carlo method; Hard-sphere interactions; Solvent primitive model electrolyte; GCMC simulation

* Corresponding author: Tel.: +48 61 8291454, Fax: +48 61 829 1505, Email: slamper@amu.edu.pl

1. Introduction

The chemical potential is perhaps the most important function describing the thermodynamic properties of a fluid. Its essential component, the activity coefficient, describes all the deviations from ideality. The activity coefficient can be determined experimentally, calculated from the statistical thermodynamics, or obtained from the numerical molecular simulations. Unfortunately the entropy-related functions like the free energy or the chemical potential cannot be estimated directly from the Monte Carlo (MC) simulation in the canonical ensemble [1] as this method samples the regions making small contribution to the configuration partition function. However, application of the Gibbs-Helmholtz equation [1] allows us to calculate the free energy from MC simulations, but the method requires many simulations at different temperatures. Widom [2] proposed a simpler and more attractive technique. It consists in inserting a particle into the simulation box, which tests the intermolecular interactions. The individual ionic activity coefficients can be calculated by this method [3,4] but the extrapolation of the results to the thermodynamic limit is required as the results depend on the number of particles in the simulation box. Another method used to calculate the activity coefficient is the Monte Carlo simulation in the grand-canonical ensemble (GCMC). This technique was applied among others by Adams [5] to calculate the chemical potential of hard-sphere fluids, and by Valleau and Cohen [6,7] to evaluate the mean activity coefficient of electrolyte. A drawback of the GCMC method is it can often be inconvenient to use as it gives the concentration of particles for a specified activity coefficient. Recently, the GCMC method has been modified by Lamperski [8], and in the modified form it allows evaluation of the activity coefficient for a specified concentration of particles. It is called the inverse grand-canonical Monte Carlo method (IGCMC) as it uses the GCMC techniques, but it works in the reverse direction. Malasics *et al.* [9] have proposed two other efficient iterative MC algorithms in the grand-canonical ensemble to determine the chemical potentials. These authors applied their method to the Lennard-Jones fluids and to the primitive electrolyte mixture.

The IGCMC technique was successfully used to calculate the individual and mean activity coefficients of the 1:1, 2:2, 2:1, 3:1 PM electrolyte with ions of the same and different size at different electrolyte concentrations [8]. The results for ions of the same size were compared with the corresponding data obtained by Valleau and Cohen [6], while the results for a 1:1 electrolyte with ions of different size were compared with those of Sloth and Sørensen [4], and with the theoretical predictions of Sloth and Sørensen [10], Molero *et al.* [11] and Outhwaite *et al.* [12,13]. The results obtained by the IGCMC method were comparable with those obtained or predicted by other methods. The advantage of IGCMC over GCMC is that the former can be used to compute also the individual ionic activity coefficients which cannot be done by the standard GCMC method.

In this paper we want to demonstrate that the IGCMC method gives correct results for hard-sphere fluids at high densities. The following three systems are the objects of our investigation here: i) one-component fluid, ii) binary mixture, and iii) solvent primitive model (SPM) electrolyte.

2. The model

The simplest model that takes into account the finite size of the molecules is the hard-sphere model. Representing the molecules of the i th kind by hard spheres of the diameter d_i , the pair potential of interaction is given by

$$u^{\text{hs}}(r_{ij}) = \begin{cases} 0, & r_{ij} > d_{ij} \\ \infty, & r_{ij} \leq d_{ij} \end{cases} \quad (1)$$

where r_{ij} is the distance between the centres of two interacting molecules (hard spheres) and d_{ij} is given by

$$d_{ij} = \frac{d_i + d_j}{2} \quad (2)$$

The SPM electrolyte model is a simple yet non-trivial model of an electrolyte that takes into account the molecular nature of solvent. We assume that the ions and solvent molecules are represented by hard spheres of diameter d_{\pm} and d_s , respectively. The ions have additionally a point electric charge $z_i e$ embedded at the centre (e is the elementary charge and z_i is the charge number of ions of the i th kind). The inter-ionic electrostatic interactions are described by

$$u^e(r_{ij}) = \frac{z_i z_j e^2}{4\pi\epsilon_0\epsilon_r r_{ij}} \quad (3)$$

where ϵ_r is the relative permittivity of the solvent and ϵ_0 is the vacuum permittivity. The importance of the SPM and hard-sphere models stems from the fact that they can be used both in molecular MC simulations and theoretical studies. Consideration of solvent molecules makes the model of electrolyte more realistic and allows investigation at densities similar to those of the real systems.

3. The method

In the grand-canonical ensemble the volume, V , temperature, T , and the chemical potential, μ , are constant, while the number of molecules, N , and the energy, E , fluctuate. In the GCMC technique the activity coefficient is used to calculate the concentration of particles. Recently Lamperski [8] has shown that this method can be made to work in the reverse direction. The new technique, called the inverse GCMC (IGCMC), can be used to estimate the activity coefficient of particles for their specified concentration. We will describe the performance of this method with an example of a one-component fluid. Extension to a multi-component solution is straightforward and gives the individual activity coefficients. The technique consists of m short GCMC simulations in each of which n number of configurations are generated. As in the GCMC technique, we input the activity coefficient ($\ln\gamma_l$) into each l -th sub-simulation and obtain the average number $\langle N \rangle_l$ of molecules. This number is compared with the number N^0 which corresponds to the specified concentration c

$$N^0 = 1000cN_A V \quad (4)$$

where N_A is the Avogadro constant and V is the volume of the simulation box. If $\langle N \rangle_l$ is lower than N^0 then the value of $\ln \gamma_l$ is increased by a small amount *e.g.* 0.2 %, otherwise it is decreased by 0.2 %. When the simulation is finished, the average value of the activity coefficients evaluated after each sub-simulation is calculated to give the final activity coefficient

$$\ln \gamma = \frac{1}{m} \sum_{l=1}^m \ln \gamma_l \quad (5)$$

The simulations were carried out for $n = 20000$ configurations and $m = 7500$ sub-simulations. The first 2500 sub-simulations were used to equilibrate the system.

In the IGCMC, as in GCMC, there are three equally probable moves: displacement, insertion and removal. In the first move a particle selected at random is displaced to a new random position in the simulation box. The acceptance of displacement is given by

$$\text{acc}(m \rightarrow n) = \min\{1, \exp[-(u_n - u_m)/kT]\} \quad (6)$$

where u_m is the potential energy of the configuration before displacement, u_n is the potential energy of the new trial configuration and k is the Boltzmann constant.

In the second move a molecule is inserted at a random position in the simulation box. The acceptance of insertion is described by the equation

$$\text{acc}(N \rightarrow N+1) = \min\left[1, \exp(-\Delta u_i/kT + \ln \frac{\gamma N^0}{N+1})\right] \quad (7)$$

where Δu_i is the potential of insertion and N is the current number of molecules.

In the last move a particle selected at random is removed from the box. The acceptance of this move is

$$\text{acc}(N \rightarrow N-1) = \min\left[1, \exp(-\Delta u_r/kT + \ln \frac{N}{\gamma N^0})\right] \quad (8)$$

where Δu_r is the potential of removal.

4. Results and discussion

4.1 Hard-sphere one-component system

At first we consider a one-component hard-sphere fluid. The IGCMC technique was applied to calculate the activity coefficient. The simulations were performed for the following reduced densities ρ^* : 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 at the diameter $d = 400$ pm ($\rho^* = \rho d^3$, ρ is the number density, $\rho = N^0/V = 1000cN_A$). The results are shown in figure 1 and collected in table 1 where they are compared with the corresponding Adams [5] data and the theoretical predictions obtained from the Ebeling and Scherwinski equation [14]. The value of the activity coefficient increases with increasing density of a system. Our results and those obtained by Adams are nearly identical, while the predictions evaluated from the Ebeling and Scherwinski's equation are also very similar, but systematically lower. For comparison with the results of Adams, our results in column 1, of table 1, are presented in terms of the reduced

density ρ^* (~~$=\rho d^3$~~ ; ρ density). The corresponding packing fractions η ($=\pi\rho^*/6$) are given in column 2 and are used in figure 1.

4.2 Binary mixture of hard spheres

The second model studied was a two-component mixture of hard spheres of different diameters, $d_1 = 300$ pm and $d_2 = 500$ pm. The simulations were carried out for the following packing fractions, $\eta : 0.05, 0.10, 0.15, 0.20, 0.25, 0.30,$ and 0.35 at different compositions described by the mole fraction x . **This time the packing fraction is given by**

$$\eta = \frac{\pi}{6} \rho \sum_i x_i d_i^3 \quad (9)$$

where ρ

$$\rho = \sum_i \rho_i \quad (10)$$

is the total number density. At constant packing fraction the total number density is changing.

The IGC MC results were compared with those calculated from the equation of Ebeling and Scherwinski [14]. The activity coefficients of the larger hard spheres are shown in figure 2A and in table 2. The simulations broke down for $x_2 = 0.1$ at $\eta = 0.30$ and for $x_2 = 0.1, 0.2$ and 0.3 at $\eta = 0.35$, because of the problem of inserting the larger molecules into the mixture with predominantly smaller molecules, arising from the lack of sufficient free space in a system. This problem does not occur in the calculation of activity coefficient for smaller molecules (figure 2B, table 3). As previously seen, the activity coefficient increases with increasing density of the system (figure 2). Here we further note that the activity coefficient of a mixture depends also on the mixture composition. The activity coefficient of larger molecules decreases (figure 2A) while that of the smaller increases (figure 2B) with increasing mole fraction of the component considered. We see that at low concentration of larger molecules their activity coefficient is greater than that of the pure component ($x_2 = 1$) (figure 2A). On the other hand, the activity coefficient of smaller spheres is lower than that of the pure component ($x_1 = 1$). This means that at a constant η the probability of insertion of a smaller sphere into a fluid of larger spheres is higher than the probability of insertion of a larger sphere into a fluid of smaller spheres.

4.3 SPM model

The next system studied was the SPM electrolyte model. Its thermodynamic properties, except for the individual activity coefficients, were intensively explored [15] by the MC simulations. We considered a 1:1 electrolyte at the total packing fraction of the solution $\eta = 0.3$ and at different electrolyte concentrations. **To keep constant the packing fraction with the variation of the electrolyte concentration, the solvent concentration was changing.** The temperature was 298.15 K and the relative permittivity $\epsilon_r = 78.5$. We considered systems with ions, d_{\pm} , and the solvent molecules, d_s of the same and different diameters. The first simulations were carried out for the same diameters of the ions and the solvent molecules. The results for $d = d_{\pm} = d_s = 300, 400$ and 500 pm are shown in figure 3 by graphic characters and collected in table 4. Additionally figure 3 presents the activity coefficients from the

1
2
3 symmetric Poisson-Boltzmann (SPB) theory [11] (solid lines) and the mean-spherical
4 approximation (MSA) [14] (dashed lines). The SPB activity coefficients were derived via the
5 charging route for the individual ion activities while the MSA equation was derived starting
6 from a uniform neutralizing background developed by Parrinello and Tosi [16]. In SPB and
7 MSA the solvent activity coefficients were calculated from the equation of Ebeling and
8 Scherwinski [14]. It is interesting to note that the activity coefficient of the solvent, $\ln\gamma_s$,
9 essentially does not depend on either the electrolyte concentration or the diameter d . The
10 mean activity coefficient, $\ln\gamma_{\pm}$, of ions in the limit of zero electrolyte concentration has the
11 same value as for the solvent molecules and decreases with increasing electrolyte
12 concentration. The effect depends on the diameter, d . Although the SPB and MSA results
13 diverge slightly from the IGCMC, their qualitative behaviour is similar. A better agreement
14 with the MC results gives the SPB theory at low electrolyte concentration while the MSA at
15 higher concentrations. The divergence of the SPB results at high electrolyte concentration is
16 perhaps due to the neglect of the fluctuation potential in the SPB theory and the choice of
17 thermodynamic route [11].

18
19
20
21 It is worth noting that the behaviour of $\ln\gamma_{\pm}$ is different than that observed for the
22 primitive model (PM) electrolyte [11]. In figure 3 we do not observe a rise in $\ln\gamma_{\pm}$ at higher
23 electrolyte concentrations characteristic of the PM electrolyte [6,8,11]. In PM this increase is
24 due to increased hard-sphere interactions at higher concentrations. In our case the hard-sphere
25 interactions are constant as the value of the packing fraction, η , is fixed and the diameters of
26 all the components are the same. These interactions do not influence the characteristics of the
27 ion activity coefficient curves, but only their values. The course depends on the inter-ionic
28 electrostatic interactions including the exclusion volume and correlation effects. It is relevant
29 to mention here that experiments have shown some electrolytes (*e.g.* aqueous KCl, KI, NaCl)
30 to exhibit a concentration dependence of the mean activity coefficient similar to that shown in
31 figure 3 [17].

32
33
34 Next we investigated a system with the ions and solvent molecules of different size
35 (the diameters of the anion and the cation being equal). At first the ion diameter was fixed at
36 $d_{\pm} = 400$ pm while the diameter of the solvent molecules d_s was 300, 400 or 500 pm. The
37 other physical parameters were the same as for the earlier case of the common diameter. The
38 **ionic** concentration dependence of the activity coefficients of ions and solvent molecules is
39 given in table 5 and illustrated in figures 4A and 4B by graphic characters. The corresponding
40 SPB and MSA results are shown by the solid and dashed lines, respectively. Better agreement
41 between the MC and the theoretical results is observed for the solvent. The mean activity
42 coefficient of ions (figure 4A) shows generally a decreasing tendency with increasing
43 electrolyte concentration, however for $d_s = 500$ pm some increase is observed at high
44 electrolyte concentrations. Even for the constant ion diameter, the mean activity coefficient
45 depends on the solvent diameter. In a mixture with smaller solvent molecules the mean
46 activity coefficient is greater than in that with larger molecules. The dependence of the
47 activity coefficient of solvent (figure 4B) on the electrolyte concentration varies: the curve for
48 $d_s = 500$ pm is increasing, for $d_s = 300$ pm it is decreasing, while for $d_s = 400$ pm it is constant
49 and this is the case discussed earlier.

50
51
52
53 The next simulations were performed for the fixed diameter of the solvent molecules,
54 $d_s = 400$ pm and for the following ion diameters: $d_{\pm} = 300, 400$ and 500 pm. The IGCMC
55 results are shown in figure 5 by graphic characters and collected in table 6. The solid and
56 dashed lines in figure 5 show the SPB and MSA results, respectively. The mean activity
57 coefficient decreases with increasing electrolyte concentration (figure 5A) although a slight
58 rise at higher concentrations is observed. Its value evidently depends on the ion diameters: the
59 larger the ions the greater the activity coefficient. The dependence of the activity coefficient
60

1
2
3 of solvent on the electrolyte concentration (figure 5B) varies again but this time the curve for
4 $d_{\pm} = 300$ pm increases while for $d_{\pm} = 500$ pm it decreases. The curve for $d_{\pm} = 400$ pm is flat.

5
6 Is it clearly seen from Figs. 2, 4 and 5 that smaller molecules in a mixture with larger
7 ones have lower activity coefficient than the larger molecules in a mixture with smaller at the
8 same **packing fraction** of a system. This effect can be explained by the fact that the probability
9 of insertion of smaller molecules into the system with a predominance of larger molecules is
10 higher than that of the reverse situation. ~~It is an entropy effect.~~

11 12 13 14 **5. Conclusions**

15
16 The principal achievement of this paper has been a successful implementation of the IGCMC
17 technique is in calculating the activity coefficient of a high density fluid. Our calculations for
18 the binary mixtures clearly indicate that smaller molecules have lower activity coefficient
19 than the larger ones. This finding allows an explanation of the unexpected adsorption
20 behaviour of Cs^+ cations from the mixed electrolyte with Li^+ or Mg^{2+} ions at the negatively
21 charged Langmuir film noticed by Shapovalov and Brezesinski [18]. They found that the
22 adsorption of Cs^+ cations of the smallest hydrated radius, in comparison with the adsorption of
23 larger Li^+ and divalent Mg^{2+} , is stronger than one could expect. Earlier this behaviour was
24 confirmed by the volume corrected Poisson–Boltzmann equation [19] and by the Monte Carlo
25 simulations [20]. Also the present results indicate that adsorption of smaller ions, which have
26 the lower activity coefficient and thus the lower chemical potential, should be enhanced.
27 **Finally we would like to notice that although we intended to use the SPB and MSA theories to
28 test the IGCMC method, we have shown that these theories could be applied successfully to
29 the SPM electrolyte with the realistic concentrations of ions and solvent molecules.**

30 31 32 33 34 **Acknowledgements**

35
36 The authors are very grateful to Professors C.W. Outhwaite and L.B. Bhuiyan for their
37 comments and suggestions and for making available their SPB program. Financial support
38 from Adam Mickiewicz University, Faculty of Chemistry, is appreciated.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- [1] M.P. Allen, D.J. Tildsley. *Computer Simulation of Liquids*, pp. 49-50, Clarendon Press, Oxford (1987).
- [2] B. Widom. Some Topics in the theory of fluids. *J. Chem. Phys.*, **39**, 2808 (1963).
- [3] P. Sloth, T.S. Sørensen. Monte Carlo simulations of single-ion chemical potentials. Preliminary results for the restricted primitive model. *Chem. Phys. Lett.*, **143**, 140 (1988).
- [4] P. Sloth, T.S. Sørensen. Monte Carlo simulations of single ion chemical potentials. Results for the unrestricted primitive model. *Chem. Phys. Lett.*, **146**, 452 (1988)
- [5] D.J. Adams. Chemical potential of hard-sphere fluids by Monte Carlo methods. *Molecular Physics*, **28**, 1241 (1974).
- [6] J.P. Valleau, L.K. Cohen. Primitive model electrolytes. I. Grand canonical Monte Carlo computations. *J. Chem. Phys.*, **72**, 5935 (1980).
- [7] J.P. Valleau, L.K. Cohen, D.N. Card. Primitive model electrolytes. II. The symmetrical electrolyte. *J. Chem. Phys.*, **72**, 5942 (1980).
- [8] S. Lamperski. The individual and mean activity coefficients of an electrolyte from the inverse GCMC simulation. *Molecular Simulation*, **33**, 1193 (2007).
- [9] A. Malasics, D. Gillespie, D. Boda. Simulating prescribed particle densities in the grand canonical ensemble using iterative algorithms. *J. Chem. Phys.*, **128**, 124102 (2008).
- [10] P. Sloth, T.S. Sørensen. Single-Ion Activity Coefficients and Structure of Ionic Fluids. Results for the primitive model of electrolyte solutions. *J. Phys. Chem.*, **94**, 2116 (1990).
- [11] M. Molero, C.W. Outhwaite, L.B. Bhuiyan. Individual ionic activity coefficients from a symmetric Poisson-Boltzmann theory. *J. Chem. Soc. Faraday Trans.*, **88**, 1541 (1992).
- [12] C.W. Outhwaite, M. Molero, L.B. Bhuiyan. Primitive model electrolytes in the modified Poisson-Boltzmann theory. *J. Chem. Soc. Faraday Trans.*, **89**, 1315 (1993).
- [13] C.W. Outhwaite, M. Molero, L.B. Bhuiyan. Corrigendum to primitive model electrolytes in the modified Poisson-Boltzmann theory. *J. Chem. Soc. Faraday Trans.*, **90**, 2002 (1994).
- [14] W. Ebeling, K. Scherwinski. On the estimation of theoretical individual activity coefficients of electrolytes. *Z. Phys. Chem. (Leipzig)*, **264**, 1 (1983).
- [15] J.Reščič, V.Vlachy, L.B.Bhuiyan, C.W.Outhwaite. Monte Carlo simulations of a mixture of an asymmetric electrolyte and a neutral species. *J.Chem.Phys.* **107**, 3611 (1997).
- [16] M. Parrinello, M.P. Tosi. Analytic solution of the mean spherical approximation for a multi-component plasma. *Chem. Phys. Letters*, **64**, 579 (1979).
- [17] D.R. Lide. *Handbook of Chemistry and Physics*, 88th Edition 2007-2008, pp. 5-79 – 5-84, Taylor & Fancis Group Press, LLC (2008).
- [18] V.L. Shapovalov, G. Brezesinski. Breakdown of the Gouy-Chapman model for highly charged Langmuir monolayers: Counterion size effect. *J. Phys. Chem. B*, **110**, 10032 (2006).
- [19] P.M. Biesheuvel, M. Soestbergen. Counterion volume effects in mixed electrical double layers. *J. Colloid Interface Sci.*, **316**, 490 (2007).
- [20] S. Lamperski, C.W. Outhwaite. Monte-Carlo simulation of mixed electrolytes next to a plane charged surface. *J. Colloid Interface Sci.*, **328**, 458 (2008).

TABLES

Table 1. Activity coefficients a of one-component hard-sphere fluid.

ρ^*	η	$\ln \gamma$		
		IGCMC	Adams [5]	Ebeling and. Scherwinski equation [14]
0.1	0.0524	0.465	0.466	0.464
0.2	0.1047	1.040	1.043	1.035
0.3	0.1571	1.764	1.770	1.747
0.4	0.2094	2.656	2.651	2.648
0.5	0.2618	3.830	3.832	3.807
0.6	0.3142	5.357	5.359	5.326
0.7	0.3665	7.392	7.391	7.359
0.8	0.4189	10.211	10.210	10.153

Table 2. Activity coefficients of larger hard spheres ($d_2 = 500$ pm) in a mixture with smaller ones ($d_1 = 300$ pm) – the IGCMC results.

x_2	$\ln\gamma$							
	η	0.05	0.10	0.15	0.20	0.25	0.30	0.35
0.1	0.8806	1.9286	3.3003	5.0560	7.4012	-	-	-
0.2	0.7129	1.6385	2.7853	4.2500	6.1683	8.8690	-	-
0.3	0.6441	1.4520	2.4593	3.7430	5.3895	7.6170	-	-
0.4	0.5927	1.3240	2.2356	3.3913	4.8747	6.8490	9.7430	-
0.5	0.5520	1.2271	2.0706	3.1371	4.4910	6.2830	8.9360	-
0.6	0.5217	1.1550	1.9442	2.9376	4.2110	5.8620	8.1040	-
0.7	0.4975	1.0973	1.8448	2.7846	3.9870	5.5440	7.6210	-
0.8	0.4718	1.0496	1.7661	2.6605	3.8060	5.2960	7.2809	-
0.9	0.4578	1.0116	1.6981	2.5584	3.6540	5.0645	6.9642	-
1.0	0.4391	0.9629	1.6431	2.4734	3.5329	4.8898	6.7373	-

Table 3. Activity coefficients of smaller hard spheres ($d_1 = 300$ pm) in a mixture with larger ones ($d_2 = 500$ pm) – the IGCMC results.

x_1	$\ln \gamma$							
	η	0.05	0.10	0.15	0.20	0.25	0.30	0.35
0.1	0.2285	0.4902	0.8003	1.1734	1.6257	2.1835	2.8880	
0.2	0.2339	0.5032	0.8238	1.2079	1.6751	2.2409	2.9810	
0.3	0.2417	0.5198	0.8514	1.2496	1.7330	2.3151	3.0890	
0.4	0.2502	0.5392	0.8844	1.3002	1.8031	2.3912	3.2210	
0.5	0.2615	0.5643	0.9273	1.3659	1.8959	2.5214	3.4050	
0.6	0.2767	0.5971	0.9832	1.4507	2.0138	2.6517	3.6260	
0.7	0.2960	0.6409	1.0593	1.5660	2.1800	2.8932	3.9420	
0.8	0.3228	0.7055	1.1680	1.7337	2.4197	3.3280	4.4250	
0.9	0.3672	0.8040	1.3380	1.9976	2.8162	3.8690	5.2520	
1.0	0.43909	0.9629	1.6431	2.4734	3.5329	4.8896	6.7374	

Table 4. The activity coefficient of ions and solvent for the SPM model
at $\eta = 0.3$ – the IGCMC results.

c / M	$d_{\pm} = d_s = 300 \text{ pm}$		$d_{\pm} = d_s = 400 \text{ pm}$		$d_{\pm} = d_s = 500 \text{ pm}$	
	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$
0.05	4.6507	4.8895	4.6645	4.8894	4.6868	4.8873
0.10	4.5681	4.8925	4.6010	4.8851	4.6214	4.8873
0.25	4.4323	4.8848	4.4935	4.8838	4.5135	4.8862
0.50	4.3288	4.8862	4.3875	4.8820	4.4354	4.8844
1.00	4.1982	4.8849	4.2861	4.8801	4.3353	4.8771
1.50	4.1147	4.8776	4.2123	4.8785	4.2765	4.8711
2.00	4.0466	4.8743	4.1634	4.8743	4.2269	4.8600
2.50	3.9965	4.8785	4.1179	4.8717	4.1934	4.8584
3.00	3.9637	4.8730	4.0912	4.8637	4.1437	4.8336

Table 5. The activity coefficients of ions and solvent for the SPM model at $\eta = 0.3$, $d_{\pm} = 400$ pm for different solvent diameters – the IGCMC results.

c / M	$d_s = 300 \text{ pm}$		$d_s = 400 \text{ pm}$		$d_s = 500 \text{ pm}$	
	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$
0.05	8.3259	4.8743	4.6645	4.8894	3.1037	4.9042
0.10	8.1975	4.8568	4.6010	4.8851	3.0417	4.9203
0.25	7.9975	4.8142	4.4935	4.8838	2.9671	4.9645
0.50	7.7784	4.7490	4.3875	4.8820	2.9182	5.0505
1.00	7.4055	4.6113	4.2861	4.8801	2.9021	5.1957
1.50	7.0683	4.4748	4.2123	4.8785	2.9461	5.3715
2.00	6.7451	4.3404	4.1634	4.8743	2.9932	5.5276
2.50	6.4551	4.2080	4.1179	4.8717	3.0497	5.6831
3.00	6.1844	4.0725	4.0912	4.8637	3.1116	5.8348

Table 6. The activity coefficient of ions and solvent for the SPM model at $\eta = 0.3$, $d_s = 400$ pm for different ion diameters – the IGCMC results.

c / M	$d_{\pm} = 300$ pm		$d_{\pm} = 400$ pm		$d_{\pm} = 500$ pm	
	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$	$\ln\gamma_{\pm}$	$\ln\gamma_s$
0.05	2.7623	4.8959	4.6645	4.8894	7.2361	4.8611
0.10	2.6706	4.9037	4.6010	4.8851	7.1516	4.8403
0.25	2.5770	4.9323	4.4935	4.8838	6.9340	4.7753
0.50	2.4832	4.9827	4.3875	4.8820	6.6554	4.6675
1.00	2.3993	5.0710	4.2861	4.8801	6.2097	4.4457
1.50	2.3658	5.1603	4.2123	4.8785	5.8025	4.2359
2.00	2.3478	5.2532	4.1634	4.8743	5.4336	4.0232
2.50	2.3519	5.3502	4.1179	4.8717	5.0290	3.8118
3.00	2.3614	5.4316	4.0912	4.8637	4.6783	3.6052

FIGURE CAPTIONS

Figure 1. The activity coefficient of one-component hard sphere fluid (solid circles – simulation, open circles – Adams results [5], solid lines – Ebeling and Scherwinski equation).

Figure 2. (COLOUR ON WEB) The individual activity coefficient of a binary hard-sphere mixture from IGCMC method (graphic characters) and from the Ebeling and Scherwinski equation (solid lines) for different x_2 (the snapshot A) and x_1 (the snapshot B) at the following η : 0.05 – solid circles, 0.10 – open circles, 0.15 – solid squares, 0.20 – open squares, 0.25 – solid triangles, 0.30 – open triangles, 0.35 – solid diamonds.

Figure 3. (COLOUR ON WEB) Dependence of the activity coefficient of ions and solvent for the SPM electrolyte at $\eta = 0.3$ on the electrolyte concentration for $d = 300, 400$ and 500 pm (graphic characters – IGCMC, solid lines – SPB, dashed lines - MSA).

Figure 4. (COLOUR ON WEB) Dependence of the activity coefficient of ions (A) and solvent (B) for the SPM electrolyte at $\eta = 0.3$ on the electrolyte concentration for $d_{\pm} = 400$ pm and $d_s = 300, 400$ and 500 pm (graphic characters – IGCMC, solid lines – SPB, dashed lines - MSA).

Figure 5. (COLOUR ON WEB) Dependence of the activity coefficient of ions (A) and molecules (B) for the SPM electrolyte at $\eta = 0.3$ on the electrolyte concentration for $d_s = 400$ pm and $d_{\pm} = 300, 400$ and 500 pm (graphic characters – IGCMC, solid lines – SPB, dashed lines - MSA).









