



**HAL**  
open science

# Fractional order kinetic equations and hypoellipticity

Radjesvarane Alexandre

► **To cite this version:**

Radjesvarane Alexandre. Fractional order kinetic equations and hypoellipticity. 2011. hal-00565013v2

**HAL Id: hal-00565013**

**<https://hal.science/hal-00565013v2>**

Preprint submitted on 11 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FRACTIONAL ORDER KINETIC EQUATIONS AND HYPOELLIPTICITY

R. ALEXANDRE

ABSTRACT. We give simple proofs of hypoelliptic estimates for some models of kinetic equations with a fractional order diffusion part. The proofs are based on energy estimates together with F. Bouchut and B. Perthame previous ideas.

## 1. INTRODUCTION

Recently, after the study initiated by Morimoto and Xu [15], the paper of Lerner and all [14] was concerned with hypoelliptic effects related to a kinetic equation similar to the following one:

$$(1.1) \quad \partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) + a(t, x, v) |D_v|^{2\beta} f = g$$

Here we assume that  $(t, x, v) \in \mathbb{R}^{1+n+n}$  for some integer  $n \geq 1$ , that  $g \in L^2$ , where  $L^2 = L^2(\mathbb{R}^{1+2n})$ . We denote by  $\|\cdot\|$  the associated norm. The usual interpretation from kinetic theory is that  $t$  plays the role of a time variable,  $x$  the position and  $v$  the velocity. The coefficient  $a$  is assumed for example to be smooth and strictly positive, see below for precise hypothesis. The parameter  $\beta$  is assumed to satisfy  $0 < \beta \leq 1$ .

As regards Fourier transformation, we shall denote by  $\tau$ ,  $k$  and  $\xi$  the Fourier variables dual to  $t$ ,  $x$  and  $v$  respectively. Other notations used in (1.1) are standard, see for example [13, 17]. Let us note immediately that the third (elliptic) term on the l.h.s. of (1.1) is not exactly similar to the one considered in [14, 15] in that the behavior therein was taken as  $a |\xi|^2$  for small frequency variables  $\xi$ , but this is not an important point from the point of view of  $L^2$  theory. In the rest of the paper, we shall *always assume that all functions such as  $f$ ,  $g$  are smooth*.

For  $\beta = 1$ , (1.1) is a well known model of Fokker Planck or Kolmogorov equation, for which one can find numerous methods for proving hypoellipticity, see for example [13, 9, 14, 15, 16] and the references therein.

We refer for example to [19] for physical motivations for this type of kinetic equations. Another motivation is linked with the study of the spatially inhomogeneous Boltzmann equation without cut-off, see for example [2, 18, 1, 3, 7, 5, 6, 11], see also the recent results of [8] and references therein.

As far as we know, the study of hypoelliptic effects for problem (1.1) was initiated by Morimoto and Xu [15] and they derive therein a partial and non optimal result. This study was then completed with optimal results by Lerner and all [14], where they proved typically that  $|D_x|^{\frac{\beta}{1+2\beta}} f \in L^2$  and a similar estimate w.r.t. time variable. In both works, the authors used  $L^2$  type methods.

While still working with an energy method, we want to show that a slight modification of the computations of Bouchut [9] can lead to the same results as Lerner and all [14], and therefore in comparison, we provide a very simple proof. One advantage is that it is very simple to keep track of the different constants depending on the given coefficients  $a$ , and furthermore, we avoid using any deep pseudo differential calculus. However, in order to study the model problem (1.1), we do use one result, namely Proposition 1.1 from Bouchut [9], whose proof is also elementary as it relies on averaging regularity type arguments. Bouchut's result is given by

---

Date: February 2011.

2000 *Mathematics Subject Classification*. 35H10, 76P05, 84C40.

*Key words and phrases*. Boltzmann equation, hypoellipticity, non-cutoff cross-sections.

**Proposition 1.1.** [Proposition 1.1 of [9]] Assume  $f \in L^2$ ,  $g \in L^2$ ,  $|D_v f|^\alpha f \in L^2$  for some  $\alpha \geq 0$  and

$$(1.2) \quad \partial_t + v \cdot \nabla_x f = g,$$

Then

$$\| |D_x|^{\frac{\alpha}{1+\alpha}} f \| \lesssim \| g \|_{1+\alpha} \| |D_v|^\alpha f \|_{1+\alpha}^{\frac{1}{\alpha}}$$

The proof done in [9] uses both Fourier transform w.r.t. time and space variables  $(t, x)$ , and arguments introduced in [10]. It might be plausible to only use Fourier transform w.r.t. variable  $x$  by using the arguments of [10]. For completeness, we give yet another proof which was used in Alexandre [1] following arguments introduced by Perthame [16] in his study of higher moments estimates. Our proof is also elementary, but uses this time both space and velocity variables Fourier transform. However, we do not use Fourier transform w.r.t. time variable, and therefore we can also deal with the Cauchy initial value problem associated with model problem (1.2). We refer also to [4] for another proof involving Fourier transform w.r.t. time and space variables and a certain kind of uncertainty principle. All in all, it is now clear that any other different and simple proof of the above Proposition would be of interest.

Once given Proposition 1.1, we can proceed to study hypoelliptic effects connected with (1.1). As usual, we shall begin to study the case of constant  $a$ , say 1, that is

$$(1.3) \quad \partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) + |D_v|^{2\beta} f = g.$$

Of course, a direct Fourier analysis is able to take care of this simple model, but recall that we are looking for energy type estimates.

Our main result is given by

**Theorem 1.2.** (1) Let  $f$  satisfy (1.3). Then one has

$$\| |D_v|^{2\beta} f \| + \| |D_x|^{\frac{2\beta}{1+2\beta}} f \| \lesssim \| g \|.$$

(2) Let  $f$  satisfy (1.1). Assume that  $a = b^2 \chi + a_-$  for some strictly positive constant  $a_-$ , a smooth positive function  $b$  and a compactly supported and positive function  $\chi$ . Then one has

$$\| |D_v|^{2\beta} f \| + \| |D_x|^{\frac{2\beta}{1+2\beta}} f \| \lesssim C_a [\| g \| + \| f \|].$$

As it will appear clearly in the proof, similar estimate w.r.t. space variable also holds for time variable. Furthermore, the proofs given below can also be adapted to take into account the initial value problem. Finally, the assumption on  $a$  might appear strange, but this is one possible choice among many others. We mention that if  $a$  is assumed to be only locally bounded from below, then all proofs adapt up to introducing cutoff functions. Finally, the proof also shows that it is not necessary to have a diffusion term as above. One might consider instead an equation such as

$$\partial_t + v \cdot \nabla_x f + \int_{\mathbb{R}^n} \frac{K(t, x, v)}{|h|^{n+2\beta}} [f(v+h) - f(v)] dh = g$$

with good assumptions on the kernel  $K$ . This example is closer to a linear version of Boltzmann operator. Details are left to the interested reader. In any case, the main issue is concerned with the multipliers introduced in the proofs with the above kernel  $K$ .

## 2. THE FREE TRANSPORT EQUATION: PROOF OF PROPOSITION 1.1

We are interested in the transport equation (1.2), under the assumptions of Proposition 1.1.

It is shown by Bouchut that we obtain  $|D_x|^{\frac{\alpha}{1+\alpha}} f \in L^2$ . The method of proof was based on velocity arguments, see [9, 10] and references therein. In fact another proof is also provided in [4] using a kind of uncertainty principle, but which is more complex. Finally, another argument based on Perthame method [16] is also possible, see for example Alexandre [1]. As mentioned by Bouchut, the

commutator method seems to require more derivatives w.r.t. variable  $v$ , but in that case, the proof is very easy (see the proof in [9]).

We shall apply Perthame's argument for the usual Fokker Planck case below, following [1]. Note that this a Fourier method, using characteristics associated on the Fourier side which is also somehow used in the paper by Lerner and all [14].

If  $\hat{\cdot}$  denotes the Fourier transform with respect to the variables  $(x, v)$  and  $(k, \xi)$  the dual variables, one has

$$\partial_t \hat{f} - k \cdot \nabla_{\xi} \hat{f} = \hat{g}.$$

Multiplying by  $\bar{f}$ , and taking the complex conjugate also, we obtain:

$$\partial_t |\hat{f}|^2 - k \cdot \nabla_{\xi} |\hat{f}|^2 = 2\mathcal{R}e(\hat{g}, \bar{f}) \lesssim |\hat{g}| |\hat{f}|.$$

Then

$$|\hat{f}(t, k, \xi)|^2 \lesssim \int_{-\infty}^t (|\hat{f}\hat{g}|(k, \xi + sk, t-s)) ds.$$

Fix  $r \geq 0$  and  $D \geq 0$ , and  $k$ . Then

$$\begin{aligned} & \int_{-\infty}^{\infty} dt \int_{\xi} |k|^r |\hat{f}(t, k, \xi)|^2 \\ &= \int_{-\infty}^{\infty} dt \int_{|\xi| \geq D} |k|^r |\hat{f}(t, k, \xi)|^2 + \int_{-\infty}^{\infty} dt \int_{|\xi| \leq D} |k|^r |\hat{f}(t, k, \xi)|^2 = A + B. \end{aligned}$$

For B, one has:

$$B \lesssim \int_{-\infty}^{\infty} dt \int_{|\xi| \leq D} |k|^r \int_{-\infty}^t (|\hat{f}\hat{g}|(k, \xi + sk, t-s)) ds.$$

Changing variable in  $\xi$ , this gives

$$B \lesssim \int_{-\infty}^{\infty} dt \int d\xi |k|^r \int_{-\infty}^{\infty} 1_{s \leq t} 1_{|\xi - sk| \leq D} (|\hat{f}\hat{g}|(k, \xi, t-s)) ds$$

Change variables in  $t$  (for fixed  $s$ ) to get

$$B \lesssim \int_0^{\infty} dt \int d\xi |k|^r \int_{-\infty}^{\infty} 1_{|\xi - sk| \leq D} (|\hat{f}\hat{g}|(k, \xi, t)) ds$$

Since  $|\xi - sk| \geq \|\xi\| - s|k|$ , it follows that

$$\begin{aligned} B &\lesssim \int_0^{\infty} dt \int d\xi |k|^{r-1} D (|\hat{f}\hat{g}|(k, \xi, t)) \\ &\lesssim \varepsilon \int_0^{\infty} dt \int d\xi |k|^{2(r-1)} D^2 |\hat{f}|^2(k, \xi, t) + C_{\varepsilon} \int_0^{\infty} dt \int d\xi |\hat{g}|^2(k, \xi, t) \end{aligned}$$

for any  $\varepsilon > 0$ .

Now for A, we get directly

$$A \lesssim \int_{-\infty}^{\infty} dt \int d\xi |\xi|^m D^{-m} |k|^r |\hat{f}(t, k, \xi)|^2.$$

Choose  $D = |k|^{\frac{r}{m}}$ . Then

$$A \lesssim \int_{-\infty}^{\infty} dt \int d\xi |\xi|^m |\hat{f}(t, k, \xi)|^2.$$

Then note that  $|k|^{2(r-1)} D^2 = |k|^{2(r-1) + \frac{2r}{m}} = |k|^r$ , if we choose the value of  $r$  such that  $r = \frac{2m}{m+2}$ . We choose  $m = 2\alpha$ . Therefore  $r = \frac{2\alpha}{1+\alpha}$ . In conclusion with all these choices, we get, for fixed  $k$ , by absorbing the right hand side with the left hand side:

$$\int_{-\infty}^{\infty} dt \int_{\xi} |k|^{\frac{2\alpha}{1+\alpha}} |\hat{f}(t, k, \xi)|^2 \lesssim \int_{-\infty}^{\infty} dt \int d\xi |\xi|^{2\alpha} |\hat{f}(t, k, \xi)|^2 + C_{\varepsilon} \int_0^{\infty} dt \int d\xi |\hat{g}|^2(k, \xi, t)$$

and therefore

$$\| |D_x|^{\frac{\alpha}{1+\alpha}} f \| \lesssim \| |D_v|^\alpha f \| + \|g\|.$$

It should be observed that we do not have the same scaling as in Bouchut. But this can be fixed easily as follows. We proceed as above, replacing  $D$  by  $\lambda D$  but use instead Cauchy Schwarz inequality for  $B$ , with the same choice of parameters:

$$B \lesssim \lambda \int_0^\infty dt \int d\xi |k|^{r-1} D(|\hat{f}\hat{g}|(k, \xi, t)) \lesssim \lambda \left\{ \int_0^\infty dt \int d\xi |k|^r |\hat{f}|^2(k, \xi, t) \right\}^{\frac{1}{2}} \left\{ \int_0^\infty dt \int d\xi |\hat{g}|^2(k, \xi, t) \right\}^{\frac{1}{2}}$$

while

$$A \lesssim \lambda^{-m} \int_{-\infty}^\infty dt \int d\xi |\xi|^m |\hat{f}(t, k, \xi)|^2.$$

Then we get an inequality such as:

$$U \lesssim \lambda U^{\frac{1}{2}} V^{\frac{1}{2}} + \lambda^{-m} W.$$

If we choose  $\lambda$  such that the two terms on the r.h.s. coincide then we get, after some computations

$$U \lesssim V^{\frac{m}{m+2}} W^{\frac{2}{m+2}}.$$

Now integrate w.r.t.  $k$  and use Holder inequality with exponent  $p = m/(m+2)$  to get

$$\int U \lesssim \left[ \int V \right]^{\frac{m}{m+2}} \left[ \int W \right]^{\frac{2}{m+2}}$$

Recalling that  $m = 2\alpha$ , we get exactly

$$\| |D_x|^{\frac{\alpha}{1+\alpha}} f \| \lesssim \|g\|^{\frac{\alpha}{1+\alpha}} \| |D_v|^\alpha f \|^{\frac{1}{1+\alpha}}.$$

**Remark 2.1.** *One can also get estimations for the initial value problem, say if we consider the transport equation (1.2) for say positive time and a given initial value at time 0,  $f_0$ . For example, in that case, the small frequency part gives an additional term which can be estimated as follows*

$$\begin{aligned} III &= \int_0^\infty dt \int_{|\xi| \leq D} |k|^r |\hat{F}_0(k, \xi + tk)|^2 = \int_0^\infty dt \int_{|\xi - tk| \leq D} |k|^r |\hat{F}_0(k, \xi)|^2 \leq \\ &\lesssim \int_0^\infty dt \int_{\mathbb{R}_\xi^3} 1_{|t - \frac{|\xi|}{|k|}| \leq \frac{D}{|k|}} |\hat{F}_0(k, \xi)|^2 \lesssim |k|^{r-1} D \int_{\mathbb{R}_\xi^3} |\hat{F}_0(k, \xi)|^2. \end{aligned}$$

### 3. PROOF OF THE FIRST PART OF THEOREM 1.2: THE CONSTANT COEFFICIENT CASE

Here we shall prove the first part of Theorem 1.2 related to  $f$  satisfying (1.3). We shall again adapt the ideas of Bouchut, except for a modification of the test multiplier in Step 4 below.

**Step 1:** We multiply (1.3) by  $\bar{f}$  and integrate over all variables. Taking into account usual symmetry cancellation, we get

$$(3.4) \quad \| |D_v|^\beta f \| \lesssim \|g\|^{\frac{1}{2}} \|f\|^{\frac{1}{2}}.$$

**Step 2:** Now having in mind that we want to prove that  $|D_v|^{2\beta} f \in L^2$  and knowing that  $g \in L^2$ , we note that

$$\partial_t f + v \cdot \nabla_x f = G \equiv -|D_v|^{2\beta} f + g.$$

Therefore applying Proposition 1.1 of Bouchut (with the parameter  $\alpha$  there replaced by  $2\beta$ ), it follows that

$$\begin{aligned} \| |D_x|^{\frac{2\beta}{1+2\beta}} f \| &\lesssim \| |D_v|^{2\beta} f \|_{\frac{1}{1+2\beta}} \cdot \| -|D_v|^{2\beta} f + g \|_{\frac{2\beta}{1+2\beta}} \\ &\lesssim \| |D_v|^{2\beta} f \|_{\frac{1}{1+2\beta}} \left[ \| |D_v|^{2\beta} f \| + \| |D_v|^{2\beta} f \|_{\frac{1}{1+2\beta}} \cdot \|g\|_{\frac{2\beta}{1+2\beta}} \right]. \end{aligned}$$

Thus

$$(3.5) \quad \||D_x|^{\frac{2\beta}{1+2\beta}} f\| \lesssim \||D_v|^{2\beta} f\| + \||D_v|^{2\beta} f\|^{\frac{1}{1+2\beta}} \cdot \|g\|^{\frac{2\beta}{1+2\beta}}.$$

**Step 3:** Now apply  $|D_x|^{\frac{\beta}{1+2\beta}}$  on (1.3), multiply by  $|D_x|^{\frac{\beta}{1+2\beta}} \bar{f}$  and integrate to get

$$(3.6) \quad \||D_v|^\beta |D_x|^{\frac{\beta}{1+2\beta}} f\| \lesssim \||D_x|^{\frac{2\beta}{1+2\beta}} f\|^{\frac{1}{2}} \|g\|^{\frac{1}{2}}.$$

**Step 4:** This step is different from Bouchut's arguments, in that we choose another multiplier, taking into account the control for large frequency variable associated with  $x$ .

Considering (1.3), multiply it by  $(|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f}$  (see the remark below for the choice of this multiplier) and integrate to get

$$\begin{aligned} & \int (|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f} \cdot |D_v|^{2\beta} f = \\ & -\operatorname{Re}((|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f}, v \cdot \nabla_x f) + \operatorname{Re}((|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f}, g) = I + II \end{aligned}$$

Using Fourier transformation for example, it follows that

$$II \lesssim (\||D_v|^{2\beta} f\| + \||D_x|^{\frac{2\beta}{1+2\beta}} f\|) \|g\|.$$

With the previous steps, we get

$$II \lesssim \||D_v|^{2\beta} f\| \|g\| + \||D_v|^{2\beta} f\|^{\frac{1}{1+2\beta}} \cdot \|g\|^{1+\frac{2\beta}{1+2\beta}}$$

On the other hand, using Parseval relation

$$\begin{aligned} I &= -\operatorname{Re}((|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f}, v \cdot \nabla_x f) = -\operatorname{Re}((|\xi|^2 + |k|^{\frac{2}{1+2\beta}})^\beta \bar{f}, k_j \partial_{\xi_j} \hat{f}) \\ &= \operatorname{Re}(\partial_{\xi_j} [ (|\xi|^2 + |k|^{\frac{2}{1+2\beta}})^\beta ] \bar{f}, k \hat{f}) + \operatorname{Re}((|\xi|^2 + |k|^{\frac{2}{1+2\beta}})^\beta \partial_{\xi_j} \bar{f}, k \hat{f}). \end{aligned}$$

Thus

$$\begin{aligned} I &= \beta \operatorname{Re}(\xi_j (|\xi|^2 + |k|^{\frac{2}{1+2\beta}})^{\beta-1} \bar{f}, k \hat{f}) = \operatorname{Re} \int \bar{f} \hat{f}(\xi, k) (|\xi|^2 + |k|^{\frac{2}{1+2\beta}})^{\beta-1} \\ &\lesssim \int \bar{f} \hat{f} |\xi| |k| |\xi|^{\beta-1} |k|^{\frac{\beta-1}{1+2\beta}} \lesssim \int |\xi|^\beta |k|^{\frac{\beta}{1+2\beta}} \bar{f} \cdot |k|^{\frac{2\beta}{1+2\beta}} \hat{f} \\ &\lesssim \||D_v|^\beta |D_x|^{\frac{\beta}{1+2\beta}} f\| \cdot \||D_x|^{\frac{2\beta}{1+2\beta}} f\|. \end{aligned}$$

Using the previous steps, it follows that

$$I \lesssim \||D_v|^{2\beta} f\|^{\frac{3}{2}} \|g\|^{\frac{1}{2}} + \||D_v|^{2\beta} f\|^{\frac{3}{2} \frac{1}{1+2\beta}} \|g\|^{\frac{3}{2} \frac{2\beta}{1+2\beta} + \frac{1}{2}}.$$

**Step 5:** In conclusion, we get

$$I \lesssim \||D_v|^{2\beta} f\|^{\frac{3}{2}} \|g\|^{\frac{1}{2}} + \||D_v|^{2\beta} f\|^{\frac{3}{2} \frac{1}{1+2\beta}} \|g\|^{\frac{3}{2} \frac{2\beta}{1+2\beta} + \frac{1}{2}}$$

and

$$II \lesssim \||D_v|^{2\beta} f\| \|g\| + \||D_v|^{2\beta} f\|^{\frac{1}{1+2\beta}} \cdot \|g\|^{1+\frac{2\beta}{1+2\beta}}$$

Using Holder inequality, we get for example, for small  $\varepsilon > 0$

$$\begin{aligned} \||D_v|^{2\beta} f\| \|g\| &\lesssim \varepsilon^2 \||D_v|^{2\beta} f\|^2 + C_\varepsilon \|g\|^2 \\ \||D_v|^{2\beta} f\|^{\frac{3}{2}} \|g\|^{\frac{1}{2}} &\lesssim \varepsilon^{\frac{4}{3}} \||D_v|^{2\beta} f\|^2 + C_\varepsilon \|g\|^2 \\ \||D_v|^{2\beta} f\|^{\frac{1}{1+2\beta}} \|g\|^{1+\frac{2\beta}{1+2\beta}} &\lesssim \varepsilon^{2(1+2\beta)} \||D_v|^{2\beta} f\|^2 + C_\varepsilon \|g\|^2 \end{aligned}$$

and therefore, it follows that

$$\||D_v|^{2\beta} f\| \lesssim C_\varepsilon \|g\|.$$

We get also from Step 2 that

$$\| |D_x|^{\frac{2\beta}{1+2\beta}} f \| \lesssim \|g\|.$$

- Remark 3.1.** (1) *To get also the same result for the derivative w.r.t.  $t$ , we repeat the above arguments, but with the multiplier  $(|D_t|^{\frac{2}{1+2\beta}} + |D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta$ . As well we could have done the computations from the beginning with this multiplier.*
- (2) *We choose a multiplier which is somehow singular near null value of the frequency variables. It would have been better to choose  $(\delta + |D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta$ , for a small  $\delta$ . Nothing is changed, except that now the upper bound involves  $\|f\|$ . In fact an even better choice would have been to choose  $(\delta + |D_v|^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta$ , whose symbol is smooth.*
- (3) *For the initial value problem, it might be interesting to consider the above multiplier by also  $(\delta + |tD_v|^2 + \langle tD_x \rangle^{\frac{2}{1+2\beta}})^\beta$ .*

#### 4. PROOF OF THE SECOND PART OF THEOREM 1.2: THE NON CONSTANT COEFFICIENT CASE

We now consider the model problem (1.1). As it should be clear now, the main issue is the estimation of the commutator of any smooth function with the operator  $(\langle D_v \rangle^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta$ , see the remarks in the previous section for the choice of this multiplier. Recall that we assume

$$a = b^2 \chi^2 + a_-$$

with  $b \geq 0$  smooth and  $\chi \geq 0$  compactly supported, and that we do not assume any lower bound on  $b$ . Then, we write

$$\begin{aligned} & \int (\langle D_v \rangle^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta \bar{f} \cdot a \cdot |D_v|^{2\beta} f \\ &= \int (\langle D_v \rangle^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta \bar{f} \cdot a_- \cdot |D_v|^{2\beta} f + \int (\langle D_v \rangle^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta \bar{f} \cdot b^2 \chi^2 \cdot |D_v|^{2\beta} f \\ &= I + II. \end{aligned}$$

The first term  $I$  is nice since it will give a lower bound as we wish. So we need to deal with  $II$  to make appear a positive term and commutators terms:

We write below  $P = (\langle D_v \rangle^2 + \langle D_x \rangle^{\frac{2}{1+2\beta}})^\beta$  and  $Q = |D_v|^{2\beta}$ . Then

$$\begin{aligned} II &= \int (|D_v|^2 + |D_x|^{\frac{2}{1+2\beta}})^\beta \bar{f} \cdot b^2 \chi^2 \cdot |D_v|^{2\beta} f = \int b \chi P f \cdot b \chi Q \bar{f} \\ &= \int \{ [b \chi, P] f + P(b \chi f) \} \{ [b \chi, Q] f + Q(b \chi f) \} \\ &= \int [b \chi, P] f \cdot [b \chi, Q] f + [b \chi, P] f \cdot Q(b \chi f) + P(b \chi f) \cdot [b \chi, Q] f + P(b \chi f) Q(b \chi f) \end{aligned}$$

The last term is positive. So we need to consider the first three terms, and in particular to study the commutator.

The easiest commutator is  $[b \chi, Q]$ , Indeed, we note that

**Lemma 4.1.** *For  $\beta \leq \frac{1}{2}$ , one has*

$$\| [b \chi, Q] f \| \lesssim c_b \|f\|$$

*and for  $\beta \geq \frac{1}{2}$ , one has*

$$\| [b \chi, Q] f \| \lesssim c_b [\| |D_v|^{\beta-1/2} f \| + \|f\|]$$

*where the constant  $c_b$  only depends on a finite number of derivatives of  $b$ .*

*Proof.* This is a well know result, so we just sketch the main arguments. One possibility is to write (see for example Stein [17]), for some constant  $c_n$

$$Qf = c_n \int_h [f(v+h) - f(v)]/|h|^{n+2\alpha}.$$

Then

$$\begin{aligned} b\chi Qf &= c_n b\chi \int_h [f(v+h) - f(v)]/|h|^{n+2\alpha} \\ &= c_n \int_h [b\chi f(v+h) - b\chi f(v)]/|h|^{n+2\alpha} + \int_h [b\chi(v) - b\chi(v+h)]f(v+h)/|h|^{n+2\alpha}. \end{aligned}$$

Therefore:

$$\begin{aligned} [b\chi, Q]f &= \int_h [b\chi(v) - b\chi(v+h)]f(v+h)/|h|^{n+2\alpha} \\ &= \int_z [(b\chi(v) - b\chi(z))]f(z)/|z-v|^{n+2\alpha} = \int_z K(v, z)f(z)dz \end{aligned}$$

Note that:

$$|K(v, z)| \lesssim |z-v|^{n+2\alpha-1} \text{ and } |K(v, z)| \lesssim 1/|z-v|^{n+2\alpha}$$

Thus if  $\beta < \frac{1}{2}$ , we can apply Shur's Lemma to see that  $[\chi, Q]$  is a  $L^2$  bounded operator. For larger values of  $\beta$  we need to use the symetrized version of the integral expression of  $Q$ . In fact another method is the following: write  $|D_v|^{2\alpha} = [|D_v|^{2\alpha} - \langle D_v \rangle^{2\alpha}]_+ + \langle D_v \rangle^{2\alpha}$ . The first factor is clearly bounded in  $L^2$  while the second one is dealt with the same method as the Lemma just below for the commutator with  $P$ . □

**Lemma 4.2.** For  $\beta \leq \frac{1}{2}$ , one has

$$\| [b\chi, P]f \| \lesssim c_b \|f\|$$

and for  $\beta \geq \frac{1}{2}$ , one has

$$\| [b\chi, P]f \| \lesssim c_b [\| |D_v|^{\beta-1/2} f \| + \|f\|]$$

where the constant  $c_b$  only depends on a finite number of derivatives of  $b$ .

*Proof.* Set  $P = p(D_x, D_v) = (|D_v|^2 + |D_x|^{1+\frac{2}{1+2\beta}})^\beta$ , with  $p(k, \xi) = (\langle \xi \rangle^2 + \langle k \rangle^{1+\frac{2}{1+2\beta}})^\beta$ . Let  $\tilde{b} = b\chi$ . Then  $[P, \tilde{b}]u = P(\tilde{b}u) - \tilde{b}(Pu)$ . Therefore

$$\begin{aligned} \widehat{[P, \tilde{b}]u}(k, \xi) &= \widehat{P(\tilde{b}u)} - \widehat{\tilde{b}(Pu)} = p(k, \xi)\widehat{\tilde{b}} * \widehat{u}(k, \xi) - \widehat{\tilde{b}} * [p\widehat{u}] \\ &= \int_{k', \xi'} [p(k, \xi) - p(k', \xi')]\widehat{\tilde{b}}(k-k', \xi-\xi')\widehat{u}(k', \xi') \end{aligned}$$

Set

$$K(k, \xi, k', \xi') = [p(k, \xi) - p(k', \xi')]\widehat{\tilde{b}}(k-k', \xi-\xi') = K_1 + K_2$$

with

$$K_1 = [p(k, \xi') - p(k', \xi')]\widehat{\tilde{b}}(k-k', \xi-\xi')$$

and

$$K_2 = [p(k, \xi) - p(k, \xi')]\widehat{\tilde{b}}(k-k', \xi-\xi')$$

It is immediately seen that the second term gives rise to a kernel for which we can apply Schur Lemma. That is we see that  $|K_2| \lesssim |\widehat{\tilde{b}}(k-k', \xi-\xi')||k-k'|$ , and then (for any small  $\delta$ )

$$\int_{k, \xi} \langle k \rangle > |\widehat{\tilde{b}}(k, \xi)| \lesssim \left[ \int \langle (k, \xi) \rangle^{2n+2+\delta} |\widehat{\tilde{b}}|^2 \right]^{\frac{1}{2}}.$$



Thus, going back to the inverse Fourier transform, we have an operator  $\tilde{K}_1$  such that  $\|\tilde{K}_1 f\| \lesssim \|f\|$ . For the part related to  $K_2$ , suppose first that  $\beta \leq 1/2$ . Then using Taylor's formulae, it follows that

$$\int_{k,\xi} |K_2(k, \xi, k', \xi')| \lesssim \int_{k,\xi} |\xi - \xi'| \widehat{b}(k - k', \xi - \xi') \lesssim \left[ \int_{k,\xi} \langle (k, \xi) \rangle^{2n+2\delta} \widehat{b}^2 \right]^{\frac{1}{2}}$$

Schur Lemma applies and yields that it is a  $L^2$  bounded operator. If  $\beta \geq \frac{1}{2}$ , we have an upper bound on  $K_2$  as  $|\hat{\alpha}||\xi - \xi'| \ll \langle \xi \rangle^{\beta-1/2} + \langle \xi' \rangle^{\beta-1/2}$ . Then, it's enough to use Petree's inequality to conclude.  $\square$

**Remark 4.3.** *The bound above depends on the norm of  $b\chi$  is  $H_{x,v}^m$ , with  $m = n + 1 + \delta$  for small  $\delta > 0$ . It is likely not optimal. When  $a$  does not depend on variable  $x$ , one can obtain better bounds.*

Putting together the two previous Lemma, we can conclude the proof of the second part of Theorem 1.2.

#### REFERENCES

- [1] R. Alexandre, Some solutions of the Boltzmann equation without angular cutoff. *J. Statist. Phys.* 104 (2001), no. 1-2, 327-358.
- [2] R. Alexandre, A review on Boltzmann equation with singular kernels. *Kinetic and Related Models*, **2-4** (2009) 541-646.
- [3] R. Alexandre, L. Desvillettes, C. Villani and B. Wennberg, Entropy dissipation and long-range interactions, *Arch. Rational Mech. Anal.*, **152** (2000) 327-355.
- [4] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu and T. Yang, Uncertainty principle and kinetic equations, *J. Funct. Anal.*, **255** (2008) 2013-2066.
- [5] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu and T. Yang, Regularizing effect and local existence for non-cutoff Boltzmann equation, *Arch. Rational Mech. Anal.*, **198** (2010), 39-123.
- [6] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu and T. Yang, Global existence and full regularity of the Boltzmann equation without angular cutoff, Preprint HAL, <http://hal.archives-ouvertes.fr/hal-00439227/fr/>
- [7] R. Alexandre and C. Villani, On the Boltzmann equation for long-range interaction, *Comm. Pure Appl. Math.* **55** (2002) 30-70.
- [8] D. Arsenio, From Boltzmann's equation to the incompressible Navier-Stokes-Fourier system with long-range interactions. Preprint.
- [9] F. Bouchut, Hypocoelliptic regularity in kinetic equations. *J. Math. Pures Appl.* (9) 81 (2002), no. 11, 1135-1159.
- [10] F. Bouchut, L. Desvillettes, Averaging lemmas without time Fourier transform and application to discretized kinetic equations. *Proc. Roy. Soc. Edinburgh Sect. A* 129 (1999), no. 1, 193-6.
- [11] P. Gressman, R. Strain, Global strong solutions of the Boltzmann equation without angular cut-off. arXiv:0912.0888.
- [12] F. Herau, K. Pravda-Starov, Anisotropic hypoelliptic estimates for Landau-type operators, Preprint, <http://arxiv.org/abs/1003.3265>
- [13] L. Hormander, The analysis of linear Partial Differential Operators. Vol. I - IV. Springer, Berlin 1990.
- [14] N. Lerner, Y. Morimoto, K. Pravda-Starov, Hypocoelliptic Estimates for a Linear Model of the Boltzmann Equation without Angular Cutoff. Preprint, <http://arxiv.org/abs/1012.4915>
- [15] Y. Morimoto and C.-J. Xu, Hypocoellipticity for a class of kinetic equations, *J. Math. Kyoto Univ.* **47** (2007), 129-152.
- [16] B. Perthame, Higher moments for kinetic equations: the Vlasov-Poisson and Fokker-Planck cases. *Math. Methods Appl. Sci.* 13 (1990), no. 5, 441-452.
- [17] E. Stein, Singular Integrals and Differentiability Properties of Functions. Princeton University Press, New York, 1970.
- [18] C. Villani, *A review of mathematical topics in collisional kinetic theory*. Handbook of Fluid Mechanics. Ed. S. Friedlander, D. Serre, 2002.
- [19] G.-M. Zaslavsky, The physics of chaos in hamiltonian systems. World Scientific Publ. Co., Imperial College Press 2007.

R. ALEXANDRE, DEPARTMENT OF MATHEMATICS, SHANGHAI JIAO TONG UNIVERSITY  
SHANGHAI, 200240, P. R. CHINA

AND

IRENAV, ARTS ET METIERS PARIS TECH, ECOLE NAVALE,  
LANVEOC POULMIC, BREST 29290 FRANCE

E-mail address: radjesvarane.alexandre@ecole-navale.fr