



HAL
open science

On logical hierarchies within FO2-definable languages

Manfred Kufleitner, Pascal Weil

► **To cite this version:**

Manfred Kufleitner, Pascal Weil. On logical hierarchies within FO2-definable languages. 2011. hal-00567171v1

HAL Id: hal-00567171

<https://hal.science/hal-00567171v1>

Preprint submitted on 18 Feb 2011 (v1), last revised 16 Mar 2012 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON LOGICAL HIERARCHIES WITHIN FO²-DEFINABLE LANGUAGES

MANFRED KUFLEITNER AND PASCAL WEIL

ABSTRACT. We consider the class of languages defined in the 2-variable fragment of the first-order logic of the linear order. Many interesting characterizations of this class are known, as well as the fact that restricting the number of alternations yields an infinite hierarchy. We show that each level of this hierarchy forms a variety of languages and hence, it admits an algebraic characterization. With this tool, we show that the quantifier alternation hierarchy over FO²[<] is decidable within one unit. For this purpose, we relate each level of the hierarchy with decidable varieties of languages, which can be defined in terms of iterated deterministic and co-deterministic products. A crucial notion in this process is that of condensed rankers, a refinement of the rankers of Weis and Immerman and the turtle languages of Schwentick, Thérien and Vollmer.

Many important properties of systems are modeled by finite automata. Frequently, the formal languages induced by these systems are definable in first-order logic. Our understanding of its expressive power is of direct relevance for a number of application fields, such as verification.

The first-order logic we are interested in, in this paper, is the first-order logic of the linear order, written FO[<], interpreted on finite words. It is well-known that the languages that are definable in this logic are exactly the star-free languages, or equivalently the regular languages whose syntactic monoid is aperiodic (that is: satisfies an identity of the form $x^{n+1} = x^n$ for some integer n) [18, 14] (see also [15, 21, 22]); and that deciding whether a finite automaton accepts such a language is PSPACE-complete [3].

Fragments of first-order logic defined by the limitation of certain resources have been studied in detail. For instance, the quantifier alternation hierarchy, with its close relation with the dot-depth hierarchy of star-free languages, offers one of the oldest open problems in formal language theory: we know that the hierarchy is infinite and that its levels are characterized algebraically (by a property of the syntactic monoids), but we do not know whether these levels (besides levels 0 and 1) are decidable. In contrast, it is known that the quantifier alternation hierarchy for the first-order logic of the successor, FO[S], collapses at level 2 [27, 16].

Date: February 12, 2011.

1998 ACM Classification: F.4.3, F.4.1.

The second author acknowledges support from ANR project 2010 BLAN 0202 01 FREC. Both authors acknowledge support from the ESF program AUTOMATHA.

Another natural limitation considers the number of variables in a formula. This limitation has attracted a good deal of attention, as the trade-off between formula size and number of variables is known to be related with the trade-off between parallel time and number of processes, see [9, 1, 8].

It is well-known that every first-order formula is equivalent to one using at most three variables. On the other hand, the first-order formulas using at most two variables, written $\text{FO}^2[<]$ are strictly less expressive. The class of languages defined by such formulas admits many remarkable characterizations [24]. To begin with, a language is $\text{FO}^2[<]$ -definable if and only if it is recognized by a monoid in the pseudovariety **DA** [26] (a precise definition will be given in Section 2). As with the characterization of $\text{FO}[<]$ -definability by aperiodic monoids, this characterization implies decidability. The $\text{FO}^2[<]$ -definable languages are also characterized in terms of unambiguous products of languages (see Section 6.3) and in terms of the unary fragment of propositional temporal logic [7] (see Section 1.3). For a survey of these properties, the reader is referred to [24, 6].

In this paper, we consider the quantifier alternation hierarchy within the two-variable fragment of first-order logic. We denote by $\text{FO}_m^2[<]$ the fragment of $\text{FO}^2[<]$ consisting of formulas using at most 2 variables and at most m alternations of quantifiers. In the sequel, we omit specifying the predicate $<$ and we write simply FO , FO^2 or FO_m^2 .

Schwentick, Thérien and Vollmer introduced the so-called *turtle programs* to characterize the expressive power of FO^2 [20]. These programs are sequences of directional instructions of the form *go to the next a to the right*, *go to the next b to the left*. More details can be found in Section 1.2 below. Turtle programs were then used, under the name of *rankers*, by Weis and Immerman [30] (first published in [29]) to characterize FO_m^2 in terms of rankers with m alternations of directions (right *vs.* left). Their subtle characterization, Theorem 1.7 below, does not yield a decidability result. It forms however the basis of our results.

Our first set of results shows that \mathcal{FO}_m^2 (the set of $\text{FO}_m^2[<]$ -definable languages) forms a variety of languages, and the same holds for the classes of languages defined by rankers having m alternations of directions. The meaning of this result is that membership of a language L in these classes depends only on the syntactic monoid of L . This justifies using algebraic methods to approach the decidability problem for \mathcal{FO}_m^2 — a technique that has proved very useful in a number of situations (see for instance [15, 24, 23, 6]).

Our results next show that rankers are actually better suited to the study of a natural hierarchy within the unary fragment of propositional temporal logic, and we introduce the notion of *condensed* rankers, which turns out to be more adapted to discuss the quantifier alternation within $\text{FO}_m^2[<]$. Condensed rankers introduce a notion of efficiency in the path they describe in a word, see Section 3. Again, we show that the classes of languages defined by rankers with at most m changes of directions define varieties of languages,

written \mathcal{R}_m and \mathcal{L}_m depending on whether the initial move is towards the right or towards the left. We show that these classes are decidable, and that they admit a neat characterization in terms of closure under alternated deterministic and co-deterministic products. Moreover, we show that

$$\mathcal{R}_m \cup \mathcal{L}_m \subseteq \mathcal{FO}_m^2 \subseteq \mathcal{R}_{m+1} \cap \mathcal{L}_{m+1}.$$

This shows that one can effectively compute, given a language L in \mathcal{FO}^2 , an integer m such that L is in \mathcal{FO}_{m+1}^2 , possibly in \mathcal{FO}_m^2 , but not in \mathcal{FO}_{m-1}^2 . That is, we can compute the quantifier alternation depth of L within one unit. As indicated above, this is much more precise than the current level of knowledge on the general quantifier alternation hierarchy in $\text{FO}[\prec]$.

Many of these results were announced in [11], with a few differences. In particular, the definition of the sets $\underline{R}_{m,n}^X$ (Section 1.2) in [11] introduced a mistake which is corrected here. And the proof of [11, Proposition 2.9] contained a gap: the correct statement is Proposition 4.3 below.

1. RANKERS AND LOGICAL HIERARCHIES

Let A be a finite alphabet. We denote by A^* the set of all words over A (that is, of sequences of elements of A), and by A^+ the set of non-empty words. If u is a length n ($n > 0$) word over A , we say that an integer $1 \leq i \leq n$ is an *a-position* of u if the i -th letter of u , written $u[i]$, is an a . If $1 \leq i \leq j \leq n$, we let $u[i; j]$ be the factor $u[i] \cdots u[j]$ of u .

FO denotes the set of first-order formulas using the unary predicates \mathbf{a} ($a \in A$) and the binary predicate $<$, and FO^2 denotes the fragment of FO consisting of formulas which use at most two variable symbols.

If u is a length n ($n > 0$) word over A , we identify the word u with the logical structure $(\{1, \dots, n\}, (\mathbf{a})_{a \in A})$, where \mathbf{a} denotes the set of a -positions in u . Formulas from FO are naturally interpreted over this structure, and we denote by $L(\varphi)$ the language *defined by* the formula $\varphi \in \text{FO}$, that is, the set of all words which satisfy φ .

1.1. Quantifier-alternation within FO^2 . We now concentrate on FO^2 -formulas and we define two important parameters concerning such formulas. To simplify matters, we consider only formulas where negation is used only on atomic formulas so that, in particular, no quantifier is negated. This is naturally possible up to logical equivalence. Now, with each formula $\varphi \in \text{FO}^2$, we associate in the natural way a parsing tree: each occurrence of a quantification, $\exists x$ or $\forall x$, yields a unary node, each occurrence of \vee or \wedge yields a binary node, and the leaves are labeled with atomic or negated atomic formulas. The *quantifier depth* of φ is the maximum number of quantifiers along a path in its parsing tree.

With each path from root to leaf in this parsing tree, we also associate its quantifier label, which is the sequence of quantifier node labels (\exists or \forall) encountered along this path. A *block* in this quantifier label is a maximal factor consisting only of \exists or only of \forall , and we define the *number of blocks*

of φ to be the maximum number of blocks in the quantifier label of a path in its parsing tree. Naturally, the quantifier depth of φ is at least equal to its number of blocks.

We let $\text{FO}_{m,n}^2$ ($n \geq m$) denote the set of first-order formulas with quantifier depth at most n and with at most m blocks and let FO_m^2 denote the union of the $\text{FO}_{m,n}^2$ for all n . We also denote by \mathcal{FO}^2 (\mathcal{FO}_m^2 , $\mathcal{FO}_{m,n}^2$) the class of FO^2 (FO_m^2 , $\text{FO}_{m,n}^2$)-definable languages.

1.2. Rankers. A *ranker* is a non-empty word on the alphabet $\{\mathsf{X}_a, \mathsf{Y}_a \mid a \in A\}$. Rankers define positions in words: given a word $u \in A^+$ and a letter $a \in A$, we denote by $\mathsf{X}_a(u)$ (resp. $\mathsf{Y}_a(u)$) the least (resp. greatest) integer $1 \leq i \leq |u|$ such that $u[i] = a$. If a does not occur in u , we say that $\mathsf{Y}_a(u)$ and $\mathsf{X}_a(u)$ are not defined. If in addition q is an integer such that $1 \leq q \leq |u|$, we let

$$\begin{aligned}\mathsf{X}_a(u, q) &= \mathsf{X}_a(u[q+1; |u|]) \\ \mathsf{Y}_a(u, q) &= \mathsf{Y}_a(u[1; q-1]).\end{aligned}$$

These definitions are extended to all rankers: if r' is a ranker, $Z \in \{\mathsf{X}_a, \mathsf{Y}_a \mid a \in A\}$ and $r = r'Z$, we let

$$r(u, q) = Z(u, r'(u, q))$$

if $r'(u, q)$ and $Z(u, r'(u, q))$ are defined, and we say that $r(u, q)$ is undefined otherwise. (In particular: rankers are processed from left to right.)

Finally, if r starts with an X - (resp. Y -) letter, we say that r defines the position $r(u) = r(u, 0)$ (resp. $r(u) = r(u, |u| + 1)$), or that it is undefined on u if this position does not exist.

Remark 1.1. Rankers were first introduced, under the name of *turtle programs*, by Schwentick, Thérien and Vollmer [20], as sequences of instructions: go to the next a to the right, go to the next b to the left, etc. These authors write (\rightarrow, a) and (\leftarrow, a) instead of X_a and Y_a . Weis and Immerman [30] write \triangleright_a and \triangleleft_a instead, and they introduced the term *ranker*. We rather follow the notation in [6, 10, 4], where X and Y refer to the future and past operators of PTL. \square

Example 1.2. The ranker $\mathsf{X}_a \mathsf{Y}_b \mathsf{X}_c$ (go to the first a starting from the left, thence to the first b towards the left, thence to the first c towards the right) is defined on bac and bca , but not on abc or cba .

We will encounter $\mathsf{X}_a \mathsf{Y}_b \mathsf{X}_c$ again in Examples 3.3 and 4.2. \square

By $L(r)$ we denote the language of all words on which r is defined. We say that the words u and v *agree on a class* R of rankers if exactly the same rankers from R are defined on u and v . And we say that two rankers r and s *coincide* on a word u if they are both defined on u and $r(u) = s(u)$.

Example 1.3. If $r = X_{a_1} \cdots X_{a_k}$ (resp. $Y_{a_k} \cdots Y_{a_1}$), then $L(r)$ is the set of words that contain $a_1 \cdots a_k$ as a subword, $L(r) = A^* a_1 A^* \cdots a_k A^*$. \square

The *depth* of a ranker r is defined to be its length (as a word). A *block* in r is a maximal factor in $\{X_a \mid a \in A\}^+$ (an X-block) or in $\{Y_a \mid a \in A\}^+$ (a Y-block). If $n \geq m$, we denote by $R_{m,n}^X$ (resp. $R_{m,n}^Y$) the set of m -block, depth n rankers, starting with an X- (resp. Y-) block, and we let $R_{m,n} = R_{m,n}^X \cup R_{m,n}^Y$ and $\underline{R}_{m,n}^X = \bigcup_{m' \leq m, n' \leq n} R_{m',n'}^X \cup \bigcup_{m' < m, n' < n} R_{m',n'}^Y$. We define $\underline{R}_{m,n}^Y$ dually and we let $\underline{R}_m^X = \bigcup_{n \geq m} \underline{R}_{m,n}^X$, $\underline{R}_m^Y = \bigcup_{n \geq m} \underline{R}_{m,n}^Y$ and $\underline{R}_m = \underline{R}_m^X \cup \underline{R}_m^Y$.

Remark 1.4. Readers familiar with [30] will notice differences between our $\underline{R}_{m,n}^X$ and their analogous $R_{m>,n}^*$; introduced for technical reasons, it creates no difference between our $\underline{R}_{m,n}$ and their $R_{m,n}^*$, the classes which intervene in crucial Theorem 1.7 below. \square

1.3. Rankers and unary temporal logic. Let us depart for a moment from the consideration of FO²-formulas, to observe that rankers are naturally suited to describe the different levels of a natural class of temporal logic. The symbols X_a and Y_a ($a \in A$) can be seen as modal (temporal) operators, with the *future* and *past* semantics respectively. We denote the resulting temporal logic (known as *unary temporal logic*) by TL: its only atomic formula is \top , the other formulas are built using Boolean connectives and modal operators. Let $u \in A^+$ and let $0 \leq i \leq |u| + 1$. We say that \top holds at every position i , $(u, i) \models \top$; Boolean connectives are interpreted as usual; and $(u, i) \models X_a \varphi$ (resp. $Y_a \varphi$) if and only if $(u, X_a(u, i)) \models \varphi$ (resp. $(u, Y_a(u, i)) \models \varphi$). In particular, $(u, i) \not\models X_a \varphi$ (resp. $Y_a \varphi$) if there is no a -position greater (resp. less) than i . We also say that $u \models X_a \varphi$ (resp. $Y_a \varphi$) if $(u, 0) \models X_a \varphi$ (resp. $(u, 1 + |u|) \models Y_a \varphi$).

TL is a fragment of *propositional temporal logic* PTL; the latter is expressively equivalent to FO and TL is expressively equivalent to FO², see the survey [10].

As in the case of FO²-formulas, one may consider the parsing tree of a TL-formula and define inductively its depth and number of alternations (between past and future operators). If $n \geq m$, the fragment $\text{TL}_{m,n}^X$ (resp. $\text{TL}_{m,n}^Y$) consists of the TL-formulas with depth n and with m alternated blocks, in which every branch (of the parsing tree) with exactly m alternations starts with future (resp. past) operators. Branches with less alternations may start with past (resp. future) operators. The fragments $\text{TL}_{m,n}$, $\underline{\text{TL}}_{m,n}^X$, $\underline{\text{TL}}_{m,n}^Y$, $\underline{\text{TL}}_m^X$, $\underline{\text{TL}}_m^Y$ and $\underline{\text{TL}}_m$ are defined according to the same pattern as in the definition of $R_{m,n}$, $\underline{R}_{m,n}^X$, $\underline{R}_{m,n}^Y$, \underline{R}_m^X , \underline{R}_m^Y and \underline{R}_m . We also denote by $\mathcal{TL}_{m,n}^X$ (\mathcal{TL}_m^X , $\underline{\mathcal{TL}}_m^X$, etc) the class of $\text{TL}_{m,n}^X$ (TL_m^X , $\underline{\mathcal{TL}}_m^X$, etc)-definable languages.

Proposition 1.5. *Let $1 \leq m \leq n$. Two words satisfy the same $\text{TL}_{m,n}^X$ formulas if and only if they agree on rankers from $R_{m,n}^X$. A language is in $\mathcal{TL}_{m,n}^X$ if and only if it is a Boolean combination of languages of the form $L(r)$, $r \in R_{m,n}^X$.*

Similar statements hold for $\text{TL}_{m,n}^Y$, $\text{TL}_{m,n}$, $\underline{\text{TL}}_{m,n}^X$, $\underline{\text{TL}}_{m,n}^Y$, $\underline{\text{TL}}_m^X$, $\underline{\text{TL}}_m^Y$ and $\underline{\text{TL}}_m$, relative to the corresponding classes of rankers.

Sketch of Proof. Since every ranker can be viewed as a TL -formula, it is easily verified that if u and v satisfy the same $\text{TL}_{m,n}^X$ -formulas, then they agree on rankers from $R_{m,n}^X$. To prove the converse, it suffices to show that a $\text{TL}_{m,n}^X$ -formula is equivalent to a Boolean combination of formulas that are expressed by a single ranker. That is: we only need to show that modalities can be brought outside the formula. This follows from the following elementary logical equivalences:

$$X_a(\varphi \wedge \psi) \equiv X_a\varphi \wedge X_a\psi \quad \text{and} \quad X_a(\neg\varphi) \equiv X_a \wedge \neg X_a\varphi.$$

□

Remark 1.6. Together with Example 1.3, this proposition confirms the elementary observation that a language is $\underline{\text{TL}}_1$ (resp. $\underline{\text{TL}}_1^X$, $\underline{\text{TL}}_1^Y$) definable if and only if it is a Boolean combination of languages of the form $A^*a_1A^* \cdots a_kA^*$ ($a_i \in A$), that is, if and only if L is piecewise testable (see [15]). It is well-known that this class of languages is a variety, whose corresponding pseudovarieties of monoids is the class \mathbf{J} of \mathcal{J} -trivial monoids (see Section 2 below), and hence $\underline{\text{TL}}_1$ -definability is decidable. □

1.4. **Rankers and FO^2 .** The connection established by Weis and Immerman [30, Theorem 4.5] between rankers and formulas in $\text{FO}_{m,n}^2$, Theorem 1.7 below, is much deeper. If x, y are integers, we let $\text{ord}(x, y)$, the *order type* of x and y , be one of the symbols $<$, $>$ or $=$, depending on whether $x < y$, $x > y$ or $x = y$.

Theorem 1.7 (Weis and Immerman [30]). *Let $u, v \in A^*$ and let $1 \leq m \leq n$. Then u and v satisfy the same formulas in $\text{FO}_{m,n}^2$ if and only if*

- (WI 1) *u and v agree on rankers from $\underline{R}_{m,n}$,*
- (WI 2) *if the rankers $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m-1,n-1}$ are defined on u and v , then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.*
- (WI 3) *if $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m,n-1}$ are defined on u and v and end with different direction letters, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.*

Corollary 1.8. *For each $n \geq m \geq 1$, $\underline{\mathcal{TL}}_{m,n} \subseteq \mathcal{FO}_{m,n}^2$ and $\underline{\mathcal{TL}}_m \subseteq \mathcal{FO}_m^2$.*

Proof. Let L be a $\underline{\text{TL}}_{m,n}$ -definable language. For each $u \in L$, let φ_u be the conjunction of the $\text{FO}_{m,n}^2$ -sentences satisfied by u and let φ be the disjunction of the formulas φ_u ($u \in L$). Since $\text{FO}_{m,n}^2$ is finite (up to logical equivalence), the conjunctions and disjunctions in the definition of φ are all finite.

A word v satisfies φ if and only if it satisfies φ_u for some word $u \in L$. Then v satisfies the same FO _{m,n} ²-sentences as u and, by comparing the statements in Proposition 1.5 and Theorem 1.7, we see that u and v satisfy the same $\underline{\text{TL}}_{m,n}$ -formulas. Since L is defined by such a formula, it follows that $v \in L$. Conversely, every word $v \in L$ satisfies φ since it satisfies φ_v , which is logically equivalent to a term in the disjunction defining φ . Thus L is defined by the FO _{m,n} ²-sentence φ . \square

1.5. FO _{m} ² and $\underline{\text{TL}}_m$ are varieties of languages. We use the ranker characterizations in Sections 1.3 and 1.4 to prove that $\underline{\text{TL}}_m$ - and FO _{m} ²-definable languages form varieties of languages. We refer the reader to [15, 23] and to Section 2 below for background and discussion on varieties of languages.

Lemma 1.9. *Let $n \geq m \geq 1$. The property $\rho_{m,n}$ (resp. $\lambda_{m,n}$), for two words $u, v \in A^*$ to agree on rankers in $\underline{R}_{m,n}^X$ (resp. $\underline{R}_{m,n}^Y$), is a finite index congruence on A^* .*

Proof. It is immediate that $\rho_{m,n}$ and $\lambda_{m,n}$ are equivalence relations, and they have finite index on A^* since $\underline{R}_{m,n}^X$ and $\underline{R}_{m,n}^Y$ are finite. We now verify that $\rho_{m,n}$ and $\lambda_{m,n}$ are stable by product.

We proceed by induction on m . The result is elementary if $m = 1$, see Example 1.3. Let us assume that $m \geq 2$, $u \rho_{m,n} v$ and $u' \rho_{m,n} v'$, and $r \in \underline{R}_{m,n}^X$ is defined on uu' . We want to show that r is defined on vv' .

If r starts with a Y-letter, then $r \in \underline{R}_{m-1,n-1}^Y$. Since $\underline{R}_{m-1,n-1}^Y \subseteq \underline{R}_{m,n}^X$, we have $u \lambda_{m-1,n-1} v$ and $u' \lambda_{m-1,n-1} v'$. Moreover, $\lambda_{m-1,n-1}$ is a congruence by induction. So $uu' \lambda_{m-1,n-1} vv'$ and hence, r is defined on vv' .

If r starts with an X-letter, the sequence of positions it defines in uu' visits sometimes u and sometimes u' . More precisely let r_1 be the longest prefix of r that is defined on u (r_1 may be empty): then $r = r_1 r'_1$ and if r'_1 is non-empty, then r'_1 starts with an X-letter. Let now r_2 be the longest prefix of r'_1 which is defined on u' . Then r_2 is not empty and if $r_2 \neq r'_1$, then $r = r_1 r_2 r'_2$ where r'_2 starts with a Y-letter. Iterating this process, we find a factorization of r as $r = r_1 r_2 \cdots r_k$, where each r_i is the longest prefix of $r_i r_{i+1} \cdots r_k$ which is defined on u if i is odd, on u' if i is even: then r_1 may be empty, r_i ($i \geq 1$) starts with an X-letter if i is even, with a Y-letter if i is odd; and the position in uu' defined by $r_1 \cdots r_i$ is the position defined by r_i in u if i is odd, in u' if i is even.

Since the r_i are all in $\underline{R}_{m,n}^X$ and since $u \rho_{m,n} v$ and $u' \rho_{m,n} v'$, the factorization $r = r_1 r_2 \cdots r_k$ has exactly the same properties with respect to v and v' instead of u and u' . In particular, r is defined on vv' .

The proof that $u \lambda_{m,n} v$ and $u' \lambda_{m,n} v'$ imply $uu' \lambda_{m,n} vv'$ is similar. \square

Lemma 1.10. *Let $n \geq m \geq 1$. The property for two words $u, v \in A^*$ to satisfy Properties (WI 1), (WI 2) and (WI 3) of Theorem 1.7 (with respect to the pair (m, n)) is a finite index congruence on A^* .*

Proof. As in Lemma 1.9, the only problem is to verify that if Properties **(WI 1)**, **(WI 2)** and **(WI 3)** hold for u and v , and also for u' and v' , then they hold for uu' and vv' . For Property **(WI 1)**, this is proved by Lemma 1.9.

Let us now consider Property **(WI 2)**, that is, let $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m-1,n-1}$ be defined on uu' and vv' .

Case 1. r and r' start with X-letters. We factor r and r' as in the proof of Lemma 1.9, say, $r = r_1 \cdots r_k$ and $r' = r'_1 \cdots r'_\ell$. As discussed in that proof, the position in uu' defined by r is the position defined by r_k in u if k is odd, in u' if k is even. Similarly, the position in uu' defined by r' is the position defined by r'_ℓ in u if ℓ is odd, in u' if ℓ is even. In view of Property **(WI 1)**, the factorizations $r = r_1 \cdots r_k$ and $r' = r'_1 \cdots r'_\ell$ have the same properties with respect to v and v' .

If k is odd and ℓ is even, then r determines a position in u (resp. v) and r' determines a position in u' (resp. v'), so both $\text{ord}(r(uu'), r'(uu'))$ and $\text{ord}(r(vv'), r'(vv'))$ are $<$. The situation is equally simple if k is even and ℓ is odd.

If both k and ℓ are odd, then $r(uu') = r_k(u)$, $r'(uu') = r_\ell(u)$, $r(vv') = r_k(v)$ and $r'(vv') = r_\ell(v)$, so $\text{ord}(r(uu'), r'(uu')) = \text{ord}(r_k(u), r'_\ell(u))$ and $\text{ord}(r(vv'), r'(vv')) = \text{ord}(r_k(v), r'_\ell(v))$. Now $r_k \in \underline{R}_{m,n}$ and $r'_\ell \in \underline{R}_{m-1,n-1}$ so Property **(WI 2)** shows that $\text{ord}(r(uu'), r'(uu')) = \text{ord}(r(vv'), r'(vv'))$.

The proof is similar if k and ℓ are both even.

Case 2. r starts with an X-letter and r' starts with a Y-letter. We consider again factorizations $r = r_1 \cdots r_k$ and $r' = r'_1 \cdots r'_\ell$ as in the proof of Lemma 1.9. The proof is the same as in Case 1, except that the roles of evenness and oddness are exchanged.

The remaining cases, where r starts with a Y-letter, are right-left duals of the previous cases, and are handled similarly.

This concludes the proof that Property **(WI 2)** holds for uu' and vv' . The handling of Property **(WI 3)** is entirely similar. \square

Corollary 1.11. *Let $n \geq m \geq 1$. The classes of $\underline{\text{TL}}_{m,n}^X$, $\underline{\text{TL}}_m^X$, $\underline{\text{TL}}_{m,n}^Y$, $\underline{\text{TL}}_m^Y$, $\underline{\text{TL}}_m$, $\text{FO}_{m,n}^2$ and FO_m^2 -definable languages are varieties of languages.*

Proof. Proposition 1.5 states that a language in A^* is $\underline{\text{TL}}_{m,n}^X$ -definable if and only if it is a Boolean combination of $\rho_{m,n}$ -classes in A^* . By Lemma 1.9, this is equivalent to being recognized by the finite monoid $A^*/\rho_{m,n}$. In particular, a language is $\underline{\text{TL}}_{m,n}^X$ -definable if and only if it is recognized by a monoid in the pseudovariety generated by the monoids of the form $A^*/\rho_{m,n}$ (A , a finite alphabet). And a language is $\underline{\text{TL}}_m^X$ -definable if and only if it is recognized by a monoid in the pseudovariety generated by the monoids of the form $A^*/\rho_{m,n}$ (A , a finite alphabet and $n \geq m$).

The same reasoning applies to the classes of $\underline{\text{TL}}_{m,n}^{\text{Y}}$, $\underline{\text{TL}}_{m^-}^{\text{Y}}$, $\underline{\text{TL}}_{m^-}$, $\text{FO}_{m,n}^2$ - and FO_m^2 -definable languages, using Theorem 1.7 and Lemma 1.10 in the last two cases. \square

This result shows that, for a given regular language L , $\underline{\text{TL}}_m^{\text{X}}$ (resp. $\underline{\text{TL}}_m$, FO_m^2 , etc) definability is characterized algebraically, that is, it depends only on the syntactic monoid of L . This justifies using the algebraic path to tackle decidability of these definability problems. Eilenberg's theory of varieties provides the mathematical framework. A limited working summary of this theory (for the purpose of this paper) is given in Section 2 below.

2. ON VARIETIES AND PSEUDOVARIETIES

We summarize in this section the information on monoid and variety theory that will be relevant for our purpose. For more detailed information and proofs, we refer the reader to [15, 2, 24, 25, 23], among other sources.

A *semigroup* is a set equipped with a binary associative operation. A *monoid* is a semigroup which contains a unit element. The set A^* of all words on alphabet A , equipped with the concatenation product, is the *free monoid* on A : it has the specific property that, if $\varphi: A \rightarrow M$ is a map into a monoid, then there exists a unique monoid morphism $\psi: A^* \rightarrow M$ which extends φ . Apart from free monoids, the semigroups and monoids which we will consider in this paper are finite.

If A is a finite alphabet and M is a finite monoid, we say that a language $L \subseteq A^*$ is *recognized* by M if there exists a morphism $\varphi: A^* \rightarrow M$ such that $L = \varphi^{-1}(\varphi(L))$.

Example 2.1. If $u \in A^*$ and $B \subseteq A$, let

$$\begin{aligned} \text{alph}(u) &= \{a \in A \mid u = vaw \text{ for some } v, w \in A^*\}, \\ [B] &= \{u \in A^* \mid \text{alph}(u) = B\} \end{aligned}$$

Let φ be the following morphism from A^* into the direct product of $|A|$ copies of the 2-element monoid $\{0, 1\}$ (multiplicative): for each letter $a \in A$, $\varphi(a)$ is the A -tuple in which every component is 0, except for the a -component. It is elementary to show that $[B] = \varphi^{-1}(\varphi([B]))$ and hence, $[B]$ is accepted by a monoid that is *idempotent* (every element is equal to its own square) and commutative. Conversely, one can show that every language recognized by an idempotent and commutative monoid is a Boolean combination of languages of the form $[B]$ ($B \subseteq A$). \square

A *pseudovariety* of monoids (resp. semigroups) is a class of finite monoids (resp. semigroups) which is closed under taking direct products, homomorphic images and submonoids (resp. subsemigroups). A class of languages \mathcal{V} is a collection $\mathcal{V} = (\mathcal{V}(A))_A$, indexed by all finite alphabets A , such that $\mathcal{V}(A)$ is a set of languages in A^* . If \mathbf{V} is a pseudovariety of monoids, we let $\mathcal{V}(A)$ be the set of all languages of A^* which are recognized by a monoid in

V. The class \mathcal{V} has important closure properties: each $\mathcal{V}(A)$ is closed under Boolean operations and under taking residuals (if $L \in \mathcal{V}(A)$ and $u \in A^*$, then Lu^{-1} and $u^{-1}L$ are in $\mathcal{V}(A)$); and if $\varphi: A^* \rightarrow B^*$ is a morphism and $L \in \mathcal{V}(B)$, then $\varphi^{-1}(L) \in \mathcal{V}(A)$. Classes of recognizable languages with these properties are called *varieties* of languages, and Eilenberg's theorem (see [15]) states that the correspondence $\mathbf{V} \mapsto \mathcal{V}$, from pseudovarieties of monoids to varieties of languages, is one-to-one and onto. Moreover, the decidability of membership in the pseudovariety \mathbf{V} , implies the decidability of the variety \mathcal{V} : indeed, a language is in \mathcal{V} if and only if its (effectively computable) syntactic monoid is in \mathbf{V} .

For every finite semigroup S , there exists an integer, usually denoted ω , such that every element of the form s^ω in S is idempotent. The *Green relations* are another important concept to describe semigroups and monoids: if S is a semigroup and $s, t \in S$, we say that $s \leq_{\mathcal{J}} t$ (resp. $s \leq_{\mathcal{R}} t$, $s \leq_{\mathcal{L}} t$) if $s = utv$ (resp. $s = tv$, $s = ut$) for some $u, v \in S \cup \{1\}$. We also say that $s \mathcal{J} t$ is $s \leq_{\mathcal{J}} t$ and $t \leq_{\mathcal{J}} s$. The relations \mathcal{R} and \mathcal{L} are defined similarly.

Pseudovarieties that will be important in this paper are the following.

- **J₁**, the pseudovariety of idempotent and commutative monoids; as discussed in Example 2.1, the corresponding variety of languages consists of the Boolean combinations of languages of the form $[B]$.

- **R**, **L** and **J**, the pseudovarieties of \mathcal{R} -, \mathcal{L} - and \mathcal{J} -trivial monoids; a monoid is, say, **R**-trivial if each of its \mathcal{R} -classes is a singleton. The variety of languages corresponding to **J**, called the piecewise testable languages, was described by Simon (see [15]): a language in A^* is piecewise testable if and only if it is a Boolean combination of languages of the form $A^*a_1A^* \cdots a_rA^*$, with $a_1, \dots, a_r \in A$. We encountered these languages in Example 1.3.

- **A**, the variety of *aperiodic* monoids, i.e., monoids in which $x^\omega = x^{\omega+1}$ for each x . Celebrated theorems of Schützenberger, McNaughton and Papert and Kamp show that the corresponding variety of languages consists of the star-free languages, the languages that are definable in FO, and the languages definable in propositional temporal logic, see for instance [15, 25, 5, 6, 22, 23].

- **DA** is the pseudovariety of all monoids in which $(xy)^\omega x(xy)^\omega = (xy)^\omega$ for all x, y . This pseudovariety has many characterizations in combinatorial, algebraic and logical terms. Of particular interest to us are the fact that the corresponding variety of languages consists of the languages that are definable in FO², and equivalently, of the languages that are defined in unary temporal logic, see [24, 25, 6, 28] among others.

- **K** (resp. **D**, **LI**) is the pseudovariety of semigroups in which $x^\omega y = x^\omega$ (resp. $yx^\omega = x^\omega$, $x^\omega yx^\omega = x^\omega$) for all x, y .

Finally we describe the operations of the *Mal'cev product on pseudovarieties*. Let \mathbf{V} be a pseudovariety of monoids, \mathbf{W} a pseudovariety of semigroups and M a finite monoid. We say that $M \in \mathbf{W} \circledast \mathbf{V}$ if there exists a finite monoid T and onto morphisms $\alpha: T \rightarrow M$ and $\beta: T \rightarrow N$ such that $N \in \mathbf{V}$ and, for each idempotent e of N , $\beta^{-1}(e) \in \mathbf{W}$. Then $\mathbf{W} \circledast \mathbf{V}$

is a pseudovariety of monoids, see [15, 2, 17]. The following equalities are well-known [15]:

$$\mathbf{K} \textcircled{m} \mathbf{J}_1 = \mathbf{K} \textcircled{m} \mathbf{J} = \mathbf{R}, \quad \mathbf{D} \textcircled{m} \mathbf{J}_1 = \mathbf{D} \textcircled{m} \mathbf{J} = \mathbf{L}, \quad \mathbf{LI} \textcircled{m} \mathbf{J}_1 = \mathbf{LI} \textcircled{m} \mathbf{J} = \mathbf{DA}.$$

We denote by $\underline{\mathbf{TL}}_{m,n}^{\mathbf{X}}$, $\underline{\mathbf{TL}}_{m,n}^{\mathbf{Y}}$, $\underline{\mathbf{TL}}_m^{\mathbf{Y}}$, $\underline{\mathbf{TL}}_m^{\mathbf{X}}$, $\underline{\mathbf{TL}}_{m,n}$, $\underline{\mathbf{TL}}_m$, $\mathbf{FO}_{m,n}^2$ and \mathbf{FO}_m^2 the pseudovarieties corresponding to the language varieties discovered in Corollary 1.11.

3. CONDENSED RANKERS

Our main tool to approach the decidability of FO_m²-definability lies in the notion of condensed rankers, a variant of rankers which was introduced implicitly by Weis and Immerman to prove Proposition 6.4 below (see [30, Theorem 4.7]). Recall that a ranker can be seen as a sequence of directional instructions (see Example 1.2). We say that a ranker r is *condensed on u* if it is defined on u , and if the sequence of positions visited *zooms in* on $r(u)$, never crossing over a position already visited. Formally, $r = Z_1 \cdots Z_n$ is condensed on u if there exists a chain of open intervals

$$(0, |u| + 1) = (i_0, j_0) \supset (i_1, j_1) \supset \cdots \supset (i_{n-1}, j_{n-1}) \ni r(u)$$

such that for all $1 \leq \ell \leq n - 1$ the following properties are satisfied:

- If $Z_\ell Z_{\ell+1} = \mathbf{X}_a \mathbf{X}_b$ then $(i_\ell, j_\ell) = (\mathbf{X}_a(u, i_{\ell-1}), j_{\ell-1})$.
- If $Z_\ell Z_{\ell+1} = \mathbf{Y}_a \mathbf{Y}_b$ then $(i_\ell, j_\ell) = (i_{\ell-1}, \mathbf{Y}_a(u, j_{\ell-1}))$.
- If $Z_\ell Z_{\ell+1} = \mathbf{X}_a \mathbf{Y}_b$ then $(i_\ell, j_\ell) = (i_{\ell-1}, \mathbf{X}_a(u, i_{\ell-1}))$.
- If $Z_\ell Z_{\ell+1} = \mathbf{Y}_a \mathbf{X}_b$ then $(i_\ell, j_\ell) = (\mathbf{Y}_a(u, j_{\ell-1}), j_{\ell-1})$.

Remark 3.1. The i_ℓ and j_ℓ are either 0 or $1 + |u|$, or positions of the form $r'(u)$ for some prefix r' of r . More precisely, if r_ℓ is the depth ℓ prefix of r ($\ell < n$), then $r_\ell(u) = i_\ell$ if $Z_{\ell+1}$ is of the form \mathbf{X}_a , and $r_\ell(u) = j_\ell$ if $Z_{\ell+1}$ is of the form \mathbf{Y}_a . \square

Remark 3.2. If $r = r_1 r_2$ is condensed on u , then $r(u) > r_1(u)$ if r_2 starts with an X-letter, and $r(u) < r_1(u)$ if r_2 starts with a Y-letter. \square

Example 3.3. The ranker $\mathbf{X}_a \mathbf{Y}_b \mathbf{X}_c$ is defined on the words bac and bca , but it is condensed only on bca .

Rankers in $\underline{\mathbf{R}}_{1,n}$ and rankers of the form $\mathbf{X}_a \mathbf{Y}_{b_1} \cdots \mathbf{Y}_{b_k}$ or $\mathbf{Y}_a \mathbf{X}_{b_1} \cdots \mathbf{X}_{b_k}$ are condensed on all words on which they are defined. \square

Condensed rankers form a natural notion, which is equally well-suited to the task of describing FO_m²-definability (see Theorem 3.15 below). With respect to \mathbf{TL} , for which Proposition 1.5 shows a perfect match with the notion of rankers, they can be interpreted as adding a strong notion of unambiguity, see Section 6.3 below and the work of Lodaya, Pandya and Shah [13].

Let us say that two words u and v *agree on condensed rankers from a set R* of rankers, if the same rankers in R are condensed on u and v . We

write $u \triangleright_{m,n} v$ (resp. $u \triangleleft_{m,n} v$) if u and v agree on condensed rankers in $\underline{R}_{m,n}^X$ (resp. $\underline{R}_{m,n}^Y$). If r is a ranker, let $L_c(r)$ be the language of all words on which r is condensed.

3.1. Technical properties of condensed rankers. Lemmas 3.4 and 3.5 admit an elementary verification. A factorization $u = u_- au_+$ of a word $u \in A^*$ is called the *a-left factorization* of u if $a \notin \text{alph}(u_-)$. Symmetrically, $u = u_- au_+$ is the *a-right factorization* of u if $a \notin \text{alph}(u_+)$. Thus, the *a-left* (resp. *a-right*) factorization of a identifies the first occurrence of a when reading u from the left (resp. the right).

Lemma 3.4. *Let s be a ranker, $a \in A$ and $r = X_a s$. Let also $u \in A^+$ and let $u = u_- au_+$ be its *a-left factorization*. Then r is condensed on u if and only if*

- s is condensed on u_+ if s starts with an X -block;
- s is condensed on u_- if s starts with a Y -block.

A dual statement holds if r is of the form $r = Y_a s$, with respect to the *a-right factorization* of u .

Lemma 3.5. *Let r be a ranker and $a \in A$. Let also $u \in A^+$ and let $u = u_- au_+$ be its *a-left factorization*.*

If r starts with an X -letter, then

- r is defined on u_- if and only if r is defined on u , r does not contain X_a or Y_a and, for every prefix p of r ending with an X -letter, pY_a is not defined on u .
- r is condensed on u_-
 - if and only if r is defined on u_- and condensed on u ,
 - if and only if r is condensed on u , r does not contain X_a or Y_a and, if p is the maximal prefix of r consisting only of X -letters, then pY_a is not defined on u ,
 - if and only if r is condensed on u , r does not contain X_a or Y_a and, if $p = X_{b_1} \cdots X_{b_k}$ ($k \geq 1$) is the initial X -block of r , then $X_a Y_{b_k} \cdots Y_{b_1}$ is defined on u .
- r is defined on u_+ if and only if $X_a r$ is defined on u and, for every prefix p of r ending with a Y -letter, $X_a p Y_a$ is defined on u .
- r is condensed on u_+ if and only if $X_a r$ is condensed on u .

If r starts with a Y -letter, then

- r is defined on u_- if and only if $X_a r$ is defined on u , r does not contain X_a or Y_a and, for every prefix p of r ending with an X -letter, $X_a p Y_a$ is not defined on u .
- r is condensed on u_- if and only if $X_a r$ is condensed on u .
- r is defined on u_+ if and only if r is defined on u and, for every prefix p of r ending with a Y -letter, pY_a is defined on u .
- r is condensed on u_+
 - if and only if r is defined on u_+ and condensed on u ,

- if and only if r is condensed on u and, if $p = Y_{b_1} \cdots Y_{b_k}$ ($k \geq 1$) is the initial Y -block of r , then pY_a is defined on u .

We also note the following, very useful characterization of the relations $\triangleright_{m,n}$ and $\triangleleft_{m,n}$.

Proposition 3.6. *The families of relations $\triangleright_{m,n}$ and $\triangleleft_{m,n}$ ($n \geq m \geq 1$) are uniquely determined by the following properties.*

- (1) $u \triangleright_{1,n} v$ if and only if $u \triangleleft_{1,n} v$, if and only if u and v have the same subwords of length at most n .
- (2) If $m \geq 2$, then $u \triangleright_{m,n} v$ if and only if $\mathbf{alph}(u) = \mathbf{alph}(v)$, $u \triangleleft_{m-1,n-1} v$ and for each letter $a \in \mathbf{alph}(u)$, the a -left factorizations $u = u_- au_+$ and $v = v_- av_+$ satisfy $u_- \triangleleft_{m-1,n-1} v_-$ and $u_+ \triangleright_{m,n-1} v_+$ ($u_+ \triangleright_{m-1,n-1} v_+$ if $n = m$).
- (3) If $m \geq 2$, then $u \triangleleft_{m,n} v$ if and only if $\mathbf{alph}(u) = \mathbf{alph}(v)$, $u \triangleright_{m-1,n-1} v$ and for each letter $a \in \mathbf{alph}(u)$, the a -right factorizations $u = u_- au_+$ and $v = v_- av_+$ satisfy $u_+ \triangleright_{m-1,n-1} v_+$ and $u_- \triangleleft_{m,n-1} v_-$ ($u_- \triangleleft_{m-1,n-1} v_-$ if $n = m$).

Proof. Statement (1) follows directly from Examples 1.3 and 3.3. Let us now assume that $m \geq 2$.

Suppose that $\mathbf{alph}(u) = \mathbf{alph}(v)$, $u \triangleleft_{m-1,n-1} v$ and for each $a \in \mathbf{alph}(u)$, the a -left factorizations $u = u_- au_+$ and $v = v_- av_+$ satisfy $u_- \triangleleft_{m-1,n-1} v_-$ and $u_+ \triangleright_{m,n-1} v_+$ if $n > m$ ($u_+ \triangleright_{m-1,n-1} v_+$ if $n = m$). Let $r \in \underline{R}_{m,n}^X$ be condensed on u . If r starts with a Y -letter, then $r \in \underline{R}_{m-1,n-1}^Y$, and hence r is condensed on v since $u \triangleleft_{m-1,n-1} v$. If instead r starts with an X -letter, say $r = X_a s$, we consider the a -left factorizations of u and v . If s starts with a Y -letter, then $s \in \underline{R}_{m-1,n-1}^Y$, s is condensed on u_- (Lemma 3.4) and hence s is condensed on v_- since $u_- \triangleleft_{m-1,n-1} v_-$, from which it follows again that r is condensed on v . Finally, if s starts with an X -letter, then s is condensed on u_+ by Lemma 3.4. Moreover, $s \in \underline{R}_{m,n-1}^X$ if $n > m$. If $n = m$, we have in fact $r \in \underline{R}_{m-1,n}^X$ (since r starts with two X -letters) and hence $s \in \underline{R}_{m-1,n-1}^X$. Since $u_+ \triangleright_{m,n-1} v_+$ if $n > m$ and $u_+ \triangleright_{m-1,n-1} v_+$ if $n = m$, it follows that s is condensed on v_+ , and hence r is condensed on v .

Conversely, let us assume that $u \triangleright_{m,n} v$, that is, u and v agree on condensed rankers in $\underline{R}_{m,n}^X$. Considering rankers in $\underline{R}_{1,1}^X \subseteq \underline{R}_{m,n}^X$ shows that $\mathbf{alph}(u) = \mathbf{alph}(v)$. Similarly, considering rankers in $\underline{R}_{m-1,n-1}^Y \subseteq \underline{R}_{m,n}^X$ shows that $u \triangleleft_{m-1,n-1} v$. Finally, let $a \in \mathbf{alph}(u)$ and let $u = u_- au_+$ and $v = v_- av_+$ be a -left factorizations.

Let $s \in \underline{R}_{m-1,n-1}^Y$ be condensed on u_- . Note that s contains neither X_a nor Y_a , since $a \notin \mathbf{alph}(u_-)$. If s starts with a Y -letter, then $r = X_a s$ is condensed on u (Lemma 3.4) and since $r \in \underline{R}_{m,n}^X$, r is condensed on v as well, which implies that s is condensed on v_- . If instead s starts with an X -letter, then s is condensed on u and hence on v . Moreover, if $p = X_{b_1} \cdots X_{b_k}$ is the maximal prefix of s consisting only of X -letters, then $X_a Y_{b_k} \cdots Y_{b_1} \in \underline{R}_{2,n}^X$ is

condensed on u (Lemma 3.5). Since $\underline{R}_{2,n}^X \subseteq \underline{R}_{m,n}^X$, it is condensed on v as well and hence, s is condensed on v_- .

Finally, assume that $s \in \underline{R}_{m,n-1}^X$ ($\underline{R}_{m-1,n-1}^X$ if $n = m$) is condensed on u_+ . The reasoning is similar: if s starts with an X-letter, then $X_a s \in \underline{R}_{m,n}^X$ is condensed on u . Therefore $X_a s$ is condensed on v and s is condensed on v_+ . If instead s starts with a Y-letter, then s is condensed on u and $s \in \underline{R}_{m-1,n-2}^Y$ ($\underline{R}_{m-2,n-2}^Y$ if $n = m$). In particular $s \in \underline{R}_{m,n}^X$ and hence, s is condensed on v as well. Moreover, if p is the initial Y-block of s , then pY_a is condensed on u . Note that $pY_a \in \underline{R}_{1,n-1}^Y \subseteq \underline{R}_{m,n}^X$, so pY_a is condensed on v and s is condensed on v_+ . \square

Lemma 3.7. *Let $n \geq m \geq 2$, $u, v \in A^*$, $a \in A$ and let $u = u_- a u_+$ and $v = v_- a v_+$ be a -left factorizations. If $u \triangleright_{m,n} v$, then $u_- \triangleright_{m,n-1} v_-$ ($u_- \triangleright_{m-1,n-1} v_-$ if $n = m$). And if $u \triangleleft_{m,n} v$, then $u_+ \triangleleft_{m,n-1} v_+$ ($u_+ \triangleleft_{m-1,n-1} v_+$ if $n = m$). Dual statements hold for the factors of the a -right factorizations of u and v if $u \triangleleft_{m,n} v$ or $u \triangleright_{m,n} v$.*

Proof. We give the proof if $n > m$; it is easily adapted to the case where $n = m$.

Assume that $u \triangleright_{m,n} v$ and $r \in \underline{R}_{m,n-1}^X$ is condensed on u_- . By Lemma 3.5, we have:

- If r starts with an X-letter, then r is condensed on u , r does not contain occurrences of X_a or Y_a , and if $p = X_{b_1} \cdots X_{b_k}$ is the initial X-block of r , then $q = X_a Y_{b_k} \cdots Y_{b_1}$ is condensed on u . Since $q \in \underline{R}_{2,k+1}^X$ and $k < n$, we have $q \in \underline{R}_{m,n}^X$ and hence r and q are condensed on v . Therefore r is condensed on v_- .

- If r starts with a Y-letter, then $X_a r$ is condensed on u . But $r \in \underline{R}_{m-1,n-2}^Y$, so $X_a r \in \underline{R}_{m,n}^X$ and hence $X_a r$ is condensed on v . It follows that r is condensed on v_- .

Assume now that $u \triangleleft_{m,n} v$ and $r \in \underline{R}_{m,n-1}^Y$ is condensed on u_+ . Then

- If r starts with an X-letter (which is possible only if $m \geq 2$), then $r \in \underline{R}_{m-1,n-2}^X$ and $X_a r$ is condensed on u . But $X_a r \in \underline{R}_{m-1,n-1}^X \subseteq \underline{R}_{m,n}^Y$, so $X_a r$ is condensed on v and r is condensed on v_+ .

- If instead r starts with a Y-letter, then r is condensed on u and if p is the initial Y-block of r , then pY_a is condensed on u . But $r, pY_a \in \underline{R}_{m,n}^Y$, so r and pY_a are condensed on v , and r is condensed on v_+ . \square

3.2. Condensed rankers determine a hierarchy of varieties. This section is the condensed ranker analogue of Section 1.5.

Lemma 3.8. *The relations $\triangleright_{m,n}$ and $\triangleleft_{m,n}$ are finite-index congruences.*

Proof. The relations $\triangleright_{m,n}$ and $\triangleleft_{m,n}$ are clearly equivalence relations, of finite index since $\underline{R}_{m,n}$ is finite. We now verify that if u and v on one hand, and u' and v' on the other hand, are $\triangleright_{m,n}$ equivalent, then so are uu' and vv' . (The proof regarding $\triangleleft_{m,n}$ is dual.)

The proof is by induction on $m + n$. The property of having the same subwords of length n is easily seen to be a congruence (and the proof of this fact can be found in [15] as it is related to Simon's theorem on piecewise testable languages). In view of Proposition 3.6 (1), this shows that $\triangleright_{1,n}$ and $\triangleleft_{1,n}$ are congruences.

Let us now assume that $n \geq m \geq 2$, $u \triangleright_{m,n} v$ and $u' \triangleright_{m,n} v'$. By Proposition 3.6 (2), we have $\text{alph}(u) = \text{alph}(v)$, $\text{alph}(u') = \text{alph}(v')$, $u \triangleleft_{m-1,n-1} v$ and $u' \triangleleft_{m-1,n-1} v'$. It follows that $\text{alph}(uu') = \text{alph}(vv')$ and $uu' \triangleleft_{m-1,n-1} vv'$ by induction.

Let now $a \in \text{alph}(uu')$. If $a \in \text{alph}(u) = \text{alph}(v)$ and $u = u_- a u_+$ and $v = v_- a v_+$ are a -left factorizations, then the a -left factorizations of uu' and vv' are $u_- a (u_+ u')$ and $v_- a (v_+ v')$. By Proposition 3.6 (2) we have $u_- \triangleleft_{m-1,n-1} v_-$ and $u_+ \triangleright_{m,n-1} v_+$ ($u_+ \triangleright_{m-1,n-1} v_+$ if $n = m$). By induction, the latter equivalence implies $u_+ u' \triangleright_{m,n-1} v_+ v'$ ($u_+ u' \triangleright_{m-1,n-1} v_+ v'$ if $n = m$).

If instead $a \notin \text{alph}(u) = \text{alph}(v)$ and $u' = u'_- a u'_+$ and $v' = v'_- a v'_+$ are a -left factorizations, then the a -left factorizations of uu' and vv' are $(u u'_-) a u'_+$ and $(v v'_-) a v'_+$. Again by Proposition 3.6 (2) we have $u'_- \triangleleft_{m-1,n-1} v'_-$ and $u'_+ \triangleright_{m,n-1} v'_+$ ($u'_+ \triangleright_{m-1,n-1} v'_+$ if $n = m$). By induction, the first equivalence implies $u u'_- \triangleleft_{m-1,n-1} v v'_-$.

It now follows (Proposition 3.6 (2) again) that $uu' \triangleright_{m,n} vv'$. \square

Now let \mathcal{R}_m (resp. \mathcal{L}_m) be the Boolean algebra generated by the languages of the form $L_c(r)$, $r \in \underline{R}_{m,n}^X$ (resp. $\underline{R}_{m,n}^Y$), $n \geq m$. Let also \mathbf{R}_m and \mathbf{L}_m be the pseudovarieties of monoids generated respectively by the $A^*/\triangleright_{m,n}$ and by the $A^*/\triangleleft_{m,n}$, $n \geq m$.

Corollary 3.9. *For each $m \geq 1$, \mathcal{R}_m and \mathcal{L}_m are varieties of languages and the corresponding pseudovarieties of monoids are \mathbf{R}_m and \mathbf{L}_m .*

Proof. By definition, a $\triangleright_{m,n}$ -class is a Boolean combination of languages of the form $L_c(r)$, $r \in \underline{R}_{m,n}^X$. Therefore every language recognized by $A^*/\triangleright_{m,n}$ is in \mathcal{R}_m , and hence so is every language recognized by a monoid in \mathbf{R}_m .

Conversely, every $L_c(r)$ ($r \in \underline{R}_{m,n}^X$) is a union of $\triangleright_{m,n}$ -classes, and hence it is recognized by $A^*/\triangleright_{m,n}$. Therefore every language in \mathcal{R}_m is recognized by a monoid in \mathbf{R}_m (and indeed, by $A^*/\triangleright_{m,n}$ for n large enough). \square

Example 3.10. It follows from Proposition 3.6 (i) that $\mathcal{R}_1 = \mathcal{L}_1$ is the variety of piecewise testable languages, and $\mathbf{R}_1 = \mathbf{L}_1 = \mathbf{J}$, the pseudovariety of \mathcal{J} -trivial monoids. \square

Remark 3.11. It follows from the definition of the \mathcal{R}_m and \mathcal{L}_m that for each m , \mathcal{R}_m and \mathcal{L}_m are contained in both \mathcal{R}_{m+1} and \mathcal{L}_{m+1} . \square

3.3. Condensed rankers, rankers and FO². We now show that, in the characterization of $\mathcal{FO}_{m,n}^2$ in Theorem 1.7, condensed rankers can be used just as well. This is done in Theorem 3.15. The first step is to relate

agreement on rankers and agreement on condensed rankers. We start with a technical lemma.

Lemma 3.12. *If a ranker $r \in \underline{R}_{m,n}^Z$ ($Z = X$ or Y) is defined but not condensed on u , and if s is the maximal prefix of r which is condensed on u , then one of the following holds, for some $\ell \geq 1$:*

- $r = sX_b t$, $s = s_0 Y_a X_{b_1} \cdots X_{b_{\ell-1}}$ and $s_0 Y_a(u) \leq s(u) < s_0(u) \leq sX_b(u)$;
- $r = sY_b t$, $s = s_0 X_a Y_{b_1} \cdots Y_{b_{\ell-1}}$ and $s_0 X_a(u) \geq s(u) > s_0(u) > sY_b(u) = s_0 Y_b(u)$.

Moreover s_0 is not empty, $s_0 \in \underline{R}_{m-1,n-\ell}^Z$; $sX_b(u) = s_0(u)$ (resp. $sY_b(u) = s_0(u)$) if the last letter of s_0 is in $\{X_b, Y_b\}$; and $sX_b(u) = s_0 X_b(u)$ (resp. $sY_b(u) = s_0 Y_b(u)$) otherwise.

Proof. Rankers in $R_{1,n}$ are condensed on each word on which they are defined (Example 3.3). Therefore we have $m \geq 2$.

By hypothesis, $s \neq r$. We consider the case where the first letter after s is an X -letter, the other case is dual. Then r is of the form $r = sX_b t$, where t may be empty. In view of Example 3.3, $s = s_0 Y_a X_{b_1} \cdots X_{b_{\ell-1}}$ for some non-empty s_0 and $\ell \geq 1$. Since s is condensed on u but sX_b is not, we have the following (see Remark 3.2):

$$s_0 Y_a(u) < s_0 Y_a X_{b_1}(u) \cdots < s_0 Y_a X_{b_1} \cdots X_{b_{\ell-1}}(u) = s(u) < s_0(u),$$

and $sX_b(u) \geq s_0(u)$. More precisely, $sX_b(u)$ is the first b -position to the right of $s(u)$, so $sX_b(u) = s_0(u)$ if $s_0(u)$ is a b -position (i.e., if s_0 ends with X_b or Y_b), and $sX_b(u) = s_0 X_b(u)$ otherwise. \square

Proposition 3.13. *Let $n \geq m \geq 1$, $u, v \in A^+$ and $Z \in \{X, Y\}$. If u and v agree on condensed rankers in $\underline{R}_{m,n}^Z$ and if $r \in \underline{R}_{m,n}^Z$ is defined on both u and v , then there exists $r' \in \underline{R}_{m,n}^Z$ which is condensed on u and v and coincides with r on both words.*

Proof. The result is trivial if $m = 1$, since rankers in $R_{1,n}$ are condensed on each word on which they are defined (Example 3.3). We now assume that $m \geq 2$.

Let p and q be positions in u and v and let $r \in \underline{R}_{m,n}^Z$ such that $r(u) = p$ and $r(v) = q$. If r is not condensed on u , then r is not condensed on v (since the two words agree on condensed rankers). With the notation of Lemma 3.12, r coincides on both u and v with $r' = s_0 t$, $s_0 X_b t$ or $s_0 Y_b t$ (depending on the last letter of s_0 and on the letter following s in r), which starts with the same letter as r . If r' is not condensed on u and v , we repeat the reasoning. This process must terminate since each iteration reduces the depth of r' . \square

Proposition 3.14. *Let $n \geq m \geq 1$, $u, v \in A^+$ and $Z \in \{X, Y\}$. If u and v agree on condensed rankers in $\underline{R}_{m,n}^Z$, then they agree on rankers from the same class.*

Proof. If u and v do not agree on rankers from $\underline{R}_{m,n}^Z$, let $r \in \underline{R}_{m,n}^Z$ be a minimum depth ranker on which u and v disagree. Without loss of generality, we may assume that $u \in L(r)$ and $v \notin L(r)$. In particular, r is not condensed on u .

Let s, s_0 and t be as in Lemma 3.12. Without loss of generality again, we may assume that the letter following s in r is X_b . Since s is condensed on u and sX_b is not, the ranker s is condensed on v and sX_b is not. Moreover, sX_b coincides on u with $s' = s_0$, or s_0X_b , depending on the last letter of s_0 . Observe that s' is shorter than r , so s' is defined on v . In particular, there exists a b -position in v to the right of s , which is not to the left of s_0 (since sX_b is not condensed on v). It follows that $sX_b(v) = s'(v)$. Let now $r' = s't$: then r' is shorter than r , it coincides with r on u , and it is not defined on v since s' coincides with s on that word. This contradicts the minimality of r . \square

We can now prove the following variant of Theorem 1.7.

Theorem 3.15. *Let $u, v \in A^*$ and let $1 \leq m \leq n$. Then u and v satisfy the same formulas in $\text{FO}_{m,n}^2$ if and only if*

- (WI 1c) *u and v agree on condensed rankers from $\underline{R}_{m,n}$,*
- (WI 2c) *if the rankers $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m-1,n-1}$ are condensed on u and v , then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.*
- (WI 3c) *if $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m,n-1}$ are condensed on u and v and end with different direction letters, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.* \blacksquare

Proof. We need to prove that together, Properties (WI 1), (WI 2) and (WI 3) are equivalent to Properties (WI 1c), (WI 2c) and (WI 3c).

Let us first assume that (WI 1), (WI 2) and (WI 3) hold. It is immediate that (WI 2c) and (WI 3c) hold. If (WI 1c) does not hold, let r be a ranker in $\underline{R}_{m,n}$ which is condensed on v and not on u . Since (WI 1) holds, r is defined on u . Let s_0, s and t be as in Lemma 3.12 and let us assume, without loss of generality, that the letter following s in r is X_b . Then s_0 and sX_b are defined on both u and v , with $s_0 \in \underline{R}_{m-1,n-1}$ and $sX_b \in \underline{R}_{m,n}$. Since r is condensed on v , we have $sX_b(v) < s_0(v)$, and since sX_b is not condensed on u , we have $s_0(v) \leq sX_b(v)$, contradicting Property (WI 2). Thus (WI 1c) holds.

Conversely, let us assume that (WI 1c), (WI 2c) and (WI 3c) hold. Then (WI 1) holds by Proposition 3.14. Let us verify Property (WI 2): suppose that $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m-1,n-1}$ are defined on u and v . In view of (WI 1c), Proposition 3.13 shows that there exist rankers $s \in \underline{R}_{m,n}$ and $s' \in \underline{R}_{m-1,n-1}$ which are condensed on u and v , and which coincide with r and r' , respectively, on both words. By (WI 2c), we have $\text{ord}(s(u), s'(u)) = \text{ord}(s(v), s'(v))$, and hence $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. Thus Property (WI 2) holds. The verification of (WI 3) is identical. \square

These results imply the following statement, which refines Corollary 1.8 and can be proved like that Corollary, using Propositions 1.5 and 3.14, and Theorem 3.15.

Corollary 3.16. *For each $m \geq 1$, we have $\underline{\mathcal{L}}_m^X \subseteq \mathcal{R}_m \subseteq \mathcal{FO}_m^2$ and $\underline{\mathcal{L}}_m^Y \subseteq \mathcal{L}_m \subseteq \mathcal{FO}_m^2$.*

Since \mathcal{FO}^2 is the variety of languages corresponding to **DA**, this yields the following containments.

Corollary 3.17. *For each $m \geq 1$, \mathbf{R}_m and \mathbf{L}_m are contained in **DA**.*

3.4. Condensed rankers and deterministic products. Recall that a product of languages $L = L_0 a_1 L_1 \cdots a_k L_k$ ($k \geq 1$, $a_i \in A$, $L_i \subseteq A^*$) is *deterministic* if, for $0 \leq i \leq k$, each word $u \in L$ has a unique prefix in $L_0 a_1 L_1 \cdots L_{i-1} a_i$. If for each i , the letter a_i does not occur in L_{i-1} , the product $L_0 a_1 L_1 \cdots a_k L_k$ is called *visibly deterministic*: this is obviously a particular case of a deterministic product.

The definition of a *co-deterministic* or *visibly co-deterministic* product is dual, in terms of suffixes instead of prefixes. If \mathcal{V} is a class of languages and A is a finite alphabet, let $\mathcal{V}^{det}(A)$ (resp. $\mathcal{V}^{vdet}(A)$, $\mathcal{V}^{codet}(A)$, $\mathcal{V}^{vcodet}(A)$) be the set of all Boolean combinations of languages of $\mathcal{V}(A)$ and of deterministic (resp. visibly deterministic, co-deterministic, visibly co-deterministic) products of languages of $\mathcal{V}(A)$.

Schützenberger gave algebraic characterizations of the operations $\mathcal{V} \mapsto \mathcal{V}^{det}$ and $\mathcal{V} \mapsto \mathcal{V}^{codet}$, see [15].

Proposition 3.18. *If \mathcal{V} is a variety of languages and if \mathbf{V} is the corresponding pseudovariety of monoids, then \mathcal{V}^{det} and \mathcal{V}^{codet} are varieties of languages and the corresponding pseudovarieties are, respectively, $\mathbf{K} \textcircled{m} \mathbf{V}$ and $\mathbf{D} \textcircled{m} \mathbf{V}$.*

This leads to the following statement.

Theorem 3.19. *For each $m \geq 1$, we have $\mathcal{R}_{m+1} = \mathcal{L}_m^{vdet} = \mathcal{L}_m^{det}$, $\mathcal{L}_{m+1} = \mathcal{R}_m^{vcodet} = \mathcal{R}_m^{codet}$, $\mathbf{R}_{m+1} = \mathbf{K} \textcircled{m} \mathbf{L}_m$ and $\mathbf{L}_{m+1} = \mathbf{D} \textcircled{m} \mathbf{R}_m$. In particular, $\mathbf{R}_2 = \mathbf{R}$ and $\mathbf{L}_2 = \mathbf{L}$.*


The proof uses the following technical property of monoids in **DA**, whose proof can be found for instance in [6, Lemma 4.2].

Fact 3.20. *Let $\sigma: A^* \rightarrow S$ be a morphism into a monoid $S \in \mathbf{DA}$. If $u, v \in A^*$, $a \in \text{alph}(v)$ and $\sigma(u) \mathcal{R} \sigma(uv)$, then $\sigma(uva) \mathcal{R} \sigma(u)$.*

Proof of Theorem 3.19. It is immediate from the definition that $\mathcal{L}_m^{vdet} \subseteq \mathcal{L}_m^{det}$.

Let $u \in A^*$ and let $B = \text{alph}(u)$. For each $a \in B$, let $u = u_-^{(a)} a u_+^{(a)}$ be the a -left factorization of u . Let $[B]$ be the language of all strings with alphabet B , $[B] = \{u \in A^* \mid \text{alph}(u) = B\}$. Observe that

$$[B] = \bigcap_{a \in B} L_c(X_a) \setminus \bigcup_{a \notin B} L_c(X_a) = \bigcap_{a \in B} L_c(Y_a) \setminus \bigcup_{a \notin B} L_c(Y_a).$$

FIGURE 1. $M \in \mathbf{K} \textcircled{m} \mathbf{L}_m$

This shows that $[B] \in \mathcal{R}_1 = \mathcal{L}_1$. (It is also well-known that $[B]$ is piecewise testable, and hence $[B] \in \mathcal{R}_1 = \mathcal{L}_1$.)

Now let $n > m \geq 1$. It follows from Proposition 3.6 that the $\triangleright_{m+1,n}$ -class of u is the intersection of $[B]$, the $\triangleleft_{m,n-1}$ -class of u and the products KaL ($a \in B$) where K is the $\triangleleft_{m,n-1}$ -class of $u_-^{(a)}$ and L is the $\triangleright_{m+1,n-1}$ -class of $u_+^{(a)}$ if $n > m + 1$, the $\triangleright_{m,n-1}$ -class of $u_+^{(a)}$ if $n = m + 1$.

By definition of an a -left factorization, each of these products is visibly deterministic and, since every $\triangleleft_{m,n-1}$ -class is a language in \mathcal{L}_m , we have shown that the $\triangleright_{m+1,n}$ -class of u is in \mathcal{L}_m^{vdet} . Thus $\mathcal{R}_{m+1} \subseteq \mathcal{L}_m^{vdet}$.

To establish the last inclusion, namely the fact that $\mathcal{L}_m^{det} \subseteq \mathcal{R}_{m+1}$, we show rather that $\mathbf{K} \textcircled{m} \mathbf{L}_m \subseteq \mathbf{R}_{m+1}$.

Let $\gamma: A^* \rightarrow M$ be a surjective morphism, onto a monoid $M \in \mathbf{K} \textcircled{m} \mathbf{L}_m$: we want to show that there exists a morphism from $A^*/\triangleright_{m+1,n}$ onto M for some $n > m$. Since $M \in \mathbf{K} \textcircled{m} \mathbf{L}_m$, there exists a monoid T and onto morphisms $\alpha: T \rightarrow M$ and $\beta: T \rightarrow N$ such that $N \in \mathbf{L}_m$ and $\beta^{-1}(e) \in \mathbf{K}$ for each idempotent $e \in T$. Using a classical property of free monoids, we can choose a morphism $\tau: A^* \rightarrow T$ such that $\gamma = \alpha \circ \tau$. Moreover, by replacing T by $\tau(A^*)$ and N by $\beta(\tau(A^*))$, we may assume that α , β and τ are onto.

By definition of \mathbf{L}_m , there exists an integer n and a morphism δ' from $A^*/\triangleleft_{m,n}$ onto N such that $\beta \circ \tau = \delta' \circ \delta$, where δ is the natural projection morphism from A^* onto $A^*/\triangleleft_{m,n}$, see Figure 1.

Let ℓ be the maximal length of a strict \mathcal{R} -chain in M , that is: if $x_n <_{\mathcal{R}} \dots <_{\mathcal{R}} x_1$ in M , then $n \leq \ell$. We show that, for any $u, v \in A^*$,

$$(1) \quad u \triangleright_{m+1, \ell|A|+n+1} v \implies \gamma(u) = \gamma(v).$$

If $n' = \ell|A| + n + 1$, this implies the existence of a morphism from $A^*/\triangleright_{m+1, n'}$ onto M , as announced.

To prove implication (1), it suffices to show that we have

$$(2) \quad u \triangleright_{m+1, \ell|\mathbf{alph}(u)|+n+1} v \implies \gamma(u) = \gamma(v),$$

which we prove by induction on $|\mathbf{alph}(u)|$. If $|\mathbf{alph}(u)| = 0$, then $u = \varepsilon$, $\mathbf{alph}(v) = \emptyset$ and $v = \varepsilon$ as well, so that $\gamma(u) = \gamma(v)$.

Now suppose that $u \neq \varepsilon$ and assume that $u \triangleright_{m+1, \ell|\mathbf{alph}(u)|+n+1} v$. Let $u = u_1 a_1 \dots a_k u_{k+1}$ be the factorization of u such that each u_i is a word, each a_i is a letter and

$$1 \mathcal{R} \gamma(u_1) >_{\mathcal{R}} \gamma(u_1 a_1) \dots >_{\mathcal{R}} \gamma(u_1 a_1 \dots u_k a_k) \mathcal{R} \gamma(u_1 a_1 \dots a_k u_{k+1}).$$

Then $k + 1 \leq \ell$, so $k < \ell$. Moreover, by Fact 3.20 (and Corollary 3.17), for each $1 \leq i \leq k$, $a_i \notin u_i$, so that each product $u_i a_i (u_{i+1} \cdots a_k u_{k+1})$ is an a_i -left factorization ($1 \leq i \leq k$).

An easy induction on k , using Lemma 3.7, shows that v can then be factored as

$$v = v_1 a_1 v_2 \cdots a_k v_{k+1},$$

where $u_i \triangleright_{m+1, \ell|\mathbf{alph}(u)|+n-i+1} v_i$ for each $1 \leq i \leq k + 1$. Moreover, for $1 \leq i \leq k$, $|\mathbf{alph}(u_i)| < |\mathbf{alph}(u)|$. Since $i \leq k < \ell$, we have $\ell|\mathbf{alph}(u)|+n-i \geq \ell|\mathbf{alph}(u_i)|+n+1$, and by induction, we have $\gamma(u_i) = \gamma(v_i)$. However, it is possible that $\mathbf{alph}(u_{k+1}) = \mathbf{alph}(u)$, so we cannot conclude that $\gamma(u_{k+1}) = \gamma(v_{k+1})$.

But we do have the following:

$$u_{k+1} \triangleright_{m+1, \ell|\mathbf{alph}(u)|+n-k} v_{k+1} \text{ and } \gamma(u') = \gamma(v'),$$

where $u' = u_1 a_1 \cdots u_k a_k$ and $v' = v_1 a_1 \cdots v_k a_k$. The first relation implies that $u_{k+1} \triangleleft_{m, \ell|\mathbf{alph}(u)|+n-k-1} v_{k+1}$. Since $k < \ell$, we have $\ell|\mathbf{alph}(u)|+n-k-1 \geq n$, and hence $u_{k+1} \triangleleft_{m, n} v_{k+1}$.

Moreover, there exists a string $x \in A^*$ such that $\gamma(u' u_{k+1} x) = \gamma(u')$. Let ω be an integer such that every ω -power is idempotent in M and N and let $y = (x u_{k+1})^{\omega-1} x$. Then we have

$$(x u_{k+1})^\omega = y u_{k+1} \triangleleft_{m, n} y v_{k+1}.$$

By the induction hypothesis, this implies that

$$\begin{aligned} \delta'(\delta(y u_{k+1})) &= \delta'(\delta(x u_{k+1})^\omega) = \delta'(\delta(y v_{k+1})), \text{ so} \\ \beta(\tau(y u_{k+1})) &= \beta(\tau(x u_{k+1})^\omega) = \beta(\tau(x v_{k+1})^\omega). \end{aligned}$$

Since $\alpha(\beta^{-1}(e)) \in \mathbf{K}$ for each idempotent e of N , we have

$$\begin{aligned} \alpha(\tau(x u_{k+1})^\omega) \alpha(\tau(y v_{k+1})) &= \alpha(\tau(x u_{k+1})^\omega), \text{ that is,} \\ \gamma((x u_{k+1})^\omega y v_{k+1}) &= \gamma(x u_{k+1})^\omega. \end{aligned}$$

It follows that

$$\begin{aligned} \gamma(u) &= \gamma(u' u_{k+1}) = \gamma(u' u_{k+1} x u_{k+1}) \\ &= \gamma(u' u_{k+1} (x u_{k+1})^\omega) \\ &= \gamma(u' u_{k+1} (x u_{k+1})^\omega y v_{k+1}) \\ &= \gamma(u' u_{k+1} (x u_{k+1})^\omega (x u_{k+1})^{\omega-1} x v_{k+1}) \\ &= \gamma(u' (u_{k+1} x)^\omega v_{k+1}) \\ &= \gamma(u' v_{k+1}) = \gamma(v' v_{k+1}) = \gamma(v). \end{aligned}$$

This concludes the proof of Formula (2), and therefore of Theorem 3.19. \square

3.5. Structure of the \mathbf{R}_m and \mathbf{L}_m hierarchies. It turns out that the hierarchies of pseudovarieties given by the \mathbf{R}_m and the \mathbf{L}_m were studied in the semigroup-theoretic literature (Trotter and Weil [28], Kufleitner and Weil [12]). In [12], they are defined as the hierarchies of pseudovarieties obtained from \mathbf{J} by alternated applications of the operations $\mathbf{X} \mapsto \mathbf{K} \circledast \mathbf{X}$ and $\mathbf{X} \mapsto \mathbf{D} \circledast \mathbf{X}$. Theorem 3.19 shows that these are the same hierarchies as those considered in this paper¹. The following results are proved in [12, Section 4].

Proposition 3.21. *The hierarchies $(\mathbf{R}_m)_m$ and $(\mathbf{L}_m)_m$ are infinite chains of decidable pseudovarieties, and their unions are equal to \mathbf{DA} . Moreover, every m -generated monoid in \mathbf{DA} lies in $\mathbf{R}_{m+1} \cap \mathbf{L}_{m+1}$.*

The results in [28, 12] go actually further, and give defining pseudoidentities for the pseudovarieties \mathbf{R}_m and \mathbf{L}_m .

4. THE \mathbf{R}_m HIERARCHY AND UNARY TEMPORAL LOGIC

We have seen in Corollary 3.16 that $\underline{\mathcal{I}\mathcal{L}}_m^X \subseteq \mathcal{R}_m$ and $\underline{\mathcal{I}\mathcal{L}}_m^Y \subseteq \mathcal{L}_m$. In Proposition 4.3 below, we prove a weak converse. Let us however make the following observation.

Proposition 4.1. *We have*

$$\begin{aligned} \underline{\mathcal{I}\mathcal{L}}_1^X &= \underline{\mathcal{I}\mathcal{L}}_1^Y = \mathcal{R}_1 = \mathcal{L}_1, \\ \underline{\mathcal{I}\mathcal{L}}_2^X &= \mathcal{R}_2, \quad \underline{\mathcal{I}\mathcal{L}}_2^Y = \mathcal{L}_2. \end{aligned}$$

Proof. The statement concerning $\underline{\mathcal{I}\mathcal{L}}_1$ was already proved in Remark 1.6. Let us now establish that $\mathcal{R}_2 \subseteq \underline{\mathcal{I}\mathcal{L}}_2^X$. We show, by induction on $n \geq 2$, that if u and v agree on rankers in $\underline{R}_{2,2n}^X$, then they agree on condensed rankers in $\underline{R}_{2,n}^X$: $u \triangleright_{2,n} v$. We use the characterization of $\triangleright_{2,n}$ in Proposition 3.6.

The consideration of 1-letter rankers shows that $\mathbf{alph}(u) = \mathbf{alph}(v)$. Moreover, since $\underline{R}_{1,n-1}^Y$ is contained in $\underline{R}_{2,2n}^X$, and since these rankers are condensed where they are defined, we find that $u \triangleleft_{1,n-1} v$. Similarly, if $u = u_- a u_+$ and $v = v_- a v_+$ are a -left factorizations, let $s \in \underline{R}_{1,n-1}^Y$. Then s is condensed on u_- if and only if s is defined on u_- , if and only if $X_a s$ is defined on u (Lemma 3.5). Since $X_a s \in \underline{R}_{2,2n}^X$ and u and v agree on such rankers, it follows that $X_a s$ is defined on v , and s is condensed on v_- . Thus $u_- \triangleleft_{1,n-1} v_-$.

Now we need to show that $u_+ \triangleright_{2,n-1} v_+$ if $n \geq 3$, $u_+ \triangleright_{1,1} v_+$ if $n = 2$. Suppose first that $n = 2$ and consider $s \in \underline{R}_{1,1}^X$, condensed on u_+ . Then $s = X_b$ for some $b \in A$ and the consideration of $r = X_a X_b$ (in $\underline{R}_{2,2}^X$) shows that s is condensed on v_+ as well. This settles the case $n = 2$.

¹More precisely, the pseudovarieties \mathbf{R}_m and \mathbf{L}_m in [12] are pseudovarieties of semi-groups, and the \mathbf{R}_m and \mathbf{L}_m considered in this paper are the classes of monoids in these pseudovarieties.

Let us now assume that $n \geq 3$ and let us show that $u_+ \triangleright_{2,n-1} v_+$. By induction, it suffices to show that u_+ and v_+ agree on rankers in $\underline{R}_{2,2n-2}^X$. So let $s \in \underline{R}_{2,2n-2}^X$ be defined on u_+ . Then for every prefix p of s ending with a Y-letter, $X_a p Y_a$ is defined on u (Lemma 3.5). Since $X_a p Y_a \in \underline{R}_{2,2n}^X$, it follows that $X_a p Y_a$ is defined on v , and hence s is defined on v_+ . This concludes the proof. \square

Example 4.2 below shows that the statement of Proposition 4.1 cannot be extended to the higher levels of the hierarchy.

Example 4.2. We show in this example that $\underline{\mathcal{TL}}_3^X$ is properly contained in \mathcal{R}_3 . More precisely, let $r_0 = X_a Y_b X_c \in R_{3,3}^X$. We show that $L_c(r_0)$, a language in \mathcal{R}_3 , is not $\underline{\mathcal{TL}}_3^X$ -definable.

Let $u_n = (bc)^n (a(bc)^n)^n$ and $v_n = (bc)^n b (a(bc)^n)^n$ ($n \geq 1$). It is easily verified that r_0 is condensed on u_n , and that it is defined and not condensed on v_n : that is, for each n , $u_n \in L_c(r_0)$ and $v_n \notin L_c(r_0)$.

We now show that u_n and v_n agree on all rankers in $\underline{R}_{3,n}^X$, so that any $\underline{\mathcal{TL}}_3^X$ -definable language contains either both u_n and v_n , or neither – and hence $L_c(r_0)$ is not $\underline{\mathcal{TL}}_3^X$ -definable.

Let $r \in \underline{R}_{3,n}^X$. If r starts with a Y-letter, then any two words ending with $(a(bc)^n)^n$ agree on r . In particular, u_n and v_n agree on r . Similarly, if r starts with an X-letter and does not contain the letters X_a or Y_a , then any two words starting with $(bc)^n$ agree on r , so u_n and v_n agree on r .

Finally, assume that r starts with an X-letter and that $r = s_0 Z_a^{(1)} s_1 \cdots Z_a^{(k)} s_k$ with $k > 0$, each $Z^{(i)} \in \{X, Y\}$ and each s_i a (possibly empty) ranker avoiding the letters X_a and Y_a . We denote by p_i the prefix $p_i = s_0 Z_a^{(1)} s_1 \cdots Z_a^{(i)}$.

Suppose first that $r \in R_{1,n}^X$. Then p_i coincides with X_a^i on u_n as well as on v_n . Therefore r is defined and coincides with $X_a^k s_k$ on both words.

Suppose now that $r \in R_{2,n}^X$, say $r = r' r''$ with r' a non-empty string of X-letters and r'' a non-empty string of Y-letters. If r' is shorter than p_1 , then $Z^{(1)} = Y$ and r is not defined on either u_n or v_n . If $Z^{(1)} = X$, let i be maximal such that p_i is a prefix of r' , say $r' = p_i s'_i$. Then $i > 0$ and p_i coincides with X_a^i on u_n , as well as on v_n .

If $i = k$, then r is defined on u_n and v_n , and it coincides with $X_a^k s_k$ on both words.

If $1 \leq i < k$, s'_i is non-empty and s_i is defined on $(bc)^n$, then p_{i+1} coincides with X_a^i on u_n and v_n . Thus r is defined on u_n (resp. v_n) if and only if $i \geq k - i$, and in that case, it coincides with $X_a^{k-2i} s_k$.

If $1 < i < k$, s'_i is non-empty and s_i is not defined on $(bc)^n$, or if s'_i is empty, then p_{i+1} coincides with X_a^{i-1} on u_n and v_n . Thus r is defined on u_n (resp. v_n) if and only if $i > k - i$, and in that case, it coincides with $X_a^{k-2i-1} s_k$.

Finally, if $1 = i < k$, s'_i is non-empty and s_i is not defined on $(bc)^n$, or if s'_i is empty, then r (and even p_{i+1}) is not defined on either u_n or v_n .

Finally, let us assume that $r \in R_{3,n}^X$, say $r = r'r''r'''$ with r' and r''' non-empty strings of X-letters and r'' a non-empty string of Y-letters. Again, let i be maximal such that p_i is a prefix of r' ($i = 0$ if p_1 is not a prefix of r') and let j be maximal such that p_j is a prefix of $r'r''$. Then $r'r'' = p_j s'_j$ for some prefix s'_j of s_j . By the previous analysis, if $i \leq 1 < j$, then $r'r''$ is not defined on u_n nor on v_n , and hence neither is r . In all other cases, $r'r''$ is defined on both words and coincides with $X_a^{2i-j} s'_j$ or $X_a^{2i-j-1} s'_j$. Since $(k-j) + (2i-j) \leq k$, r is defined on u_n and v_n , and coincides on these words with $X_a^{k-j+2i-j} s_k$ or $X_a^{k-j+2i-j-1} s_k$.

To conclude this example, note that u_n and v_n disagree on rankers in R_4^X . More precisely, the ranker $X_a Y_c X_b Y_a$ is defined on v_n but not on u_n . Further getting ahead of ourselves, we note that this example also shows (in view of Theorem 5.1) that \mathcal{TL}_3 is properly contained in \mathcal{FO}_3^2 . \square

Finally we prove a result on the containment of the \mathcal{R}_m and \mathcal{L}_m hierarchies in the \mathcal{TL}_m hierarchy.

Proposition 4.3. *Let $m \geq 1$. Then $\mathcal{R}_m \subseteq \mathcal{TL}_{2m-1}^X$ and $\mathcal{L}_m \subseteq \mathcal{TL}_{2m-1}^Y$.*

More precisely, for all $n \geq m$, $Z \in \{X, Y\}$ and $u, v \in A^$, if u and v agree on rankers in $R_{2m-1, 2n-1}^Z$, then they agree on condensed rankers in $R_{m,n}^Z$.*

Proof. Without loss of generality, we may assume $Z = X$. The proof is by induction on m . The result is trivial if $m = 1$, since $2m - 1 = 1$ and $2n - 1 \geq n$. We now assume that $m \geq 2$ and u, v agree on rankers in $R_{2m-1, 2n-1}^X$.

We use the characterization of $\triangleright_{m,n}$ in Proposition 3.6: the consideration of length 1 rankers shows that $\text{alph}(u) = \text{alph}(v)$. Since $R_{2m-3, 2n-3}^Y$ is contained in $R_{2m-1, 2n-1}^X$, we have $u \triangleleft_{m-1, n-1} v$ by induction. Now, for each letter $a \in \text{alph}(u)$, let $u = u_- a u_+$ and $v = v_- a v_+$ be the a -left factorizations. We want to show that $u_- \triangleleft_{m-1, n-1} v_-$ and $u_+ \triangleright_{m, n-1} v_+$ ($u_+ \triangleright_{m-1, n-1} v_+$ if $m = n$). By induction, it suffices to show that u_- and v_- agree on rankers in $R_{2m-3, 2n-3}^Y$, and u_+ and v_+ agree on rankers in $R_{2m-1, 2n-3}^X$ ($R_{2m-3, 2n-3}^X$ if $m = n$). In the rest of the proof we silently rely on the results of Lemma 3.5.

Let $s \in R_{2m-3, 2n-3}^Y$ be defined on u_- . If s starts with a Y-block, then $X_a s \in R_{2m-2, 2n-2}^X$ and $X_a s$ is defined on u . Moreover, if p is any prefix of s , then $X_a p Y_a \in R_{2m-1, 2n-1}^X$ is not defined on u . It follows that s is defined on v_- .

If instead s starts with an X-block, then $s \in R_{2m-4, 2n-4}^X$ and s is defined on u . If p is any prefix of s , then $p Y_a \in R_{2m-3, 2n-3}^X$ and $p Y_a$ is not defined on u . As all these rankers are in $R_{2m-1, 2n-1}^X$, the same holds on v and s is defined on v_- .

Let now $s \in R_{2m-1, 2n-3}^X$ ($s \in R_{2m-3, 2n-3}^X$ if $n = m$) be defined on u_+ . If s starts with an X-block, then $X_a s \in R_{2m-1, 2n-2}^X$ ($X_a s \in R_{2m-3, 2n-2}^X$ if $n = m$) and $X_a s$ is defined on u . Moreover, for each prefix p of s ending with

a Y-letter, $X_a p Y_a \in \underline{R}_{2m-1, 2n-1}^X$ ($X_a p Y_a \in \underline{R}_{2m-3, 2n-1}^X$ if $n = m$) and $X_a p Y_a$ is defined on u . As all these rankers are in $\underline{R}_{2m-1, 2n-1}^X$, the same holds on v and s is defined on v_+ .

If instead s starts with a Y-block, then $s \in \underline{R}_{2m-2, 2n-4}^Y$ ($s \in \underline{R}_{2m-4, 2n-4}^Y$ if $n = m$) and s is defined on u . Moreover, if p is any prefix of s ending with a Y-letter, $p Y_a \in \underline{R}_{2m-2, 2n-3}^Y$ ($p Y_a \in \underline{R}_{2m-4, 2n-3}^Y$ if $n = m$) and $p Y_a$ is defined on u . As all these rankers are in $\underline{R}_{2m-1, 2n-1}^X$, the same holds on v and s is defined on v_+ . \square

The containment of \mathcal{R}_m and \mathcal{L}_m into $\underline{\mathcal{I}\mathcal{L}}_{2m-1}^X$ and $\underline{\mathcal{I}\mathcal{L}}_{2m-1}^Y$, respectively, is not very precise, unfortunately, especially in view of Proposition 6.4 below.

5. THE \mathbf{R}_m HIERARCHY AND \mathbf{FO}_m^2

The objective of this section is to prove the following theorem.

Theorem 5.1. *Let $m \geq 1$. Every language in \mathcal{R}_m or \mathcal{L}_m is \mathbf{FO}_m^2 -definable, and every \mathbf{FO}_m^2 -definable language is in $\mathcal{R}_{m+1} \cap \mathcal{L}_{m+1}$. Equivalently, we have*

$$\mathbf{R}_m \vee \mathbf{L}_m \subseteq \mathbf{FO}_m^2 \subseteq \mathbf{R}_{m+1} \cap \mathbf{L}_{m+1},$$

where $\mathbf{V} \vee \mathbf{W}$ denotes the least pseudovariety containing \mathbf{V} and \mathbf{W} .

5.1. Are the containments in Theorem 5.1 strict? In the particular case where $m = 1$, we know that $\mathbf{R}_2 \cap \mathbf{L}_2 = \mathbf{R} \cap \mathbf{L} = \mathbf{J} = \mathbf{R}_1 \vee \mathbf{L}_1$: this reflects the elementary observation that \mathbf{FO}_1^2 -definable languages, like \mathbf{FO}_1 -definable languages, are the piecewise testable languages. However, for $m \geq 2$, we conjecture that $\mathbf{R}_m \vee \mathbf{L}_m$ is properly contained in $\mathbf{R}_{m+1} \cap \mathbf{L}_{m+1}$. The following example proves the conjecture for $m = 2$.

Example 5.2. $L = \{b, c\}^* c a \{a, b\}^*$ is \mathbf{FO}_2^2 -definable, by the following formula:

$$\begin{aligned} & \exists i \quad (\mathbf{c}(i) \wedge (\forall j (j < i \rightarrow \neg \mathbf{a}(j))) \wedge (\forall j (j > i \rightarrow \neg \mathbf{c}(j)))) \\ & \wedge \exists i \quad (\mathbf{a}(i) \wedge (\forall j (j < i \rightarrow \neg \mathbf{a}(j))) \wedge (\forall j (j > i \rightarrow \neg \mathbf{c}(j)))) \\ & \wedge \forall i \quad (\mathbf{b}(i) \rightarrow (\exists j (j < i \wedge \mathbf{a}(j)) \vee (\exists j (j > i \wedge \mathbf{c}(j)))). \end{aligned}$$

The words $u_n = (bc)^n (ab)^n$ are in L , while the words $v_n = (bc)^n b (ca)^n$ are not. Almeida and Azevedo showed that $\mathbf{R}_2 \vee \mathbf{L}_2$ is defined by the pseudo-identity $(bc)^\omega (ab)^\omega = (bc)^\omega b (ab)^\omega$ [2, Theorem 9.2.13 and Exercise 9.2.15]. In particular, for each language K recognized by a monoid in $\mathbf{R}_2 \vee \mathbf{L}_2$, the words u_n and v_n (for n large enough) are all in K , or all in the complement of K . Therefore L is not recognized by such a monoid, which proves that $\mathbf{R}_2 \vee \mathbf{L}_2$ is strictly contained in \mathbf{FO}_2^2 , and hence also in $\mathbf{R}_3 \cap \mathbf{L}_3$. It also shows that $\underline{\mathcal{I}\mathcal{L}}_2$ is properly contained in \mathcal{FO}_2^2 . \square

Finally, we formulate the following conjecture.

Conjecture 5.3. *For each $m \geq 1$, $\mathbf{FO}_m^2 = \mathbf{R}_{m+1} \cap \mathbf{L}_{m+1}$.*

5.2. Proof of Theorem 5.1. Corollary 3.16 already established that every language in \mathcal{R}_m or \mathcal{L}_m is FO_{*m*}²-definable².

In view of Theorem 3.15, to establish that \mathcal{FO}_m^2 is contained in $\mathcal{R}_{m+1} \cap \mathcal{L}_{m+1}$, it suffices to prove the following result.

For each $n \geq m \geq 1$, if $u \triangleright_{m+1,2n} v$ or $u \triangleleft_{m+1,2n} v$, then Properties (WI 1c), (WI 2c) and (WI 3c) hold for m, n .

The result is trivial if $m = 1$, since in that case, only Property (WI 1c) is non-vacuous.

So we now assume that $m \geq 2$, $u \triangleright_{m+1,2n} v$ or $u \triangleleft_{m+1,2n} v$. Property (WI 1c) holds trivially, by definition of the $\triangleright_{m+1,2n}$ and $\triangleleft_{m+1,2n}$ relations. We now concentrate on proving that Properties (WI 2c) and (WI 3c) also hold for m, n , a task that will be completed in Section 5.2.3.

5.2.1. *The case where r and r' start with opposite directions.*

Proposition 5.4. *Let $n \geq m \geq 1$, $r = Y_{a_1}s \in \underline{R}_{m,n}^Y$ and $r' = X_c$. If $u, v \in A^*$, r is condensed on u and v and $u \triangleright_{m,n+1} v$ or $u \triangleleft_{m+1,n+1} v$, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. The dual statement (involving $r = X_{a_1}s \in \underline{R}_{m,n}^X$ and $r' = Y_c$) holds as well.*

Proof. *First suppose that $u \triangleleft_{m+1,n+1} v$, that is, u and v agree on condensed rankers in $\underline{R}_{m+1,n+1}^Y$. We are in exactly one of the following three situations:*

- rY_c is defined on u , in which case $r'(u) < r(u)$;
- rY_c is undefined on u and c is the last letter to occur in r , in which case $r'(u) = r(u)$;
- rY_c is undefined on u and c is not the last letter to occur in r , in which case $r(u) < r'(u)$.

The same trichotomy holds for v . Since $rY_c \in \underline{R}_{m+1,n+1}^Y$, u and v agree on rY_c (Proposition 3.14), and hence $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

Let us now assume that $u \triangleright_{m,n+1} v$, so that u and v agree on condensed rankers in $\underline{R}_{m,n+1}^X$. If $m = 1$ then r is of the form $r = Y_{a_1} \cdots Y_{a_k}$ and we observe again that

- either $X_c X_{a_k} \cdots X_{a_1} \in \underline{R}_{1,n+1}^X$ is defined on u , and we have $r'(u) < r(u)$;
- or $X_c X_{a_k} \cdots X_{a_1}$ is undefined on u and $c = a_k$, and we have $r'(u) = r(u)$;
- or $X_c X_{a_k} \cdots X_{a_1}$ is undefined on u and $c \neq a_k$, and we have $r'(u) > r(u)$.

The same holds for v since $X_c X_{a_k} \cdots X_{a_1} \in \underline{R}_{1,n+1}^X$ and such rankers are condensed where they are defined. Therefore we have $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

We now assume that $m \geq 2$. Let $u = u_-cu_+$ and $v = v_-cv_+$ be c -left factorizations. We distinguish two cases depending on the direction of the second letter of r .

First suppose that $r = Y_{a_1}Y_{a_2}s'$. If $a_1 \notin \text{alph}(u_+)$, then $r(u) < r'(u)$ (because r is condensed on u). Since $u_+ \triangleright_{m,n} v_+$, we have $\text{alph}(u_+) =$

²Of course, the same fact can be proved by the direct construction of an FO_{*m*}²-formula for each $\triangleright_{m,n}$ -class (by induction on m and using Proposition 3.6).

$\mathbf{alph}(v_+)$, so $r(v) < r'(v)$ as well. If instead $a_1 \in \mathbf{alph}(u_+) = \mathbf{alph}(v_+)$, let $u_+ = u_0 a_1 u_1$ and $v_+ = v_0 a_1 v_1$ be the a_1 -right factorizations. Then

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(Y_{a_2} s'(u_- c u_0), r'(u_- c u_0)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(Y_{a_2} s'(v_- c v_0), r'(v_- c v_0)). \end{aligned}$$

Since $(u_- c u_0) a_1 u_1$ and $(v_- c v_0) a_1 v_1$ are a_1 -right factorizations as well, we deduce from Lemma 3.7 that $u_- c u_0 \triangleright_{m,n} v_- c v_0$ and it follows by induction on the length of r that

$$\text{ord}(Y_{a_2} s'(u_- c u_0), r'(u_- c u_0)) = \text{ord}(Y_{a_2} s'(v_- c v_0), r'(v_- c v_0)).$$

The other case is $r = Y_{a_1} X_{b_1} s'$. If $a_1 \in \mathbf{alph}(c u_+) = \mathbf{alph}(c v_+)$ then $r'(u) < r(u)$ and $r'(v) < r(v)$. If instead $a_1 \notin \mathbf{alph}(c u_+) = \mathbf{alph}(c v_+)$, we first consider the case where r has a single alternation, i.e., $r = Y_{a_1} X_{b_1} \cdots X_{b_k}$. We have $r(u) < r'(u)$ if and only if r is defined on u_- , and hence condensed (Example 3.3). Since $u_- \triangleright_{m,n} v_-$ (Lemma 3.7), this is the case if and only if r is defined on v_- . Hence, if r is defined on u_- , we have $r(u) < r'(u)$ and $r(v) < r'(v)$. If r is not defined on u_- , but $Y_{a_1} X_{b_1} \cdots X_{b_{k-1}}$ is defined on u_- and $b_k = c$, then the same holds for v and we have $r(u) = r'(u)$ and $r(v) = r'(v)$. Otherwise, we have $r(u) > r'(u)$ and $r(v) > r'(v)$.

The last situation arises if r is of the form $r = Y_{a_1} X_{b_1} \cdots X_{b_k} Y_d s''$. In particular, $m \geq 3$. If $Y_{a_1} X_{b_1} \cdots X_{b_k}$ is defined on $u_- c$, then it is defined on $v_- c$ as well (by the same reasoning as in the previous paragraph) and we have $r(u) < r'(u)$ and $r(v) < r'(v)$.

Similarly, if $Y_{a_1} X_{b_1} \cdots X_{b_{k-1}}$ is not defined on u_- and v_- , then we have $r'(u) < r(u)$ and $r'(v) < r(v)$.

Finally, let us assume that $Y_{a_1} X_{b_1} \cdots X_{b_k}$ is not defined on $u_- c$ or $v_- c$, but $Y_{a_1} X_{b_1} \cdots X_{b_{k-1}}$ is defined on u_- and v_- . Let $u_+ = u_0 b_k u_1$ and $v_+ = v_0 b_k v_1$ be b_k -left factorizations. Then

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(Y_d s''(u_- c u_0), r'(u_- c u_0)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(Y_d s''(v_- c v_0), r'(v_- c v_0)). \end{aligned}$$

Since $u \triangleright_{m,n+1} v$, we have $u_+ \triangleright_{m,n} v_+$, and by Lemma 3.7, $u_- \triangleright_{m,n} v_-$ and $u_0 \triangleright_{m,n-1} v_0$. Therefore $u_- c u_0 \triangleright_{m,n-1} v_- c v_0$. Since $Y_d s'' \in \underline{R}_{m-2,n-2}^Y$ is condensed on both $u_- c u_0$ and $v_- c v_0$, we conclude by induction on the length of r that $\text{ord}(Y_d s''(u_- c u_0), r'(u_- c u_0)) = \text{ord}(Y_d s''(v_- c v_0), r'(v_- c v_0))$ and hence $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

This concludes the proof. \square

Proposition 5.5. *Let $n > m \geq 1$, let $r = X_{a_s} \in \underline{R}_m^X$ and $r' = Y_{b_s} \in \underline{R}_m^Y$ such that $|r| + |r'| \leq n$, and let $u, v \in A^*$ such that r and r' are condensed on u and v . If $u \triangleright_{m+1,n} v$ or $u \triangleleft_{m+1,n} v$, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.*

Proof. Without loss of generality, we assume that $u \triangleright_{m+1,n} v$. We proceed by induction, first on m . If $m = 1$, then $r = X_{a_1} \cdots X_{a_k}$ and $r' = Y_{b_1} \cdots Y_{b_\ell}$ with $k + \ell \leq n$. We observe that if $p = r X_{b_\ell} \cdots X_{b_1}$ is defined on u , then $r(u) < r'(u)$; if p is not defined on u , but $a_k = b_\ell$ and $r X_{b_{\ell-1}} \cdots X_{b_1}$ is defined

on u , then $r(u) = r'(u)$; and in all other cases, $r(u) > r'(u)$. The same holds for v , and this completes the proof in case $m = 1$.

We now assume that $m \geq 2$ and proceed by induction on n . We first note that if one of r, r' has length 1, then the result was established in Proposition 5.4. We now assume that $|r|, |r'| \geq 2$ (so $|r|, |r'| \leq n - 2$).

Suppose that $n = m + 1$ and let $\beta(r)$ the number of alternating blocks in r : then $\beta(r) \leq |r| \leq n - |r'| \leq n - 2 = m - 1$. The same inequality holds for r' and we conclude by induction on m .

We must now consider the case where $n > m + 1 > 2$. In particular, we have $r \in \underline{R}_{m,n-2}^X$ and $r' \in \underline{R}_{m+1,n-1}^X$.

First case: s starts with an X-block. Let $u = u_-au_+$ and $v = v_-av_+$ be a -left-factorizations. Then s is condensed on u_+ and v_+ and $u_+ \triangleright_{m+1,n-1} v_+$, so u_+ and v_+ agree on rankers in $\underline{R}_{m+1,n-1}^X$ (Proposition 3.14). In particular, u_+ and v_+ agree on r' . If r' is defined on u_+ , then $\text{ord}(r(u), r'(u)) = \text{ord}(s(u_+), r'(u_+))$. Moreover, r' is defined on v_+ as well and $\text{ord}(r(v), r'(v)) = \text{ord}(s(v_+), r'(v_+))$, so we conclude by induction. If instead r' is not defined on u_+ or v_+ , then $r'(u) \leq X_a(u) < r(u)$ and $r'(v) \leq X_a(v) < r(v)$. ■

Second case: s' starts with a Y-block. Let $u = u_-bu_+$ and $v = v_-bv_+$ be b -right factorizations. Then $u_- \triangleright_{m+1,n-1} v_-$ by Lemma 3.7 and this case can be handled exactly like the previous one.

Third case: s starts with a Y-block and s' starts with an X-block. If $X_a(u) \leq Y_b(u)$, then $X_a(v) \leq Y_b(v)$ (by Proposition 5.4), we have $r(u) < X_a(u) \leq Y_b(u) < r'(u)$, and the same inequalities hold for v .

We now assume that $X_a(u) > Y_b(u)$ and $X_a(v) > Y_b(v)$. In particular, $a \neq b$. Identifying the first a and the last b in u and v , we get factorizations $u = u_-bu_0au_+$ and $v = v_-bv_0av_+$ such that $a \notin \text{alph}(u_-bu_0) \cup \text{alph}(v_-bv_0)$ and $b \notin \text{alph}(u_0au_+) \cup \text{alph}(v_0av_+)$. In particular, $r(u) = s(u_-bu_0)$, $r'(u)$ is the position $s'(u_0au_+)$ in the suffix u_0au_+ of u , and the same holds in v . Moreover, $u = (u_-bu_0)au_+$ is an a -left factorization, $u = u_-b(u_0au_+)$ is a b -right factorization, and the same holds in v . Therefore, and since $u \triangleright_{m+1,n} v$, we have $u_-bu_0 \triangleleft_{m,n-1} v_-bv_0$ by definition and $u_0 \triangleleft_{m,n-2} v_0$ by Lemma 3.7.

Since $s \in \underline{R}_{m-1,n-3}^Y$ and $s' \in \underline{R}_{m-1,n-3}^X \subseteq \underline{R}_{m,n-2}^Y$, Proposition 3.14 shows that, if s is not defined on u_0 , then it is not defined on v_0 either, and $r(u) \leq Y_b(u) < r'(u)$ and similarly, $r(v) < r'(v)$. Symmetrically, if s' is not defined on u_0 , then $r(u) < X_a(u) \leq r'(u)$ and $r(v) < r'(v)$.

Finally, if s and s' are defined on u_0 , then

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(s(u_0), s'(u_0)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(s(v_0), s'(v_0)), \end{aligned}$$

and we conclude by induction. □

5.2.2. *The case where r and r' start with the same direction.*

Proposition 5.6. *Let $n \geq m \geq 2$, $r \in \underline{R}_{m,n}^X$ starting with an X-letter, and $r' = X_c$. If $u, v \in A^*$, r is condensed on u and v and $u \triangleright_{m,n+1} v$, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. The dual statement (involving $r \in \underline{R}_{m,n}^Y$, $r' = Y_c$ starting with a Y-letter, and $u \triangleleft_{m,n+1} v$) holds as well.*

Proof. We proceed by induction, first on m . If $m = 2$, then either $r = X_{a_1} \cdots X_{a_k}$ or $r = X_{a_1} \cdots X_{a_k} Y_{b_1} \cdots Y_{b_\ell}$. In the first case, the order type $\text{ord}(r(u), r'(u))$ depends, as in the proof of Proposition 5.5, on whether $X_c Y_{a_k} \cdots Y_{a_1}$ is defined on u , or if it is not defined, whether $a_k = c$ and $X_c Y_{a_{k-1}} \cdots Y_{a_1}$ is defined. Since these rankers are in $\underline{R}_{2,n+1}^X$ and are condensed where they are defined (Example 3.3), we have $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

In the second case, where $r = X_{a_1} \cdots X_{a_k} Y_{b_1} \cdots Y_{b_\ell}$, three cases arise: if rY_c is defined on u , then $X_c(u) < r(u)$; if rY_c is not defined and $c = b_\ell$, then $X_c(u) = r(u)$; in all other cases, $r(u) < X_c(u)$. Since $u \triangleright_{2,n+1} v$ and $rY_c \in \underline{R}_{2,n+1}^X$, Proposition 3.14 shows that rY_c is defined on u if and only if it is defined on v , and $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

We now assume that $m \geq 3$. If r has less than m alternating blocks, we conclude by induction on m . Let us suppose now that r has m alternating blocks and let us proceed by induction on $|r| \geq m$.

Let $r = X_a s$. If s starts with a Y-letter (which includes the base case where $|r| = m$), then $s \in \underline{R}_{m-1,n-1}^Y$ is condensed on u_- and v_- . If $c \notin \text{alph}(u_-) = \text{alph}(v_-)$, then $r(u) < r'(u)$ and $r(v) < r'(v)$. In all other cases,

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(s(u_-), r'(u_-)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(s(v_-), r'(v_-)). \end{aligned}$$

Since $u_- \triangleright_{m,n} v_-$ by Lemma 3.7, these two order types are equal by Proposition 5.4.

If instead s starts with an X-letter, then $|r| > m$, $s \in \underline{R}_{m,n-1}^X$ is condensed on u_+ and v_+ (Lemma 3.4) and we distinguish two cases. If $c \in \text{alph}(u_- a) = \text{alph}(v_- a)$, then $r'(u) < r(u)$ and $r'(v) < r(v)$. Otherwise

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(s(u_+), r'(u_+)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(s(v_+), r'(v_+)). \end{aligned}$$

Since $u_+ \triangleright_{m,n} v_+$, these two order types are equal by induction on n . \square

Proposition 5.7. *Let $n \geq m \geq 2$, let $r = X_a s \in \underline{R}_m^X$ and $r' = X_b s' \in \underline{R}_{m-1}^X$ such that $|r| + |r'| \leq n$, and let $u, v \in A^*$ such that r and r' are condensed on u and v . If $u \triangleright_{m,n} v$, then $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. The dual statement (where r, r' start with Y-blocks and $u \triangleleft_{m,n} v$) holds as well.*

Proof. The proof is by induction on m , and then on n . If one of r and r' has length 1, then the result was established in Proposition 5.6. This takes

care of the cases where $n \leq 3$, including the base case $m = n = 2$. We now assume that $|r|, |r'| \geq 2$.

Let us observe that under this assumption, if $n = m$, then the number of alternating blocks in r is less than or equal to $m - 2$: indeed it is at most equal to $|r| \leq n - 2 = m - 2$. The same inequality holds for r' , so this situation is handled by induction on m . We can now assume that $n > m$.

Let $u = u_- au_+ = u'_- bu'_+$ and $v = v_- av_+ = v'_- bv'_+$ be a -left and b -left factorizations.

First case: $a = b$. If s starts with an X -block and s' starts with a Y -block, then $r'(u) < r(u)$ and $r'(v) < r(v)$. Dually, if s starts with a Y -block and s' starts with an X -block, then $r'(u) > r(u)$ and $r'(v) > r(v)$.

If s and s' both start with a Y -block (which can happen only if $m - 1 \geq 2$), then $s \in \underline{R}_{m-1}^Y$ and $s' \in \underline{R}_{m-2}^Y$ are condensed on u_- and v_- and

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(s(u_-), s'(u_-)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(s(v_-), s'(v_-)). \end{aligned}$$

Since $u_- \triangleleft_{m-1, n-1} v_-$ and $|s| + |s'| \leq n - 2$, we have $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$ by induction on m .

If instead s and s' both start with an X -block, then $s \in \underline{R}_m^X$ and $s' \in \underline{R}_{m-1}^X$ are condensed on u_+ and v_+ , and we have

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(s(u_+), s'(u_+)) \text{ and} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(s(v_+), s'(v_+)). \end{aligned}$$

Since $u_+ \triangleright_{m, n-1} v_+$ and $|s| + |s'| \leq n - 2$, we have $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$ by induction on n .

Second case: $a \neq b$, s and s' start with X -blocks. Then $s \in \underline{R}_m^X$ and $s' \in \underline{R}_{m-1}^X$ are condensed on u_+ and v_+ . Without loss of generality, $X_b(u) < X_a(u)$, so we have $r(u) = X_a s(u) = X_b X_a s(u) = X_b r(u)$. In particular, $\text{ord}(r(u), r'(u)) = \text{ord}(r(u'_+), s'(u'_+))$. By Proposition 5.6, we also have $X_b(v) < X_a(v)$, and hence $\text{ord}(r(v), r'(v)) = \text{ord}(r(v'_+), s'(v'_+))$. Since $u \triangleright_{m, n} v$, we have $u'_+ \triangleright_{m, n-1} v'_+$ and we conclude by induction on n since $|r| + |s'| \leq n - 1$.

Third case: $a \neq b$, s and s' start with Y -blocks. This can occur only if $m - 1 \geq 2$. Then $s \in \underline{R}_{m-1}^Y$ and $s' \in \underline{R}_{m-2}^Y$ are condensed on u_- and v_- , $r(u) = s(u_-)$ and $r'(u) = s'(u'_-)$, and the same equalities hold for v . Without loss of generality, we may assume that $X_b(u) < X_a(u)$, and hence $X_b(v) < X_a(v)$ (Proposition 5.6). Let u_0 and v_0 be such that $u = u'_- bu_0 au_+$ and $v = v'_- bv_0 av_+$: then u_0 is the left factor in the a -left decomposition of u'_+ and the right factor in the b -left decomposition of u_- . An analogous statement is true for v_0 . There are two cases, depending on whether s is defined on bu_0 . If this is the case, then $r'(u) < r(u)$. Moreover, we have $u'_+ \triangleright_{m, n-1} v'_+$ and $u_0 \triangleleft_{m-1, n-2} v_0$, so s is defined on bv_0 as well, by Proposition 3.14.

If instead, s is not defined on bu_0 or bv_0 , let p be the longest prefix of s which is defined on bu_0 (and hence on bv_0): then p is either empty or a Y -block and $s = pY_c t$, where c has no occurrence in $u[X_b(u); X_a p(u) - 1]$ (so Y_c is defined on u'_-).

If $Y_c t$ is defined on u'_- , then $r(u) = s(u_-) = Y_c t(u'_-)$, so that

$$\text{ord}(r(u), r'(u)) = \text{ord}(Y_c t(u_-), s'(u_-)).$$

Now $u \triangleright_{m,n} v$ implies $u'_- \triangleright_{m,n-1} v'_-$ by Proposition 3.7, so $Y_c t$ is defined on v'_- and hence we have $\text{ord}(r(v), r'(v)) = \text{ord}(Y_c t(v'_-), s'(v'_-))$ as well. Since $|Y_c t| \leq |s| < |r|$, we conclude by induction that $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

If $Y_c t$ is not defined on u'_- , then let $Y_c q$ be the longest prefix of $Y_c t$ which is defined on u'_- (and hence on v'_-). Then q is either empty or an X -block and $Y_c t = Y_c q X_d t'$. If $d = b$, then $q X_d(u'_- b) = X_b(u)$, so $r(u) = X_b t'(u)$ and similarly, $r(v) = X_b t'(v)$. We conclude by induction on m that $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$ since $X_b t'$ has 2 blocks less than r .

If $d \neq b$, then we have $X_b(u) < X_a p Y_c q X_d(u)$. If $X_d t'$ is defined on bu_0 , then $r(u)$ lies in u_0 and $r'(u)$ lies in u'_- , so $r(u) > r'(u)$. Similarly $r(v) > r'(v)$, and we are done. If instead $X_d t'$ is not defined on bu_0 , then $X_a p Y_c q(u) < X_b(u)$ and $X_a p Y_c q X_d(u) = X_b X_d(u)$, so the condensedness of $r = X_a p Y_c q X_d t'$ on u implies that $X_b X_d t'$ is condensed on u as well. The same holds for v , and we have

$$\begin{aligned} \text{ord}(r(u), r'(u)) &= \text{ord}(X_b X_d t'(u), r'(u)) \text{ and similarly,} \\ \text{ord}(r(v), r'(v)) &= \text{ord}(X_b X_d t'(v), r'(v)). \end{aligned}$$

We conclude by induction on m since $X_b X_d t'$ has 2 blocks less than r .

Fourth case: $a \neq b$, s and s' start with different directions. Without loss of generality, we may assume that s starts with an X -block and s' starts with a Y -block. Since r starts with 2 X -letters, the number of alternating blocks of r is less than $|r| - 1 \leq n - 3$. Therefore if $n = m + 1$, $r \in \underline{R}_{m-2}^X$ and $r' \in \underline{R}_{m-1}^X$, a case that can be decided by induction on m . So we now assume that $n \geq m - 2$.

If $X_b(u) < X_a(u)$, then the same inequality holds in v (by Proposition 5.6) and we have $r'(u) < r(u)$ and $r'(v) < r(v)$. If instead $X_a(u) < X_b(u)$ and $X_a(v) < X_b(v)$, then the b -left factorizations of u_+ and v_+ are of the form $u_+ = u_0 b u'_+$ and $v_+ = v_0 b v'_+$.

Several cases arise, according to whether s and s' are defined (and condensed) on u_0 or not. We have $u_+ \triangleright_{m,n-1} v_+$ and $u_0 \triangleright_{m,n-2} v_0$ by Lemma 3.7. It follows as usual that s and s' are defined on v_0 if and only if they are defined on u_0 . If s is not defined on u_0 then the order types $\text{ord}(r(u), r'(u))$ and $\text{ord}(r(v), r'(v))$ are both $>$. Therefore, from now on we can assume that s is defined on u_0 and v_0 .

If s' is defined on u_0 then we can chop off $u_- a$ from u , $v_- a$ from v , and X_a from r : $\text{ord}(r(u), r'(u)) = \text{ord}(s(u_+), r'(u_+))$ and $\text{ord}(r(v), r'(v)) =$

$\text{ord}(s(v_+), r'(v_+))$. Since $u_+ \triangleright_{m,n-1} v_+$, $\text{ord}(s(u_+), r'(u_+))$ and $\text{ord}(s(v_+), r'(v_+))$ are equal by induction on n , and hence $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. ■

If s' is not defined on u_0 , then, as in the third case, we have to split the ranker s' at those points at which it crosses the position $X_a(u)$. Let $X_b s' = p_1 q_1 \cdots p_k q_k$ such that all p_i are defined on u_0 and all p_i are starting with an X-letter followed by a (possibly empty) Y-block. The sole exception is p_k which might contain further blocks. Moreover, each p_i is the maximal prefix of $p_i q_i \cdots p_k q_k$ which is defined on u_0 . All q_i are defined on $u_- a$ and all q_i are starting with a Y-letter followed by a (possibly empty) X-block. The sole exception is q_k which might be empty or which might contain further blocks. Each q_i is the maximal prefix of $q_i p_{i+1} \cdots p_k q_k$ which is defined on $u_- a$. Since $u_- a \triangleright_{m,n-1} v_- a$ (Lemma 3.7) and $u_0 \triangleleft_{m-1,n-2} v_0$, the same definedness and maximality properties hold on $v_- a$ and v_0 .

If q_k is empty, then $k \geq 2$ and p_1 and q_1 are non-empty. We see that $\text{ord}(r(u), r'(u)) = \text{ord}(s(u_0), p_k(u_0))$ and $\text{ord}(r(v), r'(v)) = \text{ord}(s(v_0), p_k(v_0))$. ■ By induction on n , we have $\text{ord}(s(u_0), p_k(u_0)) = \text{ord}(s(v_0), p_k(v_0))$, and hence $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$.

Finally, if q_k is non-empty, then we have $r(u) > r'(u)$ and $r(v) > r'(v)$. □

5.2.3. Completing the proof. Let us (at last!) verify that, if $u \triangleright_{m+1,2n} v$ or $u \triangleright_{m+1,2n} v$, then Properties **(WI 2c)** and **(WI 3c)** hold for m, n . By symmetry, we simply handle the case where $u \triangleright_{m+1,2n} v$.

To verify Property **(WI 2c)**, we consider rankers $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m-1,n-1}$ that are condensed on u and v . If both start with X-blocks, Proposition 5.7 shows that $\text{ord}(r(u), r'(u)) = \text{ord}(r(v), r'(v))$. If both start with Y-blocks, the same proposition allows us to conclude, after observing that we have $u \triangleleft_{m,2n-1} v$. And if r and r' start with different direction blocks, we conclude by Proposition 5.5.

To verify Property **(WI 3c)**, we consider rankers $r \in \underline{R}_{m,n}$ and $r' \in \underline{R}_{m,n-1}$ that end with different directions, and that are condensed on u and v . If r and r' start with different direction blocks, we again conclude by Proposition 5.5. If both start with X-blocks, then they must have different number of alternations, so we have $r \in R_{m_1, n_1}^X$ and $r' \in R_{m_2, n_2}^X$ for some $n_1 \leq n$, $n_2 \leq n-1$ and for distinct values $m_1, m_2 \leq m$. In particular, one of m_1 and m_2 is less than or equal to $m-1$, and we can apply Proposition 5.7.

We proceed similarly if r and r' both start with Y-blocks, after observing that $u \triangleleft_{m,2n-1} v$.

6. CONSEQUENCES

6.1. Decidability results. The main consequence we draw of Theorem 5.1 and of the decidability of the pseudovarieties \mathbf{R}_m and \mathbf{L}_m is summarized in the next statement.

Theorem 6.1. *Given an FO²-definable language L , one can compute an integer m such that L is FO _{m} ²-definable, possibly FO _{$m+1$} ²-definable, but not*

FO_{m-1}^2 -definable. That is: we can decide the quantifier alternation level of L within one unit.

Proof. Let $L \in \mathcal{FO}^2$ and let M be its syntactic monoid. Since each pseudovariety $\mathbf{R}_m \cap \mathbf{L}_m$ is decidable (Proposition 3.21), we can compute the largest m such that $M \notin \mathbf{R}_m \cap \mathbf{L}_m$. By Theorem 5.1, $M \in \mathbf{R}_{m+1} \cap \mathbf{L}_{m+1} \subseteq \mathbf{FO}_{m+1}^2$ and hence L is FO_{m+1}^2 -definable. On the other hand, $M \notin \mathbf{FO}_{m-1}^2 \subseteq \mathbf{R}_m \cap \mathbf{L}_m$. \square

Let us also record the following consequences of Proposition 3.21, Proposition 4.1 and the decidability of $\mathbf{R}_2 \vee \mathbf{L}_2$ (discussed in Example 5.2).

Proposition 6.2. *The classes $\underline{\mathcal{TL}}_1^X = \underline{\mathcal{TL}}_1^Y = \underline{\mathcal{TL}}_1 = \mathcal{FO}_1^2$, $\underline{\mathcal{TL}}_2^X$, $\underline{\mathcal{TL}}_2^Y$ and $\underline{\mathcal{TL}}_2$ are decidable.*

6.2. Infinite and collapsing hierarchies. The fact that the \mathbf{R}_m and \mathbf{L}_m form strict hierarchies (Proposition 3.21), together with Theorem 5.1, proves that the \mathcal{FO}_m^2 hierarchy is infinite. Weis and Immerman had already proved this result by combinatorial means [30, Theorem 4.11], whereas our proof is algebraic. From that result on the \mathcal{FO}_m^2 , it is also possible to recover the strict hierarchy result on the \mathbf{R}_m and \mathbf{L}_m and the fact that their union is equal to \mathbf{DA} .

By the same token, Corollary 3.16 and Proposition 4.3 show that the $\underline{\mathcal{TL}}_m$ (resp. $\underline{\mathbf{TL}}_m$) hierarchy is infinite and that its union is all of \mathcal{FO}^2 (resp. \mathbf{DA}).

Proposition 6.3. *The hierarchies \mathcal{FO}_m^2 and $\underline{\mathcal{TL}}_m$ are infinite, and their union is all of \mathcal{FO}^2 .*

Similarly, the fact (stated in Proposition 3.21) that an m -generated element of \mathbf{DA} lies in $\mathbf{R}_{m+1} \cap \mathbf{L}_{m+1}$, shows that an FO^2 -definable language in A^* lies in $\mathcal{R}_{|A|+1} \cap \mathcal{L}_{|A|+1}$, and hence in $\mathcal{FO}_{|A|+1}^2$ – a fact that was already established by combinatorial means by Weis and Immerman [30, Theorem 4.7]. It also shows that such a language is in $\underline{\mathcal{TL}}_{2|A|+1}$ by Proposition 4.3.

Proposition 6.4. *A language $L \subseteq A^*$ is FO^2 -definable if and only if it is $\text{FO}_{|A|+1}^2$ -definable. And it is \mathbf{TL} -definable if and only if it is both $\underline{\mathbf{TL}}_{2|A|+1}^X$ and $\underline{\mathbf{TL}}_{2|A|+1}^Y$ -definable.*

Even though we arrived at Proposition 6.4 by algebraic means, it is interesting to note that its statement reflects the following combinatorial property (an idea that was already used by Weis and Immerman [30, Theorem 4.7]).

Lemma 6.5. *A ranker that is condensed on a word on alphabet A , has at most $|A|$ alternating blocks.*

Proof. Let u be a word and let r be a ranker that is condensed on u . Without loss of generality, we may assume that $r \in R_{m,n}^X$, say

$$r = X_{a_1} \cdots X_{a_{k_1}} Y_{a_{k_1+1}} \cdots Y_{a_{k_2}} \cdots Z_{a_{k_{m-1}+1}} \cdots Z_{a_{k_m}}$$

with $0 < k_1 < k_2 < \dots < k_m = n$ and $Z = X$ (resp. Y) if m is odd (resp. even). By definition of condensed rankers (and with the notation in that definition, see Section 3), the interval I_{k_h} is of the form $(i_{k_h-1}, X_{a_{k_h}}(u, i_{k_h-1}))$ if h is odd, of the form $(Y_{a_{k_h}}(u, j_{k_h-1}), j_{k_h-1})$ if h is even. In either case, $a_{k_{h+1}}$ occurs in u within the interval I_{k_h} but a_{k_h} does not. Since the intervals I_{k_h} are nested, it follows that the letters $a_{k_1}, a_{k_2}, \dots, a_{k_m}$ are pairwise distinct, and hence $m \leq |A|$. \square

6.3. Infinite hierarchies and unambiguous polynomials. Finally we note the following refinement on [12, Proposition 4.6]. One of the classical (and one of the earliest) results concerning the languages recognized by monoids in **DA** (namely \mathcal{FO}^2) is the following: they are exactly the disjoint unions of unambiguous products of the form $B_0^* a_1 B_1^* \dots a_k B_k^*$, where each B_i is a subset of A (Schützenberger [19], see also [24, 25]). Deterministic and co-deterministic products (see Section 3.4) are easily seen to be particular cases of unambiguous products. Propositions 3.19 and 3.21 imply the following statement.

Proposition 6.6. *The least variety of languages containing the languages of the form B^* ($B \subseteq A$) and closed under visibly deterministic and visibly co-deterministic products, is \mathcal{FO}^2 .*

More precisely, every unambiguous product of the form $B_0^ a_1 B_1^* \dots a_k B_k^*$, where each B_i is a subset of A , can be expressed in terms of the B_i^* and the a_i using only Boolean operations and at most $|A| + 1$ alternated applications of visibly deterministic and visibly co-deterministic products – starting with a visibly deterministic (resp. co-deterministic) product.*

The analogous, but weaker statement with the word *visibly* deleted was proved in [12] by algebraic means, and independently by Lodaya, Pandya and Shah using logical and combinatorial arguments [13].

REFERENCES

- [1] M. Adler and N. Immerman. An $n!$ lower bound on formula size. *ACM Transactions on Computational Logic*, 4:296–314, 2003.
- [2] Jorge Almeida. *Finite Semigroups and Universal Algebra*. World Scientific, Singapore, 1994.
- [3] Sung Cho and Dung T. Huynh. Finite automaton aperiodicity is PSPACE-complete. *Theoretical Computer Science*, 88:96–116, 1991.
- [4] Volker Diekert and Paul Gastin. Pure future local temporal logics are expressively complete for Mazurkiewicz traces. *IC*, 204:1597–1619, 2006. Conference version in LATIN 2004, LNCS 2976, 170–182, 2004.
- [5] Volker Diekert and Paul Gastin. First-order definable languages. In J. Flum, E. Grädel, and Th. Wilke, editors, *Logic and Automata: History and Perspectives*, Texts in Logic and Games, pages 261–306. Amsterdam University Press, 2008.
- [6] Volker Diekert, Paul Gastin, and Manfred Kufleitner. A survey on small fragments of first-order logic over finite words. *International Journal of Foundations of Computer Science*, 19:513–548, 2008.

- [7] Kousha Etessami, Moshe Y. Vardi, and Thomas Wilke. First-order logic with two variables and unary temporal logic. *Information and Computation*, 179(2):279–295, 2002.
- [8] M. Grohe and N. Schweikardt. The succinctness of first-order logic on linear orders. *Logical Methods in Computer Science*, 1, 2005.
- [9] Neil Immerman. *Descriptive Complexity*. Springer, 1999.
- [10] Manfred Kufleitner. Polynomials, fragments of temporal logic and the variety DA over traces. *Tcs*, 376:89–100, 2007. Special issue DLT 2006.
- [11] Manfred Kufleitner and Pascal Weil. On \mathbf{FO}^2 quantifier alternation over words. In *Mathematical Foundations of Computer Science (MFCS 2009)*, number 5734 in Lecture Notes in Computer Science, pages 513–524. Springer-Verlag, 2009.
- [12] Manfred Kufleitner and Pascal Weil. On the lattice of sub-pseudovarieties of \mathbf{DA} . *Semigroup Forum*, 81:243–254, 2010.
- [13] Kamal Lodaya, Paritosh K. Pandya, and Simoni S. Shah. Marking the chops: an unambiguous temporal logic. In *IFIP TCS 2008*, pages 461–476, 2008.
- [14] Robert McNaughton and Seymour Papert. *Counter-Free Automata*. The MIT Press, Cambridge, Mass., 1971.
- [15] Jean-Éric Pin. *Varieties of Formal Languages*. North Oxford Academic, London, 1986.
- [16] Jean-Éric Pin. Expressive power of existential first-order sentences of Büchi’s sequential calculus. *Discrete Mathematics*, 291(1-3):155–174, 2005.
- [17] Jean-Éric Pin and Pascal Weil. Profinite semigroups, Mal’cev products and identities. *Journal of Algebra*, 182:604–626, 1996.
- [18] Marcel Paul Schützenberger. On finite monoids having only trivial subgroups. *Information and Control*, 8:190–194, 1965.
- [19] Marcel Paul Schützenberger. Sur le produit de concaténation non ambigu. *Semigroup Forum*, 13:47–75, 1976.
- [20] Thomas Schwentick, Denis Thérien, and Heribert Vollmer. Partially-ordered two-way automata: A new characterization of DA. In Werner Kuich, Grzegorz Rozenberg, and Arto Salomaa, editors, *Proc. of the 5th Int. Conf. on Developments in Language Theory (DLT)*, volume 2295 of *Lecture Notes in Computer Science*, pages 239–250. Springer, 2001.
- [21] Howard Straubing. *Finite Automata, Formal Logic, and Circuit Complexity*. Birkhäuser, Boston, Basel and Berlin, 1994.
- [22] Howard Straubing and Pascal Weil. An introduction to automata theory. In Deepak D’Souza, editor, *Modern applications of automata theory*, volume 2 of *I.I.Sc. Monographs*. World Scientific, to appear.
- [23] Howard Straubing and Pascal Weil. Varieties. In Jean-Eric Pin, editor, *Handbook of Finite Automata*. European Math. Society, to appear.
- [24] Pascal Tesson and Denis Thérien. Diamonds are forever: The variety DA. In Gracinda Maria Dos Gomes Moreira Da Cunha, Pedro Ventura Alves Da Silva, and Jean-Éric Pin, editors, *Semigroups, Algorithms, Automata and Languages, Coimbra (Portugal) 2001*, pages 475–500. World Scientific, 2002.
- [25] Pascal Tesson and Denis Thérien. Logic meets algebra: the case of regular languages. *Logical Methods in Computer Science*, 3(1):1–37, 2007.
- [26] Denis Thérien and Thomas Wilke. Over words, two variables are as powerful as one quantifier alternation. In *STOC*, pages 234–240, 1998.
- [27] Wolfgang Thomas. Classifying regular events in symbolic logic. *Journal of Computing Systems and Science*, 25:360–376, 1982.
- [28] Peter Trotter and Pascal Weil. The lattice of pseudovarieties of idempotent semi-groups and a non-regular analogue. *Algebra Universalis*, 37:491–526, 1997.
- [29] Philipp Weis and Neil Immerman. Structure theorem and strict alternation hierarchy for \mathbf{FO}^2 on words. In Jacques Duparc and Thomas A. Henzinger, editors, *Computer*

Science Logic, 21st International Workshop, CSL 2007, 16th Annual Conference of the EACSL, Lausanne, Switzerland, September 11-15, 2007, Proceedings, volume 4646 of *Lecture Notes in Computer Science*, pages 343–357. Springer, 2007.

- [30] Philipp Weis and Neil Immerman. Structure theorem and strict alternation hierarchy for FO^2 on words. *Logical Methods in Computer Science*, 5:1–23, 2009.