



**HAL**  
open science

## **Medulloblastoma in childhood: revisiting intrathecal therapy in infants and children**

Sharon Conroy, Martin Garnett, Michael Vloeberghs, Richard Grundy, Ian Craven,  
David Walker

### ► **To cite this version:**

Sharon Conroy, Martin Garnett, Michael Vloeberghs, Richard Grundy, Ian Craven, et al.. Medulloblastoma in childhood: revisiting intrathecal therapy in infants and children. *Cancer Chemotherapy and Pharmacology*, 2009, 65 (6), pp.1173-1189. <10.1007/s00280-009-1127-1>. <hal-00568268>

**HAL Id: hal-00568268**

**<https://hal.science/hal-00568268v1>**

Submitted on 23 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

***Medulloblastoma in childhood***  
**- *revisiting intrathecal therapy in infants and children***

Sharon Conroy, Lecturer in Paediatric Clinical Pharmacy  
Martin Garnett, Associate Professor Pharmaceutical Sciences  
Michael Vloeberghs, Professor Paediatric Neurosurgery  
Richard Grundy, Professor Paediatric Neuro-Oncology and Cancer Cell Biology  
Ian Craven, Research student  
David Walker, Professor Paediatric Oncology

Children's Brain Tumour Research Centre,  
University of Nottingham,  
University Park,  
Nottingham  
NG7 2RD  
UK

Corresponding author: Professor David A Walker

[david.walker@nottingham.ac.uk](mailto:david.walker@nottingham.ac.uk)

Tel: +44 1158 230632.

The authors can confirm that this work is original. It was presented as a poster at the 11<sup>th</sup> International Symposium on Pediatric Neuro-Oncology and the abstract published as below:

Conroy S, Craven I, Garnett M, Barrett S, Punt J, Parker T, Walker DA. An extensive review of chemotherapy agents suitable for intrathecal (IT) therapy in primitive neuro ectodermal tumour (PNET). Abstract EXP 7 from 11<sup>th</sup> International Symposium on Pediatric Neuro-Oncology. Neuro-Oncology 2004;6:428

There are no conflicts of interest to declare. This work was conducted without any financial assistance outside of the University of Nottingham

The manuscript has been read and approved by all the authors. Each author believes that the manuscript represents honest work.

***Medulloblastoma in childhood***  
**- *revisiting intrathecal therapy in infants and children***

***Sharon Conroy, Martin Garnett, Michael Vloeberghs, Richard Grundy, Ian Craven  
and David Walker***

## ***Abstract***

***Introduction:*** Intrathecal chemotherapy is being explored in medulloblastoma in pre-school children as part of brain-sparing strategies and as an alternative to unacceptably neurotoxic cranio-spinal radiotherapy. The range of drugs suitable for this route of administration is restricted by the lack of research evidence of pharmacological suitability and efficacy of other drugs in medulloblastoma.

***Methods:*** Ideal clinical, biological, physicochemical and pharmaceutical properties for intrathecal administration were defined through literature review of pharmaceutical texts, Medline, Embase and consulting the manufacturers. 126 chemotherapy agents were assessed against these criteria by searching the academic domain of pharmaceutical texts, computer databases and consultation with manufacturers.

***Results:*** Of 126 candidate drugs, 99 were rejected because of documentation of their irritant nature, neurotoxicity and requirement for hepatic activation in standard pharmaceutical texts. Fifty were rejected for a single identifiable reason including, neurotoxicity (n=24), irritant (n=15), needs enzyme activation (n=5), clinical evidence of intrathecal neurotoxicity (n=4) and no evidence of tumour-specific efficacy (n=2). Where two reasons were cited the justifications were: neurotoxic and irritant (n=3) and needs activation and systemic administration results in equivalent concentration (n=1). Twenty seven drugs remained of which 12 were selected as eligible for further clinical investigation, and 15 were selected for further pre-clinical investigation.

***Conclusions:*** The predetermined criteria were not applicable, in their entirety, in the majority of drugs, due to lack of information in the academic domain, emphasising the importance of a more open approach to sharing basic drug information. The prioritised list of 12 candidate drugs for clinical trial and 15 for pre-clinical investigation justify that a concerted research effort in this area of practice is made. (317 words)

## **Introduction**

Medulloblastoma, the commonest malignant CNS tumour of childhood [125], has the capacity to disseminate through the lepto-meninges, presenting at diagnosis or at relapse after treatment. Leptomeningeal metastases (LM) grow at the interface of the brain and spinal fluid. They are thought to occur as a result of interplay between operative disruption of tumour cells contaminating the CSF and inherent biological capacity for tumour cells to migrate, adhere and invade neural tissue. Where LM are present at diagnosis biological factors are increasingly being associated with this dissemination and poorer cure rates with standard therapy. Systemic chemotherapy has an established role in medulloblastoma with evidence of chemosensitivity, enhanced survival rates and rising survival rates reported by trials groups and population registries [80] in recent eras.

The use of intrathecal chemotherapy in childhood leukaemia and CNS tumours has been the focus of reviews [17, 49, 108] and reports of phase 1 and 2 studies [17]. In leukaemia the intensification of systemic and intrathecal therapy has permitted exclusion of CNS radiotherapy for the majority of newly diagnosed cases with very low CNS relapse rates, acceptable late toxicity using methotrexate alone or in combination with cytosine and hydrocortisone. The intrathecal route in leukaemia has been demonstrated to bypass the blood brain barrier (BBB), enhance CSF drug concentrations reducing systemic exposure and, therefore, toxicity. Selection of drugs suitable for intrathecal administration and with evidence of sensitivity to primary CNS tumours is a challenge in medulloblastoma, therefore. Preliminary evidence of the efficacy of intrathecal therapy is emerging. Slavc reported extensive use of intrathecal therapy at relapse using etoposide, mafosfamide and others [120, 121]. Other studies report the feasibility of this route of drug administration at relapse and as part of palliative care with infrequent complications and prolonged symptom free intervals. Rutkowski reported very high survival rates in pre-school children treated with chemotherapy-only strategy including intraventricular methotrexate [109]. The proposal that intrathecal therapy be tested in a prospective European trial of medulloblastoma in this young age group is currently the focus of debate [139].

In order to further develop an enhanced evidence-base for the selection of drugs that are effective against medulloblastoma, suitable for administration by the intrathecal route, and non-toxic to the nervous system, we developed criteria for the ideal intrathecal agent. Following this a systematic literature review was conducted using these criteria to identify candidate drugs worthy of further trial for intrathecal use in medulloblastoma.

## **Method**

### *Identification of candidate drugs*

A comprehensive list of chemotherapy agents was extracted from a drug pharmacopoeia [87]. Properties of the ideal intrathecal drug were identified based upon literature review [4, 10, 13,

33, 53, 102, 108, 132, 136, 143, 150, 151] and grouped according to: "ideal clinical properties", "ideal biological properties" and "ideal physico-chemical and pharmaceutical properties".

### ***Ideal Clinical Properties***

#### *Non-irritant*

The drug would need to be non-irritant to avoid chemical arachnoiditis or meningitis. Any drug reported in any of our reference sources to cause irritation or thrombophlebitis was therefore eliminated from further consideration.

#### *Neurotoxicity*

Any drug given intrathecally has a significant risk of neurotoxicity. Evidence for neurotoxicity when given systemically or intrathecally was determined as an unacceptable risk for further evaluation of CSF administration. It was not possible to grade the neurotoxicities as they were variably reported. Some drugs which are not usually neurotoxic when given systemically at conventional doses, may be toxic if given intra-arterially, intrathecally or in very high doses [130]. Any drug reported in any of our reference sources to cause neurotoxicity was therefore eliminated from further consideration. Methotrexate and cytosine arabinoside, whilst suitable for IT delivery, have only been studied in medulloblastoma with concurrent systemic chemotherapy agents, the chemosensitivity of medulloblastoma to these drugs is, therefore, unknown.

#### *Tumour sensitivity / mechanism of action*

Evidence of activity of the drug against medulloblastoma, PNET or leptomeningeal carcinomatosis is required to warrant further investigation. A drug with a mechanism of action effective at the CSF/leptomeningeal interface would offer a therapeutic advantage.

#### *Methotrexate and Cytosine Arabinoside*

Interestingly, the strict application of these criteria led to the exclusion of methotrexate and cytarabine (in its standard formulation) from the list of potential candidates. We are aware that these agents are commonly used intrathecally. The application of 'ideal criteria' which identifies lack of chemosensitivity data and substantial evidence of neurotoxicity led to their rejection from our ideal list, for the purposes of this review.

### ***Ideal Biological properties***

#### *CSF transport system*

Some drugs are removed from the CSF by facilitated diffusion using carrier transport systems such as those for organic ions, quaternary ammonium compounds, organic bases and acids, also transporters conferring multi-drug resistance such as P-glycoprotein, multi-drug resistance protein 1, monocarboxylic acid and organic ion transporters [126]. The ideal

intrathecal agent will have no active transport system that is capable of removing it from the CSF.

#### *Cell cycle non-specific agent*

Cells in the process of dissemination may be in a non-cycling, G<sub>0</sub>, or slow cycling phase. Drugs which act at specific stages of the cell cycle will need to be in contact with the cells for sufficient time for the cells to go through active cell division, to ensure an effective cell kill. An ideal agent would therefore be cell cycle non-specific in its action.

### ***Ideal Physicochemical and pharmaceutical properties***

#### *Active in CSF*

Drugs requiring enzyme activation (e.g. cyclophosphamide) are unlikely to be active since there are very few enzyme systems present in the CSF. Drugs administered in their active form or with predictable activation (e.g. hydrolysis) are proposed. Those requiring other processes of activation are excluded.

#### *Factors affecting BBB permeability*

Drugs in the CSF are likely to be in equilibrium with plasma as determined by the characteristics of the BBB and the drug's physical characteristics. Factors that reduce a drug's capacity to diffuse across the BBB and therefore limit drug efflux include: low lipophilicity, high hydrophilicity, ionised state at CSF pH of 7.3 and molecular weight >700 Da [53]. Such properties should therefore enhance sustained CSF drug levels and are therefore preferred for intrathecal therapy.

#### *Protein binding*

Ideally drugs normally protein bound are excluded from crossing the BBB due to molecular size, direct injection would overcome this and sustain their presence in CSF. This criteria has, however, not been used as the information was not available for the majority of drugs.

#### *Formulation for CSF administration*

The drug must be soluble in the appropriate concentration for intrathecal administration, and in a vehicle which is suitable for this route, including bio-compatibility with surgical delivery systems such as ventriculostomy reservoirs.

(Insert Table 1 here - Summary of clinical, biological and physicochemical / pharmaceutical properties used to justify selection or rejection of drugs)

### ***Literature search strategy and selection criteria***

An initial search for the properties of each drug on the candidate list was undertaken by screening standard pharmaceutical texts [9, 30, 87] and by consulting the manufacturer.

Some information was obtained from these sources. This permitted immediate elimination of a number of drugs due to evidence of irritancy, neurotoxicity and the need for liver enzyme activation. However details of ionisation state and degree of protein binding for many drugs was unavailable despite direct approaches to manufacturers.

For the remaining drugs, Medline (1966 - present) and Embase (1980 - present) searches were performed. Each drug name was separately combined with the following terms: 'medulloblastoma', 'PNET' and 'leptomeningeal carcinomatosis', used in the 'exploded' form. Where hits exceeded 50 for a combination of terms, the search was further narrowed by combining that search with 'intrathecal administration'. The full text of papers with relevant abstracts were obtained and physicochemical information was extracted as well as:

- evidence of the drug's activity against medulloblastoma, PNET and leptomeningeal carcinomatosis in vitro and in vivo
- absence or presence of neurotoxicity reported from systemic or intrathecal administration
- documentation of intrathecal administration in vivo and in vitro.

## **Results**

One hundred and twenty-six drugs were initially identified for investigation. 35 were immediately rejected after consulting standard pharmaceutical texts where there was documentation of irritant qualities, neurotoxicity or the need for enzyme activation of a pro-drug. Literature searches for the remaining 91 drugs identified 33,627 hits. Screening of abstracts reduced this to 200 full papers which were read, justifying the rejection of a further 64 drugs (see Table 2a & b).

(Insert Table 2 here – List of chemotherapy drugs considered with selection / rejection status and rejection justification)

### *Rejected drugs*

Of the 126 licensed and investigational anti-cancer chemotherapy agents identified, 99 drugs were excluded (Table 2a & b). In 45 cases rejection was due solely to the lack of information. The remaining 54 drugs were rejected for a single reason (n=50) or a combination of two reasons (n=4). Where a single reason is cited the justifications were: neurotoxicity (n=24), irritant (n=15), needs activation (n=5), neurotoxic when given intrathecally (n=4), lack of activity in relevant tumour type (n=2). Where two reasons were cited, the justifications were: neurotoxic and irritant (n=3), and needs activation and systemic administration results in equivalent concentration (n=1).

### *Prioritised drugs for intrathecal use*

From our initial list of 126 drugs, 12 drugs have been identified as suitable for further testing by the intrathecal route in medulloblastoma. Fifteen other potential candidates lack sufficient information to currently justify exploration of intrathecal administration, but warrant further research. The 27 drugs are therefore candidates for further testing via the intrathecal route focussed upon in childhood medulloblastoma and other leptomeningeal malignancies.

The most promising 12 drugs for PNET/medulloblastoma with prior reported clinical experience are detailed in Table 3 with justification of their potential for future research.

(Insert Table 3 here – Drugs currently eligible for consideration for trial by intrathecal administration for medulloblastoma)

#### *Evaluation in progress*

Liposomal cytarabine and mafosfamide are currently undergoing phase 2 trials by the intrathecal route in children with brain tumours with leptomeningeal spread. Liposomal cytarabine has now been associated with significant neurotoxicity, particularly in adult patients also treated with systemic cytarabine [16]. It may be better tolerated in children. Comitant systemic corticosteroid treatment is recommended. Further studies are needed to establish optimal use of this drug.

These should be the first drugs to undergo further evaluation in phase 3 studies in order to further evaluate their efficacy and toxicity profiles and optimise dose and administration regimens.

#### *Evaluation required in Phase 2 trials*

Carboplatin, etoposide, spartaject, busulfan and nimustine are agents suitable for treatment of PNET and are therefore also suitable for further investigation by the intrathecal route. Topotecan is undergoing intrathecal phase 2 studies in children and adults with refractory neoplastic meningitis; however chemical arachnoiditis has been reported as a dose limiting toxicity. Etoposide and nimustine have already been tested clinically with some encouraging results; however refinement of drug delivery systems may be necessary to prolong exposure in order to overcome their rapid removal from the CSF, their currently observed efficacy may, therefore, be sub-optimal.

#### *Phase 1 studies required*

Floxuridine and 4-hydroperoxycyclophosphamide have been tested intrathecally in limited studies and have shown promise in relevant tumour types. The intrathecal use of these two agents should also be further explored.

#### *Pre-clinical evaluation required*

Diaziquone is lipophilic and has low ionisation at CSF pH. Together these physical properties suggest it may not remain in the CSF unless delivered in a sustained release formulation or as an infusion. It is active in PNET and leptomeningeal carcinomatosis and has been used intrathecally in human studies. Mercaptopurine has been tested by the intraventricular route in animals and children with leptomeningeal dissemination of ALL but there is no evidence of efficacy in primary brain tumours. Rubitecan has been shown to be safe & effective in a rat model against human glioblastoma multiforme neoplastic meningitis.

Temozolomide is effective in medulloblastoma when given orally. A microcrystalline preparation has been used intrathecally in animal studies justifying further study if an appropriate formulation could be developed. Its lipophilicity would favour a sustained release formulation or an infusion

#### *Pre-clinical animal evaluation*

The nine drugs in Table 4 have documented activity in the relevant tumour types but no information was found on intrathecal use. These drugs would be worthy of intrathecal testing.

(Insert Table 4 here - Drugs with documented activity in the relevant tumour types but no information on intrathecal use)

The drugs in Table 5 have been tested in studies in the relevant tumour types by the intrathecal route (some animal, some human) with mixed reports of toxicity and efficacy. With the exception of bleomycin where neurotoxicity seems to be a major problem, these may well be worthy of further research. Carmustine is licensed in a sustained release preparation designed to be implanted into tumour resection cavities as an adjunct to surgery for GBM after relapse. However, it is documented also to have serious neurotoxicity when given systemically and therefore has not been prioritised in the list of drugs for further study.

Gemcitabine has been tested by the intrathecal route in a patient with leptomeningeal carcinomatosis from non small cell lung cancer and has been used intrathecally in non-human primate models [35, 45, 67]. Based upon this, a phase 1 clinical trial of intrathecal gemcitabine has been conducted in ten patients with neoplastic meningitis including one with medulloblastoma. However, dose limiting neurotoxicity was observed in patients who had previously received CNS-directed therapies [15] The authors concluded that the potential for severe neurotoxicity precludes further development of gemcitabine for intrathecal administration.

(Insert Table 5 here - Drugs tested by the intrathecal route (some animal, some human) with mixed reports of toxicity and efficacy)

## ***Discussion***

This work helps draw three important conclusions. First, many of the proposed ideal criteria, have not been applicable because of inadequate data in the academic and commercial information domains. A more open shared strategy between pharma and academic researchers of basic pharmacological data is required. Secondly, using supplementary clinical criteria, the systematic review has identified 12 drugs suitable for immediate intrathecal use in medulloblastoma, 15 other candidate drugs were identified for further evaluation prior to human use. Finally, the extended list of agents, identified here as suitable for intrathecal use, justifies further efforts to explore their role as part of brain sparing strategies in clinical trials.

## ***Ideal criteria***

The pre-determined criteria were applicable in their entirety in only eight drugs resulting in the current drug selection being based primarily upon alternative criteria including: a) clinical evidence of tumour chemo-sensitivity, b) previously reported intrathecal use and c) reported evidence of neurotoxicity after systemic or intrathecal administration. The overwhelming reason for rejection was because of either insufficient or inconsistent information in the academic domain (Table 2). It remains unclear why this basic physicochemical data is unavailable, highlighting the importance of promoting a more open ethic between pharma and academic groupings exploring novel applications of existing cancer drugs and in particular for intrathecal administration. This large category of known, rejected drugs as well as a presumed larger number of unknown candidate agents in industrial archives constitutes an untapped resource. In contrast, drugs with inconsistent data, but promising early laboratory or clinical evidence of suitability, are a group where further efforts for their evaluation are justified. Furthermore, as new biological agents are developed which affect tumour behaviour via non cytotoxic mechanisms, the lack of publication of physico-chemical criteria or clinical pharmacological data will slow down the process of their selection or rejection for intrathecal use.

*Drug selection:* From this review, 12 agents were identified as potentially suitable for human use. The three largest previous reviews of this treatment approach do specify optimal drug criteria for intrathecal use but do not specify the method by which they developed their proposed drug selections for intrathecal use.

(Insert Table 6 here - Comparison of candidate intrathecal agents by previous published reviews)

The current review has identified four additional drugs: carboplatin, floxuridine, 4-hydroperoxycyclophosphamide and rubitecan, yet rejected cytarabine (in its standard formulation) and methotrexate because of evidence of neurotoxicity. The extensive experience with these two latter drugs for prophylactic, therapeutic and palliative strategies in acute

leukaemia and lymphoma has permitted comprehensive reporting of neurotoxicity. It has been proposed that enhanced and prolonged drug levels due to reduced permeability of the BBB arising from tumour involvement at sites of CSF efflux is a hypothesis for this observed effect [23, 24, 55, 86]. Carmustine in its wafer preparation (<sup>TM</sup>Gliadel) is used for interstitial therapy for GBM at relapse (**ref**). Prior reports of neurotoxicity after intra-arterial administration might have excluded this drug. Its lipophilicity makes it unsuitable as an intrathecal agent without adaptation to its formulation to slow its release or as an infusion. Its current application within a sustained release wafer is, therefore, compatible with the 'ideal criteria' based upon physicochemical properties but not via the intrathecal route aimed at generalised CSF distribution.

Nine drugs were identified with evidence of efficacy against medulloblastoma but insufficient information about their suitability for intrathecal use (Table 4). More information about their physicochemical / cycle specific and neurotoxicity properties is required before they could be accepted/rejected for evaluation by the ideal criteria. The information that was missing was physicochemical (7), cell cycle specificity (4) and neurotoxicity (7). The previous reviews [17, 49, 108] made no suggestions for this category of agents. Drugs, tested by the intrathecal route but with mixed reports of toxicity (n=7 Table 5), were based upon clinical case reports from both animal and human experimentation, limiting the scope of this group as a source of new drugs for testing. There were conflicting reports of neurotoxicity. In only one drug was there a complete physico-chemical dataset (dacarbazine). In three drugs (bleomycin, busulfan and carmustine) drug formulation was reported to influence neurotoxicity and efficacy. This group of drugs highlights the importance of selecting or developing drug formulations that maximise efficacy and minimise neurotoxicity.

Having identified a short list of 12 drugs suitable for human trial via the intrathecal route (table 3) an attempt to select those suitable for immediate use has been made based upon currently available data. Carboplatin, etoposide and nimustine are identified as having the strongest evidence base for intrathecal use. Liposomal cytarabine (Depocyte) and mafosfamide are currently undergoing phase 2 trials; their efficacy and toxicity will need to be judged when given without concomitant chemotherapy or radiotherapy. Mafosfamide is unlicensed and is therefore unavailable for clinical use in trial or therapy (Personal communication: Irene Slavic - [irene.slavic@akh-wien.ac.at](mailto:irene.slavic@akh-wien.ac.at); Dr Susan Blaney: [smblaney@txccc.org](mailto:smblaney@txccc.org)). Liposomal cytarabine (Depocyte) is being extensively investigated in adult studies in a variety of cancer types. Preliminary information has identified substantial evidence of neurotoxicity [65, 81]. In selecting any of these drugs for use, the more fat soluble drugs will require either a pharmaceutical formulation aimed at prolonging drug release or the use of intrathecal infusions to sustain drug levels. Such infusional techniques are well developed for intrathecal baclofen aimed at controlling spasticity and could be adapted for such an application [123].

*Limitations:* Our criteria for accepting or rejecting drugs were applied strictly therefore. Any drug for which we found documentation of it being irritant or neurotoxic was rejected. It is possible that this criteria was applied too strictly since, for example, methotrexate and cytarabine (standard formulation) were excluded and liposomal cytarabine (Depocyte) has been accepted but evidence of neurotoxicity is now emerging. However the objective of the study was to identify drugs with ideal characteristics for intrathecal use.

It is difficult to interpret many studies of intrathecal chemotherapy in terms of efficacy, as in many cases simultaneous systemic chemotherapy or radiation, or both were administered along with intrathecal treatment making it unclear which component of therapy produced the effect. Most patients recruited into such studies are refractory to other treatments and are therefore undergoing such trials as a 'last resort'. Despite this, measurable response to IT therapy, introduced after systemic chemotherapy has proved refractory, is reported as evidence of effect.

*Drug administration:* The clinical benefits of intrathecal administration need to be balanced against the risks relating to safe prescribing, clinical governance, technical and toxicity issues [38]. Furthermore, in contrast to leukaemia and lymphomas, primary CNS tumours represent anatomical challenges to drug distribution linked to the presence of ventriculo peritoneal (VP) shunts, the influence of drainage of the third ventricle to basal cistern by neuroendoscopic third ventriculostomy, distortion of CNS anatomy linked to post-resection tumour bed, post operative incarceration of the posterior fossa and spinal blocks. Finally, the ideal timing of intrathecal administration in relation to debulking surgery is unclear. Such factors will clearly influence the distribution of CSF and any drug it carries. The challenges of drug delivery to the brain have resulted in a considerable volume of research, which may overcome some of these problems, through either local delivery approaches [105, 140] or vascular delivery across the blood brain barrier [57, 79, 103]. However, relatively little research has been focussed upon delivery systems for intrathecal delivery, the main example being that of liposomal cytarabine [22]. The role of efflux inhibitors such as valspodar in combination with intrathecal therapy in order to overcome the problems of removal of drugs from the CSF by multi-drug resistance transporters could also be explored [126].

The application of this method of drug administration in medulloblastoma is timely, as cure rates with combined chemo-radiotherapy for both localised and metastatic cases are improved (>70%, 5 year survival) in school age children. Novel biological markers are identifying patients with favourable tumour types where further de-escalation of neuro-toxic cranial radiotherapy by adjustments to dose/fractionation schedules or fields. The role of systemic

chemotherapy, whilst established in this age group, will be further examined in trials focussing upon minimising toxicity.

In pre-school age children with favourable presentations (non-metastatic, completely resected, desmoplastic histology) survival rates also exceed 70% in recently reported studies [54, 109] using both low and high intensity chemotherapy schedules with or without intrathecal methotrexate and with or without involved field radiotherapy. The interaction between biological markers and these treatment strategies in this age group is yet to be studied [8, 58, 110]. The role of chemotherapy is, therefore, established. The potential for novel biological agents is yet to be explored but is anticipated.

### ***Selecting drugs for IT trial***

The most effective and safe way of delivering a drug to the leptomeningeal CSF interface will continue to be a challenge. In seeking to optimise drug delivery, drug selection is the first step. The route and method of administration will need to be studied further i.e. the intraventricular route, the lumbar route, by bolus and by infusion. Anatomical factors will require consideration including the effects of tumour location, CSF circulation, including VP shunting and post operative complications affecting the tumour bed. Once drugs have been tested in phase 1 studies, then phase 2 studies looking for efficacy in patients with either newly presenting or relapsed leptomeningeal disease will be necessary. Once a drug has been assessed as effective by intermittent or infusional techniques of IT administration, then phase 3 trials to evaluate its clinical effectiveness or preparation of sustained release formulation for intermittent IT administration would be justified. The preparatory clinical testing can only be developed within centres with the relevant neuroscience / pharmaceutical / pharmacological research expertise.

### ***Conclusion***

This review has extended the list of candidate agents suitable for further investigation of intrathecal treatment of leptomeningeal medulloblastoma. It has identified the importance of making basic physico-chemical criteria available to the academic domain by pharma and academic research groupings in order to accelerate the development of new cancer agents. It has highlighted the opportunity to replicate the success of CNS targeting of drug therapy in leukaemia and lymphoma, by proposing trials of novel intrathecal chemotherapy in medulloblastoma. It has identified the need to establish an acceptable process for selecting new drugs for trial and a suitable trial design.

### ***Acknowledgements:***

This work has been supported by funding from the Children's Brain Tumour Research Centre at the University of Nottingham.

We would like to acknowledge the assistance of Professor Imti Choonara in the development of strategy and review of the manuscript, Steve Barrett in the conduct of the literature searching and Sue Franklin in the preparation of the manuscript.

(Word count 3919)

## References:

- [1] Interactive analysis LogP and LogW Predictors. [www.logp.com](http://www.logp.com).
- [2] (1998) Gliadel wafers for treatment of brain tumors. *The Medical Letter* 40: 92
- [3] Aaron R, Elion G, Colvin O, Graham M, Keir S, Bigner D, Friedman H (1994) Busulfan therapy of central nervous system xenografts in athymic mice. *Cancer Chemotherapy & Pharmacology* 35: 127-131
- [4] Allinson R, Stach P (1978) Intrathecal drug therapy. *Drug Intelligence and Clinical Pharmacy* 12: 347-359
- [5] Archer G, Sampson J, McLendon R, Friedman A, Colvin O, Rose M, Sands H, McCullough W, Fuchs H, Bigner D, Friedman H (1999) Intrathecal busulfan treatment of human neoplastic meningitis in athymic nude rats. *Journal of Neuro-Oncology* 44: 233-241
- [6] Arita N, Ushio Y, Hayakawa T, Nagatani M, Huang T-Y, Izumoto S, Mogami H (1988) Intrathecal ACNU - a new therapeutic approach against malignant leptomeningeal tumors. *Journal of Neuro-Oncology* 6: 221-226
- [7] Arndt C, Colvin O, Balis F, Lester C, Johnson G, Poplack D (1987) Intrathecal administration of 4-hydroperoxycyclophosphamide in rhesus monkeys. *Cancer Research* 47: 5932-5934
- [8] Ashley D, Merchant T, Zhou T, Coleman L, Pollack I, Duffner P, Burger P, Miller D, Buxton A, Douglas S (2008) Systemic chemotherapy, second-look surgery and conformal radiation therapy limited to the posterior fossa and primary site for children >8 months and <3 years with non-metastatic medulloblastoma: A Children's Oncology Group phase III study. *Neuro-Oncology* 10: 436-437
- [9] Association of the British Pharmaceutical Industry (1999) ABPI Compendium of data sheets and summaries of product characteristics 1999-2000. Datapharm Publications, London.
- [10] Baker S, Heideman RL, Crom WR, Kuttesch J, Gajjar A, Stewart C (1996) Cerebrospinal fluid pharmacokinetics and penetration of continuous infusion Topotecan in Children with central nervous system tumours. *Cancer Chemotherapy & Pharmacology* 37: 195-202
- [11] Bates S, Raphaelson M, Price R, McKeever P, Cohen S, Poplack D (1985) Ascending myelopathy after chemotherapy for central nervous system acute lymphoblastic leukemia: correlation with cerebrospinal fluid myelin basic protein. *Medical & Pediatric Oncology* 13: 4-8
- [12] Bayer (1999) Information from Medical Information Department. Newbury, Berkshire.
- [13] Bender J, Grillo-Lopez A, Posada Jnr J (1983) Diaziquone (AZQ). *Investigational New Drugs* 1: 71-84
- [14] Berg S, Balis F, Zimm S, Murphy R, Holcenberg J, Sato J, Reaman G, Steinherz P, Gillespie A, Doherty K, Poplack D (1992) Phase I/II trial and pharmacokinetics of intrathecal diaziquone in refractory meningeal malignancies. *Journal of Clinical Oncology* 10: 143-148
- [15] Bernardi R, Bomgaars L, Fox E, Balis F, Egorin M, Lagattuta T, Aikin A, Whitcome P, Renbarger J, Lieberman F, Berg S, Blaney S (2008) Phase I clinical trial of intrathecal gemcitabine in patients with neoplastic meningitis. *Cancer Chemotherapy & Pharmacology* 62: 355-361
- [16] Bhojwani D, Pui C-H (2008) Intrathecal liposomal cytarabine: more friend than foe? *Leukaemia & Lymphoma* 49: 1427-1430
- [17] Blaney S, Boyett J, Friedman H, Gajjar A, Geyer R, Horowitz M, Hunt D, Kieran M, Kun L, Packer R, Phillips P, Pollack M, Prados M, Heideman R (2005) Phase I clinical trial of mafosfamide in infants and children aged 3 years or younger with newly diagnosed embryonal tumors: A Pediatric Brain Tumor Consortium study (PBTC-001). *Journal of Clinical Oncology* 23: 525-531
- [18] Blaney S, Cole D, Godwin K, Sung C, Poplack D, Balis F (1995) Intrathecal administration of topotecan in nonhuman primates. *Cancer Chemotherapy & Pharmacology* 36: 121-124
- [19] Blaney S, Heideman R, Berg S, Adamson P, Gillespie A, Geyer J, Packer R, Matthay K, Jaeckle K, Cole D, Kuttesch N, Poplack D, Ballis F (2003) Phase I clinical trial of intrathecal topotecan in patients with neoplastic meningitis. *Journal of Clinical Oncology* 21: 143-147
- [20] Blaney S, Poplack D (1996) Pharmacologic strategies for the treatment of meningeal malignancy. *Investigational New Drugs* 14: 69-85
- [21] Blaney S, Poplack D (1998) New cytotoxic drugs for intrathecal administration. *Journal of Neuro-Oncology* 38: 219-223
- [22] Bleyer W (1999) Intrathecal depot cytarabine therapy: a welcome addition to a limited armamentarium. *Clinical Cancer Research* 5: 3349-3351
- [23] Bleyer W, Coccia P, Sather H, et al (1983) Reduction in central nervous system leukemia with a pharmacokinetically derived intrathecal methotrexate dosage regimen. *Journal of Clinical Oncology* 1: 317-325
- [24] Bleyer W, Drake J, Chabner B (1973) Neurotoxicity and elevated cerebrospinal fluid methotrexate concentration in meningeal leukemia. *New England Journal of Medicine* 289: 770-773
- [25] Boiardi A, Eoli M, Pozzi A, Salmaggi A, Broggi G, Silvani A (1999) Locally delivered chemotherapy and repeated surgery can improve survival in glioblastoma patients. *Ital J Neurol Sci* 20: 43-48
- [26] Boland I, Vassal G, Morizet J, Terrier-Lacombe M-J, Valteau-Couanet D, Kalifa C, Hartmann O, Gouyette A (1999) Busulphan is active against neuroblastoma and medulloblastoma xenografts in athymic mice at clinically achievable plasma drug concentrations. *British Journal of Cancer* 79: 787-792
- [27] Bomgaars L, Geyer J, Franklin J, Dahl G, Park J, Winick N, Klenke R, Berg S, Blaney S (2004) Phase I trial of intrathecal liposomal cytarabine in children with neoplastic meningitis. *Journal of Clinical Oncology* 19: 3916-3921

- [28] Brandes A, Pasetto L, Monfardini S (2000) New drugs in recurrent high grade gliomas. *Anti-Cancer Research* 20: 1913-1920
- [29] Brem H, Piantadosi S, Burger P, Walker M, Selker R, Vick N, Black K, Sisti M, Brem S, Mohr G, Muller P, Morawetz R, Schold S (1995) Placebo-controlled trial of safety and efficacy of intraoperative controlled delivery by biodegradable polymers of chemotherapy for recurrent gliomas. *Lancet* 345: 1008-1012
- [30] British Medical Association & Royal Pharmaceutical Society of Great Britain (2000) British National Formulary. BMA & RPS (GB), London.
- [31] Castleberry R, Ragab A, Steuber C, Kamen B, Toledano S, Starling K, Norris D, Burger P, Krischer J (1990) Aziridinylbenzoquinone (AZQ) in the treatment of recurrent pediatric brain and other malignant solid tumors. *Investigational New Drugs* 8: 401-406
- [32] Chamberlain M, Khatibi S, Kim J, Howell S, Chatelut E, Kim S (1993) Treatment of Leptomeningeal Metastasis with Intraventricular Administration of Depot Cytarabine (DTC101). *Archives of Neurology* 50: 261-264
- [33] Chamberlain M, Kormanik P, Howell S, Kim S (1995) Pharmacokinetics of Intralumbar DTC-101 for the Treatment of Leptomeningeal Metastases. *Archives of Neurology* 52: 912-917
- [34] Champagne M, Silver H (1992) Intrathecal Dacarbazine Treatment of Leptomeningeal Malignant Melanoma. *Journal Of The National Cancer Institute* 84: 1203-1204
- [35] Chen Y-M, Chen M-C, Tsai C-M, Perng R-P (2003) Intrathecal gemcitabine chemotherapy for non-small cell lung cancer patients with meningeal carcinomatosis - a case report. *Lung Cancer* 40: 99-101
- [36] Crawford S, Rustin G, Bagshawe K (1986) Acute neurological toxicity of intrathecal cytosine arabinoside. *Cancer Chemotherapy & Pharmacology* 16: 306-307
- [37] Damon L, Plunkett W, Linker C (1991) Plasma and Cerebrospinal Fluid Pharmacokinetics of 1- $\beta$ - D-Arabinofuranosylcytosine and 1- $\beta$ - D-Arabinofuranosyluracil following the Repeated Intravenous Administration of High- and Intermediate-Dose  $\beta$ - D-Arabinofuranosylcytosine. *Cancer Research* 51: 4141-4145
- [38] Department of Health (2008) Updated national guidance on the safe administration of intrathecal chemotherapy. In: Health Do (ed).
- [39] Dodion P, Sanders C, Georges P, Kenis Y (1988) In vitro chemosensitivity of brain tumors to cisplatin and its analogues, iproplatin and carboplatin. *Cancer Chemotherapy & Pharmacology* 22: 80-82
- [40] Dollery C (ed) (1999) Therapeutic drugs. Churchill Livingstone, Edinburgh.
- [41] Doz F, Pinkerton R (1994) What is the Place of Carboplatin in Paediatric Oncology? *European Journal of Cancer* 30A: 194-201
- [42] Drayton C (ed) (1990) Comprehensive medicinal chemistry. Pergamon Press, Oxford.
- [43] Eagan R, Dinapoli R, Cascino T, Scheithauer B, O'Neill B, O'Fallon R (1987) Comprehensive phase II evaluation of Aziridinylbenzoquinone (AZQ, Diaziquone) in recurrent human primary brain tumors. *Journal of Neuro-Oncology* 5: 309-314
- [44] Edwards M, Levin V, Seager M, Wilson C (1981) Intrathecal Chemotherapy for Leptomeningeal Dissemination of Medulloblastoma. *Child's brain* 8: 444-451
- [45] Egorin M, Zuhowski E, McCully C, Blaney S, Kerr J, Berg S, Ballis F (2002) Pharmacokinetics of intrathecal gemcitabine in non-human primates. *Clinical Cancer Research* 8: 2437-2442
- [46] Englehard H (2000) The role of interstitial BCNU chemotherapy in the treatment of malignant glioma. *Surg Neurol* 53: 458-464
- [47] Ettinger L, Ru N, Krailo M, Ruccione K, Krivit W, Hammond G (1990) A phase II study of diaziquone in children with recurrent or progressive primary brain tumors: A report from the Children's Cancer Study Group. *Journal of Neuro-Oncology* 9: 69-76
- [48] Ewend M, Sampath P, Williams J, Tyler B, Brem H (1998) Local delivery of chemotherapy prolongs survival in experimental brain metastases from breast carcinoma. *Neurosurgery* 43: 1185-1193
- [49] Fleischhack G, Jaehde U, Bode U (2005) Pharmacokinetics following intraventricular administration of chemotherapy in patients with neoplastic meningitis. *Clin Pharmacokinet* 4: 1-31
- [50] Friedman H (2000) Temozolomide in early stages of newly diagnosed malignant glioma and neoplastic meningitis. *Seminars in Oncology* 27: 35-40
- [51] Friedman H, Bigner S, Bigner D (1995) Cyclophosphamide therapy of medulloblastoma: from the laboratory to the clinic and back again (and again and again). *Journal of Neuro-Oncology* 24: 103-108
- [52] Friedman H, Ochs J, Finlay J, Geyer R, Arndt C, Cohen B, Phillips P, Strauss L, Hochberg F, Schold S, Bigner D, Colvin O (1993) Phase I trial of intrathecal 4-hydroperoxycyclophosphamide for neoplastic meningitis. *American Association for Cancer Research*. p. 269.
- [53] Green L (1988) Choice of diluents for intrathecal drug administration. *Clinical Pharmacy* 7: 265-266
- [54] Grill J, Sainte-Rose C, Jouvet A, Gentet J-C, Lejars O, Frappaz D, Doz F, Rialland X, Pichon F, Bertozzi A-I, Chastagner P, Couanet D, Habrand J-L, Raquin M-A, Le Deley M-C, Kalifa C (2005) Treatment of medulloblastoma with postoperative chemotherapy alone: an SFOP prospective trial in young children. *Lancet Oncology* 6: 573-580
- [55] Grossman S, Trump DL, Chen D, Thompson G, Camargo E (1982) Cerebrospinal fluid flow abnormalities in patients with neoplastic meningitis. An evaluation using <sup>111</sup>Indium DTPA ventriculography. *American Journal of Medicine* 73: 641-647

- [56] Gururangan S, Petros W, Young Poussaint T, Hancock M, Phillips P, Friedman H, Bomgaars L, Blaney S, Kun L, Boyett J (2006) Phase I trial of intrathecal Spartaject Busulfan in children with neoplastic meningitis: a Pediatric Brain Tumor Consortium study (PBTC-004). *Clinical Cancer Research* 12: 1540-1546
- [57] Gutman R, Peacock G, Lu D (2000) Targeted drug delivery for brain cancer treatment. *J Controlled Release* 65: 31-41
- [58] Haberler C, Varlet P, Legoix P, Fattet S, Janoueix-Lerosey I, Lellouch-Tubiana A, Grill J, Doz F, Sainte-Rose C, Delattre O (2008) Widespread nuclear beta-catenin expression in medulloblastoma correlates with mutation status of CTNGB1 gene encoding beta-catenin. *Neuro-Oncology* 10: 470
- [59] Hare C, Elion G, Houghton P, Houghton J, Keir S, Marcelli S, Bigner D, Friedman H (1997) Therapeutic efficacy of the topoisomerase I inhibitor 7-ethyl-10-(4-[1-piperidino]-1-piperidino)-carbonyloxy-camptothecin against pediatric and adult central nervous system tumor xenografts. *Cancer Chemotherapy & Pharmacology* 39: 187-191
- [60] Heideman RL, McCully C, Balis F, Poplack D (1993) Cerebrospinal fluid pharmacokinetics and toxicology of intraventricular and intrathecal arabinosyl-5-azacytosine (fazarabine, NSC 281272) in the nonhuman primate. *Investigational New Drugs* 11: 135-140
- [61] Hori T, Tanaka S, Nishiyama M, Kamitani H, Watanabe T, Tabuchi F, Tatsuhara T, Nakajima E (1993) Distribution of intrathecally administered ACNU in mongrel dogs: Pharmacokinetics and quantitative autoradiographic study. *Surg Neurol* 40: 183-195
- [62] Houchens D, Ovejera A, Riblet S, Slagel D (1983) Human brain tumor xenografts in nude mice as a chemotherapy model. *European Journal of Cancer & Clinical Oncology* 19: 799-805
- [63] Huang T-Y, Howng S-L (1997) Intrathecal ACNU against experimental leptomeningeal tumors. *Kaohsiung J Med Sci* 13: 626-630
- [64] Hussain M, Wozniak A, Edelstein M (1993) Neurotoxicity of antineoplastic agents. *Critical Reviews in Oncology/Hematology* 14: 61-75
- [65] Jabbour E, O'Brien S, Kantarjian H, Garcia-Manero G, Ferrajoli A, Ravandi F, Cabanillas M, Thomas DA (2007) Neurologic complications associated with intrathecal liposomal cytarabine given prophylactically in combination with high-dose methotrexate and cytarabine to patients with acute lymphocytic leukemia. *Blood* 109: 3214-3218
- [66] Jaeckle K, Batchelor T, O'Day S, Phuphanich S, New P, Lesser G, Cohn A, Gilbert M, Aiken R, Heros D, Rogers L, Wong E, Fulton D, Gutheil J, Baidas S, Kennedy J, Mason W, Moots P, Russell C, Swinnen L, Howell S (2002) An open label trial of sustained-release cytarabine (DepoCyt) for the intrathecal treatment of solid tumor neoplastic meningitis. *Journal of Neuro-Oncology* 57: 231-239
- [67] Kerr J, Berg S, Dauser R, Nuchtern J, Egorin M, McGuffey L, Aleksic A, Blaney S (2001) Plasma and cerebrospinal fluid pharmacokinetics. *Cancer Chemotherapy & Pharmacology* 47: 411-414
- [68] Kim S, Chatelut E, Kim J, Howell S, Cates C, Kormanik P, Chamberlain M (1993) Extended CSF Cytarabine Exposure Following Intrathecal Administration of DTC 101. *Journal of Clinical Oncology* 11: 2186-2193
- [69] Kim S, Khatibi S, Howell S, McCully C, Balis F, Poplack D (1993) Prolongation of Drug Exposure in Cerebrospinal Fluid by Encapsulation into DepoFoam. *Cancer Research* 53: 1596-1598
- [70] Kim S, Kim D, Geyer M, Howell S (1987) Multivesicular Liposomes Containing 1- $\beta$ -D-Arabinofuranosylcytosine for Slow-Release Intrathecal Therapy. *Cancer Research* 47: 3935-3937
- [71] Kimler B, Liu C, Evans R, Morantz R (1992) Intracerebral chemotherapy in the 9L rat brain tumor model. *Journal of Neuro-Oncology* 14: 191-200
- [72] Kitamura I, Kochi M, Matsumoto Y, Ueoka R, Kuratsu J-I, Ushio Y (1996) Intrathecal chemotherapy with 1,3-Bis(2-chloroethyl)-1-nitrosourea encapsulated into hybrid liposomes for meningeal gliomatosis: An experimental study. *Cancer Research* 56: 3986-3992
- [73] Kleinschmidt-DeMasters B, Yeh M (1992) "Locked-in syndrome" after intrathecal cytosine arabinoside therapy for malignant immunoblastic lymphoma. *Cancer* 70: 2504-2507
- [74] Kochi M, Kuratsu J-I, Mihara Y, Takaki S, Inoue N, Sueyoshi N, Uemura S, Ushio Y (1990) Neurotoxicity and pharmacokinetics of intrathecal perfusion of ACNU in dogs. *Cancer Research* 50: 3119-3123
- [75] Kochi M, Kuratsu J-I, Mihara Y, Takaki S, Seto H, Uemura S, Ushio Y (1993) Ventriculolumbar Perfusion of 3-[(4-amino-2-methyl-5-pyrimidinyl)methyl]-1-(chloroethyl)-1-nitrosourea hydrochloride. *Neurosurgery* 33: 817-823
- [76] Kochi M, Takaki S, Kuratsu J-I, Seto H, Kitamura I, Ushio Y (1994) Neurotoxicity and pharmacokinetics of ventriculolumbar perfusion of methyl 6-[3-(2-chloroethyl)-3-nitrosoureido]-6-deoxy- $\alpha$ -D-glucopyranoside (MCNU) in dogs. *Journal of Neuro-Oncology* 19: 239-244
- [77] Kohn F, Malkmus S, Brownson E, Rossi S, Yaksh T (1998) Fate of the predominant phospholipid component of DepoFoam<sup>TM</sup> drug delivery matrix after intrathecal administration of sustained-release encapsulated cytarabine in rats. *Drug Delivery* 5: 143-151
- [78] Kooistra K, Rodriguez M, Powis G (1989) Toxicity of intrathecally administered cytotoxic drugs and their antitumor activity against an intrathecal Walker 256 carcinosarcoma model for meningeal carcinomatosis in the rat. *Cancer Research* 49: 977-982
- [79] Kreuter J (2001) Nanoparticulate systems for brain delivery of drugs. *Advanced Drug Delivery Reviews* 47: 65-81
- [80] Kuhl J, Doz F, Taylor R (2004) Embryonic tumors. In: Walker D, Perilongo G, Punt J, Taylor R (eds). *Brain & Spinal Tumors of Childhood*. Arnold, London. pp. 314-330.

- [81] Lassaletta A, Lopez-Ibor B, Mateos E, Gonzalez-Vicent M, Perez-Martinez A, Sevilla J, Diaz M, Madero L (2009) Intrathecal liposomal cytarabine in children under 4 years with malignant brain tumours. *Journal of Neuro-Oncology* Epub ahead of print
- [82] Levin V, Byrd D, Sikic B, Etiz B, Campbell J, Borcich J, Davis R (1985) Central Nervous System Toxicity and Cerebrospinal Fluid Pharmacokinetics of Intraventricularly Administered Bleomycin In Beagles. *Cancer Research* 45: 3810-3815
- [83] Levin V, Chamberlain M, Silver P, Rodriguez L, Prados M (1989) Phase I/II study of intraventricular and intrathecal ACNU for leptomeningeal neoplasia. *Cancer Chemotherapy & Pharmacology* 23: 301-307
- [84] Levin V, Edwards M, Gutin P, Vestnys P, Fulton D, Seager M, Wilson C (1984) Phase II evaluation of dibromodulcitol in the treatment of recurrent medulloblastoma, ependymoma and malignant astrocytoma. *Journal of Neurosurgery* 61: 1063-1068
- [85] Lind M (1993) Pharmacokinetics of alkylating agents. *Pharmacokinetics and Cancer Chemotherapy* 17: 157-179
- [86] MacDonald D (1991) Neurologic complications of chemotherapy. *Neurol Clin* 9: 955-967
- [87] Martindale (1999) The complete drug reference. Pharmaceutical Press, London.
- [88] McKeran R, Firth G, Oliver S, Uttley D, O'Laioire S (1985) A potential application for the intracerebral injection of drugs entrapped within liposomes in the treatment of human cerebral gliomas. *Journal of Neurology, Neurosurgery and Psychiatry* 48: 1213-1219
- [89] McVie J (1992) Teniposide (VM-26) in brain tumors. *Seminars in Oncology* 19: 85-88
- [90] Morantz R, Kimler B, Vats T, Henderson S (1983) Leomycin and brain tumors. *Journal of Neuro-Oncology* 1: 249-255
- [91] Nakagawa H, Maeda N, Tsuzuki T, Suzuki T, Hirayama A, Miyahara E, Wada K (2001) Intracavitary chemotherapy with 5-fluoro-2'-deoxyuridine (FdUrd) in malignant brain tumors. *Japan Journal of Clinical Oncology* 31: 251-258
- [92] Nakagawa H, Miyahara E, Suzuki T, Wada K, Tamura M, Fukushima Y (2005) Continuous intrathecal administration of 5-fluoro-2'-deoxyuridine for the treatment of neoplastic meningitis. *Neurosurgery* 57: 266-280
- [93] Nakagawa H, Yamada M, Fukushima M, Ikenaka K (1999) Intrathecal 5-fluoro-2'-deoxyuridine (FdUrd) for the treatment of solid tumor neoplastic meningitis: an in vivo study. *Cancer Chemotherapy & Pharmacology* 43: 247-256
- [94] Nakagawa H, Yamada M, Maeda N, Iwatsuki K, Hirayama Z, Ikenaka K (1999) Clinical trial of intrathecal administration of 5-fluoro-2'-deoxyuridine for treatment of meningeal dissemination of malignant brain tumors. *Journal of Neuro-Oncology* 45: 175-183
- [95] National Cancer Institute [www.nci.nih.gov](http://www.nci.nih.gov).
- [96] National Library of Medicine Special Information Services <http://chem.sis.nlm.nih.gov>.
- [97] Newlands E, Stevens M, Wedge S, Wheelhouse R, Brock C (1997) Temozolomide: a review of its discovery, chemical properties, pre-clinical development and clinical trials. *Cancer Treatment Reviews* 23: 35-61
- [98] Nishio S, Morioka T, Takeshita I, Tadahisa S, Takanori I, Fujiwara S, Fukui M (1995) Chemotherapy for progressive pilocytic astrocytomas in the chiasmo-hypothalamic regions. *Clinical Neurology and Neurosurgery* 87: 300-306
- [99] Ochiai H, Pernell C, Archer G, Chewning T, McLendon R, Friedman H, Sampson J (2006) Treatment of neoplastic meningitis with intrathecal 9-Nitro-camptothecin. *Neurol Med Chir (Tokyo)* 46: 485-490
- [100] Olivi A, Dundan L, Corden B, Lenartz D, Brem H (1993) Direct Delivery Of Platinum Based Neoplastics To The Central Nervous System : A Toxicity And Ultrastructural Study. *Cancer Chemotherapy Pharmacology* 31: 449-454
- [101] Paal C, Erdelyi-Toth V, Pap E, Csaki C, Ferencz T, Schuler D, Borsi J (1994) Pharmacokinetic studies on Elobromol in children with brain tumors. *Anti-Cancer Drugs* 5: 539-547
- [102] Pardridge W (1996) Brain Drug Delivery and Blood-Brain Barrier Transport. *Drug Delivery* 3: 99-115
- [103] Pardridge W (2002) Drug and gene targeting to the brain with molecular Trojan horses. *Nature Reviews Drug Discovery* 1: 131-139
- [104] Phillips P, Than T, Cork L, Hilton J, Carson B, Colvin O, Grochow L (1992) Intrathecal 4-hydroperoxycyclophosphamide: Neurotoxicity, cerebrospinal fluid pharmacokinetics, and antitumor activity in a rabbit model of VX2 leptomeningeal carcinomatosis. *Cancer Research* 52: 6168-6174
- [105] Read T-A, Thorsen F, Bjerkvig R (2002) Localised delivery of therapeutic agents to CNS malignancies; old and new approaches. *Current Pharmaceutical Technology* 3: 257-273
- [106] Rhone Poulenc Rorer (1999) Information from Medical Information Department. West Malling, Kent.
- [107] Riccardi R, Riccardi A, Di Rocco C, Carelli G, Tartaglia R, Lasorella A, Servidei T, Mastrangelo R (1992) Cerebrospinal fluid pharmacokinetics of carboplatin in children with brain tumors. *Cancer Chemotherapy and Pharmacology* 30: 21-24
- [108] Ruggiero A, Conter V, Milani M, Biagi E, Lazzareschi I, Sparano P, Riccardi R (2001) Intrathecal chemotherapy with antineoplastic agents in children. *Paediatric Drugs* 3: 237-246
- [109] Rutkowski S, Bode U, Deinlein F, Ottensmeier H, Warmuth-Metz M, Soerensen N, Graf N, Emser A, Pietsch T, Wolff J, Korkmann R, Kuehl J (2005) Treatment of early childhood medulloblastoma by postoperative chemotherapy alone. *New England Journal of Medicine* 352: 978-986

- [110] Rutkowski S, Von Hoff K, Emser A, Garre ML, Walker D, Grundy R, Dhall G, Finlay J, Grill J (2008) Prognostic factors and survival of young children with medulloblastoma: An international meta-analysis. *Neuro-Oncology* 10: 437
- [111] Sampson J, Archer G, Villavicencio A, McLendon R, Friedman A, Bishop W, Bigner D, Friedman H (1999) Treatment of neoplastic meningitis with intrathecal temozolomide. *Clinical Cancer Research* 5: 1183-1188
- [112] Savaraj N, Feun L, Lu K, Gray K, Wang C, Loo T (1992) Pharmacology of intrathecal VP-16-213 in dogs. *Journal of Neuro-Oncology* 13: 211-215
- [113] Savas A, Erdem A, Tun K, Kanpolat Y (2000) Fatal toxic effect of bleomycin in brain tissue after intracystic chemotherapy for a craniopharyngioma: Case report. *Neurosurgery* 46: 213-217
- [114] Schmandt S, Kuhl J (1998) Chemotherapy as prophylaxis and treatment of meningosis in children less than 3 years of age with medulloblastoma. *Journal of Neuro-Oncology* 38: 187-192
- [115] Schold Jr S, Mahaley Jr M, Vick N, Friedman H, Burger P, DeLong E, Albright Jr R, Bullard D, Khandekar D, Cairncross J, Macdonald D, Falletta J (1987) Phase II diaziquone-based chemotherapy trials in patients with anaplastic supratentorial astrocytic neoplasms. *Journal of Clinical Oncology* 5: 464-471
- [116] Schuler D, Somlo P, Borsi J, Paraicz E, Revesz T, Kardos G, Koos R (1988) New drug combination for the treatment of relapsed brain tumors in children. *Pediatric Hematology & Oncology* 5: 153-156
- [117] Schuler D, Somlo P, Koos R, Kalmancey R, Paraicz E (1992) Treatment of malignant scala posterior brain tumors in children: the chemotherapy of relapsed medulloblastoma with a dibromdulcitol-containing drug regime and pharmacokinetic studies of dibromdulcitol in children. *Medical & Pediatric Oncology* 20: 312-314
- [118] Screnci D, McKeage M, Galettis P, Hambley T, Palmer B, Baguley B (2000) Relationships between hydrophobicity, reactivity, accumulation and peripheral nerve toxicity of a series of platinum drugs. *British Journal of Cancer* 82: 966-972
- [119] Selby G, Upchurch C, Townsend J, Eyre H (1994) A phase II evaluation of fazarabine in high-grade gliomas: a Southwest Oncology Group study. *Cancer Chemotherapy & Pharmacology* 34: 179-180
- [120] Slave I, Schuller E, Czech T, Hainfellner J, Seidl R, Dieckmann K (1998) Intrathecal mafosfamide therapy for paediatric brain tumors with meningeal dissemination. *Journal of Neuro-Oncology* 38: 213-218
- [121] Slave I, Schuller E, Falger J, Gunes M, Pillwein K, Czech T, Dietrich W, Rossler K, Dieckmann K, Prayer D, Hainfellner J (2003) Feasibility of long-term intraventricular therapy with mafosfamide (n=26) and etoposide (n=11): Experience in 26 children with disseminated malignant brain tumors. *Journal of Neuro-Oncology* 64: 239-247
- [122] SmithKline Beecham Pharmaceuticals (1999) Information from Medical Information Department. Welwyn Garden City.
- [123] Stempien L, Tsai T (2000) Intrathecal baclofen pump use for spasticity: a clinical survey. *American Journal of Physical Medicine & Rehabilitation* 79: 536-541
- [124] Stevens M, Newlands E (1993) From triazines and triazines to temozolomide. *European Journal of Cancer* 29a: 1045-1047
- [125] Stiller C, Bleyer W (2004) Epidemiology. In: Walker D, Perilongo G, Punt J, Taylor R (eds). *Brain & Spinal Tumors of Childhood*. Arnold, London. pp. 35-49.
- [126] Taylor E (2002) The impact of efflux transporters in the brain on the development of drugs for CNS disorders. *Clinical Pharmacokinetics* 41: 81-92
- [127] Taylor S (1994) New agents in the treatment of primary brain tumors. *Journal of Neuro-Oncology* 20: 141-153
- [128] Tomlinson F, Lihou M, Smith P (1991) Comparison of in vitro activity of epipodophyllotoxins with other chemotherapeutic agents in human medulloblastomas. *British Journal of Cancer* 64: 1051-1059
- [129] Tsuboi K, Yoshii Y, Hyodo A, Takada K, Nose T (1995) Leukoencephalopathy associated with intra-arterial ACNU in patients with gliomas. *Journal of Neuro-Oncology* 23: 223-231
- [130] Tuxen M, Hansen S (1994) Neurotoxicity secondary to antineoplastic drugs. *Cancer Treatment Reviews* 20: 191-214
- [131] Ushio Y, Kochi M (1998) Intrathecal perfusion chemotherapy with ACNU against subarachnoid dissemination of glioma in children. *Int J Pediat Hem Oncol* 5: 55-56
- [132] Ushio Y, Kochi M, Kitamura I, Kuratsu J (1998) Ventriculolumbar perfusion of 3-[(4-amino-2-methyl-5-pyrimidinyl)-methyl]-1-(2-chloroethyl)-1-nitrosourea hydrochloride for subarachnoid dissemination of gliomas. *Journal of Neuro-Oncology* 38: 207-212
- [133] van der Gaast A, Sonneveld P, Mans D, Splinter T (1992) Intrathecal administration of etoposide in the treatment of malignant meningitis: feasibility and pharmacokinetic data. *Cancer Chemotherapy & Pharmacology* 29: 335-337
- [134] van der Graaf W, Haaxma-Reiche H, Burghouts J, Postmus P (1993) Teniposide for meningeal carcinomatosis of small cell lung cancer. *Lung Cancer* 10: 247-249
- [135] Vassal G, Boland I, Santos A, Bissery M, Terrier-Lacombe M, Morizet J, Sainte-Rose C, Lellouch-Tubiana A, Kalifa C, Gouyette A (1997) Potent Therapeutic Activity of Irinotecan (CPT-11) and its Schedule Dependency in Medulloblastoma Xenografts in Nude Mice. *International Journal of Cancer* 73: 156-163
- [136] Vassal G, Boland I, Terrier-Lacombe M, Watson A, Margison G, Venuat A, Morizet J, Parker F, Lacroix C, Lellouch-Tubiana A, Pierre-Kahn A, Poullain M, Gouyette A (1998) Activity of Fotemustine in Medulloblastoma and Malignant Glioma Xenografts in Relation to  $O^6$ -Alkylguanine-DNA Alkyltransferase and Alkylpurine-DNA *N*-Glycosylase Activity. *Clinical Cancer Research* 4: 463-468

- [137] Vassal G, Pondarre C, Boland I, Cappelli C, Santos A, Thomas C, Lucchi E, Imadalou K, Pein F, Morizet J, Gouyette A (1998) Preclinical development of camptothecin derivatives and clinical trials in pediatric oncology. *Biochimie* 80: 271-280
- [138] Vassal G, Terrier-Lacombe M-J, Bissery M, Venuat A, Gyergyay F, Benard J, Morizet J, Boland I, Ardouin P, Bressac-de-Paillerets B, Gouyette A (1996) Therapeutic activity of CPT-11, a DNA-topoisomerase I inhibitor, against peripheral primitive neuroectodermal tumour and neuroblastoma xenografts. *British Journal of Cancer* 75: 537-545
- [139] Walker D, Wilne S (2005) Treatment of medulloblastoma in young children. *Lancet Oncology* 6: 541-542
- [140] Wang P, Frazier J, Brem H (2002) Local drug delivery to the brain. *Adv Drug Del* 54: 987-1013
- [141] Ward H (1978) Central nervous system tumors of childhood treated with CCNU, vincristine and radiation. *Medical & Pediatric Oncology* 4: 315-320
- [142] Watterson J, Toogood I, Nieder M, Morse M, Frierdich S, Lee Y, Moertel C, Priest J (1994) Excessive spinal cord toxicity from intensive central nervous system-directed therapies. *Cancer* 74: 3034-3041
- [143] Weiss R, Issell B (1982) The nitrosoureas: carmustine (BCNU) and lomustine (CCNU). *Cancer Treatment Reviews* 9: 313-330
- [144] Yamada M, Nakagawa H, Fukushima M, Shimizu K, Hayakawa T, Ikenaka K (1998) In vitro study on intrathecal use of 5-fluoro-2'-deoxyuridine (FdUrd) for meningeal dissemination of malignant brain tumors. *Journal of Neuro-Oncology* 37: 115-121
- [145] Yamasaki T, Kikuchi H, Yamashita J, Asato R, Fujita M (1989) Primary spinal intramedullary malignant melanoma: case report. *Neurosurgery* 25: 117-121
- [146] Yamashita T, Yamashita J, Shoin K (1990) Neurotoxicity of local administration of two nitrosoureas in malignant gliomas. *Neurosurgery* 26: 794-800
- [147] Yoshida T, Beuls E, Shimizu K, Koulousakis A, Sturm V (1992) Intrathecal chemotherapy with ACNU for meningeal gliomatosis. *British Journal of Cancer* 66: 999-1004
- [148] Yoshida T, Shimizu K, Koulousakis A, Sturm V (1992) Intrathecal Chemotherapy With ACNU In A Meningeal Gliomatosis Rat Model. *Journal Of Neurosurgery* 77: 778-782
- [149] Yoshimura J, Nishiyama K, Mori H, Takahashi H, Fujii Y (2008) Intrathecal chemotherapy for refractory disseminated medulloblastoma. *Child's Nervous System* 24: 581-585
- [150] Zimm S, Collins J, Curt G, O'Neill D, Poplack D (1984) Cerebrospinal fluid pharmacokinetics of intraventricular and intravenous aziridinylbenzoquinone. *Cancer Research* 44: 1698-1701
- [151] Zimm S, Collins J, Miser J, Chatterji D, Poplack D (1984) Cytosine arabinoside cerebrospinal fluid kinetics. *Clinical Pharmacological Therapeutics* 35: 826-830

Table 1: Summary of ideal clinical, biological and physicochemical / pharmaceutical properties used to justify selection or rejection of drugs.

<b><i>Property Required</i></b>
<b><i>Clinical</i></b> Non-irritant Neurotoxicity low or absent Evidence of tumour sensitivity
<b><i>Biological</i></b> CSF transport system absent Cell cycle non-specific agent
<b><i>Physicochemical &amp; pharmaceutical</i></b> Active in CSF  Hydrophilic and/or ionised at CSF pH therefore low membrane permeability (to minimise diffusion out of CSF) Molecular Size (>700 Da) Suitable formulation readily available

Table 2: List of chemotherapy drugs considered with selection / rejection status and rejection justification

Table 2a - Rejected chemotherapy agents plus justification

<b><i>Irritant</i></b>	<b><i>No tumour activity</i></b>	<b><i>No information</i></b>	<b><i>Requires activation</i></b>	<b><i>Neurotoxic</i></b>	<b><i>Neurotoxic and irritant</i></b>	<b><i>Neurotoxic via IT route</i></b>
Aclarubicin	Carbetimer	AG-337 (Noltatrexed)	Capecitabine	Acivicin	Altretamine	Doxorubicin
Amsacrine	Mitoguazone	Amonafide	Cyclophosphamide	Asparaginase	Chlorambucil	Melphalan
Azathioprine		Amrubicin	Mitomycin	Carmofur		Mitozantrone
Dactinomycin		Azacitidine	Procarbazine	Cisplatin		**Fluorouracil
Daunorubicin		Bendamustine	Trofosfamide	Cladribine		
Epirubicin		Bisantrene	*Thiotepa	Docetaxel		
Idarubicin		Caracemide		Doxifluridine		
Mustine (Mechlorethamine)		Carboquone		Fludarabine		
Paclitaxel		Chlorozotocin (DCNU)		Fotemustine		
Pirarubicin		Chromomycin		Gemcitabine		
Piroxantrone		Clofarabine		Hydroxyurea		
Plicamycin		Crisnatol		Ifosfamide		
Streptozocin		Decitabine		Methotrexate		
Valrubicin		Echinomycin		Mitotane		
Zorubicin		Edatrexate		Oxaliplatin		
		Enloplatin		Pentostatin		
		Enocitabine		Sparfosic acid		
		Estramustine		Spiromustine		
		Homoharringtonine		Suramin		
		Ilmofosine		Tegafur		
		Improsulfan		Vinblastine		
		JM-216 (Satraplatin)		Vincristine		
		Lobaplatin		Vindesine		
		Miboplatin		Vinorelbine		
		Mitobronitol				

		Mitozolamide				
		Nedaplatin				
		Pemetrexed				
		Peplomycin				
		Pipobroman				
		Piritrexim				
		Porfiromycin				
		Prednimustine				
		Raltitrexed				
		Ranpirnase				
		Razoxane				
		Sebriplatin				
		Semustine				
		Sobuzoxane				
		Teloxantrone				
		Thioguanine				
		Tirapazamine				
		Treosulfan				
		Trimetrexate				
		Ubenimex				

\* Systemic administration delivers concentration to IT route

\*\* Also irritant

Table 2b – Selected drugs

1. Carboplatin	Selection	Table 3
2. Cytarabine	Selection	Table 3
3. Diaziquone	Selection	Table 3
4. Etoposide	Selection	Table 3
5. Floxuridine (FdUrd)	Selection	Table 3
6. 4-hydroperoxycyclophosphamide	Selection	Table 3
7. Mafosfamide	Selection	Table 3
8. Mercaptopurine	Selection	Table 3
9. Nimustine (ACNU)	Selection	Table 3
10. Temozolomide	Selection	Table 3
11. Topotecan	Selection	Table 3
12. Eflornithine	Selection	Table 4
13. Mitolactol (Dibromodulcitol)	Selection	Table 4
14. Didemnin B	Selection	Table 4
15. Irinotecan	Selection	Table 4
16. Lomustine (CCNU)	Selection	Table 4
17. Lonidamine	Selection	Table 4
18. Menogaril	Selection	Table 4
19. Tauromustine	Selection	Table 4
20. Teniposide	Selection	Table 4
21. Bleomycin	Rejection	Table 5 – neurotoxic when given IT
22. Busulphan	Selection	Table 5
23. Carmustine (BCNU)	Selection	Table 5
24. Dacarbazine	Selection	Table 5
25. Fazarabine	Selection	Table 5
26. Ranimustine (MCNU)	Selection	Table 5
27. Zinostatin	Selection	Table 5

Table 3: Drugs graded by eligibility for trial by intrathecal administration for medulloblastoma

**Legend**

OWPC = octanol/water partition coefficient >0 = lipophilic; <0 = hydrophilic (different values obtained from different references)

pKa = dissociation constant. pKa <5.3 or >9.3 indicate an ionisation of >99% at CSF pH of 7.3

MW = molecular weight IV = intraventricular. IL = intralumbar. IC = intracavity/tumour. IT = intrathecal

RMM - refractory meningeal malignancy, MB = medulloblastoma, LMC = leptomeningeal carcinomatosis, LMM = leptomeningeal meningitis, NM = neoplastic meningitis **3a**

**Agents under trial.**

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of medulloblastoma (MB / PNET) chemosensitivity</b>	<b>Evidence of IT use.</b>	<b>Cell cycle phase specificity</b>	<b>Comments</b>	<b>Refs</b>
Cytarabine (liposomal formulation)	-2.46	4.3	243	Evidence of MB sensitivity, Randomised Clinical Trial in NM had 13/18 responders with IT liposomal v 3/17 with free cytarabine.	Licensed in lymphomatous meningitis in adults. Phase 1 trial in children with NM established maximum tolerated dose and showed benefit in 8/14 pts. Phase 2 trial is in development.	S phase	Neurotoxic in standard formulation. Liposomal preparation used for IT is less toxic and half life in CSF is prolonged up to 40 times, though is less in children than adults. Dexametasone is required to avoid arachnoiditis.	[1, 11, 20, 27, 32, 33, 36, 37, 42, 44, 64, 66, 68-70, 73, 77, 78, 87, 95, 96, 130, 142]
Mafosfamide	0.56,-2.11		401	MB, PNET, ependymoma pts, LMC -rabbit only	IV -non-human primate, IV -rabbit. Phase 1 study in patients with RMM, IV+ IL paediatric patients showed good effect + minimal toxicity. Phase 1 study has determined maximum tolerated dose in children <3yrs with newly diagnosed embryonal tumours - phase 2 trial in progress	non	Cyclophosphamide derivative, undergoes spontaneous hydrolysis to active species	[1, 17, 21, 42, 87, 95, 96, 104, 120, 121, 128]

**Table 3b: Agents suitable for clinical trial.**

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of medulloblastoma (MB / PNET) chemosensitivity</b>	<b>Evidence of IT use.</b>	<b>Cell cycle phase specificity</b>	<b>Comments</b>	<b>Refs</b>
Carboplatin	-0.46		371	High grade gliomas, MB, PNET, ependymoblastoma	IT in rats showed this to be the least neurotoxic of the platins. Neurotoxicity not seen until lethal dose	non		[39, 41, 95, 96, 100, 107, 118, 127]
Etoposide	0.6	9.7	589	MB	IV -dogs, humans IV - metastatic MB, malignant meningitis. Used IV in paediatric patients with good effect + minimal toxicity	Late S, G2	Poor CSF distribution following IV administration, possibly due to rapid CSF clearance due to lipophilicity and/or efflux by P-glycoprotein and MDR-associated protein 1. Possible problem with IL route due to concerns of drug and preservatives causing spinal cord damage.	[1, 40, 42, 87, 95, 96, 112, 114, 121, 126, 128, 133]
Nimustine (ACNU)	0.39 Water + lipid soluble		309	Pilocytic astrocytoma, meningeal spread of MB, PNET, glioblastoma, anaplastic glioma.	MC - rat, dog, IV in meningeal spread of MB, PNET, glioblastoma, anaplastic glioma, A phase1/11 study involving IV/IL/ IT admin in 21 patients (including children) with refractory disseminated MB showed efficacy in some patients	non	When given IV, needed infusion - bolus did not give adequate subarachnoid spread as CSF half-life very short(approximately27min)  High priority for future study in appropriate delivery system	[6, 42, 61, 63, 74-76, 83, 87, 95, 96, 98, 129, 131, 132, 146-149]

**Table 3c: Drugs requiring further investigation before clinical trial.**

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of medulloblastoma (MB / PNET) chemosensitivity</b>	<b>Evidence of IT use.</b>	<b>Cell cycle phase specificity</b>	<b>Comments</b>	<b>Refs</b>
Floxuridine (FdUrd)	-1.16	7.44	246	MB, glioma LMC -animal + humans, NM	LMC, NM by continuous IT infusion. Intracavitary administration may be useful in small volume malignant brain tumours			[1, 42, 87, 91-96, 144]
4-hydroperoxy-cyclophosphamide			293	Promising phase 1 trial IT in MB, PNET	Promising phase 1 trial IT in MB, PNET. IV administration in monkeys	non		[7, 52, 95, 96]

**3d: Drugs requiring further investigation, lower priority.**

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of medulloblastoma (MB / PNET) chemosensitivity</b>	<b>Evidence of IT use.</b>	<b>Cell cycle phase specificity</b>	<b>Comments</b>	<b>Refs</b>
Diaziquone	-0.758	low ionisation at physiological pH	364	9L rat brain tumour, LMC, GBM, AA, paed brain tumours, RMM, recurrent glioma – phase 11 studies, MB xenograft in mice	Rat, IV-non-human primate studies, Human patients including children with RMM – phase 1/11 studies – good response		Low aqueous solubility has caused formulation problems. Drug shows high rate of clearance from CSF	[1, 13, 14, 21, 31, 42, 43, 47, 62, 71, 78, 87, 95, 96, 115, 127, 150]
Mercaptopurine	0.01	7.77, 11.17	170	LMD of ALL	IV - monkey then children	S - phase	Cleared from CSF at 0.63ml/min (bulk flow 0.4ml/min) suggesting an additional mechanism of elimination - possibly efflux by MDR-associated protein 1 and monocarboxylic acid transporters	(25, 59, 60, 91, 105, 107)
Rubitecan			393	safe & effective in rat model against human GBM xenograft NM	IT safe & effective in rat model of GBM NM	S-phase		
Topotecan	0.83, -0.3	6.35, 10.1	458	ependymoma, MB, high grade glioma xenografts	IV non-human primate. IV & IL to children & adults with refractory neoplastic meningitis. Phase 2 study underway.	S-phase	Chemical arachnoiditis dose limiting toxicity. Rapid elimination from CSF. Novel mechanism of action	[1, 10, 18-21, 87, 95, 96, 99, 122, 137]

**Table 3e: Drug with insufficient information to grade**

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of medulloblastoma (MB / PNET) chemosensitivity</b>	<b>Evidence of IT use.</b>	<b>Cell cycle phase specificity</b>	<b>Comments</b>	<b>Refs</b>
Temozolomide	-0.58, -1.32		194	CNS tumour xenografts, NM Phase 2 studies in MB, astrocytoma, glioma, GBM	microcrystalline prep with increased solubility - used IT in rats with NM + malig glioma sub arachnoid xenografts		Spontaneous conversion to active mitomycin in physiological conditions. Highly insoluble in aqueous solution, microcrystalline form increased solubility	[1, 42, 50, 87, 95-97, 111, 124]

Table 4: Drugs with documented activity in the relevant tumour types but no information on intrathecal use.

DRUG	OWPC	pKa	MW	Evidence of tumour sensitivity	Previous IT use.	Cell cycle phase specific	Neuro-toxicity	Comments	Refs
Didemnin B	3.173		1112	Some response seen in GBM and progressive high grade gliomas				Natural marine product, antiviral and antineoplastic activity	[95, 96, 127]
Eflornithine	-2.945		182	In combination with mitoguanzone, response seen in anaplastic astrocytoma and GBM					[95, 96, 127]
Irinotecan	0.03 (pH 1-6) 0.095 (pH9-12). At pH7-slightly more lipophilic but likely to be more hydrophilic than lipophilic	1.07, 7.89	677	peripheral PNET, neuroblastoma xenografts, MB, glioma, ependymoma xenografts. GBM, ependymoma, MB (animal models), some evidence of activity in recurrent glioma		S-phase		Requires metabolism by carboxylesterase in liver & tissues to SN-38 to provide most of its cytotoxic activity – it is not known if this would occur in the CSF	[28, 59, 87, 95, 96, 106, 135, 138]
Lomustine (CCNU)	2.629		234	MB, glioma		non		Alkylating and carbamoylating agent	[10, 42, 62, 85, 87, 95, 96, 114, 141]
Lonidamine	4.407		321	Prolongation of survival + rate of 1 year survivors (62 v 35%) supratentorial glioma			Drowsiness, weakness	Inhibits lactate production causing interference with energy metabolism of cancer cells	[87, 95, 96, 127]
Menogaril	1.066		542	High grade glioma, phase 1 and 11 trials give conflicting results		? non		Anthracycline, therefore likely to be irritant	[42, 87, 95, 96, 127]
Mitolactol (Dibromodulcitol)	-0.426		308	MB, PNET, ependymoma - single agent, moderate effect				long duration of presence in CSF with a half life of around 24 hours	[39, 42, 84, 87, 95, 96, 101, 116, 117]
Teniposide	-0.03(estimated by calculation with low confidence probably higher)	10.13	657	MB in cell culture			As etoposide	efflux by P-glycoprotein	[39, 42, 87, 89, 95, 96, 128, 134]
Tauromustine	-0.299		287	AA, glioblastoma, clinical improvement seen in 19/46 patients		? non			(59, 60, 71)

Table 5: Drugs tested by the intrathecal route (some animal, some human) with mixed reports of toxicity and efficacy:

<b>DRUG</b>	<b>OWPC</b>	<b>pKa</b>	<b>MW</b>	<b>Evidence of tumour sensitivity</b>	<b>Previous IT use IV = intra-</b>	<b>Cell cycle phase specific</b>	<b>Neurotoxicity</b>	<b>Comments</b>	<b>Refs</b>
Bleomycin	-2.57		1416	Intracerebral administration in rat 9L gliosarcoma. Administered safely in liposomes in human cerebral glioma. Given IC in GBM - safe but efficacy doubtful. <25% increase in survival in LMC rat model	phase 1 study IC - no toxicity. Rat, beagle, human (phase 1 IV), depot preparation	Cycling & non-cycling cells M & G2 phases most sensitive	Vascular necrosis after IT (beagle), death after IC administration in craniopharyngioma patient	Antibiotic - binds to DNA - strand scission	[25, 42, 78, 82, 88, 90, 95, 96, 113]
Busulphan	-0.52		246	MB/Glioma/ependyoma - mice. Glioblastoma - rat. MB-mice	Rat	Non	Confusion + seizures - high dose.	Irritant. Bifunctional alkylating agent. Poorly water soluble however water-soluble microcrystalline formulation (Spartaject) been developed. Given IT in rats, non-human primates, adults with LM disease + phase 1 trial in children with LMD from brain tumours	[3, 5, 26, 42, 51, 56, 87, 95, 96, 122, 130]

Carmustine (BCNU)	1.53		214	Intravenous preparation licensed in brainstem glioma, MB, astrocytoma, ependyoma, and metastatic brain tumours.  GLIADEL Implant licensed in high-grade malignant glioma and GBM.	IT - in hybrid liposomes - rat + dog	Non	Convulsions, cerebral oedema in patients with implants. No Adverse Drug Reactions, non-toxic in rat/dog in liposomes. Acute encephalopathy after intracarotid administration, encephalomyelopathy - coma + death with high dose, neuroretinitis	Alkylating and carbamoylating agent. DNA alkylation followed by protein carbomoylation. Causes local venous irritation	[2, 9, 29, 42, 46, 48, 64, 71, 72, 85, 87, 95, 96, 128, 130, 136, 143]
Dacarbazine	-0.24	4.42	182	Melanoma - including primary CNS	Rat. 3 case reports human patients	Non	Very toxic in rat. CNS reaction in one patient case report	Requires demethylation for activation, though dacarbazine itself shows cytotoxic activity. Irritant-extravasation causes tissue damage + severe pain	[9, 12, 34, 42, 78, 85, 87, 95-97, 145]
Fazarabine			496	Lack of activity in phase 2 trial high grade glioma. Solid tumour activity + leukaemia xenografts. Potentially useful for NM from a variety of tumours	IT + IV (Ommaya reservoir) in Rhesus monkeys	Probably s-phase	Transient changes in CSF profiles but no evidence of neurotoxicity after IV in monkey	Cytarabine analogue. Inhibits DNA synthesis + methylation. Nucleoside metabolite, intracellular activation needed by deoxycytidine kinase. Clearance from CSF five times higher than CSF bulk flow rate in monkey	[60, 95, 96, 119]
Ranimustine (MCNU)	-1.29		328	pilocytic astrocytoma	Longer half life than ACNU in dogs. IT admin in 21 patients (including children) with refractory disseminated MB showed efficacy in	Non	Mild histological changes in dog brain after IV admin. IC admin caused marked brain oedema + focal necrosis. Paraplegia + double incontinence in some patients receiving multiple	Nitrosourea	[42, 74, 87, 95, 96, 98, 146, 149]

					some.		IL bolus injections		
Zinostatin				Glioblastoma, astrocytoma	single case report		No CNS toxicity when given via carotid artery	Antineoplastic antibiotic. Does not permeate BBB in normal brain. Half life CSF=50sec	[74, 87, 95, 96, 98, 146]

Table 6: Comparison of candidate intrathecal agents by previous published reviews

Fleischack [49].	Ruggiero [108].	Blaney [17]	Conroy
Intracavitary bleomycin			
			Carboplatin
Cytarabine	Cytarabine		
Liposomal cytarabine (DepoCyte™)	Liposomal cytarabine (DepoCyte™)	Liposomal cytarabine (DepoCyte™)	Liposomal cytarabine (DepoCyte™)
Diaziquone			Diaziquone
Etoposide			Etoposide
			Floxuridine (FdUrd)
			4-Hydroxyperoxy-cyclophosphamide
Mafosfamide	Mafosfamide	Mafosfamide	Mafosfamide
Mercaptopurine	Mercaptopurine		Mercaptopurine
Methotrexate	Methotrexate		
	Monoclonal antibodies		
Nimustine (ACNU)			Nimustine (ACNU)
			Rubitecan
	Temozolomide		Temozolomide
	Thiotepa		
Topotecan		Topotecan	Topotecan

