



HAL
open science

AM1* parameters for manganese and iron

Hakan Kayi, Timothy Clark

► **To cite this version:**

Hakan Kayi, Timothy Clark. AM1* parameters for manganese and iron. *Journal of Molecular Modeling*, 2009, 16 (6), pp.1109-1126. 10.1007/s00894-009-0614-y . hal-00568333

HAL Id: hal-00568333

<https://hal.science/hal-00568333>

Submitted on 23 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Editorial Manager(tm) for Journal of Molecular Modeling
Manuscript Draft

Manuscript Number: JMMO1039R1

Title: AM1* parameters for manganese and iron

Article Type: Original paper

Keywords: AM1*; Iron parameters; Manganese parameters; Semiempirical MO-theory

Corresponding Author: Prof. Tim Clark,

Corresponding Author's Institution: Universitaet Erlangen-Nurnberg

First Author: Hakan Kayi

Order of Authors: Hakan Kayi; Tim Clark

Abstract: We report the parameterization of AM1* for the elements manganese and iron. The basis sets for both metals contain one set each of s-, p- and d orbitals. AM1* parameters are now available for H, C, N, O and F (which use the original AM1 parameters), Al, Si, P, S, Cl, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Br, Zr, Mo, I and Au. The performance and typical errors of AM1* are discussed for Mn and Fe and compared with available NDDO Hamiltonians.

Response to Reviewers: All the suggestions of the reviewer have been incorporated

AM1* parameters for manganese and iron

Received: 28.09.2009 / Accepted: 22.10.2009

Hakan Kayi and Timothy Clark✉

Computer-Chemie-Centrum and Interdisciplinary Center for Molecular Materials,
Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 25, 91052 Erlangen,
Germany

✉E-mail: Tim.Clark@chemie.uni-erlangen.de

Abstract

We report the parameterization of AM1* for the elements manganese and iron. The basis sets for both metals contain one set each of *s*-, *p*- and *d*-orbitals. AM1* parameters are now available for H, C, N, O and F (which use the original AM1 parameters), Al, Si, P, S, Cl, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Br, Zr, Mo, I and Au. The performance and typical errors of AM1* are discussed for Mn and Fe and compared with available NDDO Hamiltonians.

Keywords AM1* · Iron parameters · Manganese parameters · Semiempirical MO-theory

Introduction

We have introduced AM1* [1-7] as an extension of AM1 [8] that uses *d*-orbitals for the elements P, S, Cl, [1] Al, Si, Ti and Zr, [2] Cu and Zn, [3] Br and I, [4] V and Cr, [5] Co and Ni, [6] and Au [7]. The AM1* parameters that are available for molybdenum are a slight modification of Voityuk and Rösch's AM1(d) parameter set [9] to include the AM1* core-core parameters for interactions between Mo and new AM1* elements. The rationalization behind AM1* is to retain the advantages of AM1 (good energies for hydrogen bonds, higher rotation barriers for π -systems than MNDO [10, 11] or PM3 [12-14]) for the first-row elements and improve the performance over AM1 for compounds that contain second-row and heavier elements and to provide a published parameterization for an AM1-like method for the transition metals. As a continuation of this work, we now report AM1* parameters for manganese and iron. Both manganese and iron are important in the chemistry of organometallic and biological catalysts and they play a very important role in biochemical systems especially in the active sites of the enzymes [15, 16].

Theory

AM1* for the two new elements uses the same basic theory as outlined previously. [1, 2] As for other element-H interactions, the core-core repulsion potential for the Mn-H and Fe-H interactions use a distance-dependent term δ_{ij} , rather than the constant term used for core-core potentials for most other interactions in AM1*. [1] This distance-dependent δ_{ij} was also used for the Mo-H and interaction in AM1(d) [9] and for Ti-H, V-H, Cr-H, Co-H, Ni-H, Cu-H, Zn-H, Br-H, Zr-H, Mo-H, I-H and Au-H in AM1* [2-7]. The core-core terms for Mn-H and Fe-H are thus:

$$E^{core}(i-j) = Z_i Z_j \rho_{ss}^0 \left[1 + r_{ij} \delta_{ij} \exp(-\alpha_{ij} r_{ij}) \right] \quad (1)$$

where all terms have the same meaning as given in reference [1].

The standard MNDO/d formula is used for all other core-core interactions:

$$E^{core}(i-j) = Z_i Z_j \rho_{ss}^0 \left[1 + \delta_{ij} \exp(-\alpha_{ij} r_{ij}) \right] \quad (2)$$

The parameterization techniques were those reported in references [1] and [2] and will not be described further here.

Parameterization Data

The target values used for parameterization and their sources are defined in Tables S1, S2 and S3 of the Supplementary Material. We have used heats of formation, Koopmans' theorem ionization potentials, dipole moments and geometrical variables as we did for the Ti, Zr, Cu, Zn, Br, I, V, Cr, Co, Ni and Au parameterizations. [2-7] Because the experimental data for heats of formation of compounds of these two metals are relatively sparse, we have also extended our parameterization data using a small series of model compounds whose heats of formation we have derived from DFT calculations. [17] As before, [1-7] we checked that experimental values for heats of formation were reasonable using DFT calculations.

DFT calculations used the Gaussian 03 suite of programs [18] with the LANL2DZ basis set and standard effective core potentials [19-22] augmented by a set of polarization functions [23] (designated LANL2DZ+pol) and the B3LYP hybrid functional. [24-26]

Experimental parameterization data for manganese and iron were taken largely from the NIST Webbook, [27] but also from the OpenMopac collection [28] and the other experimental and theoretical sources given in the Supplementary Material.

The energetic parameterization data and their sources are given in Table S1 of the Supplementary Material. In addition to the energetic data, geometries, dipole moments and ionization potentials taken from the above sources, crystal structures from the Cambridge Structural Database (CSD) [29] were also used in the parameterization to ensure that not only the energetic and electronic properties for the "prototype" compounds, but also the structures of large manganese and iron compounds are well produced.

Results

Table 1 shows the optimized AM1* parameters. Geometries were optimized with the new AM1* parameterization using VAMP 10.0, [30] while the PM5 calculations used LinMOPAC2.0. [31] Since PM5 was not parameterized for manganese, it was only used for iron compounds. The calculations with PM6 used MOPAC2009. [32] We give related errors for each property, *i.e.*, heats of formation, ionization potentials, dipole moments, geometries as the mean unsigned error (MUE) and mean signed error (MSE). MUE is used to measure the scatter of the calculated results relative to the experimental values, whereas MSE indicates the direction of any systematic deviation of the predictions from experiment.

- Table 1 about here -

Manganese

Heats of formation

The calculated heats of formation for our training set of manganese compounds are shown in Table 2. We have compared our results with Stewart's published PM6 method. [33]

- Table 2 about here -

AM1* reproduces the heats of formation for the training set of manganese compounds used in parameterization better than PM6. However, we note that this comparison is not strictly valid as it is based on the current parameterization data. As can be seen from Table 2, the AM1* parameterization data set contains 78 compounds, of which 44 are taken from original PM6 parameterization data set. These data convincingly demonstrate the influence of the extent of the training data. AM1* performs significantly better for its extended training set, whereas PM6 performs better for the subset for which it was trained. This situation is unavoidable and is a direct consequence of the relative paucity of data for parameterizing semiempirical MO techniques for transition metals.

Both AM1* and PM6 tend to give positive systematic errors for manganese-containing compounds. For AM1* with a mean signed error (MSE) of 2.3 kcal mol⁻¹, this tendency is less pronounced than for PM6 (MSE of 9.8 kcal mol⁻¹).

The mean unsigned error (MUE) for the AM1* parameterization dataset is 32.3 kcal mol⁻¹, compared with 51.0 kcal mol⁻¹ for PM6. PM6 produces especially large errors for the compounds that were not included in its original training set.

The largest errors for AM1* are mainly found for Mn³⁺, Mn²⁺ and Mn⁺ (-167.3, 161.1 and 95.8 kcal mol⁻¹, respectively). Since the ionization potentials (and hence the heats of formation of the cations) of manganese are important determinants of the reactivity of manganese centers, these errors are important. However, we could not detect serious systematic trends caused by these errors. Molecules that give the largest positive errors for AM1* are MnF₃⁻ (113.7 kcal mol⁻¹), MnCl₂O (89.5 kcal mol⁻¹), MnF₂⁻ (79.5 kcal mol⁻¹), MnCl₂ (76.5 kcal mol⁻¹), MnF₂ (75.8 kcal mol⁻¹), MnC₇H₅SO₄ (Mn(Cp)(CO)₂(SO₂), CPCSMN,) (68.2 kcal mol⁻¹), MnO₂⁻ (59.1 kcal mol⁻¹), MnH (58.1 kcal mol⁻¹) and MnH⁻ (55.5 kcal mol⁻¹). The largest negative errors are found for MnC₇N₂H₁₂S₄⁻ (Mn(C₃N₂H₄)(C₂H₄S₂)₂⁻, COWHOL) (-141.9 kcal mol⁻¹), *trans*-Mn(H₂O)₄Cl₂ (-121.5 kcal mol⁻¹), MnZr (-93.2 kcal mol⁻¹), MnC₉H₁₂O (Mn(CO)(C₄H₆)₂, BUTMNC) (-65.5 kcal mol⁻¹), *trans*-Mn(H₂O)₄Br₂ (-61.2 kcal mol⁻¹) and MnF₅⁻ (-60.3 kcal mol⁻¹). AM1* uses the unchanged AM1 parameterization for the elements H, C, N, O and F, which limits the possible accuracy of the parameterization. The large errors with the compounds containing these elements are not surprising as we have pointed out in our previous parameterizations. [3-6] As found for other metals, the large errors in pure AM1* element-containing compounds are likely to be a consequence of our sequential parameterization strategy, in contrast to the simultaneous parameterization used for PM6. [33]

Ionization potentials and dipole moments

A comparison of the calculated and experimental Koopmans' theorem ionization potentials and dipole moments for AM1* and PM6 are given Table 3.

- Table 3 about here -

AM1* with an MUE of 1.66 eV performs better than PM6 (MUE of 2.07 eV) for Koopmans' theorem ionization potentials. Both AM1* and PM6 tend to underestimate ionization potentials for manganese-containing compounds. However, this tendency is more serious for PM6 with an MSE of -2.02 eV, compared to the AM1* MSE of -0.77 eV.

Large AM1* errors are found for MnF₂ (-3.06 eV), MnF₃ (-4.00 eV) and MnF₄ (-3.88 eV). This may be an indirect result of using the original AM1 parameters for fluorine. But, however, PM6 also produces very large errors for MnF₂ (-2.84 eV), MnF₃ (-4.45 eV) and MnF₄ (-6.59 eV). Since, both methods produce these large errors in the same direction, this may be a result of possible erroneous reference values given in NIST. [27]

AM1* with an MSE of 0.03 Debye shows no significant tendency to systematic errors for dipole moments of manganese compounds, whereas PM6 gives a positive systematic error of 0.14 Debye (MSE). AM1* and PM6 perform comparably, with MUEs of 1.25 and 1.27 Debye, respectively. The largest AM1* errors are found for MnOF₂ (4.03 Debye), MnOI₂ (3.36 Debye) and MnI (-2.57 Debye). The large error for MnOF₂ may be an indirect result of using original AM1 parameters for oxygen and fluorine. However, the large errors for iodine-containing compounds are not surprising and result from the known weakness [4, 5] of the AM1* parameterization in reproducing dipole moments for iodine-containing compounds.

Geometries

Table 4 shows a comparison of AM1* and PM6 results in reproducing the geometries of the manganese-containing compounds.

- Table 4 about here -

Both AM1* and PM6 overestimate bond lengths to manganese systematically by 0.02 and 0.14 Å, respectively. AM1*, with an MUE of 0.13 Å performs better than PM6 (MUE of 0.24 Å) for bond lengths. Here we note that PM6 reproduces bond lengths to be very long for the manganese-transition metal diatomic compounds. Parameterization dataset of PM6 clearly does not cover these model compounds which seriously affects statistics. For the bond angles, PM6 with an MUE of 5.0° performs quite well, compared with AM1* (MUE=12.2°). AM1* and PM6 reproduce bond angles for manganese-containing compounds that are on average 3.0° and 2.1° too small.

Iron

Heats of formation

The results obtained for heats of formation of iron-containing compounds are shown in Table 5. We have compared our results with Stewart's published PM6 method [33] and also the unpublished PM5 method implemented in LinMopac [31].

- Table 5 about here -

AM1* reproduces the heats of formation of the training set of iron compounds used in parameterization better than both PM6 and PM5. Once again, one must consider that this comparison is not strictly valid as it is based on the current parameterization data, which sometimes differ from those used in PM6 and PM5. As can be seen from Table 5, the AM1* parameterization data set contains 98 compounds, of which 55 are taken from the original PM6 parameterization data set. Since PM5 is unpublished, we do not know whether these data cover original PM5 parameterization dataset. These data demonstrate the influence of the extent of the training data. AM1* with the mean unsigned error of 26.5 kcal mol⁻¹ performs significantly better for its extended training set than PM6 and PM5 (MUEs of 31.6 and 34.5 kcal mol⁻¹, respectively), whereas PM6 performs better (MUE = 17.8 kcal mol⁻¹) than AM1* and PM5 (MUEs of 32.4 and 33.8 kcal mol⁻¹, respectively) for the subset for which it was trained. As before, we note that, this situation is unavoidable.

AM1* and PM5 underestimate heats of formation for iron compounds by only 3.1 and 5.5 kcal mol⁻¹, respectively. On the other hand, PM6 systematically predicts the heats of formation of iron compounds to be too positive with a mean signed error of 13.7 kcal mol⁻¹. The largest positive errors for AM1* are found for the compounds FeC₁₀H₁₄O₄ (bis(acetylacetonate)iron) (100.8 kcal mol⁻¹), FeC₃₆N₆H₂₄²⁺ (ferrous tris(*ortho* phenanthroline)) (84.3 kcal mol⁻¹), Fe₂I₄ (72.1 kcal mol⁻¹), Fe⁻ (68.3 kcal mol⁻¹) and FeCl₂ (67.4 kcal mol⁻¹). The largest negative errors for AM1* are found for Fe(CN)₆³⁻ (-217.8 kcal mol⁻¹), Fe(CN)₆⁴⁻ (-123.3 kcal mol⁻¹), Fe(H₂O)₅Br⁺ (-73.8 kcal mol⁻¹), Fe(H₂O)₆²⁺ (-69.1 kcal mol⁻¹) and Fe(H₂O)₅F⁺ (-67.9 kcal mol⁻¹). The large negative errors with oxygen-containing compounds are not surprising as we have pointed out in our previous parameterizations. [5, 6] Generally large errors in AM1* for iron are given by the compounds that contain original AM1 elements or AM1 elements with chlorine, bromine or iodine. We attribute this to a weakness in the

AM1* parameterization for the halogens and also general weakness of the original AM1 parameterization.

Ionization potentials and dipole moments

A comparison of the calculated and experimental ionization potentials and dipole moments of iron-containing compounds for AM1*, PM6 and PM5 are shown in Table 6.

- Table 6 about here -

AM1* and PM5 overestimate Koopmans' theorem ionization potentials of iron-containing compounds with MSEs of 0.42 and 0.52 eV, respectively for the dataset used, whereas PM6 underestimates them by 0.63 eV. AM1* with an MUE of 0.93 eV performs better than both PM5 (MUE=1.03 eV) and PM6 (1.29 eV) for ionization potentials of iron compounds. The large errors for AM1* are mainly found for iron-carbonyl compounds, which is once again an indirect result of using original AM1 parameters for carbon and oxygen.

Both AM1* and PM5 tend to give positive systematic errors for dipole moments of iron-containing compounds with MSEs of 0.31 and 0.34 Debye, respectively, whereas PM6 underestimates by 0.35 Debye. AM1* with an MUE of 0.96 Debye performs slightly better than PM5 (MUE=1.01 Debye) and far better PM6 (1.79 Debye) for dipole moments of iron-containing compounds. The large errors for AM1* are mainly obtained from the compounds containing hydrogen, carbon, oxygen and nitrogen. This is not unexpected because AM1* uses the original AM1 parameters for these elements.

Geometries

The geometrical variables used to parameterize AM1* for iron and a comparison of the AM1*, PM6 and PM5 results are shown in Table 7.

- Table 7 about here -

AM1* shows no systematic error trend for bond lengths in iron compounds, whereas both PM5 and PM6 overestimate them by 0.08 Å and 0.12 Å, respectively. AM1* with a mean unsigned error of 0.09 Å performs better than PM5 (MUE=0.15 Å) and far better than PM6 (MUE=0.27 Å) for bond lengths. Here once again we note that the large PM6 errors for bond

lengths resulted from iron-transition metal diatomic model compounds and these compounds are not covered by PM6 parameterization dataset.

All three available methods produce negative systematic errors for bond angles of iron-containing compounds. While AM1* underestimates them by 1.3°, PM5 and PM6 also underestimate by 1.7° and 2.6°, respectively. The performance of AM1* for bond angles in iron-containing compounds is comparable to PM5 and better than PM6. The MUEs for AM1* and PM5 are 5.8° and 4.0°, respectively, and for PM6 10.4°. Here the performance of relatively old method PM5 is surprisingly comparable to or better than more modern methods.

Discussion

In this work, we have presented our new AM1* parameters for manganese and iron providing important additional elements for the chemistry of organometallic and biological catalysts and for biochemical systems especially in the active sites of the enzymes. Since the experimental data for the compounds of these two metals are relatively sparse and sometimes of poor quality, we have extended the range of the parameterization dataset and made it more reliable by checking experimental data by including results from DFT calculations. We aim to produce a parameter set that is more robust and generally applicable than the ones trained only using experimental data. For our extended training set used, AM1* parameterizations for manganese and iron give very good energetic and electronic results and also perform very well for the structural properties. Both AM1* and PM6 extend the range of applicability of NDDO-based MNDO-like techniques and provide good starting points for reaction-specific local parameterizations and comparison calculations.

Acknowledgments

This work was supported by the Deutsche Forschungsgemeinschaft by an individual grant (Cl85/17-1) and as part of GK312 “*Homogeneous and Heterogeneous Electron Transfer*” and SFB583 “*Redox-Active Metal Complexes: Control of Reactivity via Molecular Architecture*”.

We thank Dr. Paul Winget, Dr. Bodo Martin, Dr. Cenk Selçuki and Anselm Horn for support with the parameterization database.

Supplementary material

The values and the sources of the parameterization data

References

1. Winget P, Horn AHC, Selçuki C, Martin B, Clark T (2003) *J Mol Model* 9:408-414
2. Winget P, Clark T (2005) *J Mol Model* 11:439–456
3. Kayi H, Clark T (2007) *J Mol Model* 13:965-979
4. Kayi H, Clark T (2009) *J Mol Model* 15:295-308
5. Kayi H, Clark T (2009) *J Mol Model* 15:1253-1269
6. Kayi H, Clark T (2009) *J Mol Model* 15 (online first) DOI: 10.1007/s00894-009-0503-4
7. Kayi H (2009) *J Mol Model* 15, in the press
8. Dewar MJS, Zoebisch EG, Healy EF, Stewart JJP (1985) *J Am Chem Soc* 107:3902-3909
9. Voityuk AA, Rösch N (2000) *J Phys Chem A* 104:4089-4094
10. Dewar MJS, Thiel W (1977) *J Am Chem Soc* 99(15):4899-4907
11. Thiel W (1998) In: Schleyer PvR, Allinger NL, Clark T, Gasteiger J, Kollman PA, Schaefer III HF, Schreiner PR (eds) *Encyclopedia of Computational Chemistry*. Wiley, Chichester, p 1599
12. Stewart JJP (1989) *J Comp Chem* 10:209-220
13. Stewart JJP (1989) *J Comp Chem* 10:221-264
14. Stewart JJP (1998), In: Schleyer PvR, Allinger NL, Clark T, Gasteiger J, Kollman PA, Schaefer III HF, Schreiner PR (eds) *Encyclopedia of Computational Chemistry*. Wiley, Chichester, p 2080
15. Rutkowska-Zbik D, Witko M (2006) *J Mol Catal A: Chem* 258:376-380
16. Carrasco R, Morgernstern-Badarau I, Cano J (2007) *Inorg Chim Acta* 360:91-101
17. Winget P, Clark T (2004) *J Comput Chem* 25:725-733
18. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Montgomery JA Jr, Vreven T, Kudin KN, Burant JC, Millam JM, Iyengar SS, Tomasi J, Barone V, Mennucci B, Cossi M, Scalmani G, Rega N, Petersson GA, Nakatsuji H, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Klene M, Li X, Knox JE, Hratchian HP, Cross JB, Bakken V, Adamo C, Jaramillo J, Gomperts R, Stratmann RE, Yazyev O, Austin AJ, Cammi R, Pomelli C, Ochterski JW, Ayala PY, Morokuma K, Voth GA, Salvador P, Dannenberg JJ, Zakrzewski VG, Dapprich S, Daniels AD, Strain MC, Farkas O, Malick DK, Rabuck AD, Raghavachari K, Foresman JB, Ortiz JV, Cui Q, Baboul AG, Clifford S, Cioslowski J, Stefanov BB, Liu G, Liashenko A, Piskorz P, Komaromi I, Martin RL, Fox DJ, Keith T,

- Al-Laham MA, Peng CY, Nanayakkara A, Challacombe M, Gill PMW, Johnson B, Chen W, Wong MW, Gonzalez C, Pople JA (2004) Gaussian 03. Gaussian Inc, Wallingford CT
19. Dunning Jr TH, Hay PJ, (1976) In: Schaefer III HF (ed) Modern Theoretical Chemistry, vol 3. Plenum, New York, pp 1-28
 20. Hay PJ, Wadt WR (1985) J Chem Phys 82(1):270-283
 21. Hay PJ, Wadt WR (1985) J Chem Phys 82(1):284-298
 22. Hay PJ, Wadt WR (1985) J Chem Phys 82(1):299-310
 23. Frisch MJ, Pople JA, Binkley JS (1984) J Chem Phys 80:3265-3269
 24. Becke AD (1988) Phys Rev A 38:3098
 25. Lee C, Yang W, Parr RG (1988) Phys Rev B 37:785
 26. Becke AD (1993) J Chem Phys 98:5648-5652
 27. NIST Chemistry WebBook, NIST Standard Reference Database Number 69, (<http://webbook.nist.gov/>), Linstrom P, Mallard W, National Institute of Standards and Technology: Gaithersburg MD, 20899, 2003
 28. Stewart JJP, <http://openmopac.net/files.html>
 29. Cambridge Structural Database, Version 5.28 (2007) Cambridge Crystallographic Data Centre, Cambridge, UK
 30. Clark T, Alex A, Beck B, Chandrasekhar J, Gedeck P, Horn AHC, Hutter M, Martin B, Rauhut G, Sauer W, Schindler T, Steinke T (2008) Computer-Chemie-Centrum, Universität Erlangen-Nürnberg: Erlangen
 31. Stewart JJP, (2002) LinMOPAC2.0, FQS Poland: Krakow
 32. Stewart JJP, (2008) MOPAC2009, Stewart Computational Chemistry, Colorado Springs, CO, USA; <http://OpenMOPAC.net>
 33. Stewart JJP (2007) J Mol Model 13:1173-1213

Table 1 AM1* parameters for the elements Mn and Fe

Parameter	Mn	Fe
U_{ss} [eV]	-33.2275445	-99.00
U_{pp} [eV]	-27.2535784	-38.200
U_{dd} [eV]	-69.8244104	-83.10
ζ_s [bohr ⁻¹]	1.9646322	2.8351398
ζ_p [bohr ⁻¹]	18.8965365	37.1808614
ζ_d [bohr ⁻¹]	1.2348158	2.2895425
β_s [eV]	-8.1782339	-20.4335137
β_p [eV]	-21.7780791	-66.1689990
β_d [eV]	-6.2336234	-26.5282533
g_{ss} [eV]	5.0059282	8.4047951
g_{pp} [eV]	13.6386827	10.4447975
g_{sp} [eV]	12.3772691	7.6833614
g_{p2} [eV]	96.0815576	13.9440172
h_{sp} [eV]	4.5810741	1.6747978
Z_{sn} [bohr ⁻¹]	26.0703955	75.1015032
Z_{pn} [bohr ⁻¹]	1.9830213	0.7086051
Z_{dn} [bohr ⁻¹]	1.9132012	1.4519328
$\rho(\text{core})$ [bohr ⁻¹]	1.2307875	1.0769040
$\Delta H_f^\circ(\text{atom})$ [kcal mol ⁻¹]	67.701	99.3
F_{sd}^0 [eV]	3.3270203	14.1017174
G_{sd}^2 [eV]	1.8796412	13.4789419
$\alpha(ij)$		
H	4.1317380	3.6162132
C	2.8147514	4.0432263
N	1.8883089	3.8755726
O	3.0560315	4.9080254
F	2.5828364	5.3397259
Al	3.5620838	2.0238999
Si	4.2623991	2.6259521
P	2.4072008	2.9339288
S	2.7156160	2.7835642
Cl	2.5923979	2.4937637
Ti	2.3993667	1.4496317
V	3.8190630	1.9484452
Cr	2.7010078	2.0835714
Mn	3.1647134	2.4029678
Fe	2.4029678	3.3225842
Co	2.2792673	2.6623880
Ni	4.5772694	4.4251916
Cu	2.2179615	4.1806484
Zn	4.5907029	4.4519730
Br	3.6890036	2.8134065
Zr	2.6907788	1.7761525
Mo	3.0996586	1.9309316
I	4.1561815	3.5275630
Au	3.0829513	2.8842586
$\delta(ij)$		
H	-12.2571050	-10.2036220
C	3.8500280	41.7471647
N	0.6216587	18.5709816

O	6.1303519	98.7945881
F	1.8592303	98.4859864
Al	32.1147597	4.4417867
Si	79.1063414	7.6467930
P	2.9611118	21.4786698
S	3.7385117	8.4757133
Cl	2.4354976	3.7093945
Ti	1.5477850	1.1499629
V	127.1273363	5.0747072
Cr	7.1369564	3.3205625
Mn	5.9226904	4.7319791
Fe	4.7319791	20.2552175
Co	2.1958330	8.1077071
Ni	265.4707317	178.9982694
Cu	1.8893296	96.7779734
Zn	87.3866849	52.0532649
Br	25.1186833	4.3023002
Zr	3.2445347	2.7554220
Mo	14.0341168	3.2515086
I	53.7439446	43.5771427
Au	10.1393842	7.0870550

Table 2 Calculated AM1* and PM6 heats of formation and errors compared with our target values for the manganese-containing compounds used to parameterize AM1* (all values kcal mol⁻¹). Errors are classified by coloring the boxes in which they appear. Green indicates errors lower than 10 kcal mol⁻¹, yellow 10-20 kcal mol⁻¹ and pink those greater than 20 kcal mol⁻¹. The codenames within parentheses indicate the CSD-names of the compounds

Compound	Target ΔH_f°	AM1*		PM6	
		ΔH_f°	Error	ΔH_f°	Error
Mn	67.7	67.7	0.0	67.7	0.0
Mn ⁻	90.2	151.8	61.6	97.9	7.7
Mn ⁺	239.1	334.9	95.8	224.8	-14.3
Mn ²⁺	599.8	760.9	161.1	566.7	-33.1
Mn ³⁺	1376.0	1208.7	-167.3	1053.2	-322.8
Mn ₂	184.3	203.7	19.4	128.2	-56.1
MnH	57.3	115.4	58.1	114.2	56.9
MnH ⁻	41.2	96.7	55.5	124.2	83.0
MnH ₂	114.1	117.0	2.9	114.6	0.5
Mn(C ₅ H ₅) ₂	66.2	100.8	34.6	95.4	29.2
MnN	123.1	121.1	-2.0	99.2	-23.9
Mn(NH ₃) ₆ ²⁺	269.5	275.4	5.9	229.9	-39.6
MnO ₂ ⁻	-62.1	-3.0	59.1	-35.3	26.8
MnO ₃ ⁻	-127.1	-97.5	29.6	-127.7	-0.6
MnO ₄ ⁻	-158.4	-149.7	8.7	-168.8	-10.4
Mn(H ₂ O) ₆ ²⁺	65.7	9.6	-56.1	50.6	-15.1
Mn(CO) ₃ ⁻	-54.8	-58.6	-3.8	-113.0	-58.1
Mn(CO) ₄ ⁻	-122.3	-131.1	-8.7	-157.7	-35.3
Mn(CO) ₅	-178.0	-152.9	25.1	-174.4	3.6
Mn(CO) ₅ H	-176.8	-180.2	-3.4	-170.4	6.4
Mn(CO) ₆ ⁺	-49.7	-67.8	-18.1	-78.8	-29.1
Mn(CO) ₅ CH ₃	-179.5	-196.5	-17.0	-179.5	0.0
Mn(CO) ₅ C ₆ H ₅	-140.8	-169.9	-29.1	-151.3	-10.5
Mn(CO) ₅ (C ₆ H ₅ CH ₂)	-151.9	-173.7	-21.8	-154.6	-2.7
Mn(CO) ₅ COCH ₃	-215.3	-228.5	-13.2	-230.3	-15.0
Mn(CO) ₅ (C ₆ H ₅ CO)	-173.2	-195.1	-21.9	-192.3	-19.1
MnC ₁₅ H ₂₁ O ₆ (Mn(acac) ₃)	-285.2	-280.0	5.2	-267.2	18.0
MnC ₆ H ₈ O ₁₀ ⁻ (KAMMND)	-543.1	-576.9	-33.8	-537.4	5.7
MnC ₉ H ₁₂ O (BUTMNC)	11.9	-53.6	-65.5	22.8	10.9
MnC ₈ H ₅ O ₃ (cyclopentadienyl manganese)	-115.2	-65.3	49.9	-81.5	33.7

tricarbonyl)					
MnC ₉ H ₇ O ₃ (HEXMNC)	-95.5	-89.4	6.1	-83.9	11.6
Mn ₂ (CO) ₉	-314.5	-310.0	4.5	-323.3	-8.8
Mn ₂ (CO) ₁₀	-385.9	-378.7	7.2	-336.8	49.1
Mn(NO)	8.5	47.9	39.4	98.1	89.6
Mn(CO)(NO) ₃	-60.6	-52.9	7.7	-23.0	37.6
MnC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (EDTMNK01)	-388.3	-407.0	-18.7	-413.1	-24.8
MnF ₂	-126.2	-50.4	75.8	-76.8	49.4
MnF ₂ ⁻	-226.0	-146.5	79.5	-106.1	119.9
MnF ₃	-188.0	-173.3	14.7	-139.0	49.0
MnF ₃ ⁻	-272.2	-158.5	113.7	-164.8	107.4
MnF ₄	-231.0	-280.9	-49.9	-170.0	61.0
MnF ₄ ⁻	-343.6	-308.5	35.1	-250.3	93.3
MnF ₅ ⁻	-373.0	-433.3	-60.3	-300.3	72.7
Mn ₂ F ₅ ⁻	-448.3	-440.7	7.6	-257.2	191.1
Mn ₂ F ₇ ⁻	-600.0	-630.6	-30.6	-392.6	207.4
MnOF ₂	-125.8	-120.5	5.3	-114.7	11.1
Mn(H ₂ O) ₄ F ₂ trans	-333.2	-366.8	-33.6	-315.4	17.8
MnC ₇ HO ₅ F ₂ (CDFVMN)	-252.2	-273.1	-20.9	-314.4	-62.2
Mn(CO) ₅ CF ₃	-330.5	-355.7	-25.2	-363.7	-33.2
Mn(CO) ₅ COCF ₃	-359.7	-378.5	-18.8	-381.9	-22.2
MnC ₇ O ₇ F ₃ (FACMNA)	-437.0	-437.3	-0.3	-431.9	5.1
MnAl	116.5	115.6	-0.9	77.3	-39.2
MnF ₃ AlF ₄ ⁻	-704.0	-734.6	-30.6	-220.0	484.0
MnSi	153.2	127.8	-25.4	-275.3	-428.5
MnSiH	93.9	108.2	14.3	-30.9	-124.8
MnSiH ₂	85.2	91.6	6.4	131.5	46.3
MnP	81.9	81.9	0.0	110.0	28.1
MnPH	67.6	77.4	9.8	82.1	14.5
MnC ₉ H ₁₁ SO ₂ (VALXAI)	-76.7	-109.1	-32.4	-54.7	22.0
MnC ₇ H ₅ SO ₄ (CPCSMN)	-125.0	-56.8	68.2	-119.5	5.5
MnC ₇ N ₂ H ₁₂ S ₄ ⁻ (COWHOL)	-19.9	-161.8	-141.9	-37.2	-17.3
MnC ₇ H ₁₀ S ₂ ClO ₃ (GEQLUJ)	-179.2	-198.5	-19.3	-161.0	18.2
MnCl	11.3	69.7	58.4	40.9	29.6
Mn(CO) ₅ Cl (ZOSWEJ)	-219.5	-240.6	-21.1	-220.0	-0.5
MnCl ₂	-63.0	13.5	76.5	-31.2	31.8
Mn(H ₂ O) ₄ Cl ₂ trans	-263.1	-384.6	-121.5	-275.3	-12.2
Mn ₂ Cl ₄	-168.2	-163.0	5.2	-30.9	137.3

Table 3 Calculated AM1* and PM6 Koopmans' theorem ionization potentials and dipole moments for manganese-containing compounds. The errors are color coded as follows: green up to 0.5 eV or 0.5 Debye; yellow between 0.5 and 1.0; pink larger than 1.0

Compound	Target	AM1*		PM6	
			Error		Error
Koopmans' Theorem Ionization Potentials for Manganese Compounds (eV)					
MnH	7.80	9.39	1.59	5.82	-1.98
MnO	8.70	10.58	1.88	7.66	-1.04
Mn(CO) ₅	8.10	9.18	1.08	7.44	-0.66
Mn(CO) ₅ H	8.85	8.71	-0.14	8.44	-0.41
MnF ₂	11.40	8.34	-3.06	8.56	-2.84
MnF ₃	12.60	8.60	-4.00	8.15	-4.45
MnF ₄	13.50	9.62	-3.88	6.91	-6.59
MnCl ₂	11.03	9.67	-1.36	8.11	-2.92
Mn(CO) ₅ Br	8.83	8.52	-0.31	8.74	-0.09
MnBr ₂	10.30	9.70	-0.60	8.78	-1.52
Mn(CO) ₅ I	8.40	8.71	0.31	8.65	0.25
		AM1*		PM6	
N=11					
MSE		-0.77		-2.02	
MUE		1.66		2.07	
Dipole Moments for Manganese Compounds (Debye)					
MnH	0.50	0.75	0.25	0.99	0.49
MnN	3.07	4.38	1.31	3.98	0.91
MnNO	5.89	4.46	-1.43	4.45	-1.44
MnO	4.72	4.57	-0.15	3.74	-0.98
MnOH	0.40	1.64	1.24	3.13	2.73
MnF	2.27	0.67	-1.60	4.23	1.96
MnOF ₂	0.53	4.56	4.03	0.61	0.08
MnAl	1.67	1.60	-0.07	0.19	-1.48
MnSi	3.27	2.85	-0.42	4.03	0.76
MnSiH	3.15	2.75	-0.40	4.03	0.88
MnSiH ₂	3.22	2.54	-0.68	3.96	0.74
MnP	2.54	2.88	0.34	2.70	0.16
MnPH	4.34	3.12	-1.22	2.97	-1.37
MnS	4.22	3.91	-0.31	1.56	-2.66

MnCl	2.36	0.81	-1.55	3.59	1.23
MnBr	4.29	3.10	-1.19	7.72	3.43
MnOBr ₂	0.51	2.12	1.61	0.56	0.05
MnI	4.25	1.68	-2.57	2.13	-2.12
MnOl ₂	0.94	4.30	3.36	0.30	-0.64
		AM1*		PM6	
N=19					
MSE		0.03		0.14	
MUE		1.25		1.27	

Table 4 Calculated AM1* and PM6 bond lengths and angles for manganese-containing compounds. The codenames within parentheses indicate the CSD-names of the compounds. The errors are color coded as follows: green up to 0.05 Å or 0.5°; yellow between 0.05-0.1 Å or 0.5-1.0°; pink larger than 0.1 Å or 1°

Compound	Variable	Target	AM1*		PM6	
				Error		Error
Mn ₂	Mn-Mn	2.29	2.40	0.11	2.89	0.60
MnH	Mn-H	1.74	1.91	0.18	1.68	-0.05
MnH ⁻	Mn-H	1.77	1.83	0.06	1.66	-0.11
Mn(C ₅ H ₅) ₂	Mn-C	2.42	2.79	0.37	2.37	-0.05
MnO ₂ ⁻	Mn-O	1.62	1.86	0.24	1.63	0.02
MnO ₃ ⁻	Mn-O	1.59	1.85	0.26	1.62	0.03
MnO ₄ ⁻	Mn-O	1.64	1.86	0.22	1.63	-0.01
Mn(H ₂ O) ₆ ²⁺	Mn-O	2.14	2.21	0.07	2.15	0.01
	Mn-O	2.19	2.23	0.04	2.17	-0.02
	Mn-O	2.20	2.27	0.07	2.17	-0.03
Mn(CH ₃)O ₃	Mn=O	1.59	1.84	0.25	1.60	0.01
	Mn-C	1.99	1.91	-0.08	1.93	-0.06
Mn(CO) ₃ ⁻	Mn-C	1.82	1.97	0.16	1.82	0.01
Mn(CO) ₅	Mn-C	1.82	1.89	0.07	1.85	0.03
	Mn-C	1.86	1.90	0.04	1.89	0.03
Mn(CO) ₅ H	Mn-C	1.86	1.93	0.07	1.84	-0.02
	Mn-C	1.85	1.92	0.07	1.86	0.01
	H-Mn-C	85.5	90.7	5.2	83.5	-2.0
Mn(CO) ₅ CH ₃	Mn-C	1.83	1.90	0.07	1.85	0.02
	Mn-C	1.85	1.94	0.09	1.85	0.00
	Mn-C	2.19	1.99	-0.21	2.08	-0.12
Mn(CO) ₆ ⁺	Mn-C	1.91	1.95	0.04	1.89	-0.02
Mn(CO) ₅ C ₆ H ₅	Mn-C	1.83	1.90	0.07	1.85	0.02
	Mn-C	1.85	1.94	0.09	1.86	0.01
	Mn-C	2.15	1.99	-0.16	1.99	-0.16
Mn(CO) ₅ (C ₆ H ₅ CH ₂)	Mn-C	1.82	1.90	0.08	1.85	0.03
	Mn-C	1.85	1.94	0.09	1.85	0.00
	Mn-C	2.26	2.02	-0.24	2.09	-0.17
Mn(CO) ₅ COCH ₃	Mn-C	1.84	1.91	0.07	1.85	0.01
	Mn-C	1.85	1.93	0.08	1.88	0.03
	Mn-C	2.17	2.02	-0.15	1.97	-0.20
Mn(CO) ₅ (C ₆ H ₅ CO)	Mn-C	1.84	1.90	0.07	1.86	0.02

	Mn-C	1.85	1.93	0.08	1.87	0.02
	Mn-C	2.18	2.03	-0.15	1.97	-0.20
MnC ₆ H ₈ O ₁₀ ⁻ (KAMMND)	Mn-O	1.90	2.07	0.17	1.93	0.03
	Mn-O	2.30	2.28	-0.02	2.44	0.14
	O-Mn-O	91.7	74.1	-17.6	88.3	-3.4
MnC ₉ H ₁₂ O (BUTMNC)	Mn-C	1.81	1.92	0.11	1.81	0.00
	Mn-C	2.15	1.98	-0.17	2.11	-0.04
Mn(CO) ₃ Cp (CPMNCO)	Mn-C(ring)	2.14	2.21	0.07	2.16	0.02
	Mn-C(O)	1.79	1.93	0.14	1.81	0.02
	C-Mn-C	91.4	101.0	9.6	92.6	1.2
MnC ₉ H ₇ O ₃ (HEXMNC)	Mn-C	2.15	2.12	-0.03	2.17	0.02
	Mn-C	1.78	1.92	0.14	1.82	0.04
Mn ₂ C ₈ H ₁₂ O ₈ (Mn ₂ (Ac) ₄)	Mn-Mn	3.31	2.43	-0.88	3.28	-0.03
Mn ₂ (CO) ₁₀	Mn-Mn	2.89	2.36	-0.53	2.70	-0.19
Mn ₂ (CO) ₁₀ C ₂ (JIPVOT)	Mn-C	1.86	1.93	0.07	1.84	-0.02
	Mn-C	1.83	1.90	0.07	1.83	0.00
	C-Mn-C	85.9	87.6	1.7	85.4	-0.5
MnN	Mn#N	1.59	1.52	-0.07	1.57	-0.02
MnNO	Mn-N	1.73	1.76	0.03	1.74	0.01
Mn(NH ₃) ₆ ²⁺	Mn-N	2.16	2.21	0.05	2.15	-0.01
Mn(CO)(NO) ₃	Mn-N	1.70	1.67	-0.03	1.64	-0.06
	Mn-C	1.87	1.94	0.06	1.93	0.06
	C-Mn-N	106.0	92.9	-13.1	107.7	1.8
MnC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (EDTMNK01)	Mn-O	2.02	2.15	0.13	1.95	-0.07
	Mn-O	1.89	2.16	0.27	1.98	0.09
	Mn-N	2.24	2.11	-0.13	2.10	-0.14
	O-Mn-O	88.3	89.7	1.4	84.2	-4.1
	N-Mn-O	77.5	82.2	4.7	83.7	6.2
MnC ₁₂ N ₂ H ₁₂ O ₆ (XOYHAU)	Mn-O	2.15	2.19	0.04	2.33	0.18
	Mn-N	2.26	1.81	-0.45	1.84	-0.42
	O-Mn-O	85.4	66.7	-18.7	62.5	-22.9
MnC ₁₂ N ₂ H ₁₆ O ₄ (Mn phenanthroline tetrahydrate)	Mn-O	2.16	2.33	0.17	3.45	1.29
MnF	Mn-F	1.82	2.14	0.32	1.87	0.05
MnF ₂	Mn-F	1.79	1.84	0.05	1.78	-0.01
MnF ₂ ⁻	Mn-F	1.86	1.88	0.03	1.83	-0.03
MnOF ₂	Mn-O	1.55	1.89	0.34	1.57	0.02
	Mn-F	1.72	1.79	0.06	1.73	0.01

	O-Mn-F	121.3	102.0	-19.3	124.1	2.8
Mn(H ₂ O) ₄ F ₂ trans	Mn-F	1.98	1.87	-0.11	1.78	-0.20
	Mn-O	2.13	2.35	0.22	2.34	0.21
MnC ₇ HO ₅ F ₂ (CDFVMN)	Mn-C	1.72	1.90	0.18	1.85	0.13
	Mn-C	1.77	1.94	0.17	1.86	0.09
	Mn-C	1.94	2.03	0.09	1.97	0.03
	C-Mn-C	93.8	87.6	-6.2	93.3	-0.5
MnF ₃	Mn-F	1.79	1.79	0.00	1.76	-0.03
	F-Mn-F	120.0	141.7	21.7	136.1	16.1
MnC ₇ O ₇ F ₃ (FACMNA)	Mn-C	1.81	1.89	0.08	1.82	0.01
	Mn-O	2.03	2.29	0.26	2.05	0.02
Mn ₂ (CO) ₈ (CF ₂) ₂ (DOFPET)	Mn-Mn	2.66	2.42	-0.24	2.73	0.07
	Mn-C(F ₂)	2.02	2.10	0.08	1.98	-0.04
	Mn-C(O)	1.88	1.96	0.08	1.87	-0.01
	C-Mn-Mn	49.4	54.8	5.4	46.4	-3.0
MnAl	Mn-Al	2.34	2.34	0.00	2.05	-0.28
Mn(AlH ₂) ₂	Mn-Al	2.41	2.42	0.00	2.15	-0.26
MnSi	Mn-Si	2.34	2.25	-0.10	2.45	0.11
MnSiH	Mn-Si	2.34	2.34	0.00	2.51	0.17
MnSiH ₂	Mn-Si	2.38	2.44	0.06	2.60	0.22
MnSi ₂ C ₉ H ₁₅ O ₅ (KIRYUF)	Mn-Si	2.34	2.39	0.05	2.34	0.00
	Mn-O	2.95	2.99	0.04	2.99	0.04
	Si-Mn-Si	71.2	81.5	10.3	68.8	-2.4
MnP	Mn-P	1.97	2.06	0.09	2.15	0.18
MnPH	Mn-P	2.22	2.21	-0.02	2.23	0.01
MnS	Mn-S	1.99	2.14	0.15	2.01	0.02
MnS ₂	Mn-S	2.20	2.00	-0.20	2.02	-0.18
MnC ₉ H ₁₁ SO ₂ (VALXAI)	Mn-S	2.27	2.25	-0.02	2.18	-0.09
	Mn-C(O)	1.77	2.01	0.24	1.79	0.02
	Mn-C(C ₄)	2.12	2.10	-0.02	2.19	0.07
MnC ₇ H ₅ SO ₄ (CPCSMN)	Mn-S	2.04	2.38	0.34	2.10	0.06
	Mn-C	1.79	1.99	0.20	1.82	0.03
	Mn-C'	2.10	2.10	0.00	2.18	0.08
	S-Mn-C	91.0	80.8	-10.1	92.6	1.6
MnC ₇ N ₂ H ₁₂ S ₄ ⁻ (COWHOL)	Mn-S	2.32	2.33	0.01	2.28	-0.04
	S-Mn-S	157.5	144.9	-12.6	156.2	-1.3
MnC ₇ H ₁₀ S ₂ ClO ₃ (GEQLUJ)	Mn-S	2.39	2.35	-0.04	2.26	-0.13
	Mn-C	1.80	1.92	0.12	1.79	-0.01

	Mn-Cl	2.38	2.17	-0.21	2.31	-0.07
	S-Mn-Cl	92.2	87.7	-4.5	88.6	-3.6
MnC ₄ N ₂ H ₈ S ₂ O ₆ (WIFSEJ)	Mn-S	2.61	2.26	-0.35	2.21	-0.40
	Mn-O	2.20	2.23	0.03	2.13	-0.07
	O-Mn-S	87.3	111.8	24.5	80.6	-6.7
MnCl	Mn-Cl	2.12	2.31	0.19	2.13	0.01
Mn(CO) ₅ Cl (ZOSWEJ)	Mn-Cl	2.37	2.13	-0.24	2.23	-0.14
	Mn-C	1.81	1.91	0.10	1.83	0.02
	Mn-C	1.89	1.94	0.05	1.86	-0.03
Mn(H ₂ O) ₄ Cl ₂ trans	Mn-Cl	2.36	2.21	-0.15	1.98	-0.38
	Mn-O	2.10	2.33	0.23	2.21	0.11
	Mn-O	2.50	2.41	-0.09	3.70	1.20
	Mn-O	2.52	2.59	0.07	3.79	1.27
Mn ₂ Cl ₄	Mn-Mn	2.29	2.38	0.09	2.49	0.20
	Mn-Cl	2.13	2.10	-0.03	1.96	-0.18
MnCl ₂ O	Mn-O	1.55	1.89	0.34	1.57	0.02
	Mn-Cl	2.10	2.08	-0.02	1.87	-0.23
	O-Mn-Cl	121.5	86.3	-35.1	119.1	-2.3
MnC ₁₂ N ₄ H ₁₂ Cl ₂ O (MUPYUR)	Mn-Cl	2.47	2.12	-0.35	2.15	-0.32
	Mn-N(py)	2.26	1.81	-0.45	1.92	-0.34
	Mn-N'	2.34	1.94	-0.40	1.94	-0.40
	N-Mn-N'	95.2	96.2	1.0	94.7	-0.5
MnTi	Mn-Ti	2.25	2.25	0.00	3.62	1.37
MnV ₂	Mn-V	2.28	2.21	-0.07	4.90	2.62
MnCr	Mn-Cr	2.35	2.35	0.00	5.90	3.55
MnCo	Mn-Co	2.44	2.27	-0.18	3.39	0.95
MnNi	Mn-Ni	2.50	2.41	-0.08	6.02	3.52
MnCu	Mn-Cu	2.50	2.50	0.00	7.67	5.17
MnZn	Mn-Zn	2.62	2.62	0.00	2.29	-0.33
MnBr	Mn-Br	2.25	2.41	0.16	2.56	0.31
MnBr ₂	Mn-Br	2.34	2.33	-0.01	2.23	-0.11
MnOBr ₂	Mn-Br	2.24	2.29	0.05	2.31	0.07
	Mn-O	1.55	1.88	0.33	1.56	0.01
	O-Mn-Br	120.7	104.2	-16.5	125.5	4.8
Mn(H ₂ O) ₄ Br ₂ trans	Mn-Br	2.50	2.40	-0.10	2.45	-0.05
	Mn-O	2.10	2.23	0.13	2.08	-0.02
Mn(CO) ₅ Br	Mn-Br	2.56	2.39	-0.17	2.49	-0.07
	Mn-C	1.82	1.90	0.09	1.82	0.01

	Mn-C'	1.88	1.94	0.07	1.86	-0.01
MnBr ₄ ²⁻ (PYDMNB)	Mn-Br	2.50	2.43	-0.07	2.44	-0.06
MnAlC ₆ Br ₃ O ₅ (CMNCXA10)	Mn-Al	3.53	3.63	0.10	3.45	-0.08
	Mn-C	1.95	1.97	0.02	1.92	-0.03
	Mn-C'	1.87	1.93	0.06	1.87	0.00
	Br-Al-Mn	140.2	117.0	-23.2	126.9	-13.3
MnZr	Mn-Zr	2.43	2.43	0.00	3.02	0.59
MnMo	Mn-Mo	2.27	2.27	0.00	4.71	2.44
MnI	Mn-I	2.46	2.49	0.04	2.29	-0.17
MnOI ₂	Mn-I	2.45	2.33	-0.12	2.44	-0.01
	Mn-O	1.55	1.89	0.34	1.55	0.00
	O-Mn-I	120.1	112.7	-7.4	116.1	-4.0
Mn(H ₂ O) ₄ I ₂ trans	Mn-I	2.73	2.44	-0.29	2.52	-0.21
	Mn-O	2.08	2.27	0.19	2.24	0.16
Mn(CO) ₅ I (VUCFAA)	Mn-I	2.69	2.39	-0.30	2.62	-0.07
	Mn-C	1.84	1.91	0.07	1.82	-0.02
MnC ₁₃ N ₂ H ₈ O ₃ I (YOMHEN)	Mn-I	2.72	2.36	-0.36	2.70	-0.02
	Mn-N	2.05	1.98	-0.07	1.92	-0.13
	Mn-C	1.77	1.93	0.16	1.84	0.07
	I-Mn-N	87.7	87.1	-0.6	83.3	-4.4
Mn ₂ (CO) ₈ I ₂ (SIZYUV)	Mn-Mn	3.98	3.98	0.00	3.96	-0.02
	Mn-I	2.70	2.51	-0.19	2.66	-0.04
	Mn-C	1.88	1.93	0.05	1.85	-0.03
	I-Mn-Mn	42.6	37.8	-4.8	43.6	1.0
MnI(SH)	Mn-I	2.41	2.36	-0.05	2.41	0.00
	Mn-S	2.10	2.27	0.17	2.05	-0.05
	I-Mn-S	115.6	146.2	30.6	101.6	-14.0
MnI(F)(PH ₂)(SH)	Mn-I	2.50	2.39	-0.12	2.46	-0.04
	Mn-F	1.75	1.79	0.04	1.73	-0.02
	Mn-P	2.27	2.25	-0.02	2.28	0.01
	Mn-S	2.16	2.22	0.06	2.16	0.00
				AM1*		PM6
N=151						
MSE bond length				0.02		0.14
MUE bond length				0.13		0.24
N=25						
MSE bond angle				-3.0		-2.1
MUE bond angle				12.2		5.0

Table 5 Calculated AM1*, PM6 and PM5 heats of formation and errors compared with our target values for the iron-containing compounds used to parameterize AM1* (all values kcal mol⁻¹). Errors are classified by coloring the boxes in which they appear. Green indicates errors lower than 10 kcal mol⁻¹, yellow 10-20 kcal mol⁻¹ and pink those greater than 20 kcal mol⁻¹. The codenames within parentheses indicate the CSD-names of the compounds

Compound	Target ΔH_f°	AM1*		PM6		PM5	
		ΔH_f°	Error	ΔH_f°	Error	ΔH_f°	Error
Fe	99.3	99.3	0.0	99.3	0.0	99.3	0.0
Fe ⁺	281.6	251.5	-30.1	262.3	-19.3	282.2	0.6
Fe ⁻	92.0	160.3	68.3	201.3	109.3	140.7	48.7
Fe ²⁺	654.6	697.5	42.9	637.4	-17.2	647.3	-7.3
Fe ³⁺	1361.0	1361.6	0.6	1358.0	-3.0	1349.7	-11.3
Fe ₂	180.0	164.3	-15.7	210.5	30.5	207.5	27.5
FeH	113.9	90.4	-23.5	145.9	32.0	123.7	9.8
FeH ⁻	91.1	100.4	9.3	193.0	101.9	102.2	11.1
FeCH ₃	71.0	51.7	-19.3	109.6	38.6	79.9	8.9
FeC ₅ H ₅	88.0	82.5	-5.5	110.7	22.7	142.6	54.6
Fe(C ₅ H ₅) ₂	57.9	67.2	9.3	49.3	-8.6	102.3	44.3
Fe(C ₅ H ₅) ₂ ⁺	210.2	250.4	40.2	252.0	41.8	294.3	84.1
FeO	60.0	32.8	-27.2	69.2	9.2	61.0	1.0
FeO ⁺	265.0	289.8	24.8	246.0	-19.0	259.3	-5.7
FeO ⁻	31.0	81.7	50.7	94.1	63.1	44.0	13.0
FeO ₂ ⁻	-30.0	-26.9	3.1	48.0	78.0	-18.8	11.2
FeOH	32.0	13.5	-18.5	28.5	-3.5	37.6	5.6
Fe(OH) ₂	-79.0	-83.3	-4.3	-56.5	22.5	-50.4	28.6
Fe(H ₂ O) ₄ (OH) ₂	-294.0	-327.1	-33.1	-334.5	-40.5	-294.5	-0.5
Fe(H ₂ O) ₅ (OH)	-162.2	-222.7	-60.5	-194.9	-32.7	-183.3	-21.1
Fe(H ₂ O) ₆ ²⁺	64.8	-4.3	-69.1	90.8	26.0	39.4	-25.4
Fe(H ₂ O) ₆ ³⁺	492.6	466.0	-26.7	455.0	-37.6	415.8	-76.8
Fe(CO) ₄ H ₂	-131.0	-129.8	1.2	-82.7	48.3	-130.9	0.1
Fe(CO)	63.9	26.1	-37.8	59.1	-4.8	60.4	-3.5
Fe(CO) ⁻	34.8	2.0	-32.8	94.1	59.3	42.4	7.6
Fe(CO) ₂	0.2	6.7	6.5	-3.4	-3.6	1.9	1.7
Fe(CO) ₂ ⁻	-27.3	-76.6	-49.3	15.9	43.2	-43.8	-16.5
Fe(CO) ₃	-55.8	-57.3	-1.5	-32.0	23.8	-41.0	14.8
Fe(CO) ₃ ⁻	-99.2	-124.1	-24.9	-50.2	49.0	-103.6	-4.4
Fe(CO) ₄	-105.1	-148.7	-43.6	-127.5	-22.4	-108.0	-2.9

Fe(CO) ₄ ⁻	-160.9	-205.1	-44.2	-139.2	21.7	-173.0	-12.1
Fe(CO) ₄ ²⁻	-115.1	-125.9	-10.8	-114.6	0.5	-127.3	-12.2
Fe(CO) ₅	-174.0	-181.1	-7.2	-188.4	-14.4	-138.1	35.8
FeC ₆ O ₁₂ ³⁻ (Fe(III)(Ox) ₃)	-506.7	-554.5	-47.8	-510.4	-3.7	-618.7	-112.0
FeC ₆ H ₃ O ₁₂ (H ₃ Fe(III)(Ox) ₃)	-516.8	-481.2	35.6	-501.4	15.4	-505.4	11.4
Fe(CO) ₄ C ₂ H ₄	-129.2	-142.1	-12.9	-133.4	-4.2	-128.5	0.7
FeC ₈ H ₂ O ₆ (FCPEN0)	-173.8	-211.7	-37.9	-170.4	3.4	-210.0	-36.2
FeC ₁₀ H ₁₄ O ₄ (bis(acetylacetonate)iron)	-198.0	-97.2	100.8	-178.2	19.8	-150.6	47.4
FeC ₁₅ H ₂₁ O ₆ ⁻ (Fe(II)(Acac)3 anion)	-340.2	-331.6	8.6	-317.9	22.3	-340.3	-0.1
Fe ₂ (CO) ₉	-319.2	-322.7	-3.5	-378.8	-59.6	-324.1	-4.9
Fe(NH ₃) ₆ ²⁺	266.4	258.1	-8.3	284.2	17.8	247.0	-19.4
Fe(CN) ₆ ³⁻	374.3	156.5	-217.8	337.0	-37.3	126.8	-247.5
Fe(CN) ₆ ⁴⁻	484.5	361.2	-123.3	494.5	10.0	318.1	-166.4
FeC ₆ N ₆ H ₂₄ ²⁺ (Fe(II)en) ₃	283.9	290.6	6.7	286.9	3.0	283.8	-0.1
FeC ₃₆ N ₆ H ₂₄ ²⁺ (Ferrous tris(ortho phenantholine))	451.2	535.5	84.3	428.1	-23.1	489.1	37.9
FeC ₈ NH ₅ O ₂ (ACODUR)	-12.1	-12.0	0.1	-9.2	2.9	-5.1	7.0
FeC ₉ NH ₈ O ₂ ⁺ (CPACFE)	113.2	147.1	33.9	96.1	-17.1	145.4	32.2
Fe(CO) ₂ (NO) ₂	-75.0	-72.8	2.2	-62.3	12.7	-2.9	72.1
FeC ₁₆ N ₅ H ₁₁ O ₄	-53.9	8.2	62.1	-54.8	-0.9	-31.2	22.7
FeC ₇ NH ₅ O ₅ (CNOFEA)	-89.2	-57.0	32.2	-94.6	-5.4	-65.6	23.6
FeC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (Iron(III)EDTA)	-405.6	-377.8	27.9	-404.1	1.5	-425.7	-20.1
FeN	174.3	139.8	-34.5	113.5	-60.8	159.5	-14.8
FeN ⁺	348.8	340.5	-8.3	294.5	-54.3	326.5	-22.3
FeN ⁻	123.7	147.8	24.1	162.4	38.7	135.0	11.3
FeF	11.4	44.6	33.2	22.1	10.7	9.9	-1.5
Fe(H ₂ O) ₅ F ⁺	-184.7	-252.6	-67.9	-214.0	-29.3	-212.6	-27.9
FeF ₂	-93.1	-97.1	-4.0	-80.7	12.4	-95.8	-2.7
FeF ₃	-196.2	-159.3	36.9	-162.1	34.1	-185.6	10.6
FeF ₆ ³⁻	-175.3	-175.3	0.0	-177.2	-1.9	-323.5	-148.2
FeAl	157.8	161.1	3.3	181.0	23.2	181.6	23.8
Fe(AlH ₂) ₂	216.8	214.8	-2.0	299.6	82.8	169.9	-46.9
FeSiH	133.7	121.6	-12.1	190.5	56.8	162.1	28.4
FeSiH ₂	125.6	122.9	-2.7	184.7	59.1	146.7	21.1
Fe(SiH ₃) ₂	126.7	133.4	6.7	234.7	108.0	136.0	9.3
FeP	156.9	157.0	0.1	29.8	-127.1	174.0	17.1
FePH	104.6	104.4	-0.2	56.8	-47.8	126.3	21.7
FeS	88.6	55.0	-33.6	137.0	48.4	101.1	12.5

FeC ₉ H ₁₀ SO ₂ (CEYTFE)	-59.3	-42.6	16.7	-39.5	19.8	-36.1	23.2
FeC ₁₂ H ₁₄ S ₄ O ₂ (CIBGAV10)	-73.1	-73.1	0.0	-88.8	-15.7	-64.0	9.1
FeCl	60.0	51.4	-8.6	52.5	-7.5	50.7	-9.3
FeCl ₂	-33.7	33.7	67.4	12.6	46.3	-24.0	9.7
FeCl ₃	-60.5	-5.7	54.8	-37.9	22.6	-59.8	0.7
FeCl ₄ ²⁻ (GOXLUA)	-105.2	-106.9	-1.7	-100.8	4.4	-187.2	-82.0
FeCl ₆ ³⁻	-56.5	-109.4	-52.9	-54.3	2.2	-177.8	-121.3
Fe(H ₂ O) ₅ Cl ⁺	-148.4	-208.0	-59.6	-146.3	2.1	-175.4	-27.0
FeC ₆ N ₂ H ₁₈ Cl ₃ (FINJIV)	-91.7	-74.4	17.3	-104.0	-12.3	-126.7	-35.0
FeTi	179.7	179.8	0.1	140.2	-39.5	153.7	-26.0
FeV	146.0	146.0	0.0	250.8	104.8	171.9	25.9
FeCr	215.2	215.2	0.0	216.4	1.2	-20.1	-235.3
FeCo	158.4	158.4	0.0	133.9	-24.5	-11.3	-169.7
FeNi	93.6	93.6	0.0	199.4	105.8	178.5	84.9
FeCu	140.7	140.7	0.0	180.0	39.3	83.8	-56.9
FeZn	119.7	129.6	9.9	113.1	-6.6	180.9	61.2
FeBr	44.7	51.9	7.2	83.7	39.0	58.6	13.9
FeBr ₂	-9.9	-16.5	-6.6	55.9	65.8	62.3	72.2
Fe ₂ Br ₄	-60.5	-60.5	0.0	-23.5	37.0	-35.2	25.3
FeC ₆ H ₅ BrO ₃ (ALCFEA)	-110.8	-122.7	-11.9	-119.6	-8.8	-132.4	-21.6
Fe(H ₂ O) ₅ Br ⁺	-137.9	-211.7	-73.8	-141.7	-3.8	-165.0	-27.1
Fe(H ₂ O) ₄ Br ₂	-248.8	-239.4	9.5	-293.3	-44.5	-281.1	-32.3
FeZr	206.1	180.3	-25.8	311.3	105.2	314.0	107.9
FeMo	251.3	251.2	-0.1	242.5	-8.8	317.4	66.1
Fe(CO) ₃ (C ₃ H ₅)I	-82.1	-86.7	-4.6	-81.0	1.1	-67.1	15.0
FeC ₆ H ₅ IO ₃ (ALCOFE10)	-97.1	-79.1	18.0	-100.2	-3.1	-108.1	-11.0
Fe(CO) ₄ I ₂	-151.9	-127.6	24.3	-140.5	11.4	-166.4	-14.5
Fe ₂ I ₄	2.0	74.1	72.1	16.5	14.5	18.8	16.8
FeI ₂	21.0	77.7	56.7	125.4	104.4	47.3	26.3
FeI	52.9	91.2	38.3	107.2	54.3	78.3	25.4
FeMn	126.6	126.7	0.1	161.6	35.0	-	-
		AM1*		PM6		PM5	
N=98							
Most positive error		100.8		109.3		107.9	
Most negative error		-217.8		-127.1		-247.5	
MSE		-3.1		13.7		-5.5	
MUE		26.5		31.6		34.5	
RMSD		41.4		43.6		57.4	

Results for the PM6 parameterization set ($N=55$)			
MSE	-6.3	1.6	-13.5
MUE	32.4	17.8	33.8
RMSD	48.9	23.5	57.6

Table 6 Calculated AM1*, PM6 and PM5 Koopmans' theorem ionization potentials and dipole moments for iron-containing compounds. The errors are color coded as follows: green up to 0.5 eV or 0.5 Debye; yellow between 0.5 and 1.0; pink larger than 1.0

Compound	Target	AM1*		PM6		PM5	
			Error		Error		Error
Koopmans' Theorem Ionization Potentials for Manganese Compounds (eV)							
Fe(CO) ₄ H ₂	9.65	9.36	-0.29	9.70	0.05	9.72	0.07
Fe(CO) ₂	6.68	8.01	1.33	8.07	1.39	8.51	1.83
Fe(CO) ₄	8.48	9.90	1.42	9.02	0.54	9.78	1.30
Fe(CO) ₅	8.60	9.76	1.16	9.23	0.63	9.21	0.61
FeBr ₂	9.70	10.05	0.35	9.23	-0.47	11.02	1.32
Fe ₂ Br ₄	12.60	11.04	-1.56	7.11	-5.49	10.53	-2.07
Fe(CO) ₄ H ₂	9.65	9.46	-0.19	9.70	0.05	9.72	0.07
FeCl ₂	10.10	11.21	1.11	8.38	-1.72	11.09	0.99
		AM1*		PM6		PM5	
N=8							
MSE		0.42		-0.63		0.52	
MUE		0.93		1.29		1.03	
Dipole Moments for Manganese Compounds (Debye)							
FeO	7.50	6.62	-0.88	4.44	-3.06	4.95	-2.55
FeCH ₃	0.90	0.67	-0.23	0.48	-0.42	1.58	0.68
Fe(CO) ₄ C ₂ H ₄	1.50	2.02	0.52	3.25	1.75	1.16	-0.34
FeO ₂ (3B1)	2.00	0.91	-1.09	3.33	1.33	0.74	-1.26
FeO ₂ (5B2)	3.40	5.06	1.66	2.10	-1.30	4.94	1.54
Fe(CO) ₃ C ₄ H ₆	2.10	0.36	-1.74	4.41	2.31	1.73	-0.37
FeC ₆ NH ₃ O ₄ (Fe(CO) ₄ acetonitrile)	5.00	7.67	2.67	5.13	0.13	6.61	1.61
FeF	4.19	3.82	-0.37	5.00	0.81	4.53	0.34
FeO ₂ F ₂	1.60	3.04	1.44	2.00	0.40	2.34	0.74
FeC ₁₀ PH ₁₅ O ₄ (Fe(CO) ₄ (PEt ₃))	5.20	5.93	0.73	1.56	-3.64	5.10	-0.10
FeCl	4.51	5.98	1.47	1.06	-3.45	4.59	0.08
FeCl ₂ O ₂	0.22	0.42	0.20	2.70	2.48	1.65	1.43
FeBr	4.18	4.41	0.23	1.12	-3.06	6.17	1.99
FeI	4.20	3.97	-0.23	5.05	0.85	5.24	1.04
		AM1*		PM6		PM5	
N=14							
MSE		0.31		-0.35		0.34	
MUE		0.96		1.79		1.01	

Table 7 Calculated AM1*, PM6 and PM5 bond lengths and angles for iron- containing compounds. The codenames within parentheses indicate the CSD-names of the compounds. The errors are color coded as follows: green up to 0.05 Å or 0.5°; yellow between 0.05-0.1 Å or 0.5-1.0°; pink larger than 0.1 Å or 1°

Compound	Variable	Target	AM1*		PM6		PM5	
				Error		Error		Error
Fe ₂	Fe-Fe	2.19	2.02	-0.17	2.35	0.17	1.83	-0.36
FeH	Fe-H	1.63	1.63	0.00	1.08	-0.55	1.70	0.07
FeH ⁻	Fe-H	1.69	1.69	0.00	1.98	0.29	1.70	0.01
FeH ⁺	Fe-H	1.60	1.53	-0.07	1.16	-0.44	1.58	-0.02
FeCH ₃	Fe-C	1.97	1.90	-0.07	2.12	0.15	1.94	-0.03
FeC ₅ H ₅	Fe-C	2.17	2.05	-0.13	2.06	-0.11	2.03	-0.14
Fe(C ₅ H ₅) ₂	Fe-C	2.06	2.12	0.06	2.06	0.00	2.22	0.16
Fe(III)Cp ₂ ⁺	Fe-C	2.06	1.99	-0.07	2.05	-0.01	2.08	0.02
FeO	Fe-O	1.62	1.76	0.14	1.66	0.04	1.63	0.00
FeO ⁻	Fe-O	1.69	1.74	0.05	1.79	0.10	1.73	0.04
FeO ⁺	Fe-O	1.56	1.75	0.19	1.65	0.09	1.65	0.09
FeO ₂ ⁻	Fe-O	1.71	1.77	0.06	1.71	0.00	1.72	0.01
FeOH	Fe-O	1.83	1.83	0.00	1.61	-0.22	1.77	-0.06
Fe(OH) ₂	Fe-O	1.78	1.85	0.08	1.81	0.03	1.79	0.02
Fe(H ₂ O) ₄ (OH) ₂	Fe-O	2.02	1.93	-0.09	1.66	-0.36	1.94	-0.08
	Fe-O	2.08	2.13	0.05	3.17	1.09	2.32	0.24
Fe(H ₂ O) ₅ (OH)	Fe-O	1.95	1.85	-0.10	1.83	-0.12	1.92	-0.03
	Fe-O	2.05	2.05	0.00	2.17	0.12	2.10	0.05
Fe(H ₂ O) ₆ ²⁺	Fe-O	2.07	1.99	-0.08	2.01	-0.06	2.15	0.08
	Fe-O	2.15	2.08	-0.07	2.53	0.38	2.22	0.07
Fe(H ₂ O) ₆ ³⁺	Fe-O	2.06	2.01	-0.05	2.05	-0.01	2.08	0.02
FeCH ₅ O ₂ (methyl iron(III)dihydroxide)	Fe-O	1.75	1.83	0.08	1.75	0.00	1.71	-0.04
	Fe-C	1.93	1.92	-0.01	1.64	-0.29	1.93	0.00
Fe(CO)	Fe-C	1.72	1.80	0.08	1.63	-0.09	1.76	0.04
Fe(CO) ⁻	Fe-C	1.79	1.85	0.06	1.80	0.01	1.19	-0.60
Fe(CO) ₂	Fe-C	1.84	1.88	0.04	1.76	-0.08	1.84	0.00
Fe(CO) ₂ ⁻	Fe-C	1.79	1.85	0.06	1.75	-0.04	1.82	0.03
Fe(CO) ₃	Fe-C	1.67	1.82	0.15	1.67	-0.01	1.82	0.14
Fe(CO) ₃ ⁻	Fe-C	1.82	1.88	0.06	1.40	-0.42	1.92	0.09
Fe(CO) ₄	Fe-C	1.78	1.89	0.11	1.71	-0.07	1.84	0.05
	Fe-C	1.82	1.89	0.08	1.80	-0.02	1.84	0.02

Fe(CO) ₄ ⁻	Fe-C	1.78	1.88	0.10	1.66	-0.13	1.84	0.06
Fe(CO) ₄ ²⁻	Fe-C	1.75	1.84	0.09	1.46	-0.29	1.78	0.03
Fe(CO) ₄ H ₂	Fe-C	1.81	1.89	0.09	1.81	0.00	1.84	0.03
	Fe-H	1.56	1.58	0.02	1.36	-0.20	1.62	0.06
	H-Fe-C	89.9	89.9	0.0	56.9	-33.0	90.0	0.1
Fe(CO) ₅	Fe-C(eq)	1.81	1.92	0.11	1.77	-0.04	1.82	0.01
	Fe-C(ax)	1.81	1.92	0.11	1.83	0.02	1.85	0.04
Fe(CO) ₄ C ₂ H ₄	Fe-C	1.81	1.89	0.08	1.75	-0.06	1.84	0.03
	Fe-C	2.13	1.98	-0.15	2.33	0.20	2.00	-0.13
FeC ₆ O ₁₂ ³⁻ (Fe(III)(Ox) ₃)	Fe-O	1.97	1.97	0.00	1.99	0.02	2.04	0.07
	O-Fe-O	83.2	80.8	-2.4	99.8	16.6	82.1	-1.1
FeC ₆ H ₃ O ₁₂ (H ₃ Fe(III)(Ox) ₃)	Fe-O	1.92	1.93	0.01	1.89	-0.03	2.02	0.10
	Fe-O	2.03	2.02	-0.01	1.98	-0.05	2.22	0.19
FeC ₇ H ₄ O ₃ (cyclobutadiene iron tricarbonyl)	Fe-C(C=O)	1.79	1.92	0.13	1.74	-0.05	1.78	-0.01
	Fe-C(C ₄ H ₄)	2.06	2.00	-0.06	2.08	0.02	2.34	0.28
	C-Fe-C	99.3	83.0	-16.3	88.5	-10.8	97.6	-1.7
FeC ₈ H ₂ O ₆ (FCPEN0)	Fe-C	1.81	1.90	0.09	1.82	0.01	1.86	0.05
	Fe-C	2.02	1.95	-0.07	1.83	-0.19	2.02	0.00
FeC ₉ H ₁₂ O (BUDFEC01)	Fe-C(O)	1.77	2.05	0.28	1.67	-0.10	1.82	0.05
	Fe-C	2.06	2.05	-0.01	2.08	0.02	2.13	0.07
	C-Fe-C	125.6	117.9	-7.7	109.5	-16.2	118.4	-7.2
FeC ₉ H ₁₂ O (FeCO(1,3-C ₄ H ₆) ₂)	Fe-C	2.11	2.00	-0.11	2.20	0.09	2.07	-0.04
	C-Fe-C	88.8	83.3	-5.5	79.2	-9.5	82.9	-5.9
Fe ₂ (CO) ₉	Fe-C	1.83	1.94	0.11	1.80	-0.03	1.85	0.02
	Fe-C	1.98	1.96	-0.02	2.02	0.04	2.03	0.05
	Fe-C#O	180.0	175.2	-4.8	177.0	-3.0	179.9	-0.1
	Fe-C#O	140.7	138.9	-1.8	141.8	1.1	139.1	-1.7
FeC ₁₀ H ₁₄ O ₄ (Fe(II)(Acac) ₂)	Fe-O	1.90	1.91	0.01	1.92	0.02	1.94	0.04
	O-Fe-O	93.9	90.9	-3.0	73.4	-20.6	92.4	-1.5
FeC ₁₅ H ₂₁ O ₆ ⁻ (Fe(II)(Acac) ₃)	Fe-O	1.97	1.98	0.01	2.03	0.06	2.02	0.05
	O-Fe-O	95.1	90.2	-4.9	91.3	-3.7	94.7	-0.4
FeN	Fe-N	1.56	1.72	0.16	1.33	-0.23	1.62	0.06
FeN ⁺	Fe-N	1.58	1.66	0.08	1.36	-0.22	1.54	-0.04
FeN ⁻	Fe-N	1.58	1.80	0.22	1.36	-0.22	1.71	0.13
Fe(NH ₃) ₆ ²⁺	Fe-N	2.10	2.07	-0.03	2.04	-0.06	2.19	0.09
Fe(CN) ₄ ⁴⁻	Fe-C	1.99	1.98	-0.01	1.98	-0.01	1.95	-0.04
	C-Fe-C	180.0	179.9	-0.1	179.9	-0.1	180.0	0.0

Fe(CO) ₂ (NO) ₂	Fe-C	1.82	1.97	0.15	1.78	-0.04	1.88	0.05
	Fe-N	1.68	1.79	0.11	1.66	-0.02	1.69	0.01
	C-Fe-C	98.7	83.8	-14.9	90.5	-8.2	86.8	-11.9
	N-Fe-N	109.0	146.7	37.8	92.5	-16.5	118.7	9.7
FeC ₆ N ₆ H ₂₄ ²⁺ (Fe(II)en) ₃	Fe-N	2.13	2.00	-0.14	2.13	0.00	2.19	0.06
FeC ₇ NH ₅ O ₅ (CNOFEA)	Fe-C	1.79	1.89	0.10	1.93	0.14	1.82	0.03
	Fe-C	2.09	2.06	-0.03	2.07	-0.02	2.27	0.18
	Fe-O	1.97	1.97	0.00	1.84	-0.13	2.06	0.09
FeC ₈ NH ₅ O ₂ (ACODUR)	Fe-C	1.91	1.90	-0.01	1.92	0.01	1.96	0.05
	Fe-C	1.78	1.90	0.12	1.76	-0.02	1.81	0.03
	Fe-C	2.19	2.11	-0.08	2.07	-0.12	2.23	0.04
	C-Fe-C	89.2	83.0	-6.2	74.2	-15.0	89.2	0.0
FeC ₉ NH ₈ O ₂ ⁺ (CPACFE)	Fe-N	1.91	1.95	0.04	1.83	-0.08	2.04	0.13
	Fe-C	1.77	1.91	0.14	1.78	0.01	1.84	0.07
FeC ₁₀ N ₄ H ₁₀ O ₆ (Diaqua-bis(pyrazinecarboxylato)-iron)	Fe-O	2.11	1.93	-0.18	1.99	-0.12	2.08	-0.03
	Fe-N	2.12	1.95	-0.17	1.92	-0.20	2.08	-0.04
	Fe-OH ₂	2.14	2.14	0.00	2.07	-0.07	2.09	-0.05
	O-Fe-N	78.1	83.4	5.3	76.5	-1.6	79.7	1.6
FeC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (Iron(III)EDTA)	Fe-N	2.03	2.21	0.18	2.19	0.16	2.32	0.29
FeC ₁₆ N ₅ H ₁₁ O ₄	Fe-O	2.12	2.60	0.48	2.08	-0.04	2.26	0.14
	Fe-O	1.92	1.87	-0.05	1.95	0.03	2.01	0.09
	Fe-N	1.89	1.83	-0.06	1.88	-0.01	2.02	0.13
	O-Fe-O	81.4	78.8	-2.6	73.4	-8.0	77.9	-3.5
FeC ₁₇ N ₂ H ₁₆ (Toluene-(2,2'-bipyridine)-iron)	Fe-N	1.90	1.84	-0.06	1.86	-0.04	1.90	0.00
	Fe-C	2.11	2.07	-0.04	2.07	-0.04	2.28	0.17
	N-Fe-N	81.9	86.3	4.4	76.5	-5.4	82.7	0.8
	N-Fe-C	122.8	114.7	-8.1	107.3	-15.5	128.2	5.4
FeC ₂₀ H ₁₂ N ₄ (iron porphyrin)	Fe-N	1.97	1.93	-0.04	1.92	-0.05	2.02	0.05
FeC ₃₆ N ₆ H ₂₄ ⁺ (ferrous tris(ortophenantholine))	Fe-N	2.05	1.97	-0.08	1.95	-0.10	2.14	0.09
FeF	Fe-F	1.76	1.80	0.04	1.75	-0.01	1.77	0.01
Fe(II)(H ₂ O) ₅ F ⁺	Fe-F	1.89	1.70	-0.19	1.82	-0.07	1.90	0.01
	Fe-O	2.05	2.04	-0.01	2.15	0.10	2.10	0.05
FeF ₂	Fe-F	1.77	1.71	-0.06	1.76	-0.01	1.79	0.02
FeO ₂ F ₂ (iron(VI)difluoride dioxide)	Fe-F	1.74	1.71	-0.03	1.74	0.00	1.81	0.07
	Fe=O	1.58	1.85	0.27	1.55	-0.03	1.66	0.08
FeF ₃	Fe-F	1.78	1.75	-0.03	1.73	-0.05	1.77	-0.01
FeF ₆ ³⁻	Fe-F	2.06	1.85	-0.21	1.94	-0.12	2.02	-0.04

FeAl	Fe-Al	2.30	2.27	-0.03	2.65	0.35	2.28	-0.03
Fe(AIH ₂) ₂	Fe-Al	2.47	2.39	-0.08	3.06	0.59	2.69	0.22
FeSiH	Fe-Si	2.30	2.30	0.00	3.96	1.66	2.48	0.18
FeSiH ₂	Fe-Si	2.26	2.31	0.05	4.24	1.98	2.48	0.22
Fe(SiH ₃) ₂	Fe-Si	2.52	2.52	0.00	4.47	1.95	2.52	0.00
FeP	Fe-P	2.23	2.20	-0.03	1.85	-0.38	2.30	0.06
FePH	Fe-P	2.20	2.30	0.10	2.05	-0.15	2.35	0.15
FeC ₁₂ P ₄ H ₃₄ S ₂ (JIYFAY)	Fe-S	2.35	2.33	-0.02	2.10	-0.25	2.33	-0.02
	Fe-P	2.23	2.64	0.41	2.26	0.03	2.49	0.26
	P-Fe-S	89.4	90.8	1.4	83.1	-6.4	70.5	-18.9
FeS	Fe-S	2.04	2.10	0.06	1.99	-0.05	2.06	0.02
Fe(SH) ₂	Fe-S	2.25	2.26	0.02	1.82	-0.43	2.26	0.01
FeC ₉ H ₁₀ SO ₂ (CEYTFE)	Fe-S	2.30	2.32	0.02	2.00	-0.30	2.34	0.04
	Fe-C(O)	1.75	1.89	0.14	1.74	-0.01	1.80	0.05
	Fe-C(C ₄)	2.12	2.12	0.00	2.04	-0.08	2.25	0.13
	C-Fe-S	90.7	85.0	-5.7	102.3	11.6	91.9	1.2
FeC ₁₆ N ₂ H ₁₈ S ₂ O ₂ (dicarbonyl ethylenediamine bis(phenylthiolato)iron)	Fe-S	2.33	2.37	0.04	2.26	-0.07	2.38	0.05
	Fe-S	2.34	2.37	0.03	2.13	-0.21	2.39	0.04
	Fe-N	2.04	1.96	-0.08	2.15	0.11	2.15	0.11
	Fe-N	2.03	1.97	-0.06	1.95	-0.08	2.15	0.12
	Fe-C	1.76	1.88	0.12	1.73	-0.03	1.79	0.03
	N-Fe-S	86.8	88.2	1.4	80.1	-6.7	82.0	-4.8
	C-Fe-S	93.2	84.9	-8.3	96.1	2.9	98.0	4.8
FeC ₉ N ₃ H ₁₈ S ₃ O ₃ (tris(N,N-dimethylthiocarbamato)iron(III))	Fe-S	2.43	2.44	0.01	2.49	0.06	2.52	0.09
	Fe-O	2.09	1.99	-0.10	1.88	-0.21	2.07	-0.02
	S-Fe-O	68.9	70.5	1.6	72.1	3.2	66.9	-2.0
	S-Fe-S	103.1	96.0	-7.1	98.0	-5.1	101.5	-1.6
Fe ₄ S ₄	Fe-S	2.27	2.27	0.00	2.28	0.01	2.37	0.10
FeC ₁₂ H ₁₄ S ₄ O ₂ (CIBGAV10)	Fe-S	2.31	2.35	0.04	2.23	-0.08	2.36	0.05
	Fe-C	1.80	1.89	0.09	1.73	-0.07	1.81	0.01
	S-Fe-S	85.2	85.7	0.4	108.8	23.6	87.8	2.6
	C-Fe-S	90.1	89.4	-0.7	83.1	-7.0	90.7	0.6
FeC ₁₄ N ₆ H ₁₆ S ₆ (diisothiocyanato-bisthiazoline iron(II))	Fe-N	2.19	2.09	-0.10	1.87	-0.32	2.06	-0.13
	Fe-N(CS)	2.08	1.87	-0.21	1.84	-0.24	1.98	-0.10
	N-Fe-N	73.9	81.3	7.4	101.9	28.0	78.1	4.2
	SCN-Fe-	97.4	94.3	-3.1	99.8	2.4	89.0	-8.4

	NCS							
FeCl	Fe-Cl	2.13	2.23	0.10	1.55	-0.58	2.16	0.02
Fe(II)(H ₂ O) ₅ Cl ⁺	Fe-Cl	2.29	2.17	-0.12	2.45	0.16	2.29	0.00
	Fe-O	2.06	2.06	0.00	2.07	0.01	2.10	0.04
FeCl ₂	Fe-Cl	2.16	2.17	0.01	1.94	-0.22	2.15	-0.01
FeCl ₃	Fe-Cl	2.16	2.17	0.02	2.18	0.03	2.17	0.02
FeC ₆ N ₂ H ₁₈ Cl ₃ (FINJIV)	Fe-Cl	2.23	2.35	0.12	2.34	0.11	2.25	0.02
	Fe-N	2.27	1.93	-0.34	1.96	-0.31	2.28	0.01
	Cl-Fe-Cl	121.0	120.1	-0.9	119.4	-1.6	119.8	-1.3
FeCl ₄ ⁻	Fe-Cl	2.19	2.22	0.03	2.11	-0.08	2.18	-0.01
FeCl ₄ ²⁻ (GOXLUA)	Fe-Cl	2.34	2.33	-0.01	2.35	0.01	2.36	0.02
FeCl ₆ ³⁻	Fe-Cl	2.53	2.54	0.01	2.51	-0.02	2.41	-0.12
FeTi	Fe-Ti	2.54	2.54	0.00	3.67	1.13	-	-
FeV	Fe-V	2.31	2.31	0.00	3.50	1.19	2.04	-0.27
FeCr	Fe-Cr	2.27	2.06	-0.21	3.65	1.38	1.60	-0.67
FeMn	Fe-Mn	2.40	2.40	0.00	5.21	2.81	-	-
FeCo	Fe-Co	2.31	2.30	-0.01	4.68	2.38	1.70	-0.61
FeNi	Fe-Ni	2.33	2.21	-0.12	4.69	2.36	2.05	-0.29
FeCu	Fe-Cu	2.31	2.31	0.00	7.36	5.05	2.05	-0.26
FeZn	Fe-Zn	2.53	2.37	-0.16	2.98	0.45	2.63	0.10
FeBr	Fe-Br	2.23	2.16	-0.07	2.17	-0.06	2.31	0.08
FeBr ₂	Fe-Br	2.31	2.13	-0.18	2.13	-0.18	2.31	0.00
FeC ₆ H ₅ BrO ₃ (ALCFEA)	Fe-Br	2.50	2.20	-0.30	2.46	-0.04	2.43	-0.07
	Fe-C(C ₂)	2.13	2.03	-0.10	2.14	0.01	2.13	0.00
	Fe-C(O)	1.79	1.91	0.12	1.83	0.04	1.85	0.06
	C-Fe-Br	88.6	87.6	-1.0	108.9	20.3	84.9	-3.7
Fe(H ₂ O) ₅ Br ⁺	Fe-Br	2.42	2.14	-0.28	2.47	0.05	2.38	-0.04
	Fe-O	2.06	2.10	0.04	2.07	0.01	2.24	0.18
Fe(H ₂ O) ₄ Br ₂	Fe-O	2.07	1.99	-0.08	2.15	0.08	2.09	0.02
	Fe-Br	2.50	2.20	-0.30	2.71	0.21	2.48	-0.02
FeC ₄ H ₈ BrO ₇ (Diaquabromo(oxydiacetato-O,O',O'')-iron(III))	Fe-Br	2.37	2.17	-0.20	2.38	0.01	2.36	-0.02
	Fe-O	2.00	2.07	0.07	1.85	-0.15	1.98	-0.02
	Br-Fe-O	105.6	94.5	-11.1	98.6	-7.0	111.5	5.9
FeBr ₂ O ₂	Fe-Br	2.27	2.15	-0.12	2.56	0.29	2.38	0.11
	Fe=O	1.58	1.74	0.16	1.55	-0.03	1.52	-0.06
	Br-Br-Fe	110.6	110.0	-0.6	129.6	19.0	98.1	-12.5
FeZr	Fe-Zr	2.59	2.59	0.00	2.84	0.25	3.04	0.46

FeMo	Fe-Mo	2.21	2.14	-0.07	3.29	1.08	11.58	9.37
FeI	Fe-I	2.44	2.46	0.02	2.70	0.26	2.38	-0.06
Fe(CO) ₃ (C ₃ H ₅)I	Fe-I	2.75	2.61	-0.14	2.80	0.05	2.48	-0.27
	Fe-C(O)	1.80	1.96	0.16	1.73	-0.07	1.84	0.04
	C-Fe-I	81.6	81.7	0.1	75.3	-6.3	76.2	-5.4
Fe(CO) ₃ (C ₃ H ₅)I (ALCOFE10)	Fe-I	2.75	2.59	-0.16	2.80	0.05	2.59	-0.16
	Fe-C	1.80	1.95	0.15	1.73	-0.07	1.83	0.03
	C-Fe-I	81.6	96.5	14.9	75.3	-6.3	81.8	0.2
FeI ₂	Fe-I	2.50	2.49	-0.01	2.55	0.05	2.43	-0.07
				AM1*	PM6		PM5	
N=153								
MSE bond length				0.00	0.12		0.08	
MUE bond length				0.09	0.27		0.15	
N=33								
MSE bond angle				-1.3	-2.6		-1.7	
MUE bond angle				5.8	10.4		4.0	

AM1* Parameters for Manganese and Iron

Hakan Kayı and Timothy Clark

Computer-Chemie-Centrum and Interdisciplinary Center for Molecular Materials,
Friedrich-Alexander-Universität Erlangen-Nürnberg, Nögelsbachstraße 25, 91052
Erlangen, Germany

E-mail: clark@chemie.uni-erlangen.de

Supplementary Material

		Page
Table S1	Sources and calculation details for the heats of formation used to parameterize.	2
Table S2	Sources and calculation details for the ionization potentials and dipole moments used to parameterize.	5
Table S3	Sources and calculation details for the geometrical variables used to parameterize.	6
References		14

Table S1: Sources and calculation details for the heats of formation used to parameterize.

Heat of Formation for Manganese Containing Compounds			
Compound	Target(kcal/mol)	Source	Total Energy (a.u.)
Mn	67.7	S1	
Mn ⁻	90.2	S2	
Mn ⁺	239.1	S2	
Mn ²⁺	599.8	S2	
Mn ³⁺	1376.0	S2	
Mn ₂	184.3	S3	-207.6091449
MnH	57.3	S3	-104.4496606
MnH ⁻	41.2	S1	
MnH ₂	114.1	S3	-104.9516412
Mn(C ₅ H ₅) ₂	66.2	S1	
MnN	123.1	S3	-158.5358747
Mn(NH ₃) ₆ ²⁺	269.5	S4	
MnO ₂ ⁻	-62.1	S1	
MnO ₃ ⁻	-127.1	S1	
MnO ₄ ⁻	-158.4	S1	
Mn(H ₂ O) ₆ ²⁺	65.7	S4	
Mn(CO) ₃ ⁻	-54.8	S1	
Mn(CO) ₄ ⁻	-122.3	S1	
Mn(CO) ₅	-178.0	S1	
Mn(CO) ₅ H	-176.8	S1	
Mn(CO) ₆ ⁺	-49.7	S4	
Mn(CO) ₅ CH ₃	-179.5	S1	
Mn(CO) ₅ C ₆ H ₅	-140.8	S1	
Mn(CO) ₅ (C ₆ H ₅ CH ₂)	-151.9	S1	
Mn(CO) ₅ COCH ₃	-215.3	S1	
Mn(CO) ₅ (C ₆ H ₅ CO)	-173.2	S1	
MnC ₁₅ H ₂₁ O ₆ (Mn(acac) ₃)	-285.2	S4	
MnC ₆ H ₈ O ₁₀ ⁻ (KAMMND)	-543.1	S4	
MnC ₉ H ₁₂ O (BUTMNC)	11.9	S4	
MnC ₈ H ₅ O ₃ (cyclopentadienyl manganese tricarbonyl)	-115.2	S1	
MnC ₉ H ₇ O ₃ (HEXMNC)	-95.5	S4	
Mn ₂ (CO) ₉	-314.5	S1	
Mn ₂ (CO) ₁₀	-385.9	S5	
Mn(NO)	8.5	S3	-233.8033629
Mn(CO)(NO) ₃	-60.6	S4	
MnC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (EDTMNK01)	-388.3	S4	
MnF ₂	-126.2	S6	
MnF ₂ ⁻	-226.0	S1	
MnF ₃	-188.0	S6	
MnF ₃ ⁻	-272.2	S1	
MnF ₄	-231.0	S6	
MnF ₄ ⁻	-343.6	S1	
MnF ₅ ⁻	-373.0	S1	
Mn ₂ F ₅ ⁻	-448.3	S1	
Mn ₂ F ₇ ⁻	-600.0	S1	
MnOF ₂	-125.8	S4	
Mn(H ₂ O) ₄ F ₂ trans	-333.2	S4	
MnC ₇ HO ₅ F ₂ (CDFVMN)	-252.2	S4	
Mn(CO) ₅ CF ₃	-330.5	S1	
Mn(CO) ₅ COCF ₃	-359.7	S1	
MnC ₇ O ₇ F ₃ (FACMNA)	-437.0	S4	
MnAl	116.5	S3	-105.8283338

MnF ₃ AlF ₄ ⁻	-704.0	S1	
MnSi	153.2	S3	-107.645974
MnSiH	93.9	S3	-108.3333477
MnSiH ₂	85.2	S3	-108.9445051
MnP	81.9	S3	-110.3663624
MnPH	67.6	S3	-110.9840119
MnC ₉ H ₁₁ SO ₂ (VALXAI)	-76.7	S4	
MnC ₇ H ₅ SO ₄ (CPCSMN)	-125.0	S4	
MnC ₇ N ₂ H ₁₂ S ₄ ⁻ (COWHOL)	-19.9	S4	
MnC ₇ H ₁₀ S ₂ ClO ₃ (GEQLUJ)	-179.2	S4	
MnCl	11.3	S6	
Mn(CO) ₅ Cl (ZOSWEJ)	-219.5	S1	
MnCl ₂	-63.0	S6	
Mn(H ₂ O) ₄ Cl ₂ trans	-263.1	S4	
Mn ₂ Cl ₄	-168.2	S6	
MnCl ₂ O	-136.0	S3	-209.1453756
MnTi	148.8	S3	-161.8591387
MnV ₂	228.3	S3	-246.2697983
MnCr	184.4	S3	-190.0795394
MnCo	158.9	S3	-248.8832029
MnNi	80.9	S3	-273.1382819
MnCu	158.2	S3	-299.9495504
MnZn	89.3	S3	-169.4568054
MnOBr ₂	-37.0	S4	
Mn(H ₂ O) ₄ Br ₂ trans	-244.3	S4	
Mn(CO) ₅ Br	-210.9	S1	
MnBr ₄ ²⁻ (PYDMNB)	-174.7	S3	-156.9399370
MnZr	173.7	S3	-150.3305742
MnMo	191.5	S3	-171.3221772
MnI	65.8	S3	-115.2527121
MnOl ₂	-7.0	S4	
Mn(H ₂ O) ₄ I ₂ trans	-213.7	S4	
Mn(CO) ₅ I (VUCFAA)	-199.4	S1	
MnC ₁₃ N ₂ H ₈ O ₃ I (YOMHEN)	-83.2	S4	

Heat of Formation for Iron Containing Compounds

Compound	Target(kcal/mol)	Source	Total Energy (a.u.)
Fe	99.3	S1	
Fe ⁺	281.6	S2	
Fe ⁻	92.0	S1	
Fe ²⁺	654.6	S2	
Fe ³⁺	1361.0	S2	
Fe ₂	180.0	S7	
FeH	113.9	S6	
FeH ⁻	91.1	S1	
FeCH ₃	71.0	S6	
FeC ₅ H ₅	88.0	S1	
Fe(C ₅ H ₅) ₂	57.9	S1	
Fe(C ₅ H ₅) ₂ ⁺	210.2	S4	
FeO	60.0	S1	
FeO ⁺	265.0	S7	
FeO ⁻	31.0	S1	
FeO ₂ ⁻	-30.0	S1	
FeOH	32.0	S7	
Fe(OH) ₂	-79.0	S1	
Fe(H ₂ O) ₄ (OH) ₂	-294.0	S4	
Fe(H ₂ O) ₅ (OH)	-162.2	S4	
Fe(H ₂ O) ₆ ²⁺	64.8	S4	
Fe(H ₂ O) ₆ ³⁺	492.6	S4	

Fe(CO) ₄ H ₂	-131.0	S1	
Fe(CO)	63.9	S6	
Fe(CO) ⁻	34.8	S1	
Fe(CO) ₂	0.2	S6	
Fe(CO) ₂ ⁻	-27.3	S1	
Fe(CO) ₃	-55.8	S6	
Fe(CO) ₃ ⁻	-99.2	S1	
Fe(CO) ₄	-105.1	S1	
Fe(CO) ₄ ⁻	-160.9	S1	
Fe(CO) ₄ ²⁻	-115.1	S4	
Fe(CO) ₅	-174.0	S1	
FeC ₆ O ₁₂ ³⁻ (Fe(III)(Ox) ₃)	-506.7	S4	
FeC ₆ H ₃ O ₁₂ (H ₃ Fe(III)(Ox) ₃)	-516.8	S4	
Fe(CO) ₄ C ₂ H ₄	-129.2	S8	
FeC ₈ H ₂ O ₆ (FCPEN0)	-173.8	S4	
FeC ₁₀ H ₁₄ O ₄ (bis(acetylacetonate)iron)	-198.0	S6	
FeC ₁₅ H ₂₁ O ₆ ⁻ (Fe(II)(Acac)3 anion)	-340.2	S1	
Fe ₂ (CO) ₉	-319.2	S1	
Fe(NH ₃) ₆ ²⁺	266.4	S4	
Fe(CN) ₆ ³⁻	374.3	S4	
Fe(CN) ₆ ⁴⁻	484.5	S4	
FeC ₆ N ₆ H ₂₄ ²⁺ (Fe(II)en) ₃	283.9	S4	
FeC ₃₆ N ₆ H ₂₄ ²⁺ (Ferrous tris(ortho phenantholine))	451.2	S4	
FeC ₈ NH ₅ O ₂ (ACODUR)	-12.1	S4	
FeC ₉ NH ₈ O ₂ ⁺ (CPACFE)	113.2	S4	
Fe(CO) ₂ (NO) ₂	-75.0	S4	
FeC ₁₆ N ₅ H ₁₁ O ₄ (FeN3O2)	-53.9	S4	
FeC ₇ NH ₅ O ₅ (CNOFEA)	-89.2	S4	
FeC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (Iron(III)EDTA)	-405.6	S4	
FeN	174.3	S3	-178.0254633
FeN ⁺	348.8	S3	-177.7462654
FeN ⁻	123.7	S3	-178.1057778
FeF	11.4	S1	
Fe(H ₂ O) ₅ F ⁺	-184.7	S4	
FeF ₂	-93.1	S1	
FeF ₃	-196.2	S1	
FeF ₆ ³⁻	-175.3	S4	
FeAl	157.8	S3	-125.3330308
Fe(AlH ₂) ₂	216.8	S3	-129.6762622
FeSiH	133.7	S3	-127.8403453
FeSiH ₂	125.6	S3	-128.4527619
Fe(SiH ₃) ₂	126.7	S3	-134.7832601
FeP	156.9	S3	-129.8171249
FePH	104.6	S3	-130.4956816
FeS	88.6	S1	
FeC ₉ H ₁₀ SO ₂ (CEYTFE)	-59.3	S4	
FeC ₁₂ H ₁₄ S ₄ O ₂ (CIBGAV10)	-73.1	S4	
FeCl	60.0	S1	
FeCl ₂	-33.7	S1	
FeCl ₃	-60.5	S1	
FeCl ₄ ²⁻ (GOXLUA)	-105.2	S4	
FeCl ₆ ³⁻	-56.5	S4	
Fe(H ₂ O) ₅ Cl ⁺	-148.4	S4	
FeC ₆ N ₂ H ₁₈ Cl ₃ (FINJIV)	-91.7	S4	
FeTi	179.7	S3	-181.3805987
FeV	146.0	S3	-194.6316283
FeCr	215.2	S3	-209.601116

FeCo	158.4	S3	-268.4543326
FeNi	93.6	S3	-292.6888573
FeCu	140.7	S3	-319.5481875
FeZn	119.7	S3	-188.9789412
FeBr	44.7	S3	-136.6278006
FeBr ₂	-9.9	S1	
Fe ₂ Br ₄	-60.5	S1	
FeC ₆ H ₅ BrO ₃ (ALCFEA)	-110.8	S4	
Fe(H ₂ O) ₅ Br ⁺	-137.9	S4	
Fe(H ₂ O) ₄ Br ₂	-248.8	S4	
FeZr	206.1	S3	-169.8300106
FeMo	251.3	S3	-190.7974445
Fe(CO) ₃ (C ₃ H ₅)I	-82.1	S1	
FeC ₆ H ₅ IO ₃ (ALCOFE10)	-97.1	S4	
Fe(CO) ₄ I ₂	-151.9	S1	
Fe ₂ I ₄	2.0	S1	
FeI ₂	21.0	S1	
FeI	52.9	S3	-134.8446233
FeMn	126.6	S3	-258.893132

Table S2: Sources and calculation details for the ionization potentials and dipole moments used to parameterize.

Koopmans' Theorem Ionization Potentials for Manganese Containing Compounds		
Compound	Target(eV)	Source
MnH	7.80	S1
MnO	8.70	S1
Mn(CO) ₅	8.10	S1
Mn(CO) ₅ H	8.85	S1
MnF ₂	11.40	S1
MnF ₃	12.60	S1
MnF ₄	13.50	S1
MnCl ₂	11.03	S1
Mn(CO) ₅ Br	8.83	S1
MnBr ₂	10.30	S1
Mn(CO) ₅ I	8.40	S1
Koopmans' Theorem Ionization Potentials for Iron Containing Compounds		
Fe(CO) ₄ H ₂	9.65	S6
Fe(CO) ₂	6.68	S6
Fe(CO) ₄	8.48	S1
Fe(CO) ₅	8.60	S1
FeBr ₂	9.70	S1
Fe ₂ Br ₄	12.60	S1
Fe(CO) ₄ H ₂	9.65	S1
FeCl ₂	10.10	S1

Dipole Moments for Manganese Containing Compounds		
Compound	Target(Debye)	Source
MnH	0.50	S3
MnN	3.07	S3

MnNO	5.89	S3
MnO	4.72	S3
MnOH	0.40	S3
MnF	2.27	S3
MnOF ₂	0.53	S4
MnAl	1.67	S3
MnSi	3.27	S3
MnSiH	3.15	S3
MnSiH ₂	3.22	S3
MnP	2.54	S4
MnPH	4.34	S3
MnS	4.22	S4
MnCl	2.36	S3
MnBr	4.29	S4
MnOBr ₂	0.51	S4
MnI	4.25	S4
MnOI ₂	0.94	S4
Dipole Moments for Iron Containing Compounds		
FeO	7.50	S6
FeCH ₃	0.90	S9
Fe(CO) ₄ C ₂ H ₄	1.50	S10
FeO ₂ (3B1)	2.00	S11
FeO ₂ (5B2)	3.40	S11
Fe(CO) ₃ C ₄ H ₆	2.10	S10
FeC ₆ NH ₃ O ₄ (Fe(CO) ₄ acetonitrile)	5.00	S10
FeF	4.19	S4
FeO ₂ F ₂	1.60	S4
FeC ₁₀ PH ₁₅ O ₄ (Fe(CO) ₄ (PET ₃))	5.20	S10
FeCl	4.51	S4
FeCl ₂ O ₂	0.22	S4
FeBr	4.18	S4
FeI	4.20	S4

Table S3: Sources and calculation details for the geometrical variables used to parameterize.

Geometrical Variables for Manganese Containing Compounds			
Compound	Variable	Target	Source
Mn ₂	Mn-Mn	2.29	S3
MnH	Mn-H	1.74	S3
MnH ⁻	Mn-H	1.77	S3
Mn(C ₅ H ₅) ₂	Mn-C	2.42	S3
MnO ₂ ⁻	Mn-O	1.62	S3
MnO ₃ ⁻	Mn-O	1.59	S3
MnO ₄ ⁻	Mn-O	1.64	S4
Mn(H ₂ O) ₆ ²⁺	Mn-O	2.14	S12
	Mn-O	2.19	S12
	Mn-O	2.20	S12
Mn(CH ₃)O ₃	Mn=O	1.59	S4

	Mn-C	1.99	S4
$\text{Mn}(\text{CO})_3^-$	Mn-C	1.82	S3
$\text{Mn}(\text{CO})_5$	Mn-C	1.82	S3
	Mn-C	1.86	S3
$\text{Mn}(\text{CO})_5\text{H}$	Mn-C	1.86	S13
	Mn-C	1.85	S13
	H-Mn-C	85.5	S13
$\text{Mn}(\text{CO})_5\text{CH}_3$	Mn-C	1.83	S3
	Mn-C	1.85	S3
	Mn-C	2.19	S3
$\text{Mn}(\text{CO})_6^+$	Mn-C	1.91	S4
$\text{Mn}(\text{CO})_5\text{C}_6\text{H}_5$	Mn-C	1.83	S3
	Mn-C	1.85	S3
	Mn-C	2.15	S3
$\text{Mn}(\text{CO})_5(\text{C}_6\text{H}_5\text{CH}_2)$	Mn-C	1.82	S3
	Mn-C	1.85	S3
	Mn-C	2.26	S3
$\text{Mn}(\text{CO})_5\text{COCH}_3$	Mn-C	1.84	S3
	Mn-C	1.85	S3
	Mn-C	2.17	S3
$\text{Mn}(\text{CO})_5(\text{C}_6\text{H}_5\text{CO})$	Mn-C	1.84	S3
	Mn-C	1.85	S3
	Mn-C	2.18	S3
$\text{MnC}_6\text{H}_8\text{O}_{10}^-$ (KAMMND)	Mn-O	1.90	S14
	Mn-O	2.30	S14
	O-Mn-O	91.7	S14
$\text{MnC}_9\text{H}_{12}\text{O}$ (BUTMNC)	Mn-C	1.81	S14
	Mn-C	2.15	S14
$\text{Mn}(\text{CO})_3\text{Cp}$ (CPMNCO)	Mn-C(ring)	2.14	S14
	Mn-C(O)	1.79	S14
	C-Mn-C	91.4	S14
$\text{MnC}_9\text{H}_7\text{O}_3$ (HEXMNC)	Mn-C	2.15	S14
	Mn-C	1.78	S14
$\text{Mn}_2\text{C}_8\text{H}_{12}\text{O}_8$ ($\text{Mn}_2(\text{Ac})_4$)	Mn-Mn	3.31	S14
$\text{Mn}_2(\text{CO})_{10}$	Mn-Mn	2.89	S14
$\text{Mn}_2(\text{CO})_{10}\text{C}_2$ (JIPVOT)	Mn-C	1.86	S14
	Mn-C	1.83	S14
	C-Mn-C	85.9	S14
MnN	Mn-N	1.59	S3
MnNO	Mn-N	1.73	S3
$\text{Mn}(\text{NH}_3)_6^{2+}$	Mn-N	2.16	S4
$\text{Mn}(\text{CO})(\text{NO})_3$	Mn-N	1.70	S4
	Mn-C	1.87	S4
	C-Mn-N	106.0	S4
$\text{MnC}_{10}\text{N}_2\text{H}_{12}\text{O}_8^-$ (EDTMNK01)	Mn-O	2.02	S14
	Mn-O	1.89	S14
	Mn-N	2.24	S14
	O-Mn-O	88.3	S14
	N-Mn-O	77.5	S14
$\text{MnC}_{12}\text{N}_2\text{H}_{12}\text{O}_6$ (XOYHAU)	Mn-O	2.15	S14
	Mn-N	2.26	S14
	O-Mn-O	85.4	S14
$\text{MnC}_{12}\text{N}_2\text{H}_{16}\text{O}_4$ (Mn	Mn-O	2.16	

phenanthroline tetrahydrate)			S12
MnF	Mn-F	1.82	S3
MnF ₂	Mn-F	1.79	S3
MnF ₂ ⁻	Mn-F	1.86	S3
MnOF ₂	Mn-O	1.55	S4
	Mn-F	1.72	S4
	O-Mn-F	121.3	S4
Mn(H ₂ O) ₄ F ₂ trans	Mn-F	1.98	S4
	Mn-O	2.13	S4
MnC ₇ HO ₅ F ₂ (CDFVMN)	Mn-C	1.72	S14
	Mn-C	1.77	S14
	Mn-C	1.94	S14
	C-Mn-C	93.8	S14
MnF ₃	Mn-F	1.79	S3
	F-Mn-F	120.0	S3
MnC ₇ O ₇ F ₃ (FACMNA)	Mn-C	1.81	S14
	Mn-O	2.03	S14
Mn ₂ (CO) ₈ (CF ₂) ₂ (DOFPET)	Mn-Mn	2.66	S14
	Mn-C(F ₂)	2.02	S14
	Mn-C(O)	1.88	S14
	C-Mn-Mn	49.4	S14
MnAl	Mn-Al	2.34	S3
Mn(AlH ₂) ₂	Mn-Al	2.41	S3
MnSi	Mn-Si	2.34	S3
MnSiH	Mn-Si	2.34	S3
MnSiH ₂	Mn-Si	2.38	S3
MnSi ₂ C ₉ H ₁₅ O ₅ (KIRYUF)	Mn-Si	2.34	S14
	Mn-O	2.95	S14
	Si-Mn-Si	71.2	S14
MnP	Mn-P	1.97	S4
MnPH	Mn-P	2.22	S3
MnS	Mn-S	1.99	S4
MnS ₂	Mn-S	2.20	S3
MnC ₉ H ₁₁ SO ₂ (VALXAI)	Mn-S	2.27	S14
	Mn-C(O)	1.77	S14
	Mn-C(C ₄)	2.12	S14
MnC ₇ H ₅ SO ₄ (CPCSMN)	Mn-S	2.04	S14
	Mn-C	1.79	S14
	Mn-C'	2.10	S14
	S-Mn-C	91.0	S14
MnC ₇ N ₂ H ₁₂ S ₄ ⁻ (COWHOL)	Mn-S	2.32	S14
	S-Mn-S	157.5	S14
MnC ₇ H ₁₀ S ₂ ClO ₃ (GEQLUJ)	Mn-S	2.39	S14
	Mn-C	1.80	S14
	Mn-Cl	2.38	S14
	S-Mn-Cl	92.2	S14
MnC ₄ N ₂ H ₈ S ₂ O ₆ (WIFSEJ)	Mn-S	2.61	S14
	Mn-O	2.20	S14
	O-Mn-S	87.3	S14
MnCl	Mn-Cl	2.12	S4
Mn(CO) ₅ Cl (ZOSWEJ)	Mn-Cl	2.37	S14
	Mn-C	1.81	S14
	Mn-C	1.89	S14

Mn(H ₂ O) ₄ Cl ₂ trans	Mn-Cl	2.36	S4
	Mn-O	2.10	S4
	Mn-O	2.50	S4
	Mn-O	2.52	S4
Mn ₂ Cl ₄	Mn-Mn	2.29	S4
	Mn-Cl	2.13	S4
MnCl ₂ O	Mn-O	1.55	S4
	Mn-Cl	2.10	S4
	O-Mn-Cl	121.5	S4
MnC ₁₂ N ₄ H ₁₂ Cl ₂ O (MUPYUR)	Mn-Cl	2.47	S14
	Mn-N(py)	2.26	S14
	Mn-N'	2.34	S14
	N-Mn-N'	95.2	S14
MnTi	Mn-Ti	2.25	S3
MnV ₂	Mn-V	2.28	S3
MnCr	Mn-Cr	2.35	S3
MnCo	Mn-Co	2.44	S3
MnNi	Mn-Ni	2.50	S3
MnCu	Mn-Cu	2.50	S3
MnZn	Mn-Zn	2.62	S3
MnBr	Mn-Br	2.25	S4
MnBr ₂	Mn-Br	2.34	S4
MnOBr ₂	Mn-Br	2.24	S4
	Mn-O	1.55	S4
	O-Mn-Br	120.7	S4
Mn(H ₂ O) ₄ Br ₂ trans	Mn-Br	2.50	S4
	Mn-O	2.10	S4
Mn(CO) ₅ Br	Mn-Br	2.56	S3
	Mn-C	1.82	S3
	Mn-C'	1.88	S3
MnBr ₄ ²⁻ (PYDMNB)	Mn-Br	2.50	S14
MnAlC ₆ Br ₃ O ₅ (CMNCXA10)	Mn-Al	3.53	S14
	Mn-C	1.95	S14
	Mn-C'	1.87	S14
	Br-Al-Mn	140.2	S14
MnZr	Mn-Zr	2.43	S3
MnMo	Mn-Mo	2.27	S3
MnI	Mn-I	2.46	S4
MnOI ₂	Mn-I	2.45	S4
	Mn-O	1.55	S4
	O-Mn-I	120.1	S4
Mn(H ₂ O) ₄ I ₂ trans	Mn-I	2.73	S4
	Mn-O	2.08	S4
Mn(CO) ₅ I (VUCFAA)	Mn-I	2.69	S14
	Mn-C	1.84	S14
MnC ₁₃ N ₂ H ₈ O ₃ I (YOMHEN)	Mn-I	2.72	S14
	Mn-N	2.05	S14
	Mn-C	1.77	S14
	I-Mn-N	87.7	S14
Mn ₂ (CO) ₈ I ₂ (SIZYUV)	Mn-Mn	3.98	S14
	Mn-I	2.70	S14
	Mn-C	1.88	S14
	I-Mn-Mn	42.6	S14

MnI(SH)	Mn-I	2.41	S4
	Mn-S	2.10	S4
	I-Mn-S	115.6	S4
MnI(F)(PH ₂)(SH)	Mn-I	2.50	S4
	Mn-F	1.75	S4
	Mn-P	2.27	S4
	Mn-S	2.16	S4
Geometrical Variables for Iron Containing Compounds			
Compound	Variable	Target	Source
Fe ₂	Fe-Fe	2.19	S3
FeH	Fe-H	1.63	S4
FeH ⁻	Fe-H	1.69	S3
FeH ⁺	Fe-H	1.60	S4
FeCH ₃	Fe-C	1.97	S3
FeC ₅ H ₅	Fe-C	2.17	S3
Fe(C ₅ H ₅) ₂	Fe-C	2.06	S4
Fe(III)Cp ₂ ⁺	Fe-C	2.06	S4
FeO	Fe-O	1.62	S4
FeO ⁻	Fe-O	1.69	S3
FeO ⁺	Fe-O	1.56	S3
FeO ₂ ⁻	Fe-O	1.71	S3
FeOH	Fe-O	1.83	S4
Fe(OH) ₂	Fe-O	1.78	S3
Fe(H ₂ O) ₄ (OH) ₂	Fe-O	2.02	S4
	Fe-O	2.08	S4
Fe(H ₂ O) ₅ (OH)	Fe-O	1.95	S4
	Fe-O	2.05	S4
Fe(H ₂ O) ₆ ²⁺	Fe-O	2.07	S4
	Fe-O	2.15	S4
Fe(H ₂ O) ₆ ³⁺	Fe-O	2.06	S12
FeCH ₅ O ₂ (methyl iron(III)dihydroxide)	Fe-O	1.75	S4
	Fe-C	1.93	S4
Fe(CO)	Fe-C	1.72	S3
Fe(CO) ⁻	Fe-C	1.79	S3
Fe(CO) ₂	Fe-C	1.84	S3
Fe(CO) ₂ ⁻	Fe-C	1.79	S3
Fe(CO) ₃	Fe-C	1.67	S3
Fe(CO) ₃ ⁻	Fe-C	1.82	S3
Fe(CO) ₄	Fe-C	1.78	S3
	Fe-C	1.82	S3
Fe(CO) ₄ ⁻	Fe-C	1.78	S3
Fe(CO) ₄ ²⁻	Fe-C	1.75	S4
Fe(CO) ₄ H ₂	Fe-C	1.81	S15
	Fe-H	1.56	S15
	H-Fe-C	89.9	S15
Fe(CO) ₅	Fe-C(eq)	1.81	S4
	Fe-C(ax)	1.81	S4
Fe(CO) ₄ C ₂ H ₄	Fe-C	1.81	S4
	Fe-C	2.13	S4
FeC ₆ O ₁₂ ³⁻ (Fe(III)(Ox) ₃)	Fe-O	1.97	S4
	O-Fe-O	83.2	S4
FeC ₆ H ₃ O ₁₂ (H ₃ Fe(III)(Ox) ₃)	Fe-O	1.92	S4
	Fe-O	2.03	S4

FeC ₇ H ₄ O ₃ (cyclobutadiene iron tricarbonyl)	Fe-C(C=O)	1.79	S4
	Fe-C(C ₄ H ₄)	2.06	S4
	C-Fe-C	99.3	S4
FeC ₈ H ₂ O ₆ (FCPEN0)	Fe-C	1.81	S14
	Fe-C	2.02	S14
FeC ₉ H ₁₂ O (BUDFEC01)	Fe-C(O)	1.77	S14
	Fe-C	2.06	S14
	C-Fe-C	125.6	S1
FeC ₉ H ₁₂ O (FeCO(1,3-C ₄ H ₆) ₂)	Fe-C	2.11	S4
	C-Fe-C	88.8	S4
Fe ₂ (CO) ₉	Fe-C	1.83	S16
	Fe-C	1.98	S16
	Fe-C#O	180.0	S16
	Fe-C#O	140.7	S16
FeC ₁₀ H ₁₄ O ₄ (Fe(II)(Acac) ₂)	Fe-O	1.90	S4
	O-Fe-O	93.9	S4
FeC ₁₅ H ₂₁ O ₆ ⁻ (Fe(II)(Acac) ₃)	Fe-O	1.97	S4
	O-Fe-O	95.1	S4
FeN	Fe-N	1.56	S3
FeN ⁺	Fe-N	1.58	S3
FeN ⁻	Fe-N	1.58	S3
Fe(NH ₃) ₆ ²⁺	Fe-N	2.10	S4
Fe(CN) ₆ ⁴⁻	Fe-C	1.99	S3
	C-Fe-C	180.0	S3
Fe(CO) ₂ (NO) ₂	Fe-C	1.82	S4
	Fe-N	1.68	S4
	C-Fe-C	98.7	S4
	N-Fe-N	109.0	S4
FeC ₆ N ₆ H ₂₄ ²⁺ (Fe(II)en) ₃	Fe-N	2.13	S4
FeC ₇ NH ₅ O ₅ (CNOFEA)	Fe-C	1.79	S14
	Fe-C	2.09	S14
	Fe-O	1.97	S14
FeC ₈ NH ₅ O ₂ (ACODUR)	Fe-C	1.91	S14
	Fe-C	1.78	S14
	Fe-C	2.19	S14
	C-Fe-C	89.2	S14
FeC ₉ NH ₈ O ₂ ⁺ (CPACFE)	Fe-N	1.91	S14
	Fe-C	1.77	S14
FeC ₁₀ N ₄ H ₁₀ O ₆ (Diaqua-bis(pyrazinecarboxylato)-iron)	Fe-O	2.11	S17
	Fe-N	2.12	S17
	Fe-OH ₂	2.14	S17
	O-Fe-N	78.1	S17
FeC ₁₀ N ₂ H ₁₂ O ₈ ⁻ (Iron(III)EDTA)	Fe-N	2.03	S4
FeC ₁₆ N ₅ H ₁₁ O ₄ (FeN3O2)	Fe-O	2.12	S4
	Fe-O	1.92	S4
	Fe-N	1.89	S4
	O-Fe-O	81.4	S4
FeC ₁₇ N ₂ H ₁₆ (Toluene-(2,2'-bipyridine)-iron)	Fe-N	1.90	S18
	Fe-C	2.11	S18
	N-Fe-N	81.9	S18
	N-Fe-C	122.8	S18

FeC ₂₀ H ₁₂ N ₄ (iron porphyrin)	Fe-N	1.97	S4
FeC ₃₆ N ₆ H ₂₄ ⁺ (ferrous tris(ortophenantholine))	Fe-N	2.05	S4
FeF	Fe-F	1.76	S4
Fe(II)(H ₂ O) ₅ F ⁺	Fe-F	1.89	S4
	Fe-O	2.05	S4
FeF ₂	Fe-F	1.77	S4
FeO ₂ F ₂ (iron(VI)difluoride dioxide)	Fe-F	1.74	S4
	Fe=O	1.58	S4
FeF ₃	Fe-F	1.78	S3
FeF ₆ ³⁻	Fe-F	2.06	S4
FeAl	Fe-Al	2.30	S3
Fe(AlH ₂) ₂	Fe-Al	2.47	S3
FeSiH	Fe-Si	2.30	S3
FeSiH ₂	Fe-Si	2.26	S3
Fe(SiH ₃) ₂	Fe-Si	2.52	S3
FeP	Fe-P	2.23	S3
FePH	Fe-P	2.20	S3
FeC ₁₂ P ₄ H ₃₄ S ₂ (JIYFAY)	Fe-S	2.35	S14
	Fe-P	2.23	S14
	P-Fe-S	89.4	S14
FeS	Fe-S	2.04	S3
Fe(SH) ₂	Fe-S	2.25	S3
FeC ₉ H ₁₀ SO ₂ (CEYTFE)	Fe-S	2.30	S4
	Fe-C(O)	1.75	S4
	Fe-C(C ₄)	2.12	S4
	C-Fe-S	90.7	S4
FeC ₁₆ N ₂ H ₁₈ S ₂ O ₂ (dicarbonyl ethylenediamine bis(phenylthiolato)iron)	Fe-S	2.33	S19
	Fe-S	2.34	S19
	Fe-N	2.04	S19
	Fe-N	2.03	S19
	Fe-C	1.76	S19
	N-Fe-S	86.8	S19
	C-Fe-S	93.2	S19
FeC ₉ N ₃ H ₁₈ S ₃ O ₃ (tris(N,N-dimethylthiocarbamato)iron(III))	Fe-S	2.43	S20
	Fe-O	2.09	S20
	S-Fe-O	68.9	S20
	S-Fe-S	103.1	S20
Fe ₄ S ₄	Fe-S	2.27	S3
FeC ₁₂ H ₁₄ S ₄ O ₂ (CIBGAV10)	Fe-S	2.31	S14
	Fe-C	1.80	S14
	S-Fe-S	85.2	S14
	C-Fe-S	90.1	S14
FeC ₁₄ N ₆ H ₁₆ S ₆ (diisothiocyanato-bisthiazoline iron(II))	Fe-N	2.19	S21
	Fe-N(CS)	2.08	S21
	N-Fe-N	73.9	S21
	SCN-Fe-NCS	97.4	S21
FeCl	Fe-Cl	2.13	S4

Fe(II)(H ₂ O) ₅ Cl ⁺	Fe-Cl	2.29	S4
	Fe-O	2.06	S4
FeCl ₂	Fe-Cl	2.16	S4
FeCl ₃	Fe-Cl	2.16	S3
FeC ₆ N ₂ H ₁₈ Cl ₃ (FINJIV)	Fe-Cl	2.23	S14
	Fe-N	2.27	S14
	Cl-Fe-Cl	121.0	S14
FeCl ₄ ⁻	Fe-Cl	2.19	S4
FeCl ₄ ²⁻ (GOXLUA)	Fe-Cl	2.34	S14
FeCl ₆ ³⁻	Fe-Cl	2.53	S4
FeTi	Fe-Ti	2.54	S3
FeV	Fe-V	2.31	S3
FeCr	Fe-Cr	2.27	S3
FeMn	Fe-Mn	2.40	S3
FeCo	Fe-Co	2.31	S3
FeNi	Fe-Ni	2.33	S3
FeCu	Fe-Cu	2.31	S3
FeZn	Fe-Zn	2.53	S3
FeBr	Fe-Br	2.23	S4
FeBr ₂	Fe-Br	2.31	S3
FeC ₆ H ₅ BrO ₃ (ALCFEA)	Fe-Br	2.50	S14
	Fe-C(C ₂)	2.13	S14
	Fe-C(O)	1.79	S14
	C-Fe-Br	88.6	S14
Fe(H ₂ O) ₅ Br ⁺	Fe-Br	2.42	S4
	Fe-O	2.06	S4
Fe(H ₂ O) ₄ Br ₂	Fe-O	2.07	S4
	Fe-Br	2.50	S4
FeC ₄ H ₈ BrO ₇ (Diaquabromo(oxydiacetato- O,O',O'')-iron(III))	Fe-Br	2.37	S4
	Fe-O	2.00	S4
	Br-Fe-O	105.6	S4
FeBr ₂ O ₂	Fe-Br	2.27	S4
	Fe=O	1.58	S4
	Br-Br-Fe	110.6	S4
FeZr	Fe-Zr	2.59	S3
FeMo	Fe-Mo	2.21	S3
FeI	Fe-I	2.44	S4
Fe(CO) ₃ (C ₃ H ₅)I	Fe-I	2.75	S4
	Fe-C(O)	1.80	S4
	C-Fe-I	81.6	S4
Fe(CO) ₃ (C ₃ H ₅)I (ALCOFE10)	Fe-I	2.75	S14
	Fe-C	1.80	S14
	C-Fe-I	81.6	S14
FeI ₂	Fe-I	2.50	S3

REFERENCES

- [S1] NIST Chemistry WebBook, NIST Standard Reference Database Number 69, (<http://webbook.nist.gov/>), Linstrom P., Mallard W. National Institute of Standards and Technology: Gaithersburg MD, 20899, 2003
- [S2] Derived from ionization energies.
- [S3] B3LYP/LANL2DZ with polarization functions, see main text for references.
- [S4] Stewart J.J.P., www.openmopac.net
- [S5] Cox J.O., Pilcher G., "Thermochemistry of Organic and Organometallic Compounds," Academic Press, New York, N.Y., 1970.
- [S6] Holder A. J. (1994) "SAM1 Semiempirical Parameters", Report, University of Missouri Kansas City Dept. of Chemistry, AFOSR-NL-94-29773.
- [S7] Lias S.G., Bartmess J.E., Liebman J.F., Holmes J.L., Levin R.D., Mallard W.G. (1988) "Gas Phase Ion and Neutral Thermochemistry", J. Phys. Chem., Ref. Data, Suppl. 1, 17.
- [S8] Pilcher G., Skinner H.A. (1982) The Chemistry of the Metal-Carbon Bond, F. R. Hartley and S. Patai, Eds., Wiley, New York, pp. 43-90.
- [S9] Bauschlicher C.W., Langhoff S.R., Partridge H., Barnes L.A. (1989) J. Chem. Phys. 91(4):2399-2411.
- [S10] L. McClellan A.L. (1974) "Tables of Experimental Dipole Moments," Vol 2, Raha Enterprises, El Cerrito.
- [S11] Gutsev G.L., Khanna S.N., Rao B.K., Jena P. (1999), J. Phys. Chem. A 103(29):5812-5822.
- [S12] Zeitschrift für Kristallographie, New Crystal Structures, <http://www.oldenbourg.de/verlag/zkristallogr/mn-ncsfi2000.htm>
- [S13] LaPlaca S.J., Hamilton W.C., Ibers J.A., Davidson A. (1969), Inorg. Chem. 8(9):1928-1935.
- [S14] Cambridge Structural Database, Version 5.28 (2007) Cambridge Crystallographic Data Centre, Cambridge, UK.
- [S15] Simoes M., Beauchamp J.L. (1990) Chem. Rev. 90:629-688.
- [S16] Cotton F.A., Troup J.M. (1974) J. Chem. Soc., Dalton Trans. 800-802.
- [S17] Klein C.L., O'Connor C.J., Majeste R.J., Trefonas L.M. (1982), J. Chem. Soc., Dalton Trans. 2419-2423.
- [S18] Radonovich L.J., Eyring M.W., Groshens T.J., Klabunde K.J. (1982) J. Am. Chem. Soc. 104:2816-2819.

- [S19] Takacs J., Soos E., Nagy-Magos Z., Marko L., Gervasio G., Hoffmann T.
(1989) *Inorg. Chim. Acta* 166:39-46.
- [S20] Ahmed J., Ibers J.A. (1977) *Inorg. Chem.* 16:935-937.
- [S21] Ozarowski A., McGarvey B.R., Sarkar A.B., Drake J.E. (1988) *Inorg. Chem.*
27:628-635