



**HAL**  
open science

## Management of post-splenectomy patients in the Netherlands

A. J. J. Lammers, D. Veninga, M. J. M. H. Lombarts, J. B. L. Hoekstra, P. Speelman

► **To cite this version:**

A. J. J. Lammers, D. Veninga, M. J. M. H. Lombarts, J. B. L. Hoekstra, P. Speelman. Management of post-splenectomy patients in the Netherlands. *European Journal of Clinical Microbiology and Infectious Diseases*, 2010, 29 (4), pp.399-405. 10.1007/s10096-009-0870-x . hal-00568346

**HAL Id: hal-00568346**

**<https://hal.science/hal-00568346v1>**

Submitted on 23 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manuscript Number: EJCMIID-D-09-00369R1

Title: Management of post-splenectomy patients in the Netherlands.

Article Type: Article

Keywords: splenectomy; immunizations; sepsis; management

Corresponding Author: Ms. Jolanda Lammers, M.D.

Corresponding Author's Institution: Academic Medical Center

First Author: Jolanda Lammers, MD

Order of Authors: Jolanda Lammers, MD; D Veninga; M J Lombarts; J B Hoekstra; P Speelman

Abstract: Purpose.

After splenectomy, patients are at increased risk of sepsis with considerable mortality. The risk of sepsis can be reduced by immunizing these patients and by prescribing antibiotic prophylaxis. The purpose of our study was to determine compliance with the international standards for the management of splenectomised patients in the Netherlands, by investigating (i) vaccination rates, (ii) prescription of antibiotics and (iii) information in discharge letters.

Methods.

A retrospective review of medical records and discharge correspondence of 609 splenectomy patients from 1997 to 2008 was performed. Data were collected from 28 hospitals. Adherence to vaccination guidelines and the prescription of antibiotics were assessed.

Results.

85.4% of post-splenectomy patients received pneumococcal vaccination, 39.4% received H. Influenzae type B and 32.3% received meningococcal group C vaccination. 12.4% of patients were discharged on prophylactic antibiotics. In less than 25% of cases adequate recommendations regarding post-splenectomy management were given to the general practitioner.

Conclusions.

In the Netherlands compliance with recommendations for management of patients after splenectomy is insufficient. Fifteen percent does not receive vaccination against pneumococci and the majority of patients does not receive antibiotic prophylaxis. The development and implementation of a national guideline for splenectomised patients is urgently required.

Response to Reviewers: Response to Reviewers:

1. The Results section is clear but perhaps the categories of splenectomy could more easily be divided into Emergency and Elective, with further sub-division of Emergency into non-iatrogenic and iatrogenic and Elective into planned pre-operative and planned peri-operative.

Answer:

At present, in the manuscript the following 4 categories are used:

- A. acute/emergency splenectomy
- B. elective/planned splenectomy
- C. iatrogenic/inadvertent splenectomy
- D. durante operatione.

The suggestion of reviewer #1 is to change the categories into:

- A. emergency splenectomy, further divided into iatrogenic vs. non-iatrogenic
- B. elective splenectomy, further divided into planned pre-operatively and planned peri-operatively.

We have considered this division during our data-analysis as well, but concluded the definitions would than not be appropriate for the following reasons:

1. In our definition, not all iatrogenic splenectomies are emergency splenectomies: in our acute/emergency category only those surgeries were included where there was spontaneous bleeding of the spleen with shock, rupture of an aneurysm of the splenic artery, or trauma to the spleen caused for example by high-energetic trauma. Splenectomies due to bleeding caused by the surgeon when operating for another indication were not included here. In our opinion, these are different categories, for example differing in setting (multiple trauma/acute operation versus scheduled bowel-surgery) and operating staff (trauma surgeons versus surgeons specialized in stomach/esophagus surgery).

2. The category of elective splenectomy cannot be divided into "planned peri-operatively", since we are investigating timing of vaccination. When a splenectomy is planned, ideally the vaccination should be given 2 weeks prior to the surgery. If it is decided during surgery to remove the spleen, pre-operative vaccination has not been performed.

2. The authors have not indicated whether there was any change in practice over time

Answer:

These data are readily available in our database. We decided to omit these data considering length of the manuscript, but we have added them as suggested by reviewer #1. See table 5.

3. nor were there any differences in management by institution (eg. Teaching versus non-Teaching).

Answer:

We acknowledge the lack of institutional data in this manuscript, as suggested by reviewer #1. However, considering the length of the manuscript, as well as relevance to the EJCMIID, we have decided to accumulate these management data in a second manuscript (including data regarding hospital teaching status, but also the influence of the presence of hospital guideline or hospital protocol on hospital performance in the prevention of infections, as well as the influence of the number of surgeons in the hospital staff on performance). This second manuscript is currently under revision at the Journal of Hospital Medicine.

4. Data on Emergency versus Elective procedures is available for vaccination but not antibiotic prescribing. Although one would predict differences in these areas confirmation of the influence of these measurables on out-come would be of some interest.

Answer:

Again, we acknowledge the lack of these data in the manuscript, as suggested by reviewer #1. Considering the low antibiotic prescription rates, we considered these numbers not to be as relevant as for vaccination status. However, these data can and have been added to the manuscript. See table 4.

1  
2  
3  
4 **Management of post-splenectomy patients in the Netherlands**  
5  
6  
7

8  
9 A.J.J. Lammers, D. Veninga, M.J.M.H. Lombarts, J.B.L. Hoekstra, P. Speelman.  
10

11  
12  
13  
14 *Corresponding author:*

15  
16 A.J.J. Lammers, Department of Infectious Diseases, Tropical Medicine and AIDS, G2-  
17  
18 105, Academic Medical Center, Meibergdreef 9, 1105 AZ, Amsterdam, Netherlands.  
19

20  
21 Tel.(+31)-20-566 7062 / 64380, fax: (+31)-20-6972286.  
22

23  
24 Email: [a.j.lammers@amc.uva.nl](mailto:a.j.lammers@amc.uva.nl)  
25  
26  
27

28  
29 *Affiliations and addresses of other authors:*  
30

31 D. Veninga, Department of Infectious Diseases, Tropical Medicine and AIDS, F4-217,  
32  
33 Academic Medical Center, Meibergdreef 9, 1105 AZ, Amsterdam, Netherlands.  
34  
35  
36  
37

38 M.J.M.H. Lombarts, PhD, Department of Quality and Process Innovation, J2-211,  
39  
40 Academic Medical Center, Meibergdreef 9, 1105 AZ, Amsterdam, Netherlands.  
41  
42  
43  
44

45 J.B.L. Hoekstra, professor, Department of Internal Medicine, F4-217,  
46  
47 Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, Netherlands  
48  
49  
50  
51  
52

53 P. Speelman, professor, Department of Infectious Diseases, Tropical Medicine and AIDS,  
54  
55 Academic Medical Center, F4-217, Meibergdreef 9, 1105 AZ Amsterdam, Netherlands.  
56  
57  
58  
59  
60  
61  
62

1  
2  
3  
4 **Abstract**  
5  
6  
7  
8

9 *Purpose.* After splenectomy, patients are at increased risk of sepsis with considerable  
10 mortality. The risk of sepsis can be reduced by immunizing these patients and by  
11 prescribing antibiotic prophylaxis. The purpose of our study was to determine compliance  
12 with the international standards for the management of splenectomised patients in the  
13 Netherlands, by investigating (i) vaccination rates, (ii) prescription of antibiotics and (iii)  
14 information in discharge letters.  
15  
16  
17  
18  
19  
20  
21  
22

23 *Methods.* A retrospective review of medical records and discharge correspondence of 609  
24 splenectomy patients from 1997 to 2008 was performed. Data were collected from 28  
25 hospitals. Adherence to vaccination guidelines and the prescription of antibiotics were  
26 assessed.  
27  
28  
29  
30  
31  
32

33 *Results.* 85.4% of post-splenectomy patients received pneumococcal vaccination, 39.4%  
34 received *H. Influenzae* type B and 32.3% received meningococcal group C vaccination.  
35 12.4% of patients were discharged on prophylactic antibiotics. In less than 25% of cases  
36 adequate recommendations regarding post-splenectomy management were given to the  
37 general practitioner.  
38  
39  
40  
41  
42  
43  
44

45 *Conclusions.* In the Netherlands compliance with recommendations for management of  
46 patients after splenectomy is insufficient. Fifteen percent does not receive vaccination  
47 against pneumococci and the majority of patients does not receive antibiotic prophylaxis.  
48 The development and implementation of a national guideline for splenectomised patients  
49 is urgently required.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 **Introduction**  
5  
6  
7

8  
9 The spleen has an important function in the host protection against invading micro-organisms [1].  
10 Splenic marginal zone macrophages filter and phagocytose bacteria and other blood-borne  
11 pathogens from the blood that flows through the splenic red pulp [2]. After splenectomy there is a  
12 diminished clearance of poorly opsonized bacterial antigens, an impaired primary humoral  
13 response to antigenic challenge and loss of the vagally mediated anti-inflammatory cholinergic  
14 response [3,4].  
15  
16  
17  
18  
19  
20

21  
22 The asplenic state thus results in a serious impairment of clearance of mainly encapsulated  
23 bacteria, which can lead to a life-threatening infection known as post-splenectomy sepsis, PSS  
24 [5,6]. The encapsulated bacterium *Streptococcus pneumoniae* is the causative pathogen in 80% of  
25 cases of PSS [7,8]. Although often preceded by nonspecific symptoms, PSS deteriorates in hours  
26 rather than days to a severe septic state that is associated with a high mortality of 50-70 %  
27 [7,9,10]. The incidence of sepsis in post-splenectomy patients is estimated to be 600 times greater  
28 than in the general population [11]. The incidence of invasive infection among children and adults  
29 after splenectomy is similar, 3.3% and 3.2% respectively [12].  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 Several relevant organizations developed recommendations to prevent this syndrome [13].

41  
42 Evidence supporting these recommendations is mostly based on clinical studies (retrospective  
43 studies, series of case-reports) and expert opinion. Unfortunately, data from randomized  
44 controlled intervention trials are not available. In the Netherlands, a guideline for the management  
45 of post-splenectomy patients has not yet been developed.  
46  
47  
48  
49  
50

51 In 1996, the British Committee for Standards in Haematology published Guidelines for the  
52 prevention and treatment of infection in patients with an absent or dysfunctional spleen [14], that  
53 are considered to reflect best-practice. These recommendations were updated in 2002 [15], and  
54 consist of the following key elements:  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1. All splenectomised patients should receive pneumococcal immunization and revaccination. If not previously immunized they should also receive *Haemophilus influenzae* type B and meningococcal C vaccine. Yearly influenza immunization is recommended.

2. Lifelong prophylactic antibiotics are recommended, or at least the first two years after splenectomy. In case of suspected or proven infection patients should be given systemic antibiotics and be admitted to a hospital.

3. All patients should be educated about the risks of infection and the risk associated with traveling (such as becoming infected by *Plasmodium falciparum*) and unusual infections (i.e. dog bites). Patient records should be labeled to indicate the risk.

It has become clear from recent literature that post-splenectomy patients are not all being managed according to best practice [5,8,9,16-21]

The goal of our study was to determine the management of post-splenectomy patients for the first time in the Netherlands. We therefore assessed compliance with the British best practice standards by investigating (i) vaccination rates, (ii) whether or not antibiotics were prescribed and (iii) if the hospital provided the general practitioner with relevant information regarding the period after the operation.



1  
2  
3  
4 **Methods**  
5  
6  
7

8 *Hospital and patient inclusion*  
9

10 Splenectomised patients were included retrospectively from a representative sample of Dutch  
11 hospitals. First, 30% of all Dutch hospitals were identified through a blind drawing, including  
12 academic, teaching and non teaching hospitals. Subsequently, patients were identified using the  
13 Dutch Pathology Registry, since spleens are routinely sent to the pathology department after  
14 removal. In the Registry, a search query *\*milt\* (spleen)* was performed, after which all  
15 splenectomies performed from 1997 to 2008 were selected and non-relevant hits such as partial  
16 splenectomies or spleen biopsies were removed. Inclusion started with the most recent  
17 splenectomy, subsequently going back in time in a consecutive matter. After hospitals and  
18 patients were identified, the medical records and all discharge correspondence were assessed on  
19 site. In case the medical file or discharge correspondence was not available, the patient was  
20 excluded. To investigate discharge correspondence, we included discharge letters as well as all  
21 other correspondence up to at least 1 year after splenectomy, for example from follow-up out-  
22 patient visits.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 *Used definitions*  
43

44 After inclusion, patients were divided into 4 categories of splenectomy: ‘acute’ in case of  
45 emergency splenectomies, ‘elective’ in case of planned splenectomies, ‘iatrogenic’ in case of  
46 inadvertent splenectomies and ‘durante operatione’ in case it was decided during the operation to  
47 perform a non-inadvertent splenectomy.  
48  
49  
50

51 Vaccination was considered to be correct if 3 immunizations were given: pneumococcal, *H.*  
52 *influenzae* B and meningococcal group C vaccination. Pneumococcal vaccination was defined as  
53 immunization with either the 23-valent pneumococcal polysaccharides vaccine (PPV-23,  
54 Pneumovax), the 7-valent pneumococcal conjugate vaccine (PCV-7, Prevenar), or both. The  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 British guidelines recommend that pneumococcal vaccination is given either 2 weeks prior to or 2  
5  
6 weeks after splenectomy. In this study, timing was considered correct if given either 1-3 weeks  
7  
8 prior to surgery in case of elective splenectomies, or 1-3 weeks after the operation.  
9

10 Prescription of antibiotics was considered to be correct if antibiotic prophylaxis was given for the  
11  
12 first 2 years after splenectomy, in combination with a prescription for on-demand (broad  
13  
14 spectrum) antibiotics to be taken in case of suspected infection.  
15  
16  
17  
18  
19

### 20 *Data collection method*

21 All data were collected by the same two investigators (DV, JL), using a standardized survey form.  
22  
23 Individual data included patient demographics, indication for splenectomy, documentation of  
24  
25 vaccine administration, timing of vaccination in relation to splenectomy and documentation of  
26  
27 post-splenectomy infections.  
28  
29

30 Furthermore, discharge correspondence was checked for mentioning of each of the following:  
31  
32 performed splenectomy, vaccination status, the need for revaccination, prescribed prophylactic  
33  
34 antibiotics, the need of urgent use of antibiotics in case of suspected infection and the advice for  
35  
36 annual flu-vaccination.  
37  
38  
39  
40  
41  
42  
43

### 44 *Statistics*

45  
46 Analysis of data was performed in SPSS 16.0. Confidence intervals were computed using CIA  
47  
48 (Confidence Interval Analysis) version 1.0, copy 5/0618 (Gardner, British Medical Journal).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 **Results**  
5  
6  
7

8  
9 In the period 1997 to 2008 a total number of 609 splenectomy cases from 28 hospitals were  
10 identified. 73 cases were excluded: 50 hospital records were not available and 23 patients could  
11 not be identified (transferred to another hospital, patient not known, no permission from patient or  
12 hospital to access the medical file). 536 patients were included (see table 1), with a mean duration  
13 of follow-up of 4.22 years. In total, 19 children under the age of 18 were included.  
14  
15  
16  
17

18 Of the 185 splenectomies of iatrogenic cause, injury to the spleen was most likely to occur in  
19 stomach/esophagus operations (32.4%), in operations of the colon descendens (18.4%) and  
20 rectum-sigmoid (11.4%) and operations of the left kidney (14.6%).  
21  
22  
23  
24  
25  
26  
27

28  
29 *Vaccination*  
30

31 To study vaccination rates, we omitted patients that died within 2 weeks after the operation  
32 (n=36). Data regarding actual vaccination was missing for 1 pneumococcal vaccination and for 2  
33 cases of the other immunizations. Overall, 426 of 499 patients (85.4%) received pneumococcal  
34 vaccination. 196 (39.4 %) patients received *H. influenzae* type B vaccination and 161 (32.3%)  
35 received the meningococcal group C vaccine. 30.5% of patients in our study received all 3  
36 recommended vaccines.  
37  
38  
39  
40  
41  
42  
43

44 Pneumococcal-vaccination rates in relation to type of operation are shown in table 2. Rates were  
45 highest for patients that underwent an acute or elective splenectomy. If the operation was  
46 performed for another indication than a spleen extirpation, such as in iatrogenic or durante-  
47 operatione splenectomies, these rates were significantly lower (CI-95% 0.158 - 0.307 for elective  
48 vs. iatrogenic splenectomy).  
49  
50  
51  
52  
53  
54

55 Of the 184 patients that underwent an elective splenectomy and received pneumococcal  
56 vaccination, 144 patients received their immunization preoperatively (78.3%) and 31  
57 postoperatively (16.7%). For 9 patients vaccination timing was not documented.  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Of all preoperatively vaccinated patients, in 12 cases (8.3%) the immunization was given 1-3  
5  
6 weeks prior to surgery, another 12 cases (8.3%) were vaccinated in the week before surgery, 13  
7  
8 (9%) were vaccinated 3 weeks to 6 months before surgery, and of 108 patients (74.5%) the exact  
9  
10 timing of vaccination was not documented.

11  
12 For immunizations given postoperatively, in 169 (62.4%) the timing of vaccination was not  
13  
14 documented. Of the remaining 107, 30 vaccinations (28%) were given 1-3 weeks after surgery, 3  
15  
16 vaccinations were given on the day of surgery, 34 within a week after surgery and 40 vaccinations  
17  
18 were given 1 – 33 months after splenectomy.  
19  
20  
21  
22

### 23 24 *Antibiotics*

25  
26 The majority of patients was discharged from the hospital without a prescription for antibiotics  
27  
28 (see table 3). The duration of antibiotic prophylaxis is shown in figure 1. Of the 77 patients  
29  
30 receiving prophylaxis, 24 patients (31.2%) received antibiotics for 2 years, 1 patient received a  
31  
32 life-long prescription because of an underlying immunodeficiency, 2 children received  
33  
34 prophylaxis until they reached 16 years of age and in 11 cases (14.3%) the duration of  
35  
36 prophylaxis was not-documented. In 6 cases (7.8%) antibiotics were given “until vaccination” (3  
37  
38 cases length unknown, 3 cases < 1 month). Table 4 shows prescription rates of antibiotics in  
39  
40 relation to type of operation.  
41  
42  
43  
44  
45  
46

### 47 *Change in time*

48  
49 In table 5, the change in vaccination rates (5A) and prescription of antibiotics (5B) over  
50  
51 time is shown. We have decided to compare the period before and after 2002, since at  
52  
53 that time the update of the British guideline was published. After 2002, there is a  
54  
55 significant increase in vaccination with *H. influenzae* and meningococci C and a  
56  
57 significant decrease in patients not receiving any antibiotics, due to more prescribed “on-  
58  
59 demand” antibiotics.  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 *Information in discharge correspondence*  
5

6 The specific information mentioned in discharge correspondence to the general practitioner (GP)  
7 is summarized in table 6. Splenectomy and given immunizations were mentioned in the majority  
8 of patients whereas other recommended advises were found in a minority of cases.  
9  
10  
11  
12  
13  
14

15 *Infectious complications*  
16

17 The number of patients that developed a documented infection after splenectomy was 295 (55%),  
18 during a mean period of follow-up of the cohort of 4.22 years (range 0-12). Infection was defined  
19 as any infectious complication found in hospital medical files and medical correspondence for  
20 each patient. Of these 295 patients, 278 (94.2 %) developed their first infection within 2 years  
21 after splenectomy.  
22  
23  
24  
25  
26  
27

28 Infectious complications were distributed over the splenectomy indications as follows: 121  
29 patients (41%) developed an infection after a splenectomy of iatrogenic cause, 90 patients (31%)  
30 after elective surgery, 60 patients (20%) after acute splenectomy and 24 (8%) if it was decided  
31 durante operatione to perform a splenectomy.  
32  
33  
34  
35  
36

37 Since asplenic hosts are particularly susceptible for blood borne infections, we investigated the  
38 incidence of bacteraemias in these patients. We found that 12.5% of patients developed a  
39 documented bacteraemic episode after splenectomy (i.e. 67 of 536 patients). A total of 96  
40 bacteraemias occurred in these 67 patients. Pathogens responsible for these bacteraemias were  
41 identified in 61 episodes and are shown in table 7. No meningococcal bacteraemia was  
42 documented. Other infections included pneumonias (155, of which 17 were proven positive for  
43 either *S. pneumoniae* or *H. influenzae*), urinary tract infections (86), fever of unknown origin (78)  
44 and miscellaneous causes (149) such as abdominal abscesses, wound infection and diarrhea.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 Patient-characteristics of documented pneumococcal and *H. influenzae* bacteraemias after  
56 splenectomy are shown in table 8.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 **Discussion**  
5  
6  
7

8  
9 Although vaccination results in the Netherlands are among the highest reported in literature, it is  
10 still of concern that 14.6% of post-splenectomy patients was not vaccinated against pneumococci,  
11 60.6% of splenectomised patients did not receive vaccination against *H. influenzae*, and 67.7%  
12 did not receive meningococcal vaccination.  
13  
14

15  
16  
17 We found only 12.4% of included patients to have received a prescription for prophylactic  
18 antibiotic therapy, of which 40.9% was of correct duration. The role and efficacy of prophylactic  
19 antibiotics after splenectomy remains unclear and their use is not generally recommended [6].  
20  
21

22  
23  
24 However, in the absence of evidence against it, the British Guidelines recommend to offer  
25 lifelong prophylaxis with oral phenoxymethylpenicillin. If there is hesitation to prescribe lifelong  
26 antibiotics, it is recommended to cover at least the first two years after splenectomy, as well as  
27 children aged up to 16. In most hospitals included in this survey, antibiotic prophylaxis is not  
28 recommended and compliance even for these first two years is estimated to be low.  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 Reviewing discharge correspondence, it is encouraging to note that in the majority of cases  
39 splenectomy status (97.7%) was documented. It is however unfortunate that of all immunized  
40 patients in 20% the vaccination was not mentioned to the general practitioner and that other  
41 recommendations were given only in a minority of cases. Primary care doctors, who are also  
42 responsible for keeping patients fully immunized, should be informed about necessary  
43 vaccinations, the importance of antibiotics and the measurements that should be taken in case of a  
44 potential PSS.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 *Infections after splenectomy*  
56

57  
58 Although very relevant, our study was not designed to describe the effectiveness of vaccination or  
59 antibiotics on reducing infections after splenectomy. In a recent review, it is described that non-  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 vaccinated patients have a higher incidence of infection than vaccinated patients [13]. In some  
5  
6 studies this difference was not significant, however overall there does appear to be both a  
7  
8 statistical and a clinically relevant difference.  
9

10 More than half of the patients in our cohort developed a documented infection after surgery. Most  
11  
12 infections occurred after an iatrogenic splenectomy. It is however likely for patients to have a  
13  
14 higher risk of infection after major abdominal surgery than after a splenectomy only. The  
15  
16 incidence of bacteraemia in splenectomised patients in this study was 12.5% during an average  
17  
18 follow up period of 4.22 years after splenectomy. There is evidence that besides the known  
19  
20 susceptibility of asplenic hosts for encapsulated bacteria, other organisms such as Gram-negative  
21  
22 bacteria and *Capnocytophaga canimorsus* cause infections and carry considerable morbidity  
23  
24 [5,10]. In a study by Holdsworth and colleagues, it is demonstrated that of 349 postsplenectomy  
25  
26 infections *E. coli* and *Pseudomonas*, although with low incidence, carried the highest mortality  
27  
28 [7]. Although the causative microorganisms in our study were not always identified, we  
29  
30 demonstrate that indeed not only encapsulated pathogens play a role in postsplenectomy  
31  
32 bacteraemia, but that asplenic hosts are also at risk for staphylococci, enterococci, *E. coli*, and  
33  
34 other pathogens. For the bacteraemias caused by coagulase-negative staphylococci however, it  
35  
36 remains unclear whether these pathogens have caused a clinical relevant infection. *N.*  
37  
38 *meningitidis* has been cited as the third most common cause for PSS [7]. Even though  
39  
40 meningococcal vaccination rates were low, we have found no bacteraemic episodes based on  
41  
42 meningococcal infection. In our study only one case of meningitis was documented, which was  
43  
44 due to *S. aureus*.  
45  
46  
47  
48  
49

50 Despite voluminous literature on post-splenectomy infection, estimates of bacteraemic risk vary  
51  
52 and precise figures remain unclear. This is in part due to variation in definitions of PSS, variation  
53  
54 in duration of follow-up, lacking documentation and heterogeneity and small sample sizes of  
55  
56 study populations.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 *Limitations*  
5

6 There are some limitations to our investigation. First, there are several aspects of our study that  
7 are likely to bias the outcome: the large variation in duration of follow-up (range from several  
8 months to 12 years), patients being lost to follow-up due to transfer to another hospital of care,  
9 (severe) infections or death under supervision of the GP thus not identified in our study, all  
10 leading to an underestimation of the true incidence of infection and death. This possible  
11 underestimation would result in unreliable conclusions on the effects of preventive measures in  
12 infection and death after splenectomy in our cohort.  
13  
14

15 Furthermore, we have not contacted patients to check for their knowledge about the need for  
16 (re)vaccination, antibiotics and precautions to be taken when travelling. Informing patients is  
17 considered to be a major factor in improving management [15]. Another consequence is that  
18 although booster vaccinations, prophylactic or on-demand antibiotics may have been  
19 recommended, we are not informed about patients' and GP's compliance. This may cause an  
20 overestimation of antibiotic use and revaccination rate.  
21  
22

23 Finally, although we demonstrate vaccination rates, it remains unclear whether these vaccinations  
24 were effective, since 5-10% of healthy adults fail to produce antibody to most or all  
25 polysaccharides in PPV23 [22]. Ideally, patients should be tested with titer-analysis for adequate  
26 antibody levels. Although there are no large scale, randomized controlled intervention trials that  
27 have tested the efficacy of bacterial vaccines in preventing PSS, data from studies that reported  
28 antibody levels measured before and after vaccination, suggest that the risk of infection in  
29 vaccinated patients is higher in those with a poor immunological response [23].  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 **Conclusion**  
54

55 We have investigated the management of post-splenectomy patients in the Netherlands for the  
56 first time. Although the pneumococcal vaccination rate after splenectomy is among the highest  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

reported in literature, hospitals could do much better at providing general practitioners with recommendations for post-splenectomy management.

Future research will have to investigate the reason for this non-compliance with international guidelines. It is important that all caregivers aim at 100% vaccination rates and all patients being discharged with antibiotics and clear instructions on how and when to use them. Therefore, we emphasize the need for a Dutch guideline for this group of patients.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## **Acknowledgements**

This study was supported by ZonMw.

We would like to thank all participating hospitals for their cooperation.

There are no potential conflicts of interest for any of the authors.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Morris DH, Bullock FD (1919) THE IMPORTANCE OF THE SPLEEN IN RESISTANCE TO INFECTION. *Ann.Surg.* 70:513-521.
2. Mebius RE, Kraal G (2005) Structure and function of the spleen. *Nat.Rev.Immunol.* 5:606-616.
3. Hosea SW, Brown EJ, Hamburger MI et al (1981) Opsonic requirements for intravascular clearance after splenectomy. *N.Engl.J.Med.* 304:245-250.
4. Huston JM, Ochani M, Rosas-Ballina M et al (2006) Splenectomy inactivates the cholinergic antiinflammatory pathway during lethal endotoxemia and polymicrobial sepsis. *J.Exp.Med.* 203:1623-1628.
5. Deodhar M, Kakkar N (2004) An audit of splenectomies in a teaching hospital in North India. Are postsplenectomy guidelines being complied with? *J.Clin.Pathol.* 57:407-410.
6. Newland A, Provan D, Myint S (2005) Preventing severe infection after splenectomy. *Bmj* 331:417-8.
7. Holdsworth RJ, Irving AD, Cuschieri A (1991) Postsplenectomy sepsis and its mortality rate: actual versus perceived risks. *Br J Surg* 78:1031-8.
8. Waghorn DJ (2001) Overwhelming infection in asplenic patients: current best practice preventive measures are not being followed. *J Clin Pathol* 54:214-8.
9. Brigden ML, Pattullo A, Brown G (2000) Pneumococcal vaccine administration associated with splenectomy: the need for improved education, documentation, and the use of a practical checklist. *Am J Hematol* 65:25-9.
10. Styrt B (1990) Infection associated with asplenia: risks, mechanisms, and prevention. *Am.J.Med.* 88:33N-42N.
11. Singer DB (1973) Postsplenectomy sepsis. *Perspect.Pediatr.Pathol.* 1:285-311.:285-311.
12. Bisharat N, Omari H, Lavi I et al (2001) Risk of infection and death among post-splenectomy patients. *J Infect* 43:182-6.
13. Mourtzoukou EG, Pappas G, Peppas G et al (2008) Vaccination of asplenic or hyposplenic adults. *Br.J.Surg.* 95:273-280.
14. (1996) Guidelines for the prevention and treatment of infection in patients with an absent or dysfunctional spleen. Working Party of the British Committee for Standards in Haematology Clinical Haematology Task Force. *BMJ.* 312:430-434.

- 1  
2  
3  
4 15. Davies JM, Barnes R, Milligan D (2002) Update of guidelines for the prevention and  
5 treatment of infection in patients with an absent or dysfunctional spleen. *Clin Med*  
6 2:440-3.  
7  
8
- 9 16. Fuentes-Ferrer ME, Cano-Escudero S, Mato-Chain G et al (2008) [Vaccination  
10 coverage against *Streptococcus pneumoniae* in splenectomized patients in a fourth-  
11 level hospital (1999-2004)]. *Enferm.Infecc.Microbiol.Clin.* 26:194-198.  
12  
13
- 14 17. Hasse B, Moll C, Oehy K et al (2005) Anti-infectious prophylaxis after splenectomy:  
15 current practice in an eastern region of Switzerland. *Swiss.Med.Wkly.* 135:291-296.  
16  
17
- 18 18. Kyaw MH, Holmes EM, Chalmers J et al (2002) A survey of vaccine coverage and  
19 antibiotic prophylaxis in splenectomised patients in Scotland. *J.Clin.Pathol.* 55:472-  
20 474.  
21
- 22 19. O'Donnell J, McGreal G, Daly P et al (2004) Management of patients undergoing  
23 splenectomy in an Irish teaching hospital: impact of guidelines. *Ir.J.Med.Sci.* 173:136-  
24 140.  
25  
26
- 27 20. Pickering J, Campbell H (2000) An audit of the vaccination and antibiotic  
28 prophylaxis practices amongst patients splenectomised in Lothian. *Health*  
29 *Bull.(Edinb.).* 58:390-395.  
30
- 31 21. Ramachandra J, Bond A, Ranaboldo C et al (2003) An audit of post-splenectomy  
32 prophylaxis--are we following the guidelines? *Ann R Coll Surg Engl* 85:252-5.  
33  
34
- 35 22. Musher DM, Ceasar H, Kojic EM et al (2005) Administration of protein-conjugate  
36 pneumococcal vaccine to patients who have invasive disease after splenectomy despite  
37 their having received 23-valent pneumococcal polysaccharide vaccine. *J.Infect.Dis.*  
38 191:1063-1067.  
39  
40
- 41 23. Cherif H, Landgren O, Konradsen HB et al (2006) Poor antibody response to  
42 pneumococcal polysaccharide vaccination suggests increased susceptibility to  
43 pneumococcal infection in splenectomized patients with hematological diseases.  
44 *Vaccine.* 24:75-81.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Tables**

Table 1. Demographics.

---

<b>Patients [n]</b>	<b>536</b>
<hr/>	
Patients per hospital type [n (hospitals)]	
Academic	40 (2)
Teaching	287 (15)
Non-teaching	209 (11)
Gender [n (%)]	
Male	276 (51,5)
Female	260 (48,5)
Median age at operation [years (range)]	64 (5-91)
Indication for splenectomy [n (%)]	
Acute	115 (21,4)
Elective	196 (36,5)
Iatrogenic lesion	185 (34,5)
Durante operatione	40 (7,4)

---

Table 2. Vaccination results for pneumococcal vaccines given to patients according to operation indication.

<b>Indication for splenectomy</b>	<b>Total</b>	<b>Vaccinated; n (%)</b>
Acute	104	96 (92.3)
Elective	193	184 (95.3)
Iatrogenic	165	119 (72.1)
Durante operatione	37	27 (73)
<b>Total <sup>a</sup></b>	<b>499</b>	<b>426 (85.4)</b>

<sup>a</sup>Total number of patients only includes patients that have survived the first two weeks after splenectomy.

Table 3. Prescribed antibiotics after splenectomy.

Antibiotics	Number of patients	%
Prophylactic <sup>a</sup>	62 /500	12.4
On-demand	41 /500	8.2
Both	15 /500	3.0
N.A. <sup>b</sup>	31 /500	6.2
No antibiotics	351 /500	70.2

<sup>a</sup> Defined as a prescription for prophylactic antibiotics given to the patient at time of discharge.

<sup>b</sup> N.A. (not applicable) refers to patients that died in the hospital before discharge, but at least 2 weeks after splenectomy.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 4. Prescription rates of antibiotics according to operation indication.

<b>Antibiotics (%)</b>	prophylactic	on-demand	both	none
<b>Indication for splenectomy</b>				
Acute	18.3	10.1	2	68.7
Elective	11.4	11	4	73.3
Iatrogenic	9.1	5.6	2.8	81.1
Durante operatione	15.8	5.6	2.8	75



Table 5. Changes in vaccination rates and antibiotic prescription over time.

5A. Vaccination rates before and after 2002.

<b>Vaccination (%)</b>	<b>1997-2001</b>	<b>2002-2008</b>	<b>CI 95%</b>
pneumococcal	88.2	84.8	-0.0422 to 0.111
<i>H. influenzae</i>	15.3	44.3	-0.380 to -0.200
meningococcal C	3.5	38.3	-0.408 to -0.286

CI = confidence interval

5B. Prescription of antibiotics before and after 2002.

<b>Antibiotics (%)</b>	<b>1997-2001</b>	<b>2002-2008</b>	<b>CI 95%</b>
prophylactic	10.6	12.8	-0.0947 to 0.0510
on-demand	3.7	9.8	-0.112 to -0.0113
both	0	3.9	0.0195 to 0.0580
none	85.4	72.6	0.0391 to 0.216

CI = confidence interval

Table 6. Information in discharge correspondence to general practitioner

<b>All correspondence</b>	Number of patients	%
Mention splenectomy	458 / 469	97.7
Mention of vaccination <sup>a</sup>	343 / 426	80.5
Advice revaccination	110 / 469	23.5
Advice annual flu-vaccination	59 / 469	12.6
Advice on-demand antibiotics	100 / 469	21.3

Numbers are minus 36 patients that died within 2 weeks after splenectomy and minus 31 patients that died before being discharged from the hospital.

<sup>a</sup> only of those cases in which pneumococcal vaccination was given

Table 7. Distribution of pathogens in post-splenectomy bacteraemia.

Pathogen	Number of positive bloodcultures
<i>S. pneumoniae</i>	6
<i>H. influenzae</i>	1
<i>S. aureus</i>	6
CNS <sup>a</sup>	16
Enterococci <sup>b</sup>	9
<i>E. coli</i>	7
<i>Pseudomonas</i>	2
<i>Klebsiella</i> <sup>c</sup>	3
<i>Proteus mirabilis</i>	2
ESBL	1
Miscellaneous <sup>d</sup>	8

<sup>a</sup> *S. epidermidis* 3, MRSE 3, CNS 6, undefined 4

<sup>b</sup> *E. faecium* 5, *E. faecalis* 2, sp. 2

<sup>c</sup> *K. pneumoniae* 2, *K. oxytoca*


<sup>d</sup> *Morganella morganii*, *Enterobacter cloacae*, *Bacteriodes fragilis*, *Streptococcus salivarius*, *Bacillus cereus*, *Serratia marcescens*, *Citrobacter freundii*, propionibacterium.

Table 8. Characteristics of patients with documented pneumococcal or *H. influenzae* bacteraemias after splenectomy.

	<i>age</i>	<i>reason for splenectomy</i>	<i>vaccination</i>	<i>antibiotics</i>	<i>PSS</i>
<b>Pt 1</b>	64	Iatrogenic (during colectomy)	23-valent	none	- <i>S. pneumoniae</i> after 5 months
<b>Pt 2</b>	78	Elective (ITP)	23-valent	none	- <i>S. pneumoniae</i> after 6 years - <i>S. pneumoniae</i> after 6.5 years
<b>Pt 3</b>	60	Elective (NHL)	none	none	- <i>S. pneumoniae</i> after 4 days - <i>H. influenzae</i>
<b>Pt 4</b>	89	Elective (AIHA)	none	Prophylactic (Amoxicillin/Clavulanate), 1 week	- <i>S. pneumoniae</i> after 2 weeks - <i>S. pneumoniae</i>

ITP; idiopathic thrombocytopenic purpura, NHL; non-Hodgkin lymphoma, AIHA; auto-immune hemolytic anemia.

Figure 1. Duration of prescribed prophylactic antibiotic therapy.



w = week, m = months, y= years.