



Zonal shear and super-rotation in a magnetized spherical Couette-flow experiment

Daniel Brito, Thierry Alboussiere, Philippe Cardin, Nadège Gagnière, Dominique Jault, Patrick La Rizza, Jean-Paul Masson, Henri-Claude Nataf, Denys Schmitt

► To cite this version:

Daniel Brito, Thierry Alboussiere, Philippe Cardin, Nadège Gagnière, Dominique Jault, et al.. Zonal shear and super-rotation in a magnetized spherical Couette-flow experiment. *Physical Review E : Statistical, Nonlinear, and Soft Matter Physics* [2001-2015], 2011, 83 (6), pp.066310. <10.1103/PhysRevE.83.066310>. <hal-00569377v2>

HAL Id: hal-00569377

<https://hal.science/hal-00569377v2>

Submitted on 16 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Zonal shear and super-rotation in a magnetized spherical Couette**
2 **flow experiment**

3 D. Brito

4 *Laboratoire des Fluides Complexes et leurs Réservoirs,*
5 *Université de Pau et des Pays de l'Adour,*
6 *CNRS, BP 1155, 64013 Pau Cedex France**

7 T. Alboussière,[†] P. Cardin, N. Gagnière, D. Jault, P.

8 La Rizza, J.-P. Masson, H.-C. Nataf, and D. Schmitt

9 *Institut des Sciences de la Terre, CNRS,*
10 *Observatoire de Grenoble, Université Joseph-Fourier,*
11 *Maison des Géosciences, BP 53, 38041 Grenoble Cedex 9, France*

12 (Dated: May 23, 2011)

Abstract

We present measurements performed in a spherical shell filled with liquid sodium, where a 74 mm-radius inner sphere is rotated while a 210 mm-radius outer sphere is at rest. The inner sphere holds a dipolar magnetic field and acts as a magnetic propeller when rotated. In this experimental set-up called *DTS*, direct measurements of the velocity are performed by ultrasonic Doppler velocimetry. Differences in electric potential and the induced magnetic field are also measured to characterize the magnetohydrodynamic flow. Rotation frequencies of the inner sphere are varied between -30 Hz and +30 Hz, the magnetic Reynolds number based on measured sodium velocities and on the shell radius reaching to about 33. We have investigated the mean axisymmetric part of the flow, which consists of differential rotation. Strong super-rotation of the fluid with respect to the rotating inner sphere is directly measured. It is found that the organization of the mean flow does not change much throughout the entire range of parameters covered by our experiment. The direct measurements of zonal velocity give a nice illustration of Ferraro's law of isorotation in the vicinity of the inner sphere where magnetic forces dominate inertial ones. The transition from a Ferraro regime in the interior to a geostrophic regime, where inertial forces predominate, in the outer regions has been well documented. It takes place where the local Elsasser number is about 1. A quantitative agreement with non-linear numerical simulations is obtained when keeping the same Elsasser number. The experiments also reveal a region that violates Ferraro's law just above the inner sphere.

* formerly member of the geodynamo team (Institut des Sciences de la Terre); daniel.brito@univ-pau.fr

† now at : Laboratoire de Sciences de la Terre, Universit de Lyon, ENS de Lyon, CNRS, Lyon, France

13 I. INTRODUCTION

14 The Earth's fluid core below the solid mantle consists of a 3480 km-radius spherical cavity
15 filled with a liquid iron alloy. A 1220 km-radius solid inner core sits in its center. It has
16 been accepted since the 1940's [1, 2] that the flows stirring the electrically conducting liquid
17 iron in the outer core produce the Earth's magnetic field by dynamo action. The fluid
18 motion is thought to originate from the cooling of the Earth's core, which results both in
19 crystallization of the inner core and in convection in the liquid outer core [3].

20 The last decade has seen enormous progress in the numerical computation of the geo-
21 dynamo problem after the first simulation of a dynamo powered by convection [4–7]. It
22 is however still unclear why many characteristics of the Earth's magnetic field are so-well
23 retrieved with simulations [8] since the latter are performed with values of important di-
24 mensionless parameters that differ much from the appropriate values for the Earth's core.
25 The main numerical difficulty is the simultaneous computation of the velocity, the magnetic
26 and the temperature fields with realistic diffusivities, respectively the fluid viscosity, the
27 magnetic and the thermal diffusivities. Those differ indeed by six orders of magnitude in
28 the outer core [9]; such a wide range is at present out of reach numerically, the simulations
29 being performed at best with two orders of magnitude difference between the values of the
30 diffusivities. An experimental approach of the geodynamo is, in that respect, promising since
31 the fluid metals used in experiments have physical properties, specifically diffusivities, very
32 close to the properties of the liquid iron alloy in the Earth's outer core. Moreover, experi-
33 ments and simulations are complementary since they span different ranges of dimensionless
34 parameters.

35 Magnetohydrodynamics experiments devoted to the dynamo study have started some 50
36 years ago (see the chapter authored by Cardin and Brito in [10] for a review). To possibly
37 induce magnetic fields, the working fluid must be liquid sodium in such experiments. Sodium
38 is indeed the fluid that best conducts electricity in laboratory conditions. A breakthrough
39 in these dynamo experiments occurred at the end of 1999 when amplification and saturation
40 of an imposed magnetic field were measured for the first time in two experiments, in Riga
41 [11] and in Karlsruhe [12]. The common property of those set-ups was to have the sodium
42 motion very much constrained spatially, in order to closely follow fluid flows well known
43 analytically to lead to a kinematic dynamo, respectively the Ponomarenko flow [13] and the

44 G.O. Roberts flow [14]. More recently, the first experimental dynamo in a fully turbulent
45 flow was obtained in a configuration where two crenelated ferromagnetic rotating discs drive
46 a von Kàrmàn swirling flow in a cylinder [15]. Earth's like magnetic field reversals were
47 also obtained in this experimental dynamo [16]. Other similar experiments have been run
48 where sodium flows are driven by propellers in a spherical geometry [17, 18]. In order to
49 emphasize the specificity of the experimental study presented in the present paper, it is
50 worth mentioning two common features of the previously mentioned sodium experiments:
51 the forcing of the sodium motion is always purely mechanical and the magnetic field is *weak*
52 in the sense that Lorentz forces are small compared to the non-linear velocity terms in the
53 equation of motion [19].

54 The experiment called *DTS* for "Derviche Tourneur Sodium" has been designed to in-
55 vestigate a supposedly relevant regime for the Earth's core, the magnetostrophic regime
56 [20–22] where the ratio of Coriolis to Lorentz forces is of the order one. The container made
57 of weakly conducting stainless steel is spherical and can rotate about a vertical axis. An
58 inner sphere consisting of a copper envelope enclosing permanent magnets is placed at the
59 center of the outer sphere; the force free magnetic field produced by those magnets enables
60 to explore dynamical regimes where Coriolis and Lorentz forces are comparable. The sodium
61 motion in the spherical gap is driven by the differential rotation between the inner sphere
62 and the outer sphere, unlike in the Earth's core where the iron motion is predominantly
63 driven by convection [23] and maybe minorly by differential rotation of the inner core [24].

64 The *DTS* experiment has not been designed to run in a dynamo regime. It has instead
65 been conceived as a small prototype of a possible future large sodium spherical dynamo
66 experiment which would benefit from its results. Note that meanwhile Daniel Lathrop and
67 collaborators have built a 3m-diameter sodium spherical experiment with an inner sphere
68 differentially rotating with respect to the outer sphere, like in *DTS*. Schaeffer, Cardin and
69 Guervilly [25, 26] have shown numerically that a dynamo could occur in a spherical Couette
70 flow at large Rm in a low magnetic Prandtl number fluid such as sodium ($Pm = \nu/\lambda$, see
71 TABLE I).

72 Numerical simulations in a *DTS*-type configuration [27–29] of Couette spherical flows
73 with an imposed magnetic field all show azimuthal flows stabilized by magnetic and rotation
74 forces. Using electric potential measurements along a meridian of the outer sphere boundary,
75 we concluded in our first report of *DTS* experimental results [30] that the amplitude of the

76 azimuthal flow may exceed the velocity of sodium in solid body rotation with the inner
77 sphere, as predicted theoretically in the linear regime [31].

78 The *DTS* experiment offers a tool to investigate non uniform rotation of an electrically
79 conducting fluid in the presence of rotation and magnetic forces. The differential rotation
80 of a body permeated by a strong magnetic field and the waves driven by the non uniform
81 rotation have received considerable attention since the work of Ferraro [32, 33]. Indeed,
82 the absence of solid envelopes makes non uniform rotation possible in stars, where it plays
83 an important role in the mixing of chemical elements [34], in contrast with the case of
84 planetary fluid cores. Ferraro found that the angular rotation in an electrically conducting
85 body permeated by a steady magnetic field symmetric about the axis of rotation tends to
86 be constant along magnetic lines of force. MacGregor and Charbonneau [35] illustrated this
87 result and showed, in a weakly rotating case, that Ferraro’s theorem holds for $Ha \gg 1$ (Ha ,
88 the Hartmann number, measures the magnetic strength, see TABLE II). An intense magnetic
89 field, probably of primordial origin, is the key actor in the transfer of angular momentum
90 from the solar radiative interior to the convection zone [36, 37]. Finally, in a geophysical
91 context, Aubert recently found, investigating zonal flows in spherical shell dynamos, that
92 Ferraro’s law of isorotation gives a good description of the geometry of the zonal flows of
93 thermal origin [38].

94 In the second study of the *DTS* experiment [39], we investigated azimuthal flows when
95 both the inner boundary and the outer boundary are rotating but at different speeds, using
96 Doppler velocimetry and electric potential measurements. Specifically, we discussed the
97 transition between the outer geostrophic region and the inner region where magnetic forces
98 dominate. Extending the asymptotic model of Kleorin et al. [40], we could explain the
99 shape of the measured azimuthal velocity profiles. We had to use a specific electric potential
100 difference as a proxy of the differential rotation between the two spheres as, unfortunately,
101 the electrical coupling between the liquid sodium and the copper casing of the interior
102 magnets was apparently both imperfect and unreliable. Finally, we reported in on our third
103 article [41] about the *DTS* experiment the presence of azimuthally traveling hydromagnetic
104 waves that we inferred mainly from electric potential measurements along parallels.

105 We investigate here again the main flows when the outer sphere is at rest. Our new study
106 benefits from a comparison with our earlier work [39] for a rotating outer sphere. There is no
107 need any more to use an indirect measure of the global rotation of the fluid as the electrical

108 coupling between liquid sodium and copper has become unimpaired. Furthermore, the *DTS*
 109 experiment has been equipped with a host of new measurement tools. The flow amplitude
 110 is measured along 7 different beams using Doppler velocimetry. Assuming axisymmetry, we
 111 have thus been able to map the azimuthal flow in most of the fluid. It turns out that the
 112 electric potential differences evolve monotonically with the inner core rotation but cannot
 113 be interpreted directly as a measure of the velocity below the outer viscous boundary layer.
 114 We have also entered a probe inside the cavity to measure the induced magnetic field in
 115 the interior. The dense measurements in the *DTS* experiment give a nice illustration of
 116 the Ferraro law of isorotation [32] in the inner region where magnetic forces dominate. In the
 117 outer region, we retrieve axially invariant azimuthal flow as the Proudman-Taylor theorem
 118 holds there, even though the outer sphere is at rest. The variation of the geostrophic velocity
 119 with the distance to the axis differs nevertheless from the case of a rotating outer sphere as
 120 recirculation in the outer Ekman layer plays an important role in the latter case.

121 The organisation of the paper is as follows. In section II, we describe the experimental set-
 122 up and the techniques that we use to measure the magnetic, electric and velocity fields; we
 123 illustrate them with a discussion of a typical experimental run. In section III, we present the
 124 governing equations and the relevant dimensionless numbers of the experiment. We devote
 125 one section of the article to the observation of differential rotation and another one to the
 126 meridional circulation. Then, the experimental measurements are compared to numerical
 127 simulations of *DTS*. We summarize and discuss the results of our study in section VII.

128 II. THE *DTS* EXPERIMENT

129 A. The experimental set-up

130 The *DTS* experimental set-up [30, 39, 41] is shown in FIG. 1. It has been installed in a
131 small building purpose-designed for sodium experiments.




FIG. 1. (Color online) Diagram and picture of the experimental set-up. A: moveable sodium reservoir, B: shielded electric slip-ring, C: electromagnetic valve, D: outer sphere, E: magnetized rotating inner sphere, F: spherical shell containing liquid sodium, G: magnetic coupling entraining the inner sphere shaft, H: crenelated belt, I: brushless electric motor driving the inner sphere, J: expansion tank for sodium, K: thermostated chamber. The total height of the set-up is 3.9 m.

132 As shown in FIG. 1, liquid sodium is contained in a spherical shell between an outer

133 sphere and an inner sphere. The radius of the outer sphere is $a = 210$ mm and that of the
 134 inner sphere $b = 74$ mm. The outer sphere is made of stainless steel and is 5 mm thick.
 135 The copper inner sphere (FIG. 2 and FIG. 3) contains magnetized Rare-Earth cobalt bricks
 136 assembled such that the resulting permanent magnetic field is very close to an axial dipole
 137 of moment intensity 700 Am^2 , with its axis of symmetry aligned with the axis of rotation.
 138 The magnetic field points upward along the rotation axis and its magnitude ranges from 345
 139 mT at the poles of the inner sphere down to 8 mT at the equator of the outer sphere.

140 Sodium is kept most of the time in the reservoir at the bottom of the set-up. When
 141 needed to run an experiment, sodium is melted and pushed up from that reservoir into the
 142 spherical shell by imposing an overpressure of Argon in the reservoir. When liquid sodium
 143 reaches the expansion tank at the top of the spherical shell, an electromagnetic valve located
 144 just below the sphere (see FIG. 1) is locked such that sodium is kept in the upper part during
 145 experiments. In case of emergency, the valve is opened and sodium pours directly into the
 146 reservoir.




FIG. 2. (Color online) (a) Picture of one hemisphere of the inner sphere. Different pieces of magnets in gray are assembled in the bulk of the inner sphere. (b) View from the side of the inner sphere and its rotating shaft. Note that the wheels at the top and bottom (only one is shown in the picture) of the rotating shaft are attached to the outer sphere.

147 The central part of the experiment is air-conditioned in a chamber maintained at around
 148 130°C during experiments: four 1 kW infrared radiants disposed around the outer sphere
 149 heat the chamber, whereas cold air pumped from outside cools the set-up when necessary.
 150 Liquid sodium is therefore usually kept some 30°C above its melting temperature during
 151 experiments. Some physical properties of sodium relevant to our study are listed in TABLE I.

TABLE I. Physical properties of pure liquid sodium at 130°C (Documents from CEA, Commissariat à l’Energie Atomique et aux énergies alternatives). *The sound velocity in sodium has been precisely measured in the present study using the UDV apparatus.

ρ	density	$9.3 \cdot 10^2 \text{ kg m}^{-3}$
σ	electric conductivity	$9 \cdot 10^6 \text{ } \Omega^{-1} \text{ m}^{-1}$
ν	kinematic viscosity	$6.5 \cdot 10^{-7} \text{ m}^2 \text{ s}^{-1}$
η	magnetic diffusivity	$8.7 \cdot 10^{-2} \text{ m}^2 \text{ s}^{-1}$
c	sound velocity*	$2.45 \cdot 10^3 \text{ m s}^{-1}$

The whole volume containing sodium, from the reservoir tank up to the expansion tank is kept under Argon pressure at all times in order to limit oxidization of sodium.

The rotation of the inner sphere, between $f = -30 \text{ Hz}$ and $f = 30 \text{ Hz}$, is driven by a crenelated belt attached to a 11 kW brushless motor (SGMH-1ADCA61 from Yaskawa Electric Corporation, Tokyo, Japan). The belt entrains a home-made magnetic coupler located around the inner sphere shaft as seen in FIG. 1. The coupler is composed of an array of magnets located outside the sodium container, another array of magnets inside the container being immersed in liquid sodium. The inner magnets are anchored to the rotating shaft of the inner sphere such that when the belt is rotated outside, the inner sphere is rotated as well. Such a coupler has the advantage of not requiring any rotating seal in liquid sodium. Torque values up to about 70 N·m have been efficiently transmitted through this coupler in the experiment.

B. Measurements

1. Ultrasonic Doppler velocimetry

We use UDV ultrasonic Doppler velocimetry [42] in order to measure liquid sodium velocities in the spherical shell. This non intrusive technique has been intensively used in our group for the last decade, in particular in rotating experiments performed either in water or in liquid metals [43–46]. The technique consists in the emission from a piezoelectric transducer of a succession of bursts of ultrasonic waves that propagate in the fluid. When

171 the wave encounters a particle with a different acoustic impedance, part of the ultrasonic
 172 wave is backscattered towards the transducer. The time elapsed between the emitted and
 173 the reflected waves and the change in that time respectively give the position of the par-
 174 ticle with respect to the transducer and the fluid velocity along the beam direction. Data
 175 processing is internal to the DOP2000 apparatus (<http://www.signal-processing.com>,
 176 Signal Processing company, Lausanne, Switzerland).




FIG. 3. (Color online) (a) 3D perspective view of the outer sphere and its interior. Caps at various latitudes hold ultrasonic velocity probes to perform UDV. The divergent ultrasonic beams emitted from each cap are shown in perspective with different colors (and numbers for the grayscale version). The five superimposed horizontal slices of magnets are assembled in the heart of the inner sphere. Differences in electric potential are measured between points from latitude $+45^\circ$ to latitude -45° , with steps of 10° (holes along a meridian at the right of the Figure). (b) Meridional view of the normalized coordinates $(s/a, z/a)$ covered by the ultrasonic trajectories numbered from 1 to 7. Some of the corresponding rays are plotted in (a) with the same color code (same numbers). The distance d from the outer sphere along the ultrasonic beam is marked by small dots drawn every 20 mm. The dotted lines are field lines of the imposed dipolar magnetic field.

177 The ultrasonic probes are held in circular stainless steel caps attached to the outer sphere,

as shown in FIG. 3(a). There are six locations with interchangeable caps on the outer sphere such that fluid velocities can be measured from any of these different positions. The thickness of the stainless steel wall between the probes and liquid sodium has been very precisely machined to 1.4 mm in order to insure the best transmission of energy from the probe to the fluid [47]. Small sodium oxides and/or gas bubbles are present and backscatter ultrasonic waves as in gallium experiments [46]. We keep the surface of the caps in contact with sodium as smooth and clean as possible to perform UDV measurements.

We use high temperature 4 MHz ultrasonic transducers (TR0405AH from Signal Processing) 10 mm long and 8 or 12 mm in diameter (piezoelectric diameter 5 or 7 mm). The measurements shown throughout the paper were performed with pulse repetition frequency (*prf*) varying from 3 kHz to 12 kHz and with a number of *prf* per profile varying from 8 to 128. A present limitation of this UDV technique is that the maximum measurable velocity obeys the following function $u_{\max} = c^2/4f_e P_{\max}$ where c is the ultrasonic velocity of the medium, f_e is the emitting frequency, and P_{\max} is the maximum measurable depth along the velocity profile. Applying this relationship to the parameters used in *DTS*, $P_{\max} \simeq 200$ mm (approximative length of the first half of the beam in Figure 3) and $f_e = 4$ Mhz, the maximum measurable velocity is of the order 2.2 m/s. In particular cases, it is possible to overcome this limitation by using aliased profiles of velocity [43] as shown later in the paper. The spatial resolution of the velocity profiles is about 1 mm, and the velocity resolution is about 0.5%, or better for the aliased profiles.

We have measured both the radial and oblique components of velocity in the bulk of the spherical shell. The radial measurements were performed from the latitudes $+10^\circ$, -20° and -40° . The oblique measurements were performed from different locations and in different planes, along rays that all deviate from the radial direction by the same angle (24°). Thus, they all have the same length in the fluid cavity. At the point of closest approach, the rays are 11 mm away from the inner sphere. The seven oblique beams used in *DTS* are sketched in FIG. 3(b). The way to retrieve the meridional and azimuthal components of the velocity field along the ultrasonic beam is detailed in the Appendix.

We use UDV measurements to confirm the strong magnetic coupling between the inner rotating sphere and sodium. In a smaller version of *DTS* performed in water, maximum angular velocities (normalized by that of the inner sphere) of the order 0.16 are obtained for a hydrodynamic Reynolds number of 10^5 in the vicinity of the equatorial plane, close

to the rotating inner sphere [26]. For similar Re in *DTS*, sodium is in super-rotation close to the inner rotating sphere and maximum measured velocities are instead around 1.2 (see FIG. 11(b) for example).

2. Magnetic field inside the sphere

The measurement technique described so far does not require probes that protrude inside the sphere. In order to measure the magnetic field inside the sphere, in the liquid, we have installed magnetometers inside a sleeve, which enters deep into the liquid. The external dimensions of the sleeve are 114 mm (length inside the sphere) and 16 mm (diameter). It contains a board equipped with high-temperature Hall magnetometers (model A1384LUA-T of Allegro Microsystems Inc). We measure the radial component of the magnetic field at radii (normalized by a the inner radius of the outer sphere) 0.93 and 0.74. The orthoradial component is measured at 0.97 and 0.78, and the azimuthal component at 0.99, 0.89, 0.79, 0.69, 0.60 and 0.50. The sleeve is mounted in place of a removable port (at a latitude of either 40° , 10° or -20°). A top view of the sleeve is shown in FIG. 6. The measured voltage is sampled at 2000 samples/second with a 16-bit 250 kHz PXI-6229 National Instruments acquisition card. The precision of the measurements (estimated from actual measurements when $f = 0$) is about $140 \mu\text{T}$, and corresponds to about 20 unit bits of the A/D converter. Magnetic fields up to 60 mT have been measured.

3. Differences in electric potentials on the outer sphere

Differences in electric potentials are measured along several meridians and along one parallel of the outer sphere [30, 39, 41]. In the present study, we are interested in the measurements performed along meridians since they are linked to the azimuthal flow velocity u_φ (we denote (r, θ, φ) the spherical coordinates). The measurements are performed between successive electrodes located from -45° to $+45^\circ$ in latitude, with electrodes 10° apart as sketched in FIG. 3(a). We note $\Delta V_{40} = V_{45} - V_{35}$ the difference between the electric potential at latitudes 45° and 35° . Electric potentials are measured by electrodes soldered to brass bolts 3 mm long, those being screwed into 1 mm-diameter, 4 mm-deep blind holes drilled in the stainless steel wall of the outer sphere. The measured voltage is filtered by an RC anti-

aliasing 215 Hz low-pass filter and then sampled at 1000 samples/second with a 16-bit 250 kHz PXI-6229 National Instruments acquisition card. The precision of the measurements (estimated from actual measurements at $f = 0$) is about $80 \mu\text{V}$, and corresponds to about 10 unit bits of the A/D converter. Electric potential differences up to 7 mV have been measured.

Denoting \mathbf{E} the electric field, we introduce the electric potential V through $\mathbf{E} = -\nabla V$, which is valid in a steady state. Then, the electric potential measurements are analysed using Ohm's law for a moving conductor, $\mathbf{j} = \sigma (\mathbf{u} \times \mathbf{B} + \mathbf{E})$ where σ is the electric conductivity, \mathbf{j} the electric current density vector, \mathbf{u} the velocity field and \mathbf{B} the magnetic field. If the meridional electric currents j_θ are small compared to $\sigma u_\varphi B_r$ in the fluid interior and away from the equatorial plane where $B_r = 0$, and if the viscous boundary layer adjacent to the outer sphere is thin, which ensures the continuity of E_θ through the layer, then the measured differences in electric potential depend on the product of the local radial magnetic field B_r by u_φ , the azimuthal fluid velocity:

$$\frac{\Delta V}{a \Delta \theta} = u_\varphi B_r, \quad (1)$$

where $\Delta \theta = 10^\circ$ is the angle between two electrodes. However, we shall question below the assumption on the smallness of j_θ , referred to as the frozen flux hypothesis.

4. Velocity and torque measured from the motor driving the inner sphere

The electronic drive of the motor entraining the inner sphere delivers an analog signal for its angular velocity and its torque. We checked and improved the velocity measurement by calibrating it using a rotation counter, which consists of a small magnet glued on the entrainment pellet and passing once per turn in front of a magnetometer. The torque signal is used to infer the power consumption in section II D.

C. A typical experiment : a complete set of measurements

A complete set of measurements performed during a typical experiment is analyzed below. The run was chosen to illustrate the various measurements but also to depict how the different observables evolve with f . During that run of 600 seconds, the inner sphere was

264 first accelerated from 0 to $f = 30$ Hz in around 120 seconds, then decelerated back to 0 during
 265 120 seconds. The inner sphere was then kept at rest for about 100 seconds and accelerated
 266 in the opposite direction to $f = -30$ Hz in 120 seconds. It returned to zero rotation in 120
 267 seconds again. That cycle of rotation is shown in FIG. 4. The torque delivered by the inner
 268 sphere motor is also shown and evolves clearly non-linearly during those cycles.




FIG. 4. (Color online) Records of the inner core rotation frequency f , torque C and differences in electric potential ΔV_{40} , ΔV_{30} , ΔV_{20} , ΔV_{10} , ΔV_{-20} , ΔV_{-30} , ΔV_{-40} as a function of time. The subscript denotes the latitude (in degrees) of the electric potential difference.

269 FIG. 4 shows electric potential records (see part II B 3) obtained during this experiment
 270 and time averaged over 0.1 s windows. The differences of potential vary in sync with the
 271 inner sphere rotation frequency as expected if the various ΔV 's measure the differential
 272 rotation between the liquid sodium and the outer sphere to which the electrodes are affixed
 273 (II B 3). However, it is also apparent that the fluid rotation as measured from the ΔV 's does
 274 not increase linearly with the inner sphere frequency. We interpret it as an indication that
 275 braking at the outer boundary, which opposes the entrainment by the inner core rotation,

varies non linearly with the differential rotation. As expected, records from electrodes pairs are anti-symmetrical with respect to the equator, since the forcing is symmetrical while the radial component of the imposed magnetic field changes sign across the equator.

FIG. 5 shows the fluid velocity $u(d)$ measured by UDV during the first half of the experiment along the ray 6 as a function of time and distance. Velocity profiles were recorded along a total distance $d \simeq 80$ mm. As demonstrated in FIG. 5(b), the velocity is aliased since the maximum measurable velocity, for the ultrasonic frequency used during the experiment, is exceeded. Since the azimuthal velocity profiles are quite simple in shape, it has been straightforward to unfold those profiles and retrieve the correct amplitudes as shown in FIG. 5(c). The evolution with f is similar to that of the electrodes, but indicates a stronger leveling-off as f increases.

FIG. 6 shows the magnetic field induced inside the fluid during the typical experiment. The measurements are taken in the sleeve placed at 40° latitude. The induced azimuthal field in FIG. 6 (a) is measured at 6 different radii (given in section II B 2). Its intensity reaches 60 mT near the inner sphere and gets larger than the imposed dipole in some locations. Note the simple evolution with f , which contrasts with that of the electric potentials and velocities in that it increases with an exponent close to 1. The induced meridional field (FIG. 6) is some 20 times weaker. It is dominated by fluctuations, and does not change sign when f does. Note that the evolution with f is not monotonic. Similar behaviors are observed at latitudes 10° and -20° .




FIG. 5. (Color online) UDV measurements performed along the ray number 6 (see FIG. 3) during the second half of the typical experiment when the inner sphere was rotated from rest to -30 Hz and then back to rest. (a) Spatio-temporal representation of the measured velocity, given by the color scale (in m/s). (b) Velocity at three distances from the probe as a function of time, extracted from the spatio-temporal shown in (a). The velocity profiles are clearly aliased since the profiles are discontinuous. (c) After applying a median time-filtering window of 0.2 s and unfolding the profiles, the correct velocities are retrieved as a continuous function of time.




FIG. 6. (Color online) (a) Azimuthal b_ϕ , (b) radial b_r and orthoradial b_θ induced magnetic field at a latitude of 40° in the sleeve at different radial positions recorded during the two triangles sequence of FIG. 4. A top view of the sleeve at the bottom of (a) gives the radial position and the orientation of the various Hall magnetometers. The intensity of the induced azimuthal field reaches 60 mT near the inner sphere and has the sign of $-f$. The fluctuations reach about 10% of the mean. The meridional components of the induced magnetic field are much weaker and dominated by fluctuations, which have been filtered out here (0.2 Hz low-pass filter).

D. Power scaling

The power dissipated by the flow is shown in FIG. 7 as a function of the rotation frequency f . It is computed from the product $\Gamma \times 2\pi f$, where Γ is the torque retrieved from the motor drive. We subtracted the power measured with an empty shell (dash-dot curve) to eliminate power dissipation in the mechanical set-up. The dissipation in the fluid reaches almost 8 kW for the highest rotation frequency of the inner sphere ($f = \pm 30$ Hz). The small spread of the data dots indicates that power fluctuations are small. The continuous line is the record of power versus f when the inner sphere is ramped from 0 to -30 Hz as in FIG. 4. The corresponding increase in kinetic energy only slightly augments power dissipation.

Power dissipation is found to scale as $f^{2.5}$, which does not differ from the scaling obtained in the laminar numerical study of section VI. There, it is explained as the result of the balance between the magnetic torque on the inner sphere and the viscous torque on the outer sphere, assuming that the fluid angular velocity below the outer viscous boundary layer is of the order of the inner sphere angular velocity. Although the outer boundary layer displays strong fluctuations, the situation is completely different from Taylor-Couette water experiments [48].




FIG. 7. Power dissipated by the flow in *DTS*. The data dots are from measurements of the motor torque for plateaus at given f . The dissipation in the mechanical set-up has been removed. It is obtained by rotating the inner sphere before filling the shell with sodium. It is drawn here upside-down in the lower panel (empty symbols) and can be fit by $\mathcal{P}_{\text{empty}}(\text{W}) = 4 \times |2\pi f| + 0.03 \times (2\pi f)^2$ (dash-dot curve). Dissipation in the flow scales as $f^{2.5}$, and is here compared with f^2 (dotted line) and f^3 (dashed line).

III. GOVERNING EQUATIONS

A spherical shell of inner radius b and outer radius a is immersed in an axisymmetric dipolar magnetic field \mathbf{B}_d :

$$\mathbf{B}_d(r, \theta, \varphi) = B_0 \left(\frac{b}{r} \right)^3 [2 \cos \theta \mathbf{e}_r + \sin \theta \mathbf{e}_\theta],$$

where (r, θ, φ) are spherical coordinates. The outer boundary is kept at rest and the inner sphere rotates with the constant angular velocity $\Omega = 2\pi f$ along the same axis as the dipole field that it carries. We assume that the electrically conducting fluid filling the cavity is homogeneous, incompressible and isothermal. We further assume that the flow inside the cavity is steady.

The inner body consists of a magnetized innermost core enclosed in an electrically conducting spherical solid envelope of finite thickness d_b . We choose b as unit length, $b\Omega$ as unit velocity, $\rho b^2 \Omega^2$ as unit pressure, and $b^2 \Omega B_0 / \eta = \text{Rm} B_0$ as unit of induced magnetic field \mathbf{b} ($\mathbf{B} = \mathbf{B}_d + \text{Rm} \mathbf{b}$). Then, the equations governing the flow \mathbf{u} and the induced magnetic field are:

$$\nabla \cdot \mathbf{u} = 0 \tag{2}$$

$$\nabla \cdot \mathbf{b} = 0 \tag{3}$$

$$(\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \Lambda ((\mathbf{B}_d \cdot \nabla) \mathbf{b} + (\mathbf{b} \cdot \nabla) \mathbf{B}_d) + \text{Re}^{-1} \nabla^2 \mathbf{u} \tag{4}$$

$$\nabla^2 \mathbf{b} = -\nabla \times (\mathbf{u} \times \mathbf{B}), \tag{5}$$

where p is a modified pressure. The notation Λ refers to the Elsasser number, classically used for rotating flows in the presence of a magnetic field. That number Λ compares the magnetic and inertial forces in the vicinity of the magnetized inner sphere. In the shell interior, the two forces are better compared by a "local" Elsasser number: $\Lambda_l = (b/r)^6 \Lambda$ (with $(b/a)^6 \simeq 1.83 \cdot 10^{-3}$). Finally, it is of interest to introduce the Hartmann number Ha that compares the magnetic and viscous forces. We have $\text{Ha} = (\Lambda \text{Re})^{1/2}$. In the shell interior, the number $(b/r)^3 \text{Ha}$ is more appropriate to compare the two forces. Typical values of these dimensionless numbers can be found in TABLE II.

The set of equations (2-5), where the non linear terms are neglected, was the subject of the analytical study of Dormy et al. [31] that described how the differential rotation

TABLE II. Typical values of the dimensionless numbers in the *DTS* experiment, computed for $f = \Omega/2\pi = 25$ Hz.

Re	$b^2\Omega/\nu$	$1.3 \cdot 10^6$
Rm	$b^2\Omega/\eta$	10
Λ	$\sigma B_0^2/\rho\Omega$	1.9
Ha	$(\text{Re}\Lambda)^{1/2}$	$1.6 \cdot 10^3$

between the fluid interior and the outer sphere drives an influx of electrical currents from the mainstream into the outer viscous Hartmann boundary layer. Electrical currents flow along the viscous boundary layer and return to the conducting inner body along a free shear layer located on the magnetic field line tangent to the outer boundary at the equator. As these electrical currents cannot flow exactly parallel to the magnetic field line, they produce a Lorentz force, which sustains "super-rotation" of the fluid. Recent studies have extended the analysis to the case of a finitely conducting outer sphere [49, 50]. On increasing the conductance of the container, Dormy et al. (2010) found that more and more electrical currents leak into the solid boundary and the super-rotation rate gets as large as $O(\text{Ha}^{1/2})$. Though the analytical results have set the stage for the interpretation of the experimental results, the neglected non linear effects are crucial in the *DTS* experiment, even for the smallest rate of rotation of the solid inner body.

Upon reversal of Ω , u_φ and b_φ change into $-u_\varphi$ and $-b_\varphi$ whilst the other components of \mathbf{u} and \mathbf{b} are kept unchanged.

347 IV. DIFFERENTIAL ROTATION

348 A. Transition between the Ferraro and geostrophic regimes

349 In this section, we use the UDV records to delve into the geometry of isorotation surfaces.
 350 The L number associated to each dipolar magnetic field line enters the equation of the
 351 surfaces spanned by dipolar lines of force:

$$r = L \sin^2 \theta . \quad (6)$$

352 Accordingly, L gives the radius of the intersection of the magnetic field line with the
 353 equatorial plane. The notation L refers to the L-value (or L-shell parameter) widely used to
 354 describe motions of low energy particles in the Earth's magnetosphere. FIG. 8 shows that,
 355 for $L \leq 2.7$, the angular velocity measured along rays 2 and 3, which are the most appropriate
 356 to map the azimuthal velocity field, is, to a large extent, a function of L only. Thus, the
 357 angular velocity does not vary along magnetic field lines near the inner sphere, where the
 358 magnetic field is the strongest. We interpret this result as a consequence of Ferraro's theorem
 359 of isorotation. The latter is written:

$$\mathbf{B}_d \cdot \nabla \left(\frac{u_\varphi}{s} \right) = 0. \quad (7)$$

360 It is obtained from the φ component of the induction equation for steady fields, ignoring
 361 magnetic diffusion. Although often invoked in the framework of ideal MHD (where magnetic
 362 diffusion is negligible), Ferraro's law does not require a large Rm [51]. It implies that there
 363 is no induced magnetic field and that, as a consequence, the magnetic force is exactly zero.
 364 More precisely, deviations from this law lead to the induction of a magnetic field, which
 365 produces a magnetic force that tends to oppose this induction process. Writing $\mathbf{u} = \mathbf{u}_0 + \mathbf{u}_1$,
 366 where \mathbf{u}_0 obeys the equation (7), we obtain $b \approx u_1$ from (5). Then, the momentum equation
 367 (4) yields $u_1 \approx (\text{Re}\Lambda)^{-1} u_0 = \text{Ha}^{-1} u_0$ (as numerically verified in [35]) when the inertial term,
 368 on the left hand side, can be neglected. Ferraro's law of isorotation, though, is not the only
 369 way to cancel the magnetic force. In the presence of electric currents parallel to the magnetic
 370 field, the magnetic force remains zero and the equation (7) can be violated [50, 51]. For the
 371 geometry of the *DTS* experiment, it cannot happen along the innermost dipolar field lines
 372 that join the two hemispheres, without touching the outer sphere. Indeed, symmetry with
 373 respect to the equatorial plane E implies that the currents do not cross E .

374 Thus, the observation of a velocity field obeying Ferraro's law is a symptom that magnetic
 375 forces predominate in that region. Note that the fact that the two legs of the profile along
 376 ray 2 show similar velocities even for large L only probes the symmetry of the flow with
 377 respect to the equatorial plane.

379 Now, FIG. 9 shows that for $s \geq 0.6$ the azimuthal velocity is largely a function of s
 380 only. There, the Proudman-Taylor theorem holds and azimuthal flows are geostrophic as
 381 the inertial forces predominate. In contrast with the case of a rotating outer sphere (see
 382 Figure 7 in [39]), there is no region of uniform rotation: zonal velocities are z -independent
 383 but vary with the distance to the z axis.

384 The transition between the Ferraro and geostrophic regimes (FIG. 10) occurs at smaller
 385 distances from the axis as the rotation frequency of the inner core increases (unfortunately,
 386 we cannot get reliable UDV data for larger f). It takes place where the local Elsasser number
 387 Λ_l , which compares the magnetic and inertial forces, is of order 1. It is noteworthy that
 388 the Elsasser number Λ defines the location (cylindrical radius) where $\Lambda_l = 1$. The surface
 389 $\Lambda_l = 1$ separates two regions of the fluid cavity. Inside this surface, the magnetic forces
 390 predominate whether outside it the rotation forces are the most important ones. Finally,
 391 the value of Λ largely defines the geometry of isorotation surfaces.

392 In the geostrophic region, magnetic stress integrated on the geostrophic cylinders remains
 393 strong enough to overcome the viscous friction at the outer boundary and to impart a rapid
 394 rotation to the fluid but becomes weaker than the Reynolds stress (which can be represented
 395 as a Coriolis force). As a result, the fluid angular velocity is still of the order of the angular
 396 velocity of the inner sphere and the velocities are predominantly geostrophic.

397 B. Inversion of velocity profiles

398 Flow velocity is constrained by its projection on the several ultrasonic rays that we
 399 shoot. We invert the Doppler velocity profiles for the large scale mean flow, assuming that
 400 the steady part of the flow is symmetric about the axis of rotation and with respect to the
 401 equatorial plane. A poloidal/toroidal decomposition,

$$\mathbf{u} = u_\varphi \mathbf{e}_\varphi + \nabla \times (u_p \mathbf{e}_\varphi) , \quad (8)$$




FIG. 8. (Color online) Rotation frequency of the fluid sodium over the inner sphere rotation frequency as a function of the magnetic field lines L for four ultrasonic velocity profiles (trajectories 1, 2, 3 and 6, with the same color code as in FIG 3) and four inner sphere rotation frequencies ($f = -1.5, -3, -6$ and -10 Hz). The dashed line is a straight line to help the eye.




FIG. 9. (Color online) Rotation frequency of the fluid sodium normalized by the inner sphere rotation frequency as a function of s , for various ultrasonic velocity profiles and four inner sphere rotation frequencies ($f = -1.5, -3, -6$ and -10 Hz). The colors of the profiles (numbers) follow the conventions laid out in FIG. 3.




FIG. 10. (Color online) Normalized cylindrical radius s/a along the UDV trajectories number 1 (blue square), 2 (red circle) and 3 (black cross) where $f_{fluid} = f$ (*i.e.* $f^* = 1$) as a function of the inner sphere rotation frequency. Pale line : $\Lambda_l = 0.5$, Dark line : $\Lambda_l = 2.5$.

is employed. We first consider the azimuthal velocity u_φ , which is expanded in associated Legendre functions with odd degree and order 1, *i.e.*

$$u_\varphi(r, \theta) = \sum_{l=0}^{l_{max}} u_\varphi^l(r) P_{2l+1}^1(\cos \theta) . \quad (9)$$

The functions $u_\varphi^l(r)$ are decomposed into a sum from $k = 0$ to k_{max} of Chebyshev polynomials of the second kind on the interval $[0, 1]$ mapped onto the interval $[b/a, 1]$, *i.e.* the fluid domain. The azimuthal velocity is not constrained to vanish at the inner and outer boundaries, in order to account for the presence of thin unresolved boundary layers.

Azimuthal velocities are more than 10 times larger than the poloidal (*i.e.* meridional) velocities. Nevertheless, the latter projects onto the ultrasound rays. We take the difference of the profiles acquired for f and $-f$ in order to eliminate this small contribution (the meridional circulation does not change sign while the azimuthal velocity does).

FIG. 11 shows the isovalues of angular frequency f^* inverted for $f = \pm 3$ Hz, with $l_{max} = 3$ and $k_{max} = 7$. A crescent of super-rotation is present near the inner sphere. There, isorotation contours roughly follow magnetic field lines, in agreement with Ferraro's theorem, as anticipated above. At larger cylindrical distance from the inner sphere, the flow becomes geostrophic: the contour lines are vertical. We note that angular velocities just above the north pole of the inner sphere do not comply with Ferraro's law. Instead, velocities decrease to quite low values inside the cylinder tangent to the inner sphere. Such violations have been shown to occur when the electric conductivity of boundaries is high [50, 51]. We speculate that we might be in this situation inside the tangent cylinder because the opening of the sphere at the top and bottom (see FIG. 3) replaces the poorly conducting stainless steel wall by sodium.

FIG. 11 compares the synthetic angular velocity profiles to the observed Doppler velocity profiles along the various rays. Note that super-rotation is clearly visible in the raw profiles. The drop in velocity just above the inner sphere is constrained by profiles 4 (green) and 6 (cyan), but its vertical extent is not.

C. f_{fluid} deduced from differences in electric potential and from UDV

As in the previous study of *DTS* with rotating outer sphere [39], we observe that the amplitudes of the differences in electric potential ΔV 's vary linearly with ΔV_{40} , the pro-




FIG. 11. (Color online) (a) Reconstructed isovalue map of fluid angular frequency f^* (the fluid angular frequency normalized by f) at $f = \pm 3$ Hz in a meridional plane, assuming axisymmetry and symmetry with respect to the equator. Three dipolar field lines (dash-dot white) are superimposed in the angular velocity maps. Super-rotation ($f^* \gtrsim 1$) is clearly visible near the inner sphere, where the Ferraro law of isorotation applies. Contours become vertical further away, where geostrophy dominates. The fluid frequency is higher than 0.4 everywhere except in thin unresolved boundary layers. The color lines are the projection in the upper half (s, z) plane of the ultrasonic rays used in the inversion (see FIG. 3). (b) Comparison between the measured ultrasonic Doppler f^* (shown by their error bars) and the synthetic profiles (solid lines) computed from the angular frequency map of (a) for $f = \pm 3$ Hz. The x -axis gives the distance along the ray (in a units). The corresponding rays are plotted in (a) with the same color code (and indicated with trajectory numbers referring to FIG. 3).

portionality factor increasing from the equator toward the poles due in particular to the increase of B_r in formula (1). We show however in the present study that measuring the electric potential does not yield a reliable indicator of the angular velocity f^* using formula (1). In FIG. 12, we compare the normalized fluid angular velocity f^* retrieved from the ΔV 's, for four different latitudes, to f^* obtained directly by UDV at the nearest measured point, around $d/a = 0.1$. The frequencies f^* obtained from ΔV and from UDV in FIG. 12, would be similar if both measurement techniques were only sensitive to u_φ in the interior below the outer viscous boundary layer. The strong discrepancy between these two sets of frequencies reveals instead that the outer boundary layer in *DTS* cannot simply be reduced to a Hartmann layer, outside of which the meridional currents j_θ can be neglected. We

440 further discuss this point in the numerical part VI.




FIG. 12. (Color online) f^* deduced from the measurements of ΔV using formula (1). Blue line : f^* value obtained with UDV measurements on the trajectory number 1 at the distance $d/a = 0.1$. Dashed red line : f^* value obtained with UDV measurements on the trajectory number 2 for $d/a = 0.1$.

441 V. MERIDIONAL CIRCULATION

442 The meridional circulation is constrained from Doppler velocity profiles of the radial ve-
 443 locity (shot along the radial direction), from profiles shot in a meridional plane, and from the
 444 projection of the meridional velocity on "azimuthal" shots. The latter is obtained by taking
 445 the sum of the profiles acquired for f and $-f$, in order to eliminate the azimuthal contribu-
 446 tion. The same is done for the radial and meridional profiles to remove any contamination
 447 from azimuthal velocities.

448 The poloidal velocity scalar u_P of equation (8) is expanded in associated Legendre func-
 449 tions with even degree and order 1, *i.e.*

$$u_P(r, \theta) = \sum_{l=0}^{l_{max}} u_P^l(r) P_{2l}^1(\cos \theta). \quad (10)$$

450 The radial u_r and orthoradial u_θ components of velocity are then obtained as:

$$u_r(r, \theta) = \sum_{l=0}^{l_{max}} \frac{u_P^l(r)}{r} \frac{1}{\sin \theta} \frac{d}{d\theta} (\sin \theta P_{2l}^1(\cos \theta)). \quad (11)$$

451

$$u_\theta(r, \theta) = - \sum_{l=0}^{l_{max}} \left(\frac{u_P^l(r)}{r} + \frac{du_P^l(r)}{dr} \right) P_{2l}^1(\cos \theta). \quad (12)$$

452 The functions $u_P^l(r)$ are decomposed into a sum of $\sin(k\pi(r - b/a)/(1 - b/a))$ from $k = 0$
 453 to k_{max} . The radial velocity is thus constrained to vanish at the inner and outer (rigid)
 454 boundaries, but the orthoradial velocity is not, in order to account for the presence of thin
 455 unresolved boundary layers. FIG. 13 shows the streamlines of the meridional circulation
 456 inverted for $f = \pm 3$ Hz, with $l_{max} = 4$ and $k_{max} = 8$. The fluid is centrifuged from the
 457 inner sphere in the equatorial plane and moves north in a narrow sheet beneath the outer
 458 boundary. It loops back to the inner sphere in a more diffuse manner. Meridional velocities
 459 are more than ten times weaker than azimuthal velocities.

460 FIG. 14 compares the synthetic radial and meridional profiles to the observed Doppler
 461 velocity profiles along the various rays. Velocities are normalized by $2\pi f a$.

462 Over a decade (from $f = 1.5$ Hz to $= 15$ Hz), radial velocities are consistently centrifugal
 463 at 10° latitude and centripetal at 40° , and are roughly proportional to f . The radial profiles
 464 at 20° are more complex and evolve with f , indicating a non-monotonic evolution of the
 465 meridional circulation, also evidenced by the records of the r and θ components of the
 466 induced magnetic field inside the fluid (see FIG. 6). FIG. 15 compiles the *rms* value of
 467 radial velocity at 20° for various f . Note that the fluctuations are larger than this value,
 468 which is almost 50 times smaller than azimuthal velocities.

469 VI. COMPARISON WITH NUMERICAL SIMULATIONS

470 Two previous numerical studies are particularly relevant to our work. Hollerbach et al.
 471 studied exactly the *DTS* configuration but for values of Λ much larger than its value in the
 472 experiment [29]. They focus their study on the modification of the linear solution by inertial




FIG. 13. (Color online) Reconstructed stream lines of the meridional circulation at $f = \pm 3$ Hz in a meridional plane, assuming axisymmetry and symmetry with respect to the equator. The interval between lines is 1.6×10^{-3} . The fluid is centrifuged away from the inner sphere in the equatorial region and moves up to the pole along the outer boundary. The color lines are the projection in the upper half (s, z) plane of the ultrasonic rays used in the inversion.




FIG. 14. (Color online) Comparison between the measured ultrasonic Doppler velocity profiles (shown by their error bars) and the synthetic profiles (solid lines) computed from the meridional circulation map of FIG. 13 for $f = \pm 3$ Hz. (a) Radial profiles along the radial direction $r1$, $r2$ and $r3$ shown in FIG. 13. (b) "Azimuthal" profiles. The contribution from the azimuthal flow has been removed by taking the sum of profiles acquired for f and $-f$. The x -axis gives the distance along the ray (in a units) and the y -axis is the velocity measured along the ray, adimensionalized by $2\pi fa$. The corresponding rays are plotted in FIG. 13 with the same color code (for the grayscale version, the trajectory numbers in (b) refers to those in FIG. 3).




FIG. 15. Compilation of the *rms* radial velocity amplitude as a function of the absolute value of f . The *rms* velocity is computed from Doppler velocimetry profiles shot at a latitude of 20° , from 3 cm beneath the outer shell down to the inner sphere (to avoid spurious values close to the outer boundary). Radial velocity is roughly proportional to f but there is a large dispersion, as the shape of the profiles changes with f . Note that for $f = 10$ Hz, the tangential velocity on the inner sphere reaches 465 cm/s.

473 effects, stressing that the magnetic field line tangent to the outer sphere at the equator loses
474 its significance in the non linear regime. As a result of the relatively large value of Λ , the
475 inertial effects remain too weak -when the outer sphere is at rest- to make a geostrophic
476 region arise at large distances from the axis. The solutions of Garaud [52] (see the figures
477 7 and 11) for a slightly different problem do show the transition between a Ferraro and a
478 geostrophic regions. In her model, which pertains to the formation of the solar tachocline,
479 a dipolar magnetic field permeates a thick spherical shell as in *DTS*, the rotation of the
480 outer boundary is imposed and the rotation of the inner boundary is a free parameter: a
481 condition of zero torque is imposed on that boundary. Numerical models [29, 39] of the
482 *DTS* experiment when the outer sphere is rotating also clearly show a Ferraro region near

the inner sphere where the magnetic field is strong and a geostrophic region in the vicinity of the equator of the outer sphere. We argue below that all these results obtained for a rotating outer sphere provide us with a useful guide to interpret the numerical solutions when the outer sphere is at rest.

A. The numerical model

The model consists of four nested spherical layers (see FIG. 16). The fluid layer is enclosed between a weakly conducting outer container and a central solid sphere comprised of an inner insulating core and of a strongly conducting outer envelope.




FIG. 16. (Color online) Geometry of the numerical model. The relative conductance of the solid outer shell is $\sigma_b \delta / \sigma a = 1/336$, with σ_b and δ respectively the conductivity and the thickness of the outer sphere. It reproduces the experimental value with σ_b chosen as the conductivity of stainless steel at 140°C . The conductivity ratio between the layers 2 and 3 reproduces the ratio (4.2) between the conductivity of copper and sodium.

The velocity field is decomposed as stated in the definitions (8) and (9). The variables $u_\varphi^l(r)$ and $u_p^l(r)$ are then discretized in radius. Analogous decompositions of variables denoted $b_\varphi^l(r)$ and $b_p^l(r)$ are employed to represent the induced magnetic field. The truncation level l_{max} (see (9)) is 120 and at least 450 unevenly spaced points are used in the radial

direction. Specifically, the density of points strongly increases close to the boundaries in order to resolve the viscous boundary layers.

The equations (4) and (5), modified to include all the non linearities and the time derivatives of \mathbf{u} and \mathbf{b} , are transformed into equations for u_φ^l , u_p^l , b_φ^l and b_p^l . We treat the non linear terms explicitly. To advance from one time step to the next, we use an Adams-Bashforth method. Diffusive terms, however, are treated implicitly. Finally, Laplace's equation in spherical coordinates separates which makes it easy to write the magnetic boundary conditions.

The dimensionless numbers Re and Λ are chosen so that steady solutions exist and are stable, with $Pm \ll 1$ (Pm enters the definition of the unit induced field). We strive to reproduce the experimental values of Λ and R_m . Solutions are obtained after time-stepping the equations until a stationary or periodic state is reached. They have been successfully compared to solutions obtained with another numerical code PARODY, which is not restricted to axisymmetric variables [26, 53].

It is not possible to simulate the Reynolds number of the experiment, which is about 10^6 . For the experimental range of Λ , steady solutions are obtained with $Re \sim 10^3$.

B. Steady axisymmetric solutions

FIG. 17 displays a typical solution for the angular and meridional velocities that illustrates well the experimental results. The fluid rotates faster than the magnetized inner body in its vicinity. There, the angular velocity is constant along magnetic field lines of force. Further away of the inner core, the zonal shear becomes almost geostrophic. In addition to these features that we have retrieved from the experimental results, the numerical solution displays recirculation in the outer boundary layer at high latitude. There, the interior flow largely consists in rigid rotation and the boundary layer has the characteristics of a Bödewadt layer with a region of enhanced angular rotation.

For large enough Re (*e.g.* $(a/b)^2 Re = 10^4$ with $(b/a)^2 Ha = 20$), circular waves are present in the Bödewadt layer, above 60° of latitude. They propagate towards the axis. Similar waves had been reported before in simulations of the flow between a rotating and a stationary disk in the absence of a magnetic field [54]. There, they eventually die out. Thus, the persistence of propagation of circular waves in the boundary layer attached to the




FIG. 17. (Color online) (a) Angular and (b) meridional velocity in a meridional plane for $\text{Re} = 9.5 \cdot 10^2$, $\text{Ha} = 163$, and $\text{Pm} = 10^{-3}$. (c) angular velocity estimated from V , using (1). Two dipolar field lines (white) are superimposed in the angular velocity maps, and the thick black contour line is where the angular velocity is unity.

sphere at rest may be attributed to the presence of a magnetic field. On the other hand, these waves arise for larger Re as Ha is augmented. Their emergence delimits the domain of steady solutions.

We have checked that the thickness of the outer boundary layer in the numerical solution scales as $\Omega^{-1/2}$. Note that it corresponds to 3 mm for $\Omega = 1.5 \text{ s}^{-1}$ and the viscosity of liquid sodium. The fluid rotation is driven by the electromagnetic torque acting at the inner boundary against the viscous torque at the outer boundary. We have found that both the viscous torque on the inner surface and the electromagnetic torque on the outer surface are negligible. Comparing different simulations, we have also checked that the main viscous torque scales as $\sim \Omega^{3/2}$, as expected from the thickness of the Bödewadt layer. Thus, the power required to drive the fluid rotation scales as $\Omega^{5/2}$, as does the experimentally measured power, and torque measurements do not give indications on turbulence (see section IID).

The angular rotation just below the outer viscous layer scaled by the inner core angular rotation decreases with Re in agreement with the experimental results. On the other hand, the angular rotation that would be inferred from the electric potential differences calculated at the outer surface using expression (1) increases with Re . FIG. 17(c) displays the angular velocity as estimated from the electric potential, according to equation (1). It can be compared to FIG. 17(a). The actual shear is well retrieved where the magnetic force predominates, in the region where Ferraro's law of isorotation holds. There, the electric current

density \mathbf{j} is limited by the strength of the magnetic force, which needs to be balanced by another force. That restriction makes it possible to neglect \mathbf{j} in Ohm's law. Then, predictions made from (1) are correct. On the other hand, the actual shear is not well recovered in the geostrophic region where the electric current density is not limited by the strength of the magnetic field. There, the frozen-flux relation (1) can be violated. We thus explain why the electric potential measurements at the surface of the *DTS* experiment do not yield a good prediction of the angular velocity immediately below the outer viscous boundary layer.

Our first discussion [30] of the electric potential measurements was based on a numerical model calculated for the experimental values of Ha and thus for too large values of Λ . As a result, the magnetic force, in the numerical model, was dominant in the entire fluid layer and the frozen-flux relationship (1) was verified, at least away from the equator where $B_r = 0$. However, equation (1), becomes less and less valid as Re is increased and Λ decreased, in agreement with the divergence that has been experimentally observed (see the FIG. 12) between the angular velocity calculated from (1) and the actual velocity.

Incidentally, cranking up the rotation of the magnetized inner sphere stabilizes the fluid circulation, at least within a certain parameter range. We have calculated the time-averaged solution (not shown) for the same parameters as the steady solution illustrated by FIG. 17, but for a lower Re . Both the flow and the induced magnetic field are periodic for this set of parameters. A second meridional roll, which is centripetal in the equatorial plane, turns up in the outer region. There, it creates a disk-shaped region where the rotation is slow and the solution is strikingly different from the almost geostrophic solution (FIG. 17) obtained for a slightly larger value of Re .

C. Comparison between numerical simulations and experimental results

We find that reproducing the Elsasser number Λ , rather than a combination of Λ and Re such as the Hartmann number $\text{Ha} = (\text{Re}\Lambda)^{1/2}$, is the key factor to recover the experimental results. The parameters for the solution displayed in FIG. 17 correspond to $\Lambda = 28$, which is the appropriate value for experiments with $\Omega = 1.5 \text{ s}^{-1}$. With $\text{Pm} = 10^{-3}$, the value of the magnetic Reynolds number is about right. It remains too small for the poloidal field to be much different from the imposed dipole field (again for the parameters of FIG. 17).

FIG. 18 shows that numerical solutions are able to satisfactorily reproduce the ultrasonic measurements of angular velocity, obtained for the same values of Λ , as expected from the similitude of the angular velocity maps 11 and 17. The simulated velocities have weaker amplitude than the measured ones in much of the fluid though. We have checked that increasing Re , whilst keeping Λ constant, favours enhanced corotation between the fluid and the inner core. As our calculations are for much smaller Re than the values realized in the experiment, that result may explain the remaining discrepancy between measured and simulated velocities.




FIG. 18. (Color online) Angular velocity along the ultrasonic rays as a function of the distance from the probe: measured (solid lines, 3 Hz, $\Lambda = 16$) and retrieved from a time-averaged numerical solution (dashed lines, $\text{Re} = 1.5 \cdot 10^3$, $\text{Ha} = 163$, $\text{Pm} = 10^{-3}$, $\Lambda = 18$). The color lines refers to those used to define the ultrasonic beams in the FIG. 3 (the numbers also refers to the ultrasonic beams numbers defined in the FIG. 3). The error bars of the experimental data are shown in FIG. 11.

581 VII. DISCUSSION AND CONCLUSION

582 In the presence of an imposed magnetic field, which favors solid body rotation, the inertial
 583 forces largely reduce to a Coriolis force, even for large Reynolds numbers. Experimental
 584 results can thus be interpreted using a single dimensionless number, the Elsasser number.
 585 In that respect, experimental results obtained with global rotation [39] provide a better
 586 guide to interpreting the present results than the linear situation studied by Dormy et al.
 587 [28, 31]. We estimate that, in *DTS*, the rotation frequency f should be less than 0.1 Hz for
 588 the latter to be approached.

589 Experiments have been conducted with the inner sphere rotating in the range -30 Hz
 590 $\leq f \leq 30$ Hz. We have been able to map extensively the shear in the fluid cavity from
 591 ultrasonic Doppler velocimetry for $|f| \leq 10$ Hz. Our observations provide a very clear
 592 experimental illustration of Ferraro’s law of isorotation, demonstrating the predominance
 593 of magnetic forces near the inner sphere. They also exhibit a strong super-rotation: in the
 594 region where magnetic forces dominate, the fluid angular velocity gets 30% larger than that
 595 of the inner sphere. This contrasts with the results obtained by Dormy et al. [28] when
 596 global rotation is present, which indicate that the phenomenon of super-rotation is hindered
 597 by the Coriolis force. The experimental results obtained in our previous study with global
 598 rotation [39] could not address this issue and we plan to run additional experiments for that
 599 purpose.

600 The experiments also display a clear violation to Ferraro’s law: quite low angular velocities
 601 are observed just above the inner sphere, where the magnetic field is strongest (see FIG. 11).
 602 We suspect that this is due to the presence of sodium at rest at the top and bottom of the
 603 cylinder tangent to the inner sphere. Indeed, such violations have been shown to occur when
 604 the electric conductivity of boundaries is high [50, 51].

605 We could follow the evolution of induced magnetic field, electric potentials and power
 606 across the full range of forcing. In a first approximation, all observables associated with the
 607 azimuthal flow (which dominates) can be described by a universal solution, both velocities
 608 and induced magnetic field scaling with f . In a second approximation, the increase of the
 609 dimensional fluid velocity with f thins the viscous boundary layer at the outer sphere and
 610 increases friction accordingly, thus reducing the adimensional velocity of the fluid inside
 611 the sphere. At the same time, the effective Coriolis force that results from the non-linear

612 $(\mathbf{u} \cdot \nabla)\mathbf{u}$ term increases with respect to the (linear) Lorentz force: the geostrophic region
 613 extends further towards the inner sphere. This explains that the fluid velocity increases with
 614 f less rapidly than f (FIG. 5) at large f whilst the torque instead increases more rapidly
 615 than f (FIG. 4) (the electric potentials follow an intermediate trend). The outer friction
 616 torque is balanced by the magnetic torque at the inner boundary. This is consistent with
 617 an increase of the induced magnetic field, near the solid inner body, that is steeper than f
 618 (see FIG. 6). On the other hand, the description of Nataf and Gagnière [55] pertains to the
 619 region where the shear is geostrophic. There, the increased torque at the outer boundary is
 620 balanced by the magnetic torque on the geostrophic cylinders in the interior, which results
 621 from the shearing of the imposed dipolar field. The direct measurement of the velocity (up to
 622 10 Hz, see FIG. 9) shows that the adimensionalized shear does not change significantly with
 623 f even though the velocity itself decreases. In addition, the induced azimuthal magnetic
 624 field that we measure inside the sphere (FIG. 6), for the whole range of f , increases more
 625 rapidly than f . At large f , we observe that b_φ gets larger than the imposed dipolar field in
 626 much of the fluid layer. Eventually, this induced field is large enough to modify the overall
 627 magnetic field, and the resulting flow.

628 This last regime, only achieved because the magnetic Reynolds number is large enough,
 629 is probably the most interesting one. Unfortunately, we cannot directly measure the flow
 630 velocities with the ultrasound technique at these very large f . Less direct techniques are
 631 now required to investigate the zonal shear for $f > 10$ Hz. Inertial waves modified in the
 632 presence of the dipolar and the induced magnetic fields have been inferred from records of
 633 the electric potential along parallels at the surface [41] and of the magnetic field along a
 634 meridian. Both their period and their wavenumber vary with the geometry of the differential
 635 rotation in the cavity. Hopefully, it will be possible to invert the zonal shear from the records
 636 of magneto-inertial waves.

637 Guided by the numerical model, we find that electric field measurements are difficult to
 638 interpret, particularly in the equatorial region where the radial magnetic field B_r vanishes.
 639 The frozen-flux approximation (1) holds when there is a mechanism that keeps under control
 640 the strength of the electrical currents [56]. This is the reason why the magnetic Reynolds
 641 number Rm is not relevant to discuss the validity of the frozen-flux approximation in our
 642 quasi-steady experiment. That approximation has predictive power, instead, in regions
 643 where the magnetic force is dominating. In the *DTS* experiment, it corresponds to the

inner region close to the magnet where $\Lambda \geq 1$.

In a geophysical context, a similar approach is routinely used [57] to invert the velocity field at the Earth’s core surface from models of the time changes of the geomagnetic field, the so-called secular variation. Taking the example of a quasi steady state, this geophysical application has been criticized from a strictly kinematic standpoint [58]. We reckon instead that it is necessary to consider the balance of forces to decide whether the frozen-flux hypothesis holds, at least for a quasi steady state as illustrated by the *DTS* experiment.

Features of the experiment that only depend upon dimensionless numbers that do not involve diffusivities have been simulated numerically. An analogous explanation has been put forward to explain the intriguing successes of geodynamo simulations [5].

ACKNOWLEDGMENTS

The *DTS* project has been supported by Fonds National de la Science, Agence Nationale de la Recherche (Research program VS-QG, grant number BLAN06-2.155316), Institut National des Sciences de l’Univers, Centre National de la Recherche Scientifique, and Université Joseph-Fourier. We are thankful to Dominique Grand and his colleagues from *SERAS* who conducted the design study of the mechanical set-up. The magnetic coupler was computed by Christian Chillet. We thank two anonymous referees for useful comments.

661 Appendix: Angular and meridional velocity along the ultrasonic oblique rays

662 The seven oblique ultrasonic rays shot in *DTS* are sketched in FIG. 3. We define the
 663 declination D as the angle between the beam and the meridional plane (D counted positively
 664 eastwards), the inclination I as the angle between the projected beam in the meridional plane
 665 and the radial direction (I counted positively upwards) and λ as the latitude of the ultrasonic
 666 probe. Using those definitions, TABLE III give the characteristics of the beams.

TABLE III. Latitude λ , inclination I , and declination D (in degrees) at the origin of the shots
 (on the outer sphere) of the oblique ultrasonic beams in *DTS*.

Trajectory number and color	λ	I	D
1, blue	40	21.1	11.7
2, red	10	2.2	23.9
3, black	10	12.5	-20.6
4, green	-20	20	-13.5
5, yellow	-20	21.1	-11.7
6, cyan	-40	21.1	11.7
7, magenta	-40	-24	0

667 1. Angular velocity

668 Along these oblique beams, the projection $u(d)$ (d is the distance from the probe) of the
 669 velocity is a combination of the components u_r , u_θ and u_φ of the total velocity field. Velocity
 670 $u(d)$ is counted positive in the shooting direction. We assume that the mean fluid flow is
 671 axisymmetric, and also $(u_r, u_\theta) \ll u_\varphi$, the meridional velocities amplitude in *DTS* being less
 672 than 10% the amplitude of the azimuthal velocities. Using projections along the beam, we
 673 retrieve the angular velocity $\omega(d)$ along trajectories 1 to 6 using the following relationship

$$\omega(d) = -\frac{u(d)}{a \cos \lambda \sin D}. \quad (\text{A.1})$$

674 2. Meridional velocity

675 We have also exploited the observation that the meridional velocity does not change sign
 676 when the rotation of the inner sphere is reversed - it remains centrifugal in the equatorial
 677 plane - whereas the angular velocity does change sign. Thus, combining measurements
 678 obtained with two opposite rotation rates of the inner core, we can separate azimuthal and
 679 meridional velocities.

680 Assuming now that the mean meridional velocity is axisymmetric and using projections,
 681 we can retrieve the radial velocity

$$u_r(d) = \frac{u(d)r(d)}{d - a \cos D \cos I}, \quad (\text{A.2})$$

682 and the orthoradial velocity

$$u_\theta(d) = \frac{u(d)r(d)s(d)}{a[a \cos D \cos \lambda \sin I - d \cos^2 D \cos(\lambda + I) \sin I + d \sin^2 D \sin \lambda]}, \quad (\text{A.3})$$

683 where $r(d) = \sqrt{x^2 + y^2 + z^2}$ is the spherical radius and $s(d) = \sqrt{x^2 + y^2}$ is the cylindrical
 684 radius at the measurement point. They (x, y, z) coordinates of the measurement point are
 685 given by:

$$x(d) = a \cos \lambda - d \cos D \cos(\lambda + I) \quad (\text{A.4})$$

$$y(d) = -d \sin D \quad (\text{A.5})$$

$$z(d) = a \sin \lambda - d \cos D \sin(\lambda + I) \quad (\text{A.6})$$

-
- 686 [1] W. M. Elsasser, Phys. Rev., **69**, 106 (1946).
 687 [2] W. M. Elsasser, Phys. Rev., **70**, 202 (1946).
 688 [3] J. Verhoogen, *Energetics of the Earth* (National Academy of Sciences, Washington, 1980).
 689 [4] G. A. Glatzmaier and P. H. Roberts, Nature, **377**, 203 (1995).
 690 [5] U. Christensen and J. Aubert, Geophys. J. Int., **166**, 97 (2006).
 691 [6] F. Takahashi, M. Matsushima, and Y. Honkura, Phys. Earth Planet. Inter., **167**, 168 (2008).
 692 [7] A. Sakuraba and P. Roberts, Nature Geoscience (2009).
 693 [8] E. Dormy, J.-P. Valet, and V. Courtillot, Geochem. Geophys. Geosyst., **1** (2000).

- [9] J.-P. Poirier, *Introduction to the Physics of the Earth's Interior* (Cambridge University Press, Cambridge, UK, 2000).
- [10] E. Dormy and A. M. Soward, *Mathematical aspects of natural dynamos* (CRC Press/Taylor Francis, 2007).
- [11] A. Gailitis, O. Lielausis, E. Platacis, S. Dement'ev, A. Cifersons, G. Gerbeth, T. Gundrum, F. Stefani, M. Christen, and G. Will, Phys. Rev. Lett., **86**, 3024 (2001).
- [12] R. Stieglitz and U. Müller, Phys. Fluids, **13**, 561 (2001).
- [13] Y. Ponomarenko, J. Appl. Mech. Tech. Phys., **14**, 775 (1973).
- [14] G. O. Roberts, Phil. Trans. R. Soc. London, Ser. A, **271**, 411 (1972).
- [15] M. Berhanu, R. Monchaux, S. Fauve, N. Mordant, F. Pétrélis, A. Chiffaudel, F. Daviaud, B. Dubrulle, L. Marié, F. Ravelet, M. Bourgoin, P. Odier, J.-F. Pinton, and R. Volk, Europhys. Lett., **77**, 59001 (2007).
- [16] R. Monchaux, M. Berhanu, M. Bourgoin, M. Moulin, P. Odier, J.-F. Pinton, R. Volk, S. Fauve, N. Mordant, F. Pétrélis, A. Chiffaudel, F. Daviaud, B. Dubrulle, C. Gasquet, L. Marié, and F. Ravelet, Phys. Rev. Lett., **98**, 044502 (2007).
- [17] D. R. Sisan, N. Mujica, W. A. Tillotson, Y.-M. Huang, W. Dorland, A. B. Hassam, T. M. Antonsen, and D. P. Lathrop, Phys. Rev. Lett., **93**, 114502 (2004).
- [18] M. D. Nornberg, E. J. Spence, R. D. Kendrick, C. M. Jacobson, and C. B. Forest, Phys. Rev. Lett., **97**, 044503 (2006).
- [19] F. Petrelis, N. Mordant, and S. Fauve, Geophys. Astrophys. Fluid Dyn., **101**, 289 (2007).
- [20] J. Taylor, Proc. R. Soc. Lond. A, **274**, 274 (1963).
- [21] P. Cardin, D. Brito, D. Jault, H.-C. Nataf, and J.-P. Masson, Magnetohydrodynamics, **38**, 177 (2002).
- [22] D. Jault, Phys. Earth Planet. Inter., **166**, 67 (2008).
- [23] F. H. Busse, J. Fluid Mech., **44**, 441 (1970).
- [24] X. Song and P. Richards, Nature, **382**, 221 (1996).
- [25] N. Schaeffer and P. Cardin, Earth Planet. Sci. Lett., **245**, 595 (2006).
- [26] C. Guervilly and P. Cardin, Geophys. Astrophys. Fluid Dyn., **104**, 221 (2010).
- [27] R. Hollerbach, Proc. R. Soc. Lond. A, **444**, 333 (1994).
- [28] E. Dormy, P. Cardin, and D. Jault, Earth Planet. Sci. Lett., **160**, 15 (1998).
- [29] R. Hollerbach, E. Canet, and A. Fournier, Eur. J. Mech. B, **26**, 729 (2007).

- [30] H.-C. Nataf, T. Alboussière, D. Brito, P. Cardin, N. Gagnière, D. Jault, J.-P. Masson, and D. Schmitt, *Geophys. Astrophys. Fluid Dyn.*, **100**, 281 (2006).
- [31] E. Dormy, D. Jault, and A. M. Soward, *J. Fluid Mech.*, **452**, 263 (2002).
- [32] V. Ferraro, *Mon. Not. Roy. Astron. Soc.*, **97**, 458 (1937).
- [33] H. Spruit, *Astron. Astrophys.*, **349**, 189 (1999).
- [34] C. Charbonnel and S. Talon, *Science*, **309**, 2189 (2005).
- [35] K. MacGregor and P. Charbonneau, *Astrophys. J.*, **519**, 911 (1999).
- [36] L. Mestel and N. Weiss, *Mon. Not. Roy. Astron. Soc.*, **226**, 123 (1987).
- [37] D. Gough, in *Magnetic Coupling between the Interior and Atmosphere of the Sun*, edited by S. Hasan and R. Rutten (Springer, 2010) pp. 68–85.
- [38] J. Aubert, *J. Fluid Mech.*, **542**, 53 (2005).
- [39] H.-C. Nataf, T. Alboussière, D. Brito, P. Cardin, N. Gagnière, D. Jault, and D. Schmitt, *Phys. Earth Planet. Inter.*, **170**, 60 (2008).
- [40] N. Kleeorin, I. Rogachevskii, A. Ruzmaikin, A. M. Soward, and S. Starchenko, *J. Fluid Mech.*, **344**, 213 (1997).
- [41] D. Schmitt, T. Alboussière, D. Brito, P. Cardin, N. Gagnière, D. Jault, and H.-C. Nataf, *J. Fluid Mech.*, **604**, 175 (2008).
- [42] Y. Takeda, *Nucl. Technol.*, **79**, 120 (1987).
- [43] D. Brito, H.-C. Nataf, P. Cardin, J. Aubert, and J.-P. Masson, *Exp. Fluids*, **31**, 653 (2001).
- [44] J. Noir, D. Brito, K. Aldridge, and P. Cardin, *Geophys. Res. Lett.*, **28**, 3785 (2001).
- [45] J. Aubert, D. Brito, H.-C. Nataf, P. Cardin, and J.-P. Masson, *Phys. Earth Planet. Inter.*, **128**, 51 (2001).
- [46] N. Gillet, D. Brito, D. Jault, and H.-C. Nataf, *J. Fluid Mech.*, **580**, 83 (2007).
- [47] S. Eckert and G. Gerbeth, *Exp. Fluids*, **32**, 542 (2002).
- [48] D. Lathrop, J. Fineberg, and H. Swinney, *Physical review letters*, **68**, 1515 (1992), ISSN 1079-7114.
- [49] K. Mizerski and K. Bajer, *Phys. Earth Planet. Inter.*, **160**, 245 (2007).
- [50] A. Soward and E. Dormy, *J. Fluid Mech.*, **645**, 145 (2010).
- [51] J. Allen, P. Auer, and V. Endean, *Plasma physics*, **18**, 143 (1976).
- [52] P. Garaud, *Mon. Not. Roy. Astron. Soc.*, **329**, 1 (2002).
- [53] J. Aubert, J. Aurnou, and J. Wicht, *Geophys. J. Int.*, **172**, 945 (2008).

- 756 [54] J. Lopez, F. Marques, A. Rubio, and M. Avila, Phys. Fluids, **21**, 114107 (2009).
- 757 [55] H.-C. Nataf and N. Gagnière, C.R. Physique, **9**, 702 (2008).
- 758 [56] D. Jackson, *Classical Electrodynamics* (John Wiley, New York, 1975).
- 759 [57] R. Holme, *Treatise on Geophysics, Vol. 8 Core Dynamics*, edited by P. Olson and G. Schubert
760 (Elsevier, 2007) Chap. 8.04, pp. 107–129.
- 761 [58] J. J. Love, Geophys. J. Int., **138**, 353 (1999).