



HAL
open science

Transitionless quantum drivings for the harmonic oscillator

J G Muga, X Chen, S Ibáñez, I Lizuain, A Ruschhaupt

► **To cite this version:**

J G Muga, X Chen, S Ibáñez, I Lizuain, A Ruschhaupt. Transitionless quantum drivings for the harmonic oscillator. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 2010, 43 (8), pp.85509. 10.1088/0953-4075/43/8/085509 . hal-00569912

HAL Id: hal-00569912

<https://hal.science/hal-00569912>

Submitted on 25 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Transitionless quantum drivings for the harmonic oscillator

J. G. Muga¹, X. Chen^{1,2}, S. Ibáñez¹, I. Lizuain¹, A. Ruschhaupt³

¹ Departamento de Química-Física, UPV-EHU, Apdo 644, 48080 Bilbao, Spain

² Department of Physics, Shanghai University, 200444 Shanghai, P. R. China

³ Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany

Abstract. Two methods to change a quantum harmonic oscillator frequency without transitions in a finite time are described and compared. The first method, a transitionless-tracking algorithm, makes use of a generalized harmonic oscillator and a non-local potential. The second method, based on engineering an invariant of motion, only modifies the harmonic frequency in time, keeping the potential local at all times.

PACS numbers: 37.10.De, 42.50.-p, 37.10.Vz

1. Introduction

Changing the external parameters of the Hamiltonian is a fundamental and standard operation to probe, control, or prepare a quantum system. In many cases it is desirable to go from an initial parameter configuration to a final one without inducing transitions, as in the expansions performed in fountain clocks [1], the release of atoms from optical lattices [2], or compressions to increase the spatial density [2]. In fact most of the current experiments with cold atoms are based on a cooling stage and then an adiabatic drive of the system to some desired final trap or regime [3]. These “transitionless” [4], or “frictionless” [5] adiabatic processes may require exceedingly large times and become impractical [2], even impossible [3], or quite simply a faster process is desirable, e.g. to increase the repetition rate of a cycle, or a signal-to-noise ratio. This motivates the generic objective of achieving the same final state as the slow adiabatic processes, possibly up to phase factors, but in a much shorter time. One may try to fulfill that goal in two different ways: (a) designing appropriate “parameter trajectories” of the Hamiltonian from the initial to the final times, or (b) applying entirely new interactions that modify the Hamiltonian beyond a simple parameter evolution of the original form, for example by adding different terms to it. In this paper we shall analyze and discuss, for the harmonic oscillator, two recently proposed methods, whose relation had not been investigated, which actually implement these two different routes. While most of the treatment is applicable to an “abstract” harmonic oscillator, we shall discuss physical

implementations specific for ultracold atoms or ions. Indeed, harmonic traps and their manipulation are basic working horses of this field.

For the harmonic oscillator the parameter we consider is the trap frequency, which should go from ω_0 to ω_f in a time t_f , preserving the populations of the levels, $P_n(t_f) = P_n(0)$. “ n ” labels the instantaneous n -th eigenstate of the initial and final harmonic oscillator Hamiltonians,

$$\begin{aligned}\hat{H}_0(0)|n(0)\rangle &= \hbar\omega_0(n + 1/2)|n(0)\rangle, \\ \hat{H}_0(t_f)|n(t_f)\rangle &= \hbar\omega_f(n + 1/2)|n(t_f)\rangle.\end{aligned}\tag{1}$$

One of the methods we shall discuss here relies on a general framework set by Kato in a proof of the adiabatic theorem [6], and has been formulated recently by Berry [4]. We shall term it “transitionless-tracking” approach, or TT for short; the other one [7, 8] engineers the Lewis-Riesenfeld invariant [9] by an inverse method [10] to satisfy the desired boundary conditions; we shall call this method “inverse-invariant”, or II for short. In the basic version of TT the dynamics is set to follow at all intermediate times the adiabatic path defined by an auxiliary Hamiltonian $\hat{H}_0(t)$ (in our case a regular harmonic oscillator with frequency $\omega(t)$ and boundary conditions $\omega(0) = \omega_0$ and $\omega_f = \omega(t = t_f)$), and its instantaneous eigenvectors $|n(t)\rangle$, up to phase factors. Instead, in the II approach the auxiliary object is an engineered Lewis-Riesenfeld invariant $I(t)$ set to commute with $\hat{H}_0(0)$ at $t = 0$ and with $\hat{H}_0(t_f)$ at t_f . In both cases intermediate states may be highly non-adiabatic with respect to the instantaneous eigenstates of the Hamiltonians actually applied, $\hat{H}_{TT}(t)$ and $\hat{H}_{II}(t)$.

We shall provide first the equations characterizing the two approaches and then comment on possible physical implementations.

2. Transitionless tracking algorithm

2.1. General formalism

For the general formalism we follow [4] closely. Assume a time-dependent Hamiltonian $\hat{H}_0(t)$ with initial and final values (1), instantaneous eigenvectors $|n(t)\rangle$ and eigenvalues $E_n(t)$,

$$\hat{H}_0(t)|n(t)\rangle = E_n(t)|n(t)\rangle.\tag{2}$$

A slow change would preserve the eigenvalue and eigenvector along the dynamical evolution times a phase factor, and the adiabatic approximation of the wavefunction is

$$|\psi_n(t)\rangle = \exp\left\{-\frac{i}{\hbar}\int_0^t dt' E_n(t') - \int_0^t dt' \langle n(t')|\partial_{t'} n(t')\rangle\right\}|n(t)\rangle.\tag{3}$$

We now seek a Hamiltonian $\hat{H}(t)$ such that the adiabatic approximation $|\psi_n(t)\rangle$ represents the *exact* dynamics,

$$i\hbar\partial_t|\psi_n(t)\rangle = \hat{H}(t)|\psi_n(t)\rangle.\tag{4}$$

$\hat{H}(t)$ (which is \hat{H}_{TT} if distinction with the other method is needed) is related to the corresponding unitary operator by

$$i\hbar\partial_t\hat{U}(t) = \hat{H}(t)\hat{U}(t), \quad (5)$$

$$\hat{H}(t) = i\hbar(\partial_t\hat{U}(t))\hat{U}^\dagger(t). \quad (6)$$

Choosing

$$\hat{U}(t) = \sum_n \exp\left\{-\frac{i}{\hbar}\int_0^t dt' E_n(t') - \int_0^t dt' \langle n(t')|\partial_{t'}n(t')\rangle\right\} |n(t)\rangle\langle n(0)|, \quad (7)$$

we find from (6),

$$\hat{H}(t) = \sum_n |n\rangle E_n \langle n| + i\hbar \sum_n (|\partial_t n\rangle\langle n| - \langle n|\partial_t n\rangle|n\rangle\langle n|) \equiv \hat{H}_0 + \hat{H}_1, \quad (8)$$

where we have simplified the notation, $|n\rangle = |n(t)\rangle$. It is also possible to choose other phases in (3) [4]. The simplest case is $\hat{U}(t) = \sum |n(t)\rangle\langle n(0)|$, without phase factors, which leads to $\hat{H}(t) = i\hbar \sum |\partial_t n\rangle\langle n|$. Note that with this choice $\hat{H}_0(t)$ has been formally suppressed in $\hat{H}(t)$ but still plays a role through its eigenfunctions $|n(t)\rangle$.

2.2. Application to the harmonic oscillator

We now apply the above to the harmonic oscillator

$$\hat{H}_0(t) = \hat{p}^2/2m + \omega(t)^2\hat{x}^2/2m = \hbar\omega(t)(\hat{a}_t^\dagger\hat{a}_t + 1/2), \quad (9)$$

where \hat{a}_t and \hat{a}_t^\dagger are the (Schrödinger picture!) annihilation and creation operators at time t ,

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega(t)}}(\hat{a}_t^\dagger + \hat{a}_t), \quad (10)$$

$$\hat{p} = i\sqrt{\frac{\hbar m\omega(t)}{2}}(\hat{a}_t^\dagger - \hat{a}_t), \quad (11)$$

$$\hat{a}_t = \sqrt{\frac{m\omega(t)}{2\hbar}}\left(\hat{x} + \frac{i}{m\omega(t)}\hat{p}\right), \quad (12)$$

$$\hat{a}_t^\dagger = \sqrt{\frac{m\omega(t)}{2\hbar}}\left(\hat{x} - \frac{i}{m\omega(t)}\hat{p}\right). \quad (13)$$

This time dependence may be misleading and a bit unusual at first so we insist: since the frequency depends on time the “instantaneous” ladder operators \hat{a}_t , \hat{a}_t^\dagger create or annihilate different “instantaneous” states, adapted to the corresponding frequency. Thus, ladder operators with different time labels do not commute in general, although some combinations, e.g. those equivalent to powers of \hat{x} and/or \hat{p} , do commute, as we shall see later.

The instantaneous eigenstates $|n(t)\rangle$ can be written in coordinate representation as

$$\langle x|n(t)\rangle = \frac{1}{\sqrt{2^n n!}} \left(\frac{m\omega(t)}{\pi\hbar}\right)^{1/4} \exp\left(-\frac{1}{2}\frac{m\omega(t)}{\hbar}x^2\right) H_n\left(\sqrt{\frac{m\omega(t)}{\hbar}}x\right), \quad (14)$$

and their derivative with respect to t is

$$\langle x|\partial_t n(t)\rangle = \left(\frac{1}{4} - \frac{m\omega(t)}{2\hbar}x^2\right) \frac{\dot{\omega}}{\omega(t)}|n\rangle + \sqrt{\frac{m\omega(t)}{2\hbar}}x \frac{\dot{\omega}}{\omega(t)}\sqrt{n}|n-1\rangle. \quad (15)$$

We find, using the recursion relation of Hermite polynomials and their orthogonality,

$$\langle k|\partial_t n\rangle = \begin{cases} \frac{1}{4}\sqrt{n(n-1)}\frac{\dot{\omega}}{\omega(t)} & k = n-2 \\ -\frac{1}{4}\sqrt{(n+1)(n+2)}\frac{\dot{\omega}}{\omega(t)} & k = n+2 \\ 0 & (\text{otherwise}) \end{cases}, \quad (16)$$

so that $\hat{H}_1(t)$ can be written as

$$\begin{aligned} \hat{H}_1(t) &= i\hbar \sum_n |\partial_t n\rangle \langle n| \equiv i\hbar \frac{\dot{\omega}}{\omega(t)} \sum_n \left[\left(\frac{1}{4} - \frac{m\omega(t)}{2\hbar}\hat{x}^2\right) |n\rangle \langle n| \right. \\ &\quad \left. + \sqrt{\frac{m\omega(t)}{2\hbar}}\hat{x}\sqrt{n}|n-1\rangle \langle n| \right]. \end{aligned} \quad (17)$$

Using $\hat{a}_t = \sum_n \sqrt{n}|n-1(t)\rangle \langle n(t)|$, and the relations between \hat{x} , \hat{p} , \hat{a}_t and \hat{a}_t^\dagger written above,

$$\begin{aligned} \hat{H}_1(t) &= i\hbar \frac{\dot{\omega}}{\omega(t)} \sum_n \left[\frac{1}{4} - \frac{m\omega(t)}{2\hbar}\hat{x}^2 + \sqrt{\frac{m\omega(t)}{2\hbar}}\hat{x}\hat{a}_t \right] \\ &= \frac{i\hbar}{4} \frac{\dot{\omega}}{\omega(t)} - \frac{1}{2} \frac{\dot{\omega}}{\omega(t)} \hat{x}\hat{p}. \end{aligned} \quad (18)$$

With $[\hat{x}, \hat{p}] = i\hbar$, we finally write the Hamiltonian $\hat{H}_1(t)$ in the following simple forms

$$\hat{H}_1(t) = -\frac{\dot{\omega}}{4\omega}(\hat{x}\hat{p} + \hat{p}\hat{x}) = i\hbar \frac{\dot{\omega}}{4\omega}[\hat{a}^2 - (\hat{a}^\dagger)^2]. \quad (19)$$

In the last expression the subscript t in \hat{a} and \hat{a}^\dagger has been dropped because the squeezing combination $\hat{a}^2 - (\hat{a}^\dagger)^2$ is actually independent of time, so it may be evaluated at any convenient time, e.g. at $t = 0$. The connection with squeezing operators is worked out in the appendix.

\hat{H}_1 is therefore a non-local operator, and does not have the form of a regular harmonic oscillator potential with an \hat{x}^2 term. Nevertheless the final Hamiltonian $\hat{H} = \hat{H}_0 + \hat{H}_1$ is still quadratic in \hat{x} and \hat{p} , so it may be considered a generalized harmonic oscillator [11].

2.3. Physical realization

The nonlocality of \hat{H}_1 , with a constant prefactor, can be realized in a laboratory by means of 2-photon Raman transitions for trapped ions [12, 13, 14] or, as discussed below, for neutral atoms in tight dipole traps. Since we have to evaluate as well the possibility of making the prefactor in \hat{H}_1 time dependent, we need to provide the derivation with some detail, first for a time-independent ω .

2.3.1. Raman two-photon transition in a trapped ion or neutral atom Let us consider a harmonically trapped two-level system in 1D driven by two different lasers (with coupling strengths Ω_j and frequencies ω_j , $j = 1, 2$), see Fig. 1 and Refs. [13, 14]. The time dependent ‘‘Raman’’ Hamiltonian in the Schrödinger picture will be given by

$$\hat{H}_R(t) = \hat{H}_T + \hat{H}_A + \hat{H}_{int}, \quad (20)$$

with ‘‘trap’’ (T), ‘‘atomic’’ (A), and interaction (int) terms

$$\hat{H}_T = \hbar\omega\hat{a}^\dagger\hat{a}, \quad (21)$$

$$\hat{H}_A = \hbar\omega_e|e\rangle\langle e|, \quad (22)$$

$$\hat{H}_{int} = \sum_{j=1}^2 \hbar\Omega_j \cos(\omega_j t - k_j x + \phi_j) (|g\rangle\langle e| + |e\rangle\langle g|), \quad (23)$$

where $\hbar\omega_e$ is the energy of the excited state $|e\rangle$ and $\mathbf{k}_j = k_j\hat{\mathbf{x}}$ the wavevector of each laser which are assumed to be pointing along the principal trap direction, the x -direction. A basic assumption in this Hamiltonian is that the same vibrational frequency, i.e., the same external potential for atomic motion is applicable to the ground and excited states $|g\rangle$ and $|e\rangle$. This is the standard case for ions in Paul traps, whereas for neutral atoms it may be realized in optical traps too, at least for suitably chosen species and transitions (for example alkali-earth atoms and spin forbidden transitions), by means of ‘‘magic wavelength’’ compensating techniques by which the light shifts, and thus the potentials for atomic motion in the $|g\rangle$ and $|e\rangle$ states can be controlled and made identical [15, 16].

2.3.2. Interaction picture Let us now write the above Hamiltonian in an interaction picture defined by the Hamiltonian $\hat{h}_0 = \hat{H}_T + \hbar\tilde{\omega}_L|e\rangle\langle e|$, where $\tilde{\omega}_L = (\omega_1 + \omega_2)/2$ has been introduced. The interaction Hamiltonian $\hat{H}_I = e^{i\hat{h}_0 t/\hbar}(\hat{H}_R - \hat{h}_0)e^{-i\hat{h}_0 t/\hbar}$ reads

$$\begin{aligned} \hat{H}_I(t) = & -\hbar\tilde{\Delta}|e\rangle\langle e| \\ & + \sum_{j=1}^2 \frac{\hbar\Omega_j}{2} \left(e^{i\eta_j[\hat{a}(t)+\hat{a}^\dagger(t)]} e^{-i(\omega_j-\tilde{\omega}_L)t} e^{-i\phi_j}|e\rangle\langle g| + H.c. \right), \end{aligned} \quad (24)$$

where $\tilde{\Delta} = \tilde{\omega}_L - \omega_e$, and now $\hat{a}(t) = \hat{a}e^{-i\omega t}$, $\hat{a}^\dagger(t) = \hat{a}^\dagger e^{i\omega t}$ are the time dependent Heisenberg annihilation and creation operators respectively. Note also that fast oscillating off-resonant $e^{\pm i(\omega_j+\tilde{\omega}_L)t}$ terms have been neglected in the rotating wave approximation (RWA). The parameter $\eta_j = k_j x_0$ is known as the Lamb-Dicke (LD) parameter, where $x_0 = \sqrt{\hbar/2m\omega}$ is the extension (square root of the variance) of the ion’s ground state, i. e., $\hat{x} = x_0(\hat{a} + \hat{a}^\dagger)$.

2.3.3. Adiabatic elimination and effective Hamiltonian For a general wavefunction (in the corresponding interaction picture) such as

$$|\psi_I(t)\rangle = \sum_{n=0}^{\infty} [g_n(t)|g, n\rangle + e_n(t)|e, n\rangle] \quad (25)$$




Figure 1. Schematic electronic and vibrational level structure for a two-photon transition in an ion trapped with frequency ω . ω_1 and ω_2 are the laser frequencies, and ω_e the transition frequency between ground and excited states. See the text for further details.

the differential equations of motion for the probability amplitudes $g_n(t)$ and $e_n(t)$ are obtained from the Schrödinger equation $i\hbar\partial_t|\psi_I(t)\rangle = \hat{H}_I|\psi_I(t)\rangle$,

$$i\dot{g}_n(t) = \frac{1}{2} \sum_{j=1}^2 \sum_{n'=0}^{\infty} \Omega_j e^{i(\theta_j t + \phi_j)} \langle n | e^{-in_j[\hat{a}(t) + \hat{a}^\dagger(t)]} | n' \rangle e_{n'}(t), \quad (26)$$

$$i\dot{e}_n(t) = -\tilde{\Delta}e_n(t) + \frac{1}{2} \sum_{j=1}^2 \sum_{n'=0}^{\infty} \Omega_j e^{-i(\theta_j t + \phi_j)} \langle n | e^{in_j[\hat{a}(t) + \hat{a}^\dagger(t)]} | n' \rangle g_{n'}(t), \quad (27)$$

where $\theta_j = \omega_j - \tilde{\omega}_L$. For large detunings, i. e., for $|\tilde{\Delta}| \gg \Omega_j, \omega$, see Fig. 1, and for an ion initially in the ground state one may assume that the excited state $|e\rangle$ is scarcely populated and it may be adiabatically eliminated. Then, setting $\dot{e}(t) = 0$, $e_n(t)$ may be written as a function of the $g_{n'}(t)$ from Eq. (27), and substituting this result into (26) there results a differential equation for the ground state probability amplitude,

$$i\dot{g}_n(t) = s g_n(t) + \frac{\tilde{\Omega}}{2} \sum_{n'=0}^{\infty} \mathcal{F}_{n,n'}(t) g_{n'}(t), \quad (28)$$

where

$$s = \frac{\Omega_1^2 + \Omega_2^2}{4\tilde{\Delta}}, \quad (29)$$

$$\mathcal{F}_{n,n'}(t) = \langle n | e^{-i\tilde{\eta}[\hat{a}(t) + \hat{a}^\dagger(t)]} | n' \rangle e^{i(\tilde{\delta}t + \tilde{\phi})} + \langle n | e^{i\tilde{\eta}[\hat{a}(t) + \hat{a}^\dagger(t)]} | n' \rangle e^{-i(\tilde{\delta}t + \tilde{\phi})}, \quad (30)$$

and where the effective two-photon Raman parameters, denoted by tildes, are given by

$$\begin{aligned} \tilde{\delta} &= \omega_1 - \omega_2, \\ \tilde{\eta} &= \eta_1 - \eta_2, \\ \tilde{\phi} &= \phi_1 - \phi_2, \\ \frac{\tilde{\Omega}}{2} &= \frac{\Omega_1 \Omega_2}{4\tilde{\Delta}}. \end{aligned} \quad (31)$$

The equation for the ground state probability amplitude corresponds to an effective Hamiltonian

$$\hat{H}_{eff} = \hbar s |g\rangle\langle g| + \frac{\hbar\tilde{\Omega}}{2} \left(e^{i\tilde{\eta}[\hat{a}(t)+\hat{a}^\dagger(t)]} e^{-i(\tilde{\delta}t+\tilde{\phi})} + H.c \right) |g\rangle\langle g|. \quad (32)$$

Note that the Stark-Shift produced by off resonant driving is included in s , which is a constant of motion and produces no effect on the Raman coupling between sidebands. We have thus adiabatically eliminated the excited state $|e\rangle$ ending with a Hamiltonian of the same form as (24) where the transitions between electronic levels are not present.

2.3.4. Two-photon Jaynes-Cummings Hamiltonian in the Raman Scheme: Vibrational RWA Using the Baker-Campbell-Hausdorff (BCH) identity, the exponential in the effective Hamiltonian (32) may be expanded in power series of $\tilde{\eta}$ [17, 18],

$$\hat{H}_{eff} = \frac{\hbar\tilde{\Omega}}{2} \left[e^{-\tilde{\eta}^2/2} \sum_{nn'} \frac{(i\tilde{\eta})^{n+n'}}{n!n'} \hat{a}^{\dagger n} \hat{a}^{n'} e^{i(n-n')\omega t} e^{-i\tilde{\delta}t} e^{-i\tilde{\phi}} + H.c \right]. \quad (33)$$

If the effective detuning is $\tilde{\delta} = \omega_1 - \omega_2 = 2\omega$, the second blue sideband becomes resonant, and we may neglect rapidly oscillating terms in a second or vibrational RWA [18]. The above Hamiltonian is then simplified to a two-photon Jaynes-Cummings-like Hamiltonian without electronic transitions. To leading order in $\tilde{\eta}$ it takes the form

$$\hat{H}_{eff} \approx \tilde{\eta}^2 \frac{\hbar\tilde{\Omega}}{4} \left(\hat{a}^{\dagger 2} e^{i\tilde{\phi}} + \hat{a}^2 e^{-i\tilde{\phi}} \right) = i\hbar \frac{\tilde{\eta}^2 \tilde{\Omega}}{4} \left(\hat{a}^2 - \hat{a}^{\dagger 2} \right), \quad (34)$$

where, in the last step, a relative phase between the applied fields $\tilde{\phi} = \phi_1 - \phi_2 = -\pi/2$ has been assumed.

2.3.5. Validity for time-dependent ω Unfortunately the above formal manipulations and approximations cannot be carried out in general for a time dependent ω . The interaction picture performed in 2.3.2, in particular, assumes a constant \hat{h}_0 . A time dependent one would require a more complex approach with time-ordering operators [19]. Similarly, the vibrational rotating wave approximation requires the stability of the frequency for times larger than a period to avoid off-resonant couplings. One may still obtain (34) for a sufficiently slowly varying ω , the criterion being that the change of the time-dependent trapping frequency in one time period T has to be much smaller than the frequency itself. We can write this condition as $\dot{\omega}(t)T \ll \omega(t)$ or

$$\frac{\dot{\omega}(t)}{\omega(t)^2} \ll 1, \quad (35)$$

which turns out to be the adiabaticity condition for the harmonic oscillator. Of course, if satisfied, the whole enterprise of applying the TT method would be useless. These arguments are far from constituting a proof that the TT method cannot be implemented for the harmonic oscillator. They simply leave this as an unresolved issue.

3. Engineering the Lewis-Riesenfeld invariant

In this section we describe a different method for transitionless dynamics of the harmonic oscillator [7]. A harmonic oscillator such as $\hat{H}_0(t)$ in Eq. (9) has the following time dependent invariant [9]

$$\hat{I}(t) = \frac{1}{2} \left(\frac{\hat{x}^2}{b^2} m \omega_0^2 + \frac{1}{m} \hat{\pi}^2 \right), \quad (36)$$

where $\hat{\pi} = b(t)\hat{p} - m\dot{b}\hat{x}$ plays the role of a momentum conjugate to \hat{x}/b , the dots are derivatives with respect to time, and ω_0 is in principle an arbitrary constant. The scaling, dimensionless function $b = b(t)$ satisfies the subsidiary condition

$$\ddot{b} + \omega(t)^2 b = \omega_0^2 / b^3, \quad (37)$$

an Ermakov equation where real solutions must be chosen to make \hat{I} Hermitian. ω_0 is frequently rescaled to unity by a scale transformation of b [9]. Other common and convenient choice, which we shall adopt here, is $\omega_0 = \omega(0)$. The eigenstates of $\hat{I}(t)$ become, with appropriate phase factors, solutions of the time-dependent Schrödinger equation,

$$\Psi_n(t, x) = \left(\frac{m\omega_0}{\pi\hbar} \right)^{1/4} \frac{1}{(2^n n! b)^{1/2}} \exp \left[-i(n + 1/2) \int_0^t dt' \frac{\omega_0}{b(t')^2} \right] \quad (38)$$

$$\times \exp \left[i \frac{m}{2\hbar} \left(\frac{\dot{b}}{b(t)} + \frac{i\omega_0}{b^2} \right) x^2 \right] H_n \left[\left(\frac{m\omega_0}{\hbar} \right)^{1/2} \frac{x}{b} \right], \quad (39)$$

and form a complete basis to expand any time-dependent state, $\psi(x, t) = \sum_n c_n \Psi_n(x, t)$, with the amplitudes c_n constant. A method to achieve frictionless, population preserving processes is to leave $\omega(t)$ undetermined first, and set b so that $\hat{I}(0) = \hat{H}_0(0)$ and $[\hat{I}(t_f), \hat{H}_0(t_f)] = 0$. This guarantees that the eigenstates of \hat{I} and \hat{H}_0 are common at initial and finite times and can be done by setting

$$\begin{aligned} b(0) &= 1, \dot{b}(0) = 1, \ddot{b} = 0, \\ b(t_f) &= \gamma = [\omega_0/\omega_f]^{1/2}, \dot{b}(t_f) = 0, \ddot{b}(t_f) = 0, \end{aligned} \quad (40)$$

and interpolating $b(t)$ with some real function that satisfies these boundary condition. The simplest choice is a polynomial,

$$b(t) = \sum_{j=0}^5 a_j t^j. \quad (41)$$

Once the a_j are determined from (40), $\omega(t)$ is calculated from the Ermakov equation (37), and the transitionless Hamiltonian $\hat{H}_{II}(t) = \hat{H}_0(t)$ is defined in terms of a local, ordinary, harmonic potential, but note that $\omega(t)^2$ may become negative during some time interval if the process is very fast, making the potential an expulsive parabola [7, 20]. This is illustrated in Figs. 2-4 for a transition from $\omega(0) = 2\pi \times 200$ Hz to $\omega(t_f) = 2\pi \times 2$ Hz in $t_f = 5$ ms.

The II method is thus clearly distinct from from TT and implements a different Hamiltonian. Note also, by comparing the coefficients, that the invariant operator \hat{I}




Figure 2. b (poynomial ansatz) versus t/t_f for an expansion from $\omega(0) = 2\pi \times 200$ Hz to $\omega(t_f) = 2\pi \times 2$ Hz.




Figure 3. Square of the frequency versus time for an expansion from $\omega(0) = 2\pi \times 200$ Hz to $\omega(t_f) = 2\pi \times 2$ Hz in $t_f = 5$ ms. Solid line: II method, polynomial b ; Dashed line: TT method with a linear-in-time $\omega(t)$ in \hat{H}_0 .




Figure 4. Average energy versus time. The initial state is the ground state of the initial trap and $E_0(0)$ the corresponding energy. Same processes and conditions as in Figure 3.

corresponding to \hat{H}_{II} is different from \hat{H}_{TT} , although they are both generalized harmonic oscillators.

3.1. Physical realization

The TT method only requires the time variation of a parabolic potential. Effective harmonic optical traps for neutral atoms may be formed by magnetic and/or optical means and their frequencies are routinely varied in time as part of many cold atom experiments. In magnetic traps, for example the frequency is modulated harmonically to

look for collective excitation modes of a condensate [21], and ramped down adiabatically to change its conditions (critical temperature, particle number, spatial extension) [22, 21], or as a preliminary step to superimpose an optical lattice [23]. Some experiments involve both time-dependent magnetic and optical traps or antitraps [24]. Purely optical traps are also manipulated in time, e.g. for adiabatic cooling of single neutral atoms [25]. In particular laser beams detuned with respect to the atomic transition form effective potentials for the ground state depending on Rabi frequency Ω and detuning Δ as $\Omega^2/4\Delta$ by adiabatic elimination of the excited state, thus forming attractive or repulsive potentials. This effective interaction can be made time dependent by varying the laser intensity, the frequency, or both [1], since the optical frequencies are many orders of magnitude larger than Rabi frequencies or detunings, and the changes will be slowly varying in the scale of optical periods. To vary the sign of $\omega^2(t)$, two dipole beams locked respectively on the blue and red side of the line may be superimposed. The intensity of a dipole trap can be changed by three or four orders of magnitude in 100 ns using acousto-optics or electro-optics modulators. This affects the trap radius, which can as well be controlled [2].

4. Discussion

We have compared and distinguished two different methods: a “transitionless-tracking” (TT) algorithm, and an “inverse-invariant” (II) method, to achieve transitionless dynamics for a fast frequency change of a quantum harmonic oscillator. They imply different driving Hamiltonians \hat{H}_{TT} and \hat{H}_{II} . The one in the II method can be implemented for ultracold atoms or ions by varying the trap frequency in time so as to enforce at initial and final times the commutativity between the Hamiltonian and an invariant of motion \hat{I} with the form of a generalized harmonic oscillator. In this method the states are at all times eigenstates of the invariant if initially so. Extending this method to other (anharmonic) potentials or systems may be difficult and remains an open question: in principle one should find in each case the corresponding state-carrying invariant and be able to engineer it. A generalization to Bose Einstein condensates has been worked out [8] without invoking the invariant concept, but it relies on scaling laws applicable to the harmonic oscillator. The TT method does not explicitly make use of invariants and provides a Hamiltonian for which the state dynamics follows exactly the adiabatic approximation of a time-dependent process defined by a reference Hamiltonian \hat{H}_0 . In this respect it has the advantage of being, at least formally, more generally applicable than the II approach. The feasibility of the actual realization is quite another matter and has to be studied in each case. An application example for two-level systems is discussed in [4]. For the harmonic oscillator studied here \hat{H}_{TT} becomes, as \hat{I} , a generalized harmonic oscillator too, but it is different from \hat{I} and plays a different role. In particular the “carrying states” are not instantaneous eigenstates of \hat{H}_{TT} but of the reference, harmonic oscillator Hamiltonian \hat{H}_0 . \hat{H}_{TT} includes, in addition to a harmonic term in \hat{H}_0 with a time dependent frequency going from the

initial to the final values, a time-dependent (squeezing) non-local potential with the symmetrized product of position and momentum operators. We have found a realization for cold atoms that, unfortunately, requires a time scale larger than the times needed for an adiabatic process. Therefore the applicability of the method remains an open question. Trap expansions and compressions play such an important role in cold atom experiments that investigating further the fundamental properties of both approaches and their technical realization is definitely worthwhile.

Acknowledgments

We thank M. V. Berry for discussions and acknowledge funding by Projects No. GIU07/40, FIS2009-12773-C02-01, 60806041, 08QA14030, 2007CG52, S30105, and Juan de la Cierva Program.

Appendix A. Relation to the squeezing operator

The evolution operator takes a particularly simple form when using the simplified case $E_n(t) = 0$, so that $\hat{H}(t) = \hat{H}_1(t)$. Taking into account that $[\hat{H}_1(t), \hat{H}_1(t')] = 0$ we can write

$$\hat{U}(t) = e^{-i \int_0^t \hat{H}_1(t) dt / \hbar}. \quad (\text{A.1})$$

This may be evaluated explicitly with (19) fixing the time of the creation and annihilation operators to 0,

$$\hat{U}(t) = e^{\frac{1}{2} \ln \left(\sqrt{\frac{\omega(t)}{\omega(0)}} \right) [\hat{a}_0^2 - (\hat{a}_0^\dagger)^2]} = \hat{S}[r(t)], \quad (\text{A.2})$$

which is a squeezing operator with real argument $r(t) = \ln \left(\sqrt{\frac{\omega(t)}{\omega(0)}} \right)$. It is unitary with inverse $[\hat{S}(r)]^{-1} = \hat{S}(-r)$. Using the relations

$$\begin{aligned} \hat{a}_t^\dagger + \hat{a}_t &= \sqrt{\frac{\omega(t)}{\omega(0)}} (\hat{a}_0^\dagger + \hat{a}_0), \\ \hat{a}_t^\dagger - \hat{a}_t &= \sqrt{\frac{\omega(0)}{\omega(t)}} (\hat{a}_0^\dagger - \hat{a}_0), \end{aligned} \quad (\text{A.3})$$

and the formal properties of \hat{S} , see e.g. [26], it is easy to prove that

$$\begin{aligned} \hat{S}(r) \hat{a}_0 \hat{S}(-r) &= \hat{a}_t, \\ \hat{S}(r) \hat{a}_0^\dagger \hat{S}(-r) &= \hat{a}_t^\dagger. \end{aligned} \quad (\text{A.4})$$

In fact any combination of powers of \hat{a}_0 and \hat{a}_0^\dagger is mapped to the same combination of powers of \hat{a}_t and \hat{a}_t^\dagger by this unitary transformation. $|0_t\rangle \equiv \hat{S}|0_0\rangle$ is indeed the vacuum at time t , as

$$\hat{a}_t |0_t\rangle = \hat{S}(r) \hat{S}(-r) \hat{a} \hat{S}(r) |0_0\rangle = \hat{S}(r) \hat{a}_0 |0_0\rangle = 0. \quad (\text{A.5})$$

In a similar way we note that, consistently,

$$\begin{aligned}\hat{S}(r)|n(0)\rangle &= \hat{S}(r)\frac{1}{\sqrt{n!}}(\hat{a}_0^\dagger)^n|0_0\rangle = \frac{1}{\sqrt{n!}}\hat{S}(r)(\hat{a}_0^\dagger)^n\hat{S}(-r)\hat{S}(r)|0_0\rangle \\ &= \frac{1}{\sqrt{n!}}(\hat{a}_t^\dagger)^n|0_t\rangle = |n(t)\rangle.\end{aligned}\tag{A.6}$$

References

- [1] Bize S et al. 2005 *J. Phys. B: At. Mol. Opt. Phys.* **38** S449
- [2] Han D J, DePue M T and Weiss D S 2001 *Phys. Rev. A* **63** 023405
- [3] Polkovnikov A and Gritsev V 2008 *Nature Physics* **4** 477
- [4] Berry M V 2009 *J. Phys. A: Math. Theor.* **42** 365303
- [5] Rezek Y and Kosloff R 2006 *N. J. Phys* **8** 83
- [6] Kato T 1950 *J. Phys. Soc. Japan.* **5** 435
- [7] Chen X, Ruschhaupt A, Schmidt S, del Campo A, Guéry-Odelin D and Muga J G 2009 arXiv:0910:0709; *Phys. Rev. Lett.*, to appear
- [8] Muga J G, Chen Xi, Ruschhaupt A and Guéry-Odelin D 2009 *J. Phys. B: At. Mol. Opt. Phys.* **42** 241001
- [9] Lewis H R and Riesenfeld W B 1969 *J. Math. Phys.* **10** 1458
- [10] Palao J P, Muga J G and Sala R 1998 *Phys. Rev. Lett.* **80** 5469
- [11] Berry M V 1985 *J. Phys. A: Math. Gen.* **18** 15
- [12] Meekhof D M, Monroe C, King B E, Itano W M and Wineland D J 1996 *Phys. Rev. Lett.* **76** 1796
- [13] Zeng H and Lin F 1995 *Phys. Lett. A* **201** 139
- [14] Zeng H *Phys. Lett. A* 1998 **247** 273
- [15] Katori H, Ido T and Kuwata-Gonokami M 1999 *J. Phys. Soc. Japan* **68** 2479
- [16] Ido T and Katori H 2003 *Phys. Rev. Lett.* **91** 053001
- [17] Orszag M 2000 *Quantum Optics* (Springer, Berlin)
- [18] Lizuain I, Muga J G and Eschner J 2008 *Phys. Rev. A* **77** 053817
- [19] Schrade G, Bardroff P J, Glauber R J, Leichtle C, Yakovlev V and Schleich W P 1997 *App. Phys. B* **64** 181
- [20] Khaykovich L et al. 2002 *Science* **296** 1290
- [21] Jin D S, Ensher J R, Matthews M R, Wieman C E and Cornell E A, 1996 *Phys. Rev. Lett.* **77** 420
- [22] Miesner H J, Stamper-Kurn D M, Andrews M R, Durfee D S, Inouye S and Ketterle W 1998 *Science* **279** 1005
- [23] Tuchman A K, Li W, Chien H, Dettmer S and Kasevich M A 2006 *New Journal of Physics* **8** 311
- [24] Tin-Lun Ho and Qi Zhou 2007 *Phys. Rev. Lett.* **99** 120404
- [25] Tuchendler C, Lance A M, Browaeys A, Sortais Y R P and Grangier P 2008 *Phys. Rev. A* **78** 033425
- [26] Barnett A M and Radmore P M 1997 *Methods in Theoretical Quantum Optics* (Oxford: Oxford U. Press)