



HAL
open science

Psychological approaches to treatment of post-concussion syndrome: a systematic review

Amal Al Sayegh, David Sandford, Alan J Carson

► **To cite this version:**

Amal Al Sayegh, David Sandford, Alan J Carson. Psychological approaches to treatment of post-concussion syndrome: a systematic review. *Journal of Neurology, Neurosurgery and Psychiatry*, 2010, 81 (10), pp.1128. 10.1136/jnnp.2008.170092 . hal-00570134

HAL Id: hal-00570134

<https://hal.science/hal-00570134>

Submitted on 27 Feb 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Psychological approaches to treatment of post-concussion syndrome: a systematic review

Al Sayegh A, Sandford D, Carson A J

Corresponding author

Amal Al Sayegh
Specialty Registrar
Royal Edinburgh Hospital
Morningside Terrace
Edinburgh
EH10 4HF
United Kingdom
E-mail: amal.alsayegh@nhs.net
Tel: 0131 537 6248
Fax: 0131 537 6140

Co-authors

David Sandford
Cognitive behavioural therapist
Lancashire Traumatic Stress Service
Lancashire NHS Foundation Trust
Chorley
PR7 1PS
United Kingdom

Alan J Carson
Consultant Neuropsychiatrist
Department of Psychiatry
University of Edinburgh
Royal Edinburgh Hospital
EH10 5HF
United Kingdom

Keywords

Post-concussion syndrome
Brain injury
Cognitive behavioural therapy
Psychotherapy
Psychological treatment

Word Count 3, 237 (excluding tables and figures)

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in Journal of Neurology, Neurosurgery & Psychiatry and any other BMJPGJ products to exploit all subsidiary rights, as set out in our licence (<http://jnnp.bmj.com/fora/licence.pdf>).

ABSTRACT

Background & Aim: Post-concussion syndrome is a term used to describe the complex, and controversial, constellation of physical, cognitive and emotional symptoms associated with mild brain injury. At the current time, there is a lack of clear, evidence-based treatment strategies. In this systematic review we aimed to evaluate the potential efficacy of CBT and other psychological treatments in post-concussion symptoms.

Methods: Four electronic databases were searched to November 2008 for studies of psychological approaches to treatment or prevention of post-concussion syndrome or symptoms.

Results: The search identified 7, 763 citations and 42 studies were included. This paper reports the results of 17 randomised controlled trials for psychological interventions which fell into 4 categories: CBT for PCS or specific PCS symptoms; information, reassurance and education; rehabilitation with a psychotherapeutic element and mindfulness/relaxation. Due to heterogeneity of methodology and outcome measures meta-analysis was not possible. The largest limitation to our findings was the lack of high quality studies.

Conclusion: There was evidence that CBT may be effective in the treatment of PCS. Information, education and reassurance alone may not be as beneficial as previously thought. There was limited evidence that multi-faceted rehabilitation programmes that include a psychotherapeutic element or mindfulness/relaxation benefit those with persisting symptoms. Further, more rigorous trials of CBT for post-concussion symptoms are required.

INTRODUCTION

Most clinicians are familiar with the complex constellation of physical, cognitive and emotional symptoms complained of by patients in the aftermath of a mild traumatic brain injury (MTBI). Almost every aspect of the syndrome is controversial, including rates, mechanisms and even the name. MTBI has a high incidence with 100-300 hospital-treated cases/100,000 population per year in most industrialised countries, however, a large number of MTBI cases are not treated in hospitals and the actual rate of all MTBI may be in excess of 600/100,000¹. In the immediate aftermath of injuries many patients describe a cluster of troubling symptoms (see table 1) but there is considerable controversy over the prognosis of such symptoms with some authors arguing complete recovery within weeks² and others suggesting highly disabling symptoms over years³. The World Health Organisation helpfully conducted a high quality systematic review of the epidemiological evidence and suggested that there are no MTBI attributable, objectively measured cognitive deficits beyond 1-3 months post-injury in the majority of cases⁴.

There is also disagreement over the aetiological mechanism of these putative symptoms with some authors believing that the presentation can be explained in terms of acquired neuropathological damage⁵ although much of the evidence cited to support this is problematic⁶. Many take the view that the mechanism involves a complex interplay of biological, psychological and social factors which include prior health, life stressors and compensation/litigation issues⁴. Such a view on aetiology certainly explains why similar symptoms are described after orthopaedic injuries such as long bone fractures; why there is a highly variable rate of presentation from country to country; and why financial compensation is a significant risk factor¹. This debate is translated into the actual name for such symptoms. For many years they

have been referred to as post-concussion syndrome (PCS) and this tradition is continued in ICD-10 and DSM-IV. However the WHO cautioned against this saying that such a mechanism was at the current time unproven and reminded us of the age old epidemiological rule that association was not proof of causation⁴. We agree with the WHO's logic but have continued to use the term PCS in this review to describe those with persistent symptoms as it remains the accepted term within ICD-10 and DSM-IV, the term that most clinicians are familiar with, and the term most commonly used in the studies we were systematically reviewing.

Perhaps the only area that clinicians do agree upon is that there is a lack of clear, evidence-based treatment strategies to guide our clinical management of such patients. We consider that the development of these symptoms after MTBI appears to have much in common with a number of functional symptom syndromes such as chronic fatigue syndrome⁷. We have noted the beneficial effects of cognitive behavioural therapy (CBT) in functional disorders⁸ and were interested in CBT's potential as a treatment for symptoms after MTBI. We noted with encouragement the suggestion that patients responded positively to appropriate information and reassurance given shortly after injury⁹. This was in keeping with our view that there is a significant psychological component to more persistent complaints. However, we were unaware of any definitive randomised controlled trials of CBT in this group of patients. The purpose of this systematic review was to evaluate what, if any, clinical trial evidence existed on the efficacy of psychological therapies for the treatment of PCS.

Table 1

METHODS

The sources of literature were the electronic databases Medline (1950-); Embase (1980-); PsycINFO (1967-) and CINAHL (1982-) up to the end of November 2008. The search strategy included the use of the following thesaurus terms: 'post-concussion syndrome'; 'brain concussion' and 'brain injuries'. In addition, we used the following keywords: 'brain contusion'; 'concussion'; 'post-concussion'; 'brain injury'; 'brain damage' and 'head injury'. In order to ensure that all relevant studies that used a psychological approach were identified, we combined the search strategy with keywords and, where available, subject headings including 'psychotherapy'; 'cognitive therapy'; 'cognitive behavioural therapy'; 'CBT'; 'behavioural therapy'; 'psychological therapy'; 'psychological treatment'; 'psychological techniques'; 'psychoeducation'; 'psychosocial'; 'biopsychosocial'; 'bibliotherapy'; 'computer-assisted therapy'; 'talking therapy'; 'rational emotive'; 'self-instruction'; 'self-management'; 'self-attribution' and 'non-surgical interventions'. Using the inclusion and exclusion criteria below, we reviewed the titles of all citations and retrieved relevant abstracts for more detailed evaluation. Where there was uncertainty, the full paper was studied. We also hand searched the reference list of relevant studies to aid identification of further studies.

At the outset we believed that there was only limited research in this field and we therefore included data from pilot studies and case series as well as randomised controlled trials (RCT). We also decided to include studies that described a range of severities of head injury (including moderate and severe) if it seemed that the psychological intervention was addressing chronic problems in keeping with PCS.

Finally we included studies of patients with brain injuries due to non-traumatic causes.

Inclusion criteria

- Studies examining a psychological approach to treatment or prevention of post-concussion syndrome, post-concussion symptoms or other psychiatric or psychological problems after mild acquired brain injury
- Studies that included participants with moderate and severe head injuries if they also included MTBI
- Adult participants only
- English language reports

Exclusion criteria

- Letters to editors and editorials without data
- Studies outside the timescales above as these were not available electronically
- Studies excluding MTBI or those with Glasgow Coma Scale score of 13 or above
- Studies using purely neuro-psychological/cognitive re-training
- Studies of rehabilitation programmes with no detail of psychotherapeutic elements.

The quality of each randomised controlled trial was assessed using the 22-item CONSORT statement 2001 checklist¹⁰ by AAS and DS. Where there was disagreement, AJC adjudicated.

RESULTS

The search strategy identified 7,763 references and 42 were included in the systematic review. The inclusion and exclusion of papers is shown in Figure 1. Psychological interventions fell into one of four categories:

1. Use of CBT in post-concussion syndrome or with specific post-concussion symptoms.
2. Information, reassurance and education.
3. Rehabilitation programmes with a psychotherapeutic element.
4. Mindfulness-based interventions and effects of stress/relaxation.

Figure 1

We found more randomised trial evidence than we expected and have therefore concentrated this report on data from RCTs but have presented our review of the remaining evidence as supplemental web files. The 17 RCTs discussed in this paper are presented in Table 2.

Table 2

The participants are a heterogeneous sample of head injuries of various severity (including severe) in addition to those with post-concussion syndrome or symptoms from MTBI. The non-RCT studies and ratings of the RCTs using the CONSORT checklist are published as supplemental material on-line. For ease of reference, a summary of the RCTs is shown in Table 3 showing the number of CONSORT items

met, the intervention used, number of participants entering and at follow-up, length of follow-up, whether only participants with MTBI were included, the definition of MTBI used and whether a benefit was shown.

Table 3

Due to heterogeneity of methodology and outcome measures, meta-analysis of outcome was not possible. We gave consideration to calculating the effect sizes of these trials but opted not to as we felt this would encourage a numerical comparison between studies which should not be compared in such a fashion.

Evidence from Randomised Controlled Trials

Table 3 briefly describes a summative quality measure for each trial based on the number of 'CONSORT items' met. However, we caution that comparison of such total 'scores' is not necessarily informative as each of the 22 items is given equivalent weight, whereas certain aspects of trials design, such as randomisation techniques, will have considerably more influence than, say, the structure of the discussion. A full qualitative assessment of each trial is detailed in supplemental web material.

In general the RCTs reviewed performed well on giving the scientific background and rationale; eligibility criteria for participants; details of intervention intended for each group; specific objectives and hypotheses and defined primary and secondary outcome measures. However, they were poor on reporting how sample size was determined; random allocation sequence was generated; allocation concealment was implemented; who generated the allocation sequence, enrolled participants and assigned them to groups; blinding was ascertained; the flow of participants; the dates defining periods of recruitment and follow-up; the use of intention-to-treat analysis and reporting important adverse events.

The definition of MTBI used varied between trials (Table 3) and trials tended to use widely different outcome measures (Table 2).

Cognitive behavioural therapy

There were 3 randomised controlled trials examining CBT and all concluded some form of benefit¹¹⁻¹³. Two of these met 11 of the 22 items on the CONSORT checklist^{11, 12}. One trial randomised consecutive hospital admissions after MTBI (GCS 13-15, no PTA) to one session of CBT and gave out a 10-page manual¹¹. In comparison to routine care, the CBT group reported reduced frequency, intensity and duration of symptoms at 6-month follow-up. The second trial randomised referrals from local brain injury units and community services and delivered CBT to the treatment group adapted to account for difficulties with attention, concentration, fatigue and memory¹². Compared to the waiting list controls, the CBT group showed improvement in anxiety and depression at 1-month follow-up. The third trial had a more robust methodology meeting 19 of the 22 CONSORT items¹³. They recruited participants with mild-moderate TBI and delivered thrice-weekly CBT (with thrice-weekly cognitive remediation) and concluded significant improvement in psychosocial functioning (especially anxiety and depression) but little change in cognitive measures¹³. It is, perhaps, unfortunate that stand-alone efficacy of CBT was not examined.

Information, reassurance & education

There were 10 papers that tested the efficacy of information, reassurance and education¹⁴⁻²³. Generally this involved the early provision of information about diagnosis and possible post-concussion symptoms; reassurance about prognosis; education on ways of coping and resumption of activities. Some incorporated multi-disciplinary management tailored for individual needs. Two papers were considered as one trial as they detailed different follow-up points on the same cohort of participants^{18,20}.

Three studies demonstrated a benefit^{15, 19, 21}. The first performed very poorly on the CONSORT checklist meeting only 7 items and based its conclusions on a follow-up rate of 22%¹⁵. The second¹⁹ repeated an earlier methodology¹⁷ and concluded that patients with *moderate* head injury *requiring admission* benefit from a routinely offered early intervention service but that no benefit was seen when such an intervention was offered to all patients presenting with a head injury. These studies were of higher quality meeting 16¹⁷ and 17¹⁹ CONSORT items. The third trial met 10 CONSORT items and randomised emergency department discharges after MTBI and undertook neuropsychological assessments on the treatment group and gave them an information booklet²¹. Compared to 'no intervention' they found improved sleep and anxiety, reduced distress but no difference in neuropsychological measures.

There were 6 RCTs in this category that concluded no benefit or reported inconclusive findings^{14, 16-18, 20, 22, 23}. The quality of these trials varied greatly, meeting between 7 and 19 CONSORT items and included the lowest and highest performers (Table 3). These RCTs examined inpatient information, encouragement, physiotherapy and reassurance¹⁴; inpatient information and reassurance¹⁶; early intervention, information and further treatment as needed for those discharged from the emergency department¹⁷; single session education and support in the emergency department^{18,20}; follow-up within 1 week of injury for education and multi-disciplinary treatment as needed²² and reassurance, information, telephone and outpatient reviews²³.

Rehabilitation programmes with a psychotherapeutic element

There were 2 RCTs that examined the efficacy of rehabilitation programmes that included psychotherapy^{24, 25}. One concluded no difference between groups following the addition of a problem-solving intervention, but it was of low quality demonstrating only 8 CONSORT items²⁴. The other reported mixed findings following the addition of individual and group support. It met 14 CONSORT items²⁵.

Mindfulness-based interventions & relaxation

One trial tested the effects of relaxation on PCS symptoms and found the severity of symptoms increased with stress in those reporting symptoms regardless of history of head injury and the effects were reduced by relaxation²⁶. One trial examined a mindfulness-based intervention but found no difference between groups²⁷. Both trials met between 8²⁶ and 10²⁷ CONSORT items performing at the lower end of the range compared to the other trials (Table 3).

Evidence from Non-randomised Trials

The details of interventions and main findings of studies of designs other than RCTs are available as on-line supplemental material but are summarised here.

Cognitive behavioural therapy

There were 7 studies. One was a controlled trial which showed initial benefit but this was not maintained at follow-up²⁸. Limitations included lack of power calculations to determine sample size and follow-up data not being analysed statistically²⁸. Three studies examined the use of CBT with the specific PCS symptoms of headache²⁹ and insomnia^{30, 31} and all concluded an improvement in the symptom investigated. Limitations include inadequate²⁹ or no control group³¹ and high drop-out rates²⁹. A case study examined the efficacy of CBT in treating anxiety and OCD after moderate traumatic brain injury and revealed significant improvements in most measures³². Generalizability is limited as the patient received concurrent cognitive rehabilitation. While the remaining 2 papers also concluded a positive outcome for CBT in the treatment of PCS, there were substantive methodological weaknesses including a potentially biased sample, no information on how subjects were selected from other referrals, no control group and lack of detail on how the investigators excluded a diagnosis of depression^{33, 34}.

Information, reassurance & education

There were 3 retrospective studies³⁵⁻³⁷ and 1 single case study³⁸. Two of the 3 retrospective studies considered a control group in the form of “little/no treatment”³⁵ or outcomes in those treated before a change in treatment was implemented³⁶. Both concluded a benefit. The third retrospective study had no control group and reported no difference in outcome³⁷. The single-case study reported a benefit³⁸.

Rehabilitation programmes with a psychotherapeutic element

Thirteen studies examined the efficacy of rehabilitation programmes that included psychotherapy. Two papers were treated as one study as they were published in 2 parts^{39, 40}. Interventions and outcome measures varied greatly. The psychotherapeutic interventions were part of multi-disciplinary rehabilitation and so stand-alone efficacy was not studied. Generally there was little detail about the psychotherapy undertaken. Almost all the studies concluded a benefit but had no control group³⁹⁻⁴⁴, had no randomisation⁴⁵ or were case studies with a sample size of 1 or 2⁴⁶⁻⁵¹. One retrospective study revealed inconclusive findings⁵².

Mindfulness-based interventions

In a pre-post design study with drop-outs as controls; no follow-up; high attrition rates and no control for medication, an improvement in measures of quality of life was concluded⁵³.

DISCUSSION

Our systematic review on studies of the potential efficacy of CBT as a treatment for post-concussion symptoms found 10 studies of which 3 had a randomised controlled design. All 10 studies concluded a benefit. However they had relatively small numbers and short durations of follow-up and do not allow robust conclusions about the efficacy of CBT to be drawn.

In general the 17 RCTs we described in this review had methodological weaknesses, in particular, a failure to predetermine sample size, failure to detail the randomisation procedure, failure to ensure blinding (if indicated in the methodology) and failure to undertake intention-to-treat analyses. It is important that further trials in this field address these concerns. It is unlikely that further small scale, methodologically

limited studies will add any scientifically valuable information on treatment efficacy and such investigations should be confined to pilot studies of procedural and methodological issues for definitive trials.

The benefits of information, education and reassurance in the treatment and prevention of PCS and PCS symptoms is generally endorsed in the MTBI literature^{2, 9, 54}. Yet our systematic review identified 6 randomised controlled trials that concluded no benefit versus 3 that demonstrated an improvement in symptoms. It may be argued that elements of this intervention may be justified as a cost-effective intervention to prevent the development of PCS, perhaps in selected patients, such as those whose head injury warranted admission^{17, 19}, but we concluded that evidence to support its usefulness had been perhaps overstated.

The studies of rehabilitation programmes with a psychotherapeutic element were diverse in delivery, setting and content. The psychotherapeutic elements ranged from counselling and unspecified 'psychotherapy' to CBT. Outcome measures varied greatly and meaningful comparison was not possible. Generally, participants were those with persisting problems. Although most studies in this category showed a positive outcome, the studies with inconclusive findings or showing no difference were of more robust methodology^{24, 25}. Generally there was little information about the details of the psychotherapeutic component of the programme and so it is difficult to draw conclusions as stand-alone efficacy was not addressed. There was insufficient evidence to recommend these multi-faceted programmes in the treatment of persisting PCS.

Based on the limited evidence found on mindfulness-based interventions and relaxation, these techniques cannot be currently recommended for prevention or treatment of post-concussion symptoms.

Limitations to our systematic review included publication bias with the assumption of a tendency to publish positive small trials but not small studies of no effect. Time constraints restricted contacting experts and researchers in the field for unpublished material. The inclusion criteria for individual studies were judged by only one author. However, we think the largest limitation to our conclusions was not our review methodology but the lack of high quality studies examining the treatment and prevention of PCS. We had however anticipated this at the time of designing the study and the aim was to describe the current, albeit limited evidence.

We believe that similarities can be drawn between PCS and complex functional symptom syndromes such as chronic fatigue syndrome. Most patients with fatigue are managed in primary care but some may require referral to specialist care. Only a small proportion will be found to be suffering from a recognised medical disease⁵⁵. Patients may be worried that the fatigue is a symptom of severe but undiagnosed disease. This may lead to repeated presentations to health services and impairment in physical and social functioning. NICE guidelines recommend CBT and graded exercise as the most effective specialist treatment approaches⁵⁶. Focusing on symptoms and improving function tends to be more productive than engaging in a debate about the presence or absence of disease or undergoing repeated investigations and instrumentation. Parallels can be drawn with the experience of a patient disabled by PCS. We hope that a similar approach to managing PCS might be effective. The

available data was not robust enough to make any firm conclusion in this regard. However, there is data to suggest that CBT shows some promise and that a definitive trial would be a useful contribution.

In the interim, we would also tentatively suggest that for the majority that present to emergency departments and primary care with MTBI, brief information and explanation should be provided and it may be sufficient to manage any anxieties. For those that require further investigation or admission for observation, more tailored and specific information, education and reassurance is perhaps warranted to help prevent the development of PCS. This should include reassurance that cognitive difficulties are common and usually resolve by 3 months. There was support for the use of CBT but it is likely to be a treatment for those with persisting problems or disability.

In conclusion, there was promising evidence that CBT may be effective in the treatment of PCS. Information, education and reassurance alone may not be as beneficial as previously thought. There was limited evidence that multi-faceted rehabilitation programmes that include a psychotherapeutic element are of benefit in the management of persisting symptoms. Further and more rigorous randomised controlled trials of CBT for PCS are needed.

Acknowledgements

We would like to thank Dr. Brian Pentland for valuable comments on earlier drafts; Alan Mill and Elizabeth Killean for assistance in retrieving papers and Cat Graham for advice on statistics.

Contributors

AAS did the search. AJC generated the study hypothesis and design. AAS and DS reviewed the original papers and rated quality of trials. AAS and AJC contributed to drafting the report.

Competing Interests

None


Funding

None

Table 1: Post-concussion symptoms

Physical	Cognitive	Emotional
Headache Dizziness Fatigue Visual disturbances Noise sensitivity Light sensitivity Insomnia	Memory deficits Attention/concentration deficits Executive function deficits	Irritability Depression Anxiety

Figure 1: Selection of papers



*Non-randomised controlled trials, non-controlled trials, retrospective studies, case series, case studies presented as on-line supplemental material.

Table 2: Randomised Controlled Trials (n=17)

Authors	Participants	Intervention	Outcome measures	Main results [Standard deviation]
Relander et al (1972) ¹⁴	Hospital admissions following cerebral concussion	Information, continuity of care, encouragement, physiotherapy, reassurance (n=82, 34 at follow-up) Routine care (n=92, 25 at follow-up)	Time in bed in hospital; time in hospital; time off work	Time in bed and hospital: no difference Time off work reduced by 2 weeks No difference at 12 months in number or severity of symptoms
Hinkle et al (1986) ¹⁵	Minor head injury admitted for 24-hour observation	Routine care (n=75) Information (n=no info) Info. and reassurance (n=no info) (n=166 for group 2 + 3)	Days before return to work or social activity	No difference between 3 groups but those in the 2 treatment groups returned to work 1 week earlier than Routine Care group.
Alves et al (1993) ¹⁶	Hospital admissions mild uncomplicated head injury	Routine care (n=210) Information only (n=176) Information & reassurance (n=201)	Post-concussion symptoms (relative risk of being symptomatic at follow-up)	No significant difference between groups
Mittenberg et al (1996) ¹¹	Consecutive hospital admissions after mild head trauma, GCS 13-15 and no PTA	10-page manual and one session CBT; routine care and discharge information for control group	Frequency (number of PCS symptoms), intensity (scale of 1-10) and duration (days) 6 months after discharge	CBT group significantly reduced frequency (mean 3.10[3.19] to 1.62[2.04]), intensity (mean 1.72[1.93] to 0.80[1.13]) and duration (mean 51.19[45.10] to 33.18[35.62]) of symptoms
Wade et al (1997) ¹⁷	A&E attendances and hospital admissions for head injury of any severity	Early intervention, information, advice, further intervention as required (including CBT) (n=252); standard care (n=226)	RPQ RHFUQ	No difference between groups. Subgroup analysis suggested some benefit for those with moderate or severe injury
Paniak et al (1998) ¹⁸	Volunteers with MTBI from	Single session education & support	Problem Checklist	Both groups improved, no

	consecutive admissions to emergency department.	(n=58) Treatment as needed (with neuropsychological and personality assessment (n=53)	Community Questionnaire SF-36	Integration	significant difference
Wade et al (1998) ¹⁷	Hospital admissions for head injury of any severity	Early intervention, information, advice, further intervention as required (including CBT) (n=132); standard care (n=86)	RPQ RHFUQ		Significant difference in RPQ mean 9.8[11.7] (trial) vs 13.9[13.6] (control); RHFUQ mean 5.36[7.81] (trial) vs 8.23[8.75] (control)
Paniak et al (2000) ²⁰	Volunteers with MTBI from consecutive admissions to emergency department at one year follow-up.	Single session education & support (n=53) Treatment as needed (with neuropsychological and personality assessment (n=52)	Problem Checklist Community Questionnaire SF-36	Integration	Both groups improved, no significant difference. Improvements at 3 months maintained at 12 months.
Hanna-Pladdy et al (2001) ²⁶	Undergraduate students screened for history of MTBI and PCS symptoms	Symptomatic/uninjured (n=22) Symptomatic/MTBI (n=22) Asymptomatic/uninjured (n=22) Asymptomatic/MTBI (n=22) Half group: high stress, other half: relaxation	Physiological measures; Neuropsychological measures; Self-reported measures (post-concussion symptoms and stress)		Post-concussion symptoms increase with stress in symptomatic groups regardless of history of injury, effects decreased with relaxation. No difference in symptom reporting between MTBI and uninjured.
McMillan et al (2002) ²⁷	Neurosurgical patients with traumatic brain injury with attentional problems on neuropsychological testing	Attentional Control Training (n=44) Physical exercise (n=38) Control/no intervention (n=48)	Cognitive function (objective + self report); HADS; General Health Questionnaire, RPQ		No significant differences between the 3 groups on these measures post-treatment or 6 or 12 month follow-up
Ponsford et al (2002) ²¹	Discharges from emergency department after mild head injury	Neuropsychological assessment & information booklet (n=79) No intervention (n=123)	SCL-90-R HRSRE (stress) PCS Checklist Neuropsychological measures		No means or SD reported. Improved sleep (p=0.01) and anxiety (p=0.04) No difference in neuropsychological measures

Rath et al (2003) ²⁴	High-functioning TBI (various severity) outpatients attending a neuropsychological rehabilitation programme with variety of post-concussion complaints	Problem-solving group (n=27, 18 at follow-up) Conventional treatment group (n=19, 13 at follow-up)	Measures of cognitive skills, psychosocial functioning, problem-solving and significant-other reports	Both groups improved; inconclusive
Hodgson et al (2005) ¹²	Referrals from local brain injury units and community services	CBT adapted to account for difficulties with attention, concentration, fatigue and memory, weekly sessions for 9-14 weeks (n=6) Wait-list control (n=6)	Social Phobia and Anxiety Inventory HADS Coopersmith Self-esteem Inventory Profile of Mood States	Anxiety mean 9.5[3.9] to 5.3[2.7] vs 10.9[2.6] to 11.3[5.4] in controls. Depression mean 8.9[4.9] to 5.2[4.4] vs 8.0[2.3] to 8.3[2.3] in controls Social Phobia mean 80.5[23.6] to 40.5[16.6] vs 78.1[30.0] to 64.8[37.1] in controls
Tiersky et al (2005) ¹³	Recruitment of subjects with mild-moderate TBI	CBT and cognitive remediation (50mins each, 3x per week for 11 weeks) versus waiting list control	SCL-90R PASAT CRI (problem solving) Attention questionnaire	Significant improvement SCL-90R, mean 8.06[0.41] vs 1.71[1.00]; depression subscale, mean 1.12[0.45] to 2.11[1.14]; anxiety subscale, mean 0.72[0.42] vs 1.53[1.02] and PASAT 135.55[30.71] vs 110.88[60.28]. No difference in CRI or attention
Ghaffer et al (2006) ²²	Consecutive presenters to emergency dept. with mild TBI	Follow-up within 1 week of injury for education and multi-disciplinary treatment as needed for 6 months thereafter (n=86) Control (n=84)	RPQ RHFUQ GHQ Psychometric battery	No significant treatment effects, except in improvement of depression in those with past psychiatric history (no means reported, p=0.01)
Elgmark Andersson et al (2007) ²³	Selected subjects from patients diagnosed with MTBI in emergency department	Reassurance, information, telephone contact and outpatient reviews plus specialist referral as	PCS questionnaire Life Satisfaction questionnaire CIQ	No difference other than improvement in one aspect of life satisfaction (physical

Ownsworth et al (2008) ²⁵	Acquired brain injury, various aetiology, convenience sample attending outpatient brain injury units	<p>needed (n=246) Treatment as usual (n=109)</p> <p>Individual intervention (occupation-based support) (n=10) Group intervention (self-awareness and compensatory strategies) (n=11) Combined intervention (n=10)</p>	<p>SF-36</p> <p>Canadian Occupational Performance Measure (COPM). Patient Competency Rating Scale (PCRS). The Brain Injury Community Rehabilitation Outcome 39 (BICRO-39)</p>	<p>health)</p> <p>No summary data COPM: slight improvement in goal attainment PCRS: improvement in individual and group but not combined intervention BICRO-39: mixed findings</p>
--------------------------------------	--	---	---	--

Abbreviations

BAI= Beck Anxiety Inventory; BDI= Beck Depression Inventory; BDI-II= Beck Depression Inventory II; CIQ= Community Integration Questionnaire; CES-D= Centre for Epidemiological Studies-Depression; CRI= Coping Response Inventory GHQ= General Health Questionnaire; GCS= Glasgow Coma Scale; HADS= Hospital Anxiety and Depression Scale; HRSRE= Holmes Rahe survey of recent experiences; MOCI= Maudsley Obsessive-Compulsive Inventory; MHLC= Multidimensional Health Locus of Control Scale; MTBI= Mild traumatic brain injury; PASAT= Paced auditory serial addition task; PCS= post concussive syndrome; PSS= Perceived Stress Scale; PTA= Post-Traumatic amnesia; RHFUQ= Rivermead head injury follow up questionnaire; RPQ= Rivermead Post-concussion symptoms questionnaire; SF36= Medical Outcome Study 36-item Short-Form Health Survey; SCL-90-R=Symptom checklist 90 revised; TBI= Traumatic brain injury.

Table 3: Summary of trials

	Number of CONSORT items met (max 22)	Intervention	Number entering	Number at follow-up	Follow-up length	Only participants with MTBI	MTBI definition	Benefit shown
Relander 1972	7	IER	178	59	1 year	Yes	Excluded those requiring neurosurgery	No
Hinkle 1986	7	IER	1092	241	3 months	Yes	GCS 13-15, some alteration of consciousness	Yes
Alves 1993	7	IER	1710	587	1 year	Yes	GCS 13-15, PTA <24hours	No
Mittenberg 1996	11	CBT	58	No info	6 months	Yes	GCS 13-15, PTA <24hours	Yes
Wade 1997	16	IER	1136	478	6 months	No	Head injury of any severity	No
Paniak 1998	11	IER	119	111	3 months	Yes	ACR 1993 MTBI definition	No
Wade 1998	17	IER	314	218	6 months	No	Head injury requiring admission	Yes
Paniak 2000	11	IER	119	105	1 year	Yes	ACRM 1993 MTBI definition	No
HannaPladdy2001	8	Relaxation	88	88	None	Yes	Self-report closed head injury, PTA <24hours	Yes
McMillan 2002	10	Mindfulness	145	110	1 year	No	None	No
Ponsford 2002	10	IER	262	202	3 months	Yes	Trauma to head, LOC <30mins, PTA <24hours	Yes
Rath 2005	8	Rehab	60	31	6 months	No	Geffen 1998 classification	No
Hodgson 2005	11	CBT	16	12	1 month	No	ABI at least 12 months previously	Yes
Tiersky 2005	19	CBT	29	18	3 months	No	ACRM 1993 definition	Yes
Ghaffar 2006	14	IER	191	170	6 months	Yes	ACRM 1993 definition	No
Elgmark 2007	19	IER	395	355	1 year	Yes	ACRM 1993 definition	No
Owensworth 2008	14	Rehab	35	31	3 months	No	ABI convenience sample	No

Abbreviations

ABI= Acquired brain injury; ACRM= American Congress of Rehabilitation Medicine; CBT= Cognitive-behavioural therapy; GCS= Glasgow coma scale; IER= Information, education and reassurance; LOC= Loss of consciousness; PTA= Post traumatic amnesia

REFERENCES

1. Cassidy JD, Carroll LJ, Peloso PM, Borg J, von Holst H, Holm L, et al. Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med* 2004; Feb;(43 Suppl):28-60.
2. King NS. Post-concussion syndrome: Clarity amid the controversy?. *British Journal of Psychiatry* 2003; 01 Oct;183(OCT):276-8.
3. Ryan LM, Warden DL. Post concussion syndrome. *International Review of Psychiatry* 2003; Nov;15(4):310-6.
4. Carroll LJ, Cassidy JD, Peloso PM, Borg J, von Holst H, Holm L, et al. Prognosis for mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med* 2004; Feb;(43 Suppl):84-105.
5. Doezema D, King JN, Tandberg D, Espinosa MC, Orrison WW. Magnetic resonance imaging in minor head injury. *Ann Emerg Med* 1991; Dec;20(12):1281-5.
6. Carroll LJ, Cassidy JD, Holm L, Kraus J, Coronado VG, WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. Methodological issues and research recommendations for mild traumatic brain injury: the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med* 2004; Feb;(43 Suppl):113-25.
7. Wessely S, Nimnuan C, Sharpe M. Functional somatic syndromes: one or many?[see comment]. *Lancet* 1999; Sep 11;354(9182):936-9.

8. Kroenke K, Swindle R. Cognitive-behavioral therapy for somatization and symptom syndromes: a critical review of controlled clinical trials. *Psychotherapy & Psychosomatics* 2000; Jul-Aug;69(4):205-15.
9. Comper P, Bisschop SM, Carnide N, Tricco A. A systematic review of treatments for mild traumatic brain injury. *Brain Injury* 2005; Oct;19(11):863-80.
10. Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomised trials. *Lancet* 2001; Apr 14;357(9263):1191-4.
11. Mittenberg W, Tremont G, Zielinski RE, Fichera S, Rayls KR. Cognitive-behavioral prevention of postconcussion syndrome. *Archives of Clinical Neuropsychology* 1996;11(2):139-45.
12. Hodgson J, McDonald S, Tate R, Gertler P. A Randomised Controlled Trial of a Cognitive-Behavioural Therapy Program for Managing Social Anxiety After Acquired Brain Injury. *Brain Impairment* 2005; Dec;6(3):169-80.
13. Tiersky LA, Anselmi V, Johnston MV, Kurtyka J, Roosen E, Schwartz T, et al. A trial of neuropsychologic rehabilitation in mild-spectrum traumatic brain injury. *Archives of Physical Medicine & Rehabilitation* 2005; Aug;86(8):1565-74.
14. Relander M, Troupp H, Af Bjorkesten G. Controlled trial of treatment for cerebral concussion. *Br Med J* 1972; Dec 30;4(5843):777-9.
15. Hinkle JL, Alves WM, Rimell RW, Jane JA. Restoring social competence in minor head-injury patients. *Journal of Neuroscience Nursing* 1986; Oct;18(5):268-71.

16. Alves W, Macciocchi SN, Barth JT. Postconcussive symptoms after uncomplicated mild head injury. *J Head Trauma Rehabil* 1993;8(3):48-59.
17. Wade DT, Crawford S, Wenden FJ, King NS, Moss NE. Does routine follow up after head injury help? A randomised controlled trial. *Journal of Neurology, Neurosurgery & Psychiatry* 1997; May;62(5):478-84.
18. Paniak C, Toller-Lobe G, Durand A, Nagy J. A randomized trial of two treatments for mild traumatic brain injury. *Brain Injury* 1998; Dec;12(12):1011-23.
19. Wade DT, King NS, Wenden FJ, Crawford S, Caldwell FE. Routine follow up after head injury: a second randomised controlled trial. *Journal of Neurology, Neurosurgery & Psychiatry* 1998; Aug;65(2):177-83.
20. Paniak C, Toller-Lobe G, Reynolds S, Melnyk A, Nagy J. A randomized trial of two treatments for mild traumatic brain injury: 1 year follow-up. *Brain Injury* 2000; Mar;14(3):219-26.
21. Ponsford J, Willmott C, Rothwell A, Cameron P, Kelly AM, Nelms R, et al. Impact of early intervention on outcome following mild head injury in adults. *Journal of Neurology, Neurosurgery & Psychiatry* 2002; Sep;73(3):330-2.
22. Ghaffar O, McCullagh S, Ouchterlony D, Feinstein A. Randomized treatment trial in mild traumatic brain injury. *J Psychosom Res* 2006; Aug;61(2):153-60.
23. Elgmark Andersson E, Emanuelson I, Bjorklund R, Stalhammar DA. Mild traumatic brain injuries: the impact of early intervention on late sequelae. A randomized controlled trial. *Acta Neurochir* 2007; discussion 160; Feb;149(2):151-9.

24. Rath JF, Simon D, Langenbahn DM, Sherr RL, Diller L. Group treatment of problem-solving deficits in outpatients with traumatic brain injury: A randomised outcome study. *Neuropsychological Rehabilitation* 2003; Sep;13(4):461-88.
25. Ownsworth T, Fleming J, Shum D, Kuipers P, Strong J. Comparison of individual, group and combined intervention formats in a randomized controlled trial for facilitating goal attainment and improving psychosocial function following acquired brain injury. *J Rehabil Med* 2008; Feb;40(2):81-8.
26. Hanna-Pladdy B, Berry ZM, Bennett T, Phillips HL, Gouvier WD. Stress as a diagnostic challenge for postconcussive symptoms: sequelae of mild traumatic brain injury or physiological stress response. *Clin Neuropsychol* 2001; Aug;15(3):289-304.
27. McMillan T, Robertson IH, Brock D, Chorlton L. Brief mindfulness training for attentional problems after traumatic brain injury: A randomised control treatment trial. *Neuropsychological Rehabilitation* 2002;12(2):117-25.
28. Anson K, Ponsford J. Evaluation of a coping skills group following traumatic brain injury. *Brain Injury* 2006; Feb;20(2):167-78.
29. Gurr B, Coetzer BR. The effectiveness of cognitive-behavioural therapy for post-traumatic headaches. *Brain Injury* 2005; Jul;19(7):481-91.
30. Ouellet MC, Morin CM. Cognitive behavioral therapy for insomnia associated with traumatic brain injury: a single-case study. *Archives of Physical Medicine & Rehabilitation* 2004; Aug;85(8):1298-302.

31. Ouellet MC, Morin CM. Efficacy of cognitive-behavioral therapy for insomnia associated with traumatic brain injury: a single-case experimental design. *Archives of Physical Medicine & Rehabilitation* 2007; Dec;88(12):1581-92.
32. Williams WH, Evans JJ, Fleminger S. Neurorehabilitation and cognitive-behaviour therapy of anxiety disorders after brain injury: An overview and a case illustration of obsessive-compulsive disorder. *Neuropsychological Rehabilitation* 2003; Mar;13(1-2):133-48.
33. Miller LJ, Mittenberg W. Brief cognitive behavioral interventions in mild traumatic brain injury. *Appl Neuropsychol* 1998;5(4):172-83.
34. Rees RJ, Bellon ML. Post concussion syndrome ebb and flow: longitudinal effects and management. *Neurorehabilitation* 2007;22(3):229-42.
35. Kelly R. The post-traumatic syndrome: an iatrogenic disease. *Forensic Sci* 1975; Aug-Oct;6(1-2):17-24.
36. Minderhoud JM, Boelens ME, Huizenga J, Saan RJ. Treatment of minor head injuries. *Clinical Neurology & Neurosurgery* 1980;82(2):127-40.
37. Gronwall D. Rehabilitation programs for patients with mild head injury: Components, problems, and evaluation. *J Head Trauma Rehabil* 1986; Jun;1(2):53-62.
38. Conboy TJ, Barth J, Boll TJ. Treatment and rehabilitation of mild and moderate head trauma. *Rehabilitation Psychology* 1986;31(4):203-15.

39. Cope DN, Cole JR, Hall KM, Barkan H. Brain injury: analysis of outcome in a post-acute rehabilitation system. Part 1: General analysis. *Brain Injury* 1991; Apr-Jun;5(2):111-25.
40. Cope DN, Cole JR, Hall KM, Barkan H. Brain injury: analysis of outcome in a post-acute rehabilitation system. Part 2: Subanalyses. *Brain Injury* 1991; Apr-Jun;5(2):127-39.
41. Christensen AL, Pinner EM, Moller Pedersen P, Teasdale TW, Trexler LE. Psychosocial outcome following individualized neuropsychological rehabilitation of brain damage. *Acta Neurol Scand* 1992; Jan;85(1):32-8.
42. Ownsworth TL, McFarland K, Young RM. Self-awareness and psychosocial functioning following acquired brain injury: An evaluation of a group support programme. *Neuropsychological Rehabilitation* 2000;10(5):465-84.
43. Vickery CD, Gontkovsky ST, Wallace JJ, Caroselli JS. Group psychotherapy focusing on self-concept change following acquired brain injury: a pilot investigation. *Rehabil Psychol* 2006; Feb;51(1):30-5.
44. Geurtsen GJ, Martina JD, Van Heugten CM, Geurts ACH. A prospective study to evaluate a new residential community reintegration programme for severe chronic brain injury: The Brain Integration Programme. *Brain Injury* 2008; Jul;22(7-8):545-54.
45. Sarajuuri JM, Kaipio M, Koskinen SK, Niemela MR, Servo AR, Vilkki JS. Outcome of a comprehensive neurorehabilitation program for patients with traumatic brain injury. *Arch Phys Med Rehabil* 2005; Dec;86(12):2296-302.

46. Cicerone KD. Psychotherapy after mild traumatic brain injury: Relation to the nature and severity of subjective complaints. *J Head Trauma Rehabil* 1991;6(4):30-43.
47. Montgomery GK. A multi-factor account of disability after brain injury: implications for neuropsychological counselling. *Brain Injury* 1995; Jul;9(5):453-69.
48. Mukherjee D, Reis JP, Heller W. Women living with traumatic brain injury: social isolation, emotional functioning and implications for psychotherapy. *Women Ther* 2003;26(1/2):3-26.
49. Mateer CA, Sira CS, O'Connell ME. Putting Humpty Dumpty together again: the importance of integrating cognitive and emotional interventions. *J Head Trauma Rehabil* 2005; Jan-Feb;20(1):62-75.
50. Coetzer R. Psychotherapy following traumatic brain injury: integrating theory and practice. *J Head Trauma Rehabil* 2007; Jan-Feb;22(1):39-47.
51. Arco L. Neurobehavioural treatment for obsessive-compulsive disorder in an adult with traumatic brain injury. *Neuropsychological Rehabilitation* 2008; Jan;18(1):109-24.
52. Cicerone KD, Smith LC, Ellmo W, Mangel HR, Nelson P, Chase RF, et al. Neuropsychological rehabilitation of mild traumatic brain injury. *Brain Injury* 1996; Apr;10(4):277-86.
53. Bedard M, Felteau M, Mazmanian D, Fedyk K, Klein R, Richardson J, et al. Pilot evaluation of a mindfulness-based intervention to improve quality of life among

individuals who sustained traumatic brain injuries. *Disability & Rehabilitation* 2003; Jul 8;25(13):722-31.

54. Mittenberg W, Canyock EM, Condit D, Patton C. Treatment of post-concussion syndrome following mild head injury. *Journal of Clinical & Experimental Neuropsychology: Official Journal of the International Neuropsychological Society* 2001; Dec;23(6):829-36.

55. Sharpe M, Wilks D. Fatigue. *BMJ* 2002; Aug 31;325(7362):480-3.

56. National Institute for Health and Clinical Excellence (NICE). *Chronic fatigue syndrome/myalgic encephalomyelitis (or encephalopathy): diagnosis and management of chronic fatigue syndrome/myalgic encephalomyelitis (or encephalopathy) in adults and children*. (NICE guideline CG53) ed. London: Royal College of General Practitioners; 2007.