



HAL
open science

Estimating the closed skew-normal distributions parameters using weighted moments

C. Flecher, P. Naveau, D. Allard

► **To cite this version:**

C. Flecher, P. Naveau, D. Allard. Estimating the closed skew-normal distributions parameters using weighted moments. *Statistics and Probability Letters*, 2009, 79 (19), pp.1977. 10.1016/j.spl.2009.06.004 . hal-00573468

HAL Id: hal-00573468

<https://hal.science/hal-00573468>

Submitted on 4 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Estimating the closed skew-normal distributions parameters using weighted moments

C. Flecher, P. Naveau, D. Allard

PII: S0167-7152(09)00216-8
DOI: 10.1016/j.spl.2009.06.004
Reference: STAPRO 5445

To appear in: *Statistics and Probability Letters*

Received date: 8 September 2008
Revised date: 4 June 2009
Accepted date: 4 June 2009

Please cite this article as: Flecher, C., Naveau, P., Allard, D., Estimating the closed skew-normal distributions parameters using weighted moments. *Statistics and Probability Letters* (2009), doi:10.1016/j.spl.2009.06.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



* Manuscript

[Click here to view linked References](#)

Estimating the closed skew-normal distributions parameters using weighted moments

by C. Flecher^{1,2}, P. Naveau¹ and D. Allard²

¹ Laboratoire des Sciences du Climat et de l'Environnement,
CNRS Gif-sur-Yvette, France

² Biostatistics and Spatial Processes,
INRA, Agroparc 84914 Avignon, France

June 4, 2009

Abstract

Skewness is often present in a wide range of applied problems. One possible approach to model this skewness is based on the class of skew normal distributions. Fitting such distributions remains an inference challenge in various cases. In this paper, we propose and study novel estimators using weighted moments for the closed multivariate skew-normal distribution.

1 Introduction

Multivariate skew-normal distributions are extensions of the normal distribution which admit skewness whilst retaining most of the interesting properties of the Gaussian distri-

bution. The book edited by Genton (2004) provides an overview of theoretical and applied developments related to skewed distributions. More recently, a special issue in the journal *Communications in statistics* edited by Pewsey and González-Farías (2007) was centered on the different aspects of skew distributions. In the large class of skew distributions, we focus on the multivariate closed skew-normal distribution proposed by González-Farías *et al.* (2004). Such a distribution has many interesting properties inherited from the Gaussian distribution. Concerning its definition, an k -dimensional random vector \mathbf{Y} is said to have a multivariate closed skew-normal distribution, denoted by $\text{CSN}_{k,l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta})$, if its density function is of the form

$$f_{k,l}(\mathbf{y}) = c_l \phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) \Phi_l(\mathbf{D}^t(\mathbf{y} - \boldsymbol{\mu}); \boldsymbol{\nu}, \boldsymbol{\Delta}), \text{ with } c_l^{-1} = \Phi_l(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}), \quad (1)$$

where $\boldsymbol{\mu} \in \mathbb{R}^k$, $\boldsymbol{\nu} \in \mathbb{R}^l$, $\boldsymbol{\Sigma} \in \mathbb{R}^{k \times k}$ and $\boldsymbol{\Delta} \in \mathbb{R}^{l \times l}$ are both covariance matrices, $\mathbf{D} \in \mathbb{R}^{k \times l}$, $\phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $\Phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma})$ are the probability distribution function (pdf) and cumulative distribution function (cdf), respectively, of the k -dimensional normal distribution with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$, and \mathbf{D}^t is the transpose of the matrix \mathbf{D} . If $\mathbf{D} = \mathbf{0}$, the density (1) reduces to the multivariate normal one. When $l = 1$, the density of Azzalini (2005) is obtained, i.e. the variable \mathbf{Y} follows a $\text{CSN}_{k,1}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\alpha}, 0, 1)$ distribution, where $\boldsymbol{\alpha}$ is a vector of length k .

Concerning the inference of the Azzalini skew normal distribution parameters, Azzalini and Capitanio (1999) studied the classical maximum likelihood estimation (mle) approach. In sections 5 and 6 of their paper, these authors noticed that they are numerous statistical issues with the mle procedure. They wrote that *an alternative estimation method is called for*. This is one motivation for this work.

One of the advantages of parametrization (1) resides in the additive stability of the distribution. González-Farías *et al.* (2004) proved the closure under linear transformation of independent CSN random vectors. In particular, the sum of independent CSN vectors is still a CSN vector.

They also derived the moments of \mathbf{Y} . The first moment equals to

$$\mathbb{E}(\mathbf{Y}) = \boldsymbol{\mu} + \frac{\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}, \quad (2)$$

where

$$\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}) = \sum_{i=1}^k \sum_{j=1}^l (\mathbf{D}^t \boldsymbol{\Sigma})_{ij} \Phi_l^{\{j\}}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}) \mathbf{e}_i, \quad (3)$$

and \mathbf{e}_i is a $(k \times 1)$ vector with one in the i^{th} position and zero elsewhere, and

$$\Phi_l^{\{j\}}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}) = \phi([\mathbf{D}^t \boldsymbol{\mu}]_j; \boldsymbol{\nu}_j, [\boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}]_{jj}) \Phi_{l-1}([\mathbf{D}^t \boldsymbol{\mu}]_{-j}; \boldsymbol{\nu}_{-j}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D} | [\mathbf{D}^t \boldsymbol{\mu}]_j).$$

where $[\mathbf{D}^t \boldsymbol{\mu}]_j$ is the j^{th} element of $\mathbf{D}^t \boldsymbol{\mu}$ and $[\mathbf{D}^t \boldsymbol{\mu}]_{-j}$ is the vector $\mathbf{D}^t \boldsymbol{\mu}$ without the j^{th} element. The variance can be expressed as

$$\text{Var}(\mathbf{Y}) = \boldsymbol{\Sigma} + \frac{\Phi_l^{**}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} - \mathbb{E}(\mathbf{Y} - \boldsymbol{\mu}) \mathbb{E}(\mathbf{Y} - \boldsymbol{\mu})^t, \quad (4)$$

where

$$\Phi_l^{**}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}) = \frac{\partial}{\partial \mathbf{t} \partial \mathbf{t}^t} \Phi_l(\mathbf{D}^t(\boldsymbol{\mu} + \boldsymbol{\Sigma} \mathbf{t}); \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}) |_{\mathbf{t}=\mathbf{0}}. \quad (5)$$

These expressions can quickly become cumbersome in practice. The main reason for this difficulty is the absence of analytical representations for $\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})$ and its derivatives, even when $\boldsymbol{\mu} = \mathbf{0}$. To apply the CSN to real case studies, the practitioner needs to be able to easily implement an estimation method that provides accurate estimators of the CSN parameters. In its most general form, it is unlikely that such an objective can be reached for the CSN because of its large number of parameters. This aim can be achieved in some specific cases, basically when (2) and (4) can be simplified. Such cases still can offer an added flexibility and skewness with respect to the classical Gaussian distribution. To reach this goal in such cases our strategy is to develop a novel Method of Moments (MOM) approach that will be described in Section 2.

Before closing this section, we derive a lemma that will help us to compute the mean and variance of a CSN. All our proofs can be found in the Appendix.

Lemma 1. *Let $\Phi_l(\cdot; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})$ be a l -dimensional Gaussian cdf with mean $\boldsymbol{\nu} \in \mathbb{R}^l$ and covariance matrix $\boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D}$. Then,*

$$\frac{\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} = \frac{\Phi_l^*(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}. \quad (6)$$

and

$$\frac{\Phi_l^{**}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} = \frac{\Phi_l^{**}(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}, \quad (7)$$

where $\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})$ and $\Phi_l^{**}(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})$ are defined by (3) and (5), respectively.

In practice, this lemma indicates that, although $\boldsymbol{\mu}$ has still to be estimated, it can be set equal to zero in the ratios (6) and (7).

2 Our inference method

The key concept for any MOM approach is to be able to explicitly express moments in terms of the unknown parameters. From a CSN vector, there are at least three quantities to estimate: the covariance matrix, the mean vector and the skewness parameters. Hence, working with the first two moments does not provide enough equations. A possible path is to look at higher moments but this strategy is associated with at least two drawbacks. The mathematical expression of the third and fourth CSN moments are too complex to be of any practical use. In addition, the estimation of the third and fourth CSN moments are classically tainted by large variances. To avoid these hurdles, we modify a MOM approach used in Extreme Value Theory.

Probability Weighted Moments are a class of MOM estimators introduced by Hosking *et al.* (1985). Such approaches have been extended by Diebolt *et al.* (2008). The basic idea is to compute and estimate moments of the type $\mathbb{E}(Z^s F^r(Z))$ where F corresponds to the cumulative distribution function of the univariate random variable Z and r, s are

integers, usually equal to one or two. For the extreme value distributions, such quantities have explicit analytical expressions (e.g. Hosking *et al.*, 1985; Diebolt *et al.*, 2008). In the case of the CSN, the computation of these so-called Probability Weighted Moments seems very difficult to derive because the “weight” $F^r(Z)$ is very complex for a CSN. However, replacing $F^r(\cdot)$ by the cumulative Gaussian distribution $\Phi^r(\cdot; \mu, \sigma^2)$ greatly simplifies the problem. This is even true for the multivariate case. The following proposition summarizes our findings on this topic.

Proposition 2. *Let \mathbf{Y} be a $CSN_{k,l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}, \mathbf{0}, \boldsymbol{\Delta})$ defined by (1) and $h(\mathbf{y}) = h(y_1, \dots, y_k)$ be any real valued function such that $\mathbb{E}(h(\mathbf{Y}))$ is finite, then*

$$\mathbb{E}(h(\mathbf{Y})\Phi_k^r(\mathbf{Y}; \mathbf{0}, \mathbf{I}_k)) = \frac{\Phi_{rk+l}(\mathbf{0}; \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+ + \mathbf{D}_+^t \boldsymbol{\Sigma} \mathbf{D}_+)}{\Phi_l(\mathbf{0}; \mathbf{0}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} \mathbb{E}(h(\mathbf{Y}_+)), \quad (8)$$

where \mathbf{Y}_+ follows a $CSN_{k,rk+l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}_+, \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+)$ with $\boldsymbol{\Delta}_+ = \begin{bmatrix} \mathbf{I}_{rk} & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\Delta} \end{bmatrix}$, $\mathbf{D}_+ =$

$\begin{bmatrix} \mathbf{D}_*^t \\ \mathbf{D}^t \end{bmatrix}^t$, \mathbf{D}_* a $k \times rk$ matrix defined by $\mathbf{D}_* = \begin{bmatrix} \mathbf{I}_k \\ \dots \\ \mathbf{I}_k \end{bmatrix}^t$ with \mathbf{I}_k the identity matrix of size k , and $\boldsymbol{\nu}_+$ a $rk + l$ vector defined by $\boldsymbol{\nu}_+^t = (-\boldsymbol{\mu}, \dots, -\boldsymbol{\mu}, \mathbf{0}_l)$.

To understand Equation (8) we consider two special cases. First, suppose that $r = 1$ and $h(y_1, \dots, y_k) = y_i$ for some $i = 1, \dots, k$. Then, using Lemma 1 and Equation (2) within (8), we can write

$$\mathbb{E}(Y_i \Phi_k(\mathbf{Y}; \mathbf{0}, \mathbf{I}_k)) = \mu_i \frac{\Phi_{k+l}(\mathbf{0}; \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+ + \mathbf{D}_+^t \boldsymbol{\Sigma} \mathbf{D}_+)}{\Phi_l(\mathbf{0}; \mathbf{0}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} + \frac{[\Phi_{k+l}^*(\mathbf{0}; \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+ + \mathbf{D}_+^t \boldsymbol{\Sigma} \mathbf{D}_+)]_i}{\Phi_l(\mathbf{0}; \mathbf{0}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}. \quad (9)$$

The left hand side of (9) can be viewed as a type of weighted moments. Concerning the right hand side, it basically means that, if \mathbf{D} , $\boldsymbol{\Sigma}$ and $\boldsymbol{\Delta}$ are chosen such as the first moment (2) and second moment (4) have explicit expressions or can be reasonably approximated,

then the weight moment defined by (9) can also be derived. In other words, if one can compute or approximate the first and second moments expressions, then a weighted moment can also be obtained and a MOM approach with a third equation can be implemented.

Our second case concerns the bivariate sub vector $(Y_i, Y_j)^t$ for some $i, j = 1, \dots, k$ that follows a $CSN_{2,l}(\mathbf{0}, \mathbf{\Sigma}_{ij}, \mathbf{D}_{ij}, \mathbf{0}, \mathbf{\Delta}_{ij})$ with obvious notations for the matrices. To apply Proposition 2 we choose $h(y_i, y_j) = y_i y_j$ and $r = 2$. If $\Phi(x)$ denotes a shortcut notation for the standardized cumulative Gaussian distribution $\Phi_1(x; 0, 1)$, we can then write

$$\mathbb{E}(Y_i Y_j \Phi(Y_i) \Phi(Y_j)) = \frac{\Phi_{4+l}(\mathbf{0}; \mathbf{0}, \mathbf{\Delta}_{+ij} + \mathbf{D}_{+ij}^t \mathbf{\Sigma} \mathbf{D}_{+ij})}{\Phi_l(\mathbf{0}; \mathbf{0}, \mathbf{\Delta}_{ij} + \mathbf{D}_{ij}^t \mathbf{\Sigma}_{ij} \mathbf{D}_{ij})} \mathbb{E}(Y_{+i} Y_{+j}), \quad (10)$$

where $(Y_{+i}, Y_{+j})^t$ follows a $CSN_{2,2+l}(\mathbf{0}, \mathbf{\Sigma}_{ij}, \mathbf{D}_{+ij}, \mathbf{0}, \mathbf{\Delta}_{+ij})$ and these elements can be obtained from (4). Equation (10) will be useful to derive the variance properties of our proposed estimators. In Equation (10) we took $(\mu_i, \mu_j) = (0, 0)$. When this assumption does not hold, we can take advantage of Lemma 1.

If $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ are both assumed to be unknown, then they cannot be identified in (1) because the expression $\Phi_l(\mathbf{D}^t(\mathbf{y} - \boldsymbol{\mu}); \boldsymbol{\nu}, \mathbf{\Delta})$ is over-parametrized in this case. Following González-Farías *et al.* (2004) we decide to set $\boldsymbol{\nu} = \mathbf{0}$ in the rest of the paper. In our applications, we also suppose that the matrix $\mathbf{\Delta}$ is a function of the other three unknown parameters $\boldsymbol{\mu}$, $\mathbf{\Sigma}$ and \mathbf{D} . Conceptually, it may be possible to deal with the estimation of $\mathbf{\Delta}$ by working with a moment of the type $\mathbb{E}(Y_i \Phi_k^2(\mathbf{Y}; \mathbf{0}, \mathbf{I}_k))$. But it becomes burdensome in terms of notations and complicated in terms of optimization.

3 Examples and estimation procedures

3.1 The univariate case

Assume that X_1, \dots, X_n are n independent and identically distributed random variables(iid) $CSN_{1,1}(\mu, \sigma^2, \delta, 0, \Delta)$ with μ, σ^2 and δ unknown. Define $\lambda = \sigma\delta/\sqrt{\Delta + \sigma^2\delta^2}$, the first and

second moment expressions are easily derived from (2) and (4) with $k = l = 1$

$$\mathbb{E}(X_i) = \mu + 2\Phi_1^*(0; 0, \Delta + \delta^2\sigma^2) = \mu + \frac{2}{\sqrt{2\pi}}\sigma\lambda,$$

and

$$\text{Var}(X_i) = \sigma^2 + 2\Phi_1^{**}(0; 0, \Delta + \delta^2\sigma^2) - (2\Phi_1^*(0; 0, \Delta + \delta^2\sigma^2))^2 = \sigma^2 \left(1 - \frac{2}{\pi}\lambda^2\right).$$

A direct application of Equation (9) gives

$$\mathbb{E}(X_i\Phi(X_i)) = 2\mu \Phi_2(\mathbf{0}; \nu_{2+}, \mathbf{A}_2) + 2\Phi_2^*(\mathbf{0}; \nu_{2+}, \mathbf{A}_2) \quad (11)$$

where $\Phi_2^*(\mathbf{0}; \nu_{2+}, \mathbf{A}_2) = \sigma^2 \left\{ \Phi_2^{\{1\}}(\mathbf{0}; \nu_{2+}, \mathbf{A}_2) + \delta \Phi_2^{\{2\}}(\mathbf{0}; \nu_{2+}, \mathbf{A}_2) \right\}$ and

$$\nu_{2+} = \begin{bmatrix} -\mu \\ 0 \end{bmatrix}, \quad \mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \text{and} \quad \mathbf{A}_2 = \begin{bmatrix} 1 + \sigma^2 & \sigma^2\delta \\ \sigma^2\delta & \Delta + \sigma^2\delta^2 \end{bmatrix}. \quad (12)$$

In terms of estimation, the sample mean denoted $\bar{X}_n = (X_1 + \dots + X_n)/n$ and the sample variance $s_n^2 = \sum_{i=1}^n (X_i - \bar{X}_n)^2 / (n-1)$ can be plugged in the above moment expressions. The weighted moment $m = \mathbb{E}(X_i\Phi(X_i))$ can be estimated by the unbiased statistic $m_n = n^{-1} \sum_{i=1}^n X_i\Phi(X_i)$. The following system of three equations with unknowns $\hat{\mu}$, $\hat{\delta}$ and $\hat{\sigma}$ follows

$$\begin{cases} \bar{X}_n = \hat{\mu} + \sqrt{\frac{2}{\pi}}\hat{\sigma}\hat{\lambda} \\ s_n^2 = \hat{\sigma}^2 \left(1 - \frac{2}{\pi}\hat{\lambda}^2\right) \\ m_n = 2\hat{\mu} \Phi_2(\mathbf{0}; \hat{\nu}_{2+}, \hat{\mathbf{A}}_2) + 2\Phi_2^*(\mathbf{0}; \hat{\nu}_{2+}, \hat{\mathbf{A}}_2) \end{cases}, \quad (13)$$

where $\hat{\nu}_{2+}$ and $\hat{\mathbf{A}}_2$ are defined by (12). Concerning the asymptotic properties of our moments, they can be described by the following proposition.

Proposition 3. *Assume that X_1, \dots, X_n are n iid $CSN_{1,1}(\mu, \sigma^2, \delta, 0, \Delta)$ variables where μ, σ^2 and δ are unknown and define $\lambda = \sigma\delta/\sqrt{\Delta + \sigma^2\delta^2}$. The unbiased estimator vector $\sqrt{n}(\bar{X}_n - \mathbb{E}(X), s_n^2 - \text{Var}(X), m_n - \mathbb{E}(X\Phi(X)))^t$ converges in distribution, as n goes to*

infinity, to a zero-mean Gaussian vector with a covariance matrix whose elements c_{ij} equal

$$\begin{aligned}
c_{1,1} &= \sigma^2 \left(1 - \frac{2}{\pi} \lambda^2 \right), \\
c_{1,2} &= \sigma^3 \frac{2}{\sqrt{2\pi}} \left(\frac{4}{\pi} - 1 \right) \lambda^3, \\
c_{2,2} &= \sigma^4 \left(2 - \frac{16}{2\pi} \lambda^2 + \frac{16}{\pi} \left(\frac{1}{2} - \frac{1}{\pi} \right) \lambda^4 \right), \\
c_{3,3} &= \mathbb{E}(X^2 \Phi^2(X)) - \mathbb{E}^2(X \Phi(X)), \\
c_{1,3} &= c_{3,1} = \mathbb{E}(X^2 \Phi(X)) - \mathbb{E}(X) \mathbb{E}(X \Phi(X)), \\
c_{2,3} &= c_{3,2} = \mathbb{E}(X^3 \Phi(X)) - \text{Var}(X) \mathbb{E}(X \Phi(X)) - 2\mathbb{E}(X) \mathbb{E}(X^2 \Phi(X)) + \mathbb{E}^2(X) \mathbb{E}(X \Phi(X)),
\end{aligned}$$

where $\mathbb{E}(X^p)$, $p \geq 0$ can be found in Martinez (2008) and

$$\begin{aligned}
\mathbb{E}(X^2 \Phi^2(X)) &= 2(\sigma^2 + \mu^2) \Phi_3(\mathbf{0}; \boldsymbol{\nu}_{3+}, \mathbf{A}_3) + 4\mu \Phi_3^*(\mathbf{0}; \boldsymbol{\nu}_{3+}, \mathbf{A}_3) + 2\Phi_3^{**}(\mathbf{0}; \boldsymbol{\nu}_{3+}, \mathbf{A}_3), \\
\mathbb{E}(X^2 \Phi(X)) &= 2\mu \Phi_2(\mathbf{0}; \boldsymbol{\nu}_{2+}, \mathbf{A}_2) + 2\Phi_2^*(\mathbf{0}; \boldsymbol{\nu}_{2+}, \mathbf{A}_2), \\
\mathbb{E}(X^3 \Phi(X)) &= 2\Phi_2(\mathbf{0}; \boldsymbol{\nu}_{2+}, \mathbf{A}_2) \mathbb{E}(X_+^3),
\end{aligned}$$

with $\boldsymbol{\nu}_{2+}$ and \mathbf{A}_2 defined by (12),

$$\boldsymbol{\nu}_{3+} = \begin{bmatrix} -\mu \\ -\mu \\ 0 \end{bmatrix}, \text{ and } \mathbf{A}_3 = \begin{bmatrix} 1 + \sigma^2 & \sigma^2 & \sigma^2 \delta \\ \sigma^2 & 1 + \sigma^2 & \sigma^2 \delta \\ \sigma^2 \delta & \sigma^2 \delta & \Delta + \sigma^2 \delta^2 \end{bmatrix}.$$

To assess our MOM estimation scheme with simulations we focus on the Skew-Normal model described by Azzalini and Capitanio (1999) and denoted $SN(\mu, \sigma, \alpha)$. The latter corresponds to a $CSN_{1,1}(\mu, \sigma^2, \lambda\sigma^{-1}, 0, 1 - \lambda^2)$ for some $0 \leq |\lambda| < 1$ where $\lambda = \alpha/\sqrt{1 + \alpha^2}$.

In this case, the system (13) can be written as

$$\begin{cases} \bar{X}_n = \hat{\mu} + \sqrt{\frac{2}{\pi}} \hat{\sigma} \hat{\lambda} \\ s_n^2 = \hat{\sigma}^2 \left(1 - \frac{2}{\pi} \hat{\lambda}^2 \right) \\ m_n = 2\hat{\mu} \Phi_2(\mathbf{0}; \hat{\boldsymbol{\nu}}_+, \hat{\mathbf{A}}_2) + 2\Phi_2^*(\mathbf{0}; \hat{\boldsymbol{\nu}}_+, \hat{\mathbf{A}}_2) \end{cases}, \quad (14)$$

Figure 1: Density of $CSN_{1,1}(0, 1, \lambda, 0, 1 - \lambda^2)$ for λ in $\{0, 0.2, 0.71, 0.89, 0.97\}$.

where $\Phi_2(\mathbf{0}; \hat{\boldsymbol{\nu}}_+, \hat{\mathbf{A}}_2) = \Phi_2\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}; \begin{bmatrix} -\frac{\hat{\mu}}{\sqrt{1+\hat{\sigma}^2}} \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & \hat{\lambda}\frac{\hat{\sigma}}{\sqrt{1+\hat{\sigma}^2}} \\ \hat{\lambda}\frac{\hat{\sigma}}{\sqrt{1+\hat{\sigma}^2}} & 1 \end{bmatrix}\right)$ and the probability $\Phi_2^*(\mathbf{0}; \hat{\boldsymbol{\nu}}_+, \hat{\mathbf{A}}_2)$ equals

$$\hat{\sigma}^2 \phi(\hat{\mu}; 0, 1 + \hat{\sigma}^2) \Phi\left(-\frac{\hat{\sigma}}{1 + \hat{\sigma}^2} \hat{\lambda} \hat{\mu}; 0, \frac{1 + \hat{\sigma}^2(1 - \hat{\lambda}^2)}{1 + \hat{\sigma}^2}\right) + \frac{\hat{\sigma} \hat{\lambda}}{\sqrt{2\pi}} \Phi\left(\hat{\mu}; 0, 1 + \hat{\sigma}^2(1 - \hat{\lambda}^2)\right).$$

For our simulations, 1000 samples of sizes $n = 50$ and $n = 100$ are generated from a $CSN_{1,1}(0, 1, \lambda, 0, 1 - \lambda^2)$ with λ in $\{0.20, 0.71, 0.89, 0.97\}$. These values correspond to Azzalini's Skew-Normal shape parameters α in $\{0.2, 1, 2, 4\}$. Table 1 compares the Mean Squared Errors for each parameter obtained from our weighted moment approach with the ones derived with the mle method studied by Azzalini and Capitanio (1999). The lowest MSE value for the two methods is represented in bold letters. This table indicates that our approach performs better than the mle for the small sample size $n = 50$, especially for the parameter σ . As the sample size increases, both methods improve and become comparable (although the mle has still difficulties for σ). In contrast to larger λ 's (see Table 1), the weighted moments and maximum likelihood methods have both difficulties to estimate values of λ near zero. For the latter estimation procedure, this difficulty can be explained by the singularity of the Information of Fisher matrix at $\lambda = 0$. For the weighted moment method, the asymptotic variance of \bar{X}_n and s_n^2 is maximal for $\lambda = 0$. The covariance matrix of the moments does not present any singularity when $\lambda = 0$. As an example, for $\mu = 0$, $\sigma = 1$ and $\lambda = 0$ its determinant is equal to $0.00808 > 0$.

Another way to look at the poor estimators performance for small values of λ is to display the density distributions for various λ 's, see Figure 1 below. The black and blue lines correspond to $\lambda = 0$ (Gaussian) and $\lambda = 0.2$, respectively. Except if the sample size is very large, it is extremely difficult to detect any skewness from a sample drawn from a skew-normal with $\lambda = 0.2$. In comparison, the skewness becomes more visible for $\lambda > 0.7$.

Table 1: Mean Squared Error computed from 1000 samples of sizes $n = 50$ and $n = 100$ generated from a $CSN_{1,1}(0, 1, \lambda, 0, 1 - \lambda^2)$ with λ in $\{0.20, 0.71, 0.89, 0.97\}$

To obtain the asymptotic distribution of our estimators, the classical approach is to apply the delta method in conjunction with Proposition 3 that provides the trivariate-normal asymptotic distribution of our triplet of moments. Still, the complexity of those weighted moment expressions makes it impossible to derive explicit formulas of the transformed estimators via the delta-method.

The covariance matrix of the moments established in Proposition 3 allows us to build approximate confidence intervals using a parametric bootstrap approach (Efron and Tibshirani, 1993), briefly outlined as follows. From proposition 3, sample N repetition of triplets of moments $(\bar{X}_n(i), s_n^2(i), m_n(i))$, $i = 1, \dots, N$ using the estimated parameters $(\hat{\mu}, \hat{\sigma}, \hat{\lambda})$. For each triplet, solve (13) to find a triplet $(\tilde{\mu}(i), \tilde{\sigma}(i), \tilde{\lambda}(i))$. From these one obtains an approximate variance-covariance matrix associated to $(\hat{\mu}, \hat{\sigma}, \hat{\lambda})$ as well as confidence intervals. For example we computed the confidence interval for the case $\lambda = 0.89$ and $n = 100$. With $N = 10000$, the 90% centered confidence interval was estimated to be $[0.685, 0.968]$.

3.2 A multivariate example

Following our univariate example, we study the following k -dimensional multivariate distribution $CSN_{k,k}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \lambda \boldsymbol{\Sigma}^{-1/2}, 0, (1 - \lambda^2) \mathbf{I}_k)$, where $\boldsymbol{\mu}$ is a location vector of size k , $\boldsymbol{\Sigma}$ a $k \times k$ variance covariance matrix and λ a scalar in $] - 1, 1[$. Such a parametrization corresponds to $\boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D} = \mathbf{I}_k$ and the moments are

$$\mathbb{E}(\mathbf{Y}) = \boldsymbol{\mu} + \frac{2}{\sqrt{2\pi}} \lambda \boldsymbol{\Sigma}^{1/2} \mathbf{1}_k, \quad \text{Var}(\mathbf{Y}) = \boldsymbol{\Sigma} \left(1 - \frac{2}{\pi} \lambda^2 \right) \text{ and}$$

$$\mathbb{E}(\Phi_k(\mathbf{Y}, \mathbf{0}, \mathbf{I}_k)) = 2^k \Phi_{2k} \left(\mathbf{0}; \begin{bmatrix} -\boldsymbol{\mu} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma} + \mathbf{I}_k & \lambda \boldsymbol{\Sigma}^{1/2} \\ \lambda \boldsymbol{\Sigma}^{1/2} & \mathbf{I}_k \end{bmatrix} \right)$$

Figures 2 and 3 summarize, with boxplots and histograms, our MOM estimations obtained

with 1000 samples of size 500 generated from a bivariate $CSN_{2,2}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \lambda \boldsymbol{\Sigma}^{-1/2}, \mathbf{0}, (1 - \lambda^2) \mathbf{I}_2)$ random vector with $\mu_1 = \mu_2 = 0$, $\sigma_1 = \sigma_2 = 1$, $\lambda = 0.89$ and $\rho = 0.8$. The boxplots correspond to the parameters μ and σ and the histograms to ρ and λ . These graphs indicate that our approach provides reasonable results in this bivariate case.

Figure 2: Estimated values of $\hat{\mu}_1$, $\hat{\mu}_2$, $\hat{\sigma}_1$ and $\hat{\sigma}_2$ obtained from 1000 samples of size 500 generated from a bivariate $CSN_{2,2}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \lambda \boldsymbol{\Sigma}^{-1/2}, \mathbf{0}, (1 - \lambda^2) \mathbf{I}_2)$ with $\mu_1 = \mu_2 = 0$, $\sigma_1 = \sigma_2 = 1$, $\lambda = 0.89$ and $\rho = 0.8$. Solid lines represent true values.

Figure 3: Histograms of estimated values for $\hat{\rho}$ (left) and $\hat{\lambda}$ (right) from the same simulations described in Figure 2

4 Concluding remarks

In this paper, we have introduced a new inference approach for closed skew normal distribution by modifying a MOM approach used in Extreme Value Theory. In the simulation study, we compare our estimators with mle ones in the univariate case. Our approach seems to outperform the mle for small sample sizes. As illustrated by our bivariate example, the basic principles of our method can be applied to some multivariate cases. In conclusion, our goal was not to show the superiority of our approach over the mle. Instead and as advocated by Azzalini and Capitanio (1999), we aimed at providing an alternative inference scheme for skew normal distributions. This method could be used to initialize mle algorithms or whenever the mle provides unreasonable values.

As shown in Section 3.1, the asymptotic covariance matrix of the moments allows us to build confidence intervals for the estimates using a parametric bootstrap procedure. This raises the question of the testing $\lambda = 0$ versus $\lambda \neq 0$. We showed that it was possible for high values of λ because estimates are reliable in this case. The results shown in Table 1

for small values of λ does not encourage us to recommend formal tests when the amount of skewness is low.

In this context, our advice when analyzing a sample of observations is rather to perform a simple Gaussianity test and basic visual checks (histogram, density, qq-plots, etc) before imposing a skew-normal model. If there is no strong evidence of a departure from normality, the assumption of a skew-normal distribution should be discouraged because neither mle nor our method can accurately estimate the skew-normal parameters in this instance. In other words, a skew normal model should only be assumed when some skewness is apparent in the data or when the sample size is very large. In this latter case, our method-of-moment should be performed adequately.

Acknowledgments. The authors wish to express their gratitude to the editor and referee for their very helpful comments and suggestions. This work was supported by the ANR-CLIMATOR project, the NCAR Weather and Climate Impact Assessment Science Initiative and the ANR-AssimilEx project. The authors would also like to credit the contributors of the R project.

References

- Allard, D. and Naveau, P., 2007. A new spatial skew-normal random field model. *Communication in Statistics*, **36**, 1821–1834.
- Arellano-Valle, R.B. and Azzalini, A., 2008. The centred parametrization for the multivariate skew-normal distribution. *J. Multiv. Analysis*, **99**, 1362-1382.
- Azzalini, A. and Capitanio, A., 1999. Statistical applications of the multivariate skew-normal distribution. *Journal of the Royal Statistical Society, B.*, **61, part 3**, 579–602.
- Azzalini, A., 2005. The skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, **32**, 159–188.
- Diebolt, J., A. Guillou, Naveau, P. and Ribereau, P., 2008. Improving probability-weighted

- moment methods for the generalized extreme value distribution. *REVSTAT - Statistical Journal*, **6–1**, 3350.
- Efron, B. and Tibshirani, R.J., 1993. *An Introduction to the Bootstrap*. New-York, Chapman & Hall.
- Genton, M.G. (Ed.), 2004. *Skew-Elliptical Distributions and Their Applications: A Journey Beyond Normality*. Boca Raton, FL: Chapman & Hall/CRC.
- González-Farías, G., Domínguez-Molina, J. and Gupta, A., 2004. Additive properties of skew normal random vectors. *Journal of Statistical Planning and Inference*, **126**, 521–534.
- Hosking, J.R.M., Wallis, J.R., and Wood, E.F., 1985. Estimation of the generalized extreme-value distribution by the method of probability-weighted moments, *Technometrics*, **27**, 251-261.
- Martinez, E.H., Varela, H., Gomez, H.W and Bolfarine H., 2008. A note on the likelihood and moments of the skew normal distribution. *SORT*, **32**(1), 57–66.
- Pewsey A. and González-Farías, G., 2007. Special issue preface. *Communications in Statistics - Theory and Methods*, **36–9**.

5 Appendix

Proof of Lemma 1. According to González-Farías *et al.* (2004), if \mathbf{Y} follows a $\text{CSN}_{k,l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta})$, then $\mathbf{Y} + \mathbf{c}$ follows a

$$\text{CSN}_{k,l}(\boldsymbol{\mu} + \mathbf{c}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta}) \quad (15)$$

for any real vector \mathbf{c} of length k . This implies the following equalities

$$\begin{aligned}
\frac{\Phi_l^*(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{0}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} &= \mathbb{E}(\text{CSN}_{k,l}(\mathbf{0}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta})), \text{ from (2),} \\
&= \mathbb{E}(\text{CSN}_{k,l}(\mathbf{0}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta}) + \boldsymbol{\mu}) - \boldsymbol{\mu}, \\
&= \mathbb{E}(\text{CSN}_{k,l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}, \boldsymbol{\nu}, \boldsymbol{\Delta})) - \boldsymbol{\mu}, \text{ because of (15),} \\
&= \left[\frac{\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})} + \boldsymbol{\mu} \right] - \boldsymbol{\mu}, \text{ from (2),} \\
&= \frac{\Phi_l^*(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}{\Phi_l(\mathbf{D}^t \boldsymbol{\mu}; \boldsymbol{\nu}, \boldsymbol{\Delta} + \mathbf{D}^t \boldsymbol{\Sigma} \mathbf{D})}.
\end{aligned}$$

With a similar argument, this time using (4), Equation (7) can be derived.

Proof of Proposition 2 . We write that

$$\begin{aligned}
\mathbb{E}(h(\mathbf{Y}) \Phi_k^*(\mathbf{Y}; \mathbf{0}, \mathbf{I}_k)) &= c_l \int \dots \int h(\mathbf{y}) \Phi_k^*(\mathbf{y}; \mathbf{0}, \mathbf{I}_k) \phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) \Phi_l(\mathbf{D}^t(\mathbf{y} - \boldsymbol{\mu}); \mathbf{0}, \boldsymbol{\Delta}) d\mathbf{y}, \\
&= c_l \int \dots \int h(\mathbf{y}) \phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) [\Phi_{rk}(\mathbf{D}_*^t(\mathbf{y} - \boldsymbol{\mu}); -[\boldsymbol{\mu} \dots \boldsymbol{\mu}]^t, \mathbf{I}_{rk}) \Phi_l(\mathbf{D}^t(\mathbf{y} - \boldsymbol{\mu}); \mathbf{0}, \boldsymbol{\Delta})] d\mathbf{y}, \\
&= c_l \int \dots \int h(\mathbf{y}) \phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) [\Phi_{rk+l}(\mathbf{D}_+^t(\mathbf{y} - \boldsymbol{\mu}); \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+)] d\mathbf{y}.
\end{aligned}$$

To complete the computation of $\mathbb{E}(h(\mathbf{Y}) \Phi_k(\mathbf{Y}; \mathbf{0}, \mathbf{I}_k))$, we remark that the density of \mathbf{Y}_+ follows a $\text{CSN}_{k,rk+l}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \mathbf{D}_+, \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+)$ defined in Proposition 2 is

$$f_{k,rk+l}(\mathbf{y}) = \frac{1}{\Phi_{rk+l}(\mathbf{0}; \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+ + \mathbf{D}_+^t \boldsymbol{\Sigma} \mathbf{D}_+)} \phi_k(\mathbf{y}; \boldsymbol{\mu}, \boldsymbol{\Sigma}) \Phi_{rk+l}(\mathbf{D}_+^t(\mathbf{y} - \boldsymbol{\mu}); \boldsymbol{\nu}_+, \boldsymbol{\Delta}_+).$$

The expected result follows.

Figure




Figure 1: Density of $CSN_{1,1}(0, 1, \lambda, 0, 1 - \lambda^2)$ for λ in $\{0, 0.2, 0.71, 0.89, 0.97\}$.

Figure




Figure 2: Estimated values of $\hat{\mu}_1$, $\hat{\mu}_2$, $\hat{\sigma}_1$ and $\hat{\sigma}_2$ obtained from 1000 samples of size 500 generated from a bivariate $CSN_{2,2}(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \lambda \boldsymbol{\Sigma}^{-1/2}, \mathbf{0}, (1 - \lambda^2) \mathbf{I}_2)$ with $\mu_1 = \mu_2 = 0$, $\sigma_1 = \sigma_2 = 1$, $\lambda = 0.89$ and $\rho = 0.8$. Solid lines represent true values.

Figure




Figure 3: Histograms of estimated values for $\hat{\rho}$ (left) and $\hat{\lambda}$ (right) from the same simulations described in Figure 2.

Table

Table 1: Mean Squared Error computed from 1000 samples of sizes $n = 50$ and $n = 100$ generated from a $CSN_{1,1}(0, 1, \lambda, 0, 1 - \lambda^2)$ with λ in $\{0.20, 0.71, 0.89, 0.97\}$

	Weighted Moment Method			Maximum Likelihood Method		
λ	$\hat{\mu}$	$\hat{\sigma}$	$\hat{\lambda}$	$\hat{\mu}$	$\hat{\sigma}$	$\hat{\lambda}$
$n = 50$						
0.20	0.997	0.209	0.702	0.781	0.878	0.638
0.71	0.086	0.043	0.031	0.091	0.24	0.034
0.89	0.032	0.024	0.010	0.049	0.14	0.024
0.97	0.024	0.022	0.0061	0.028	0.088	0.012
$n = 100$						
0.20	0.935	0.180	0.688	0.576	0.433	0.547
0.71	0.060	0.027	0.025	0.052	0.12	0.023
0.89	0.025	0.016	0.0092	0.030	0.071	0.015
0.97	0.013	0.012	0.0021	0.012	0.054	0.0012