



HAL
open science

A note on the equation $-\Delta u + e^u - 1 = 0$

Laurent Veron

► **To cite this version:**

| Laurent Veron. A note on the equation $-\Delta u + e^u - 1 = 0$. 2004. hal-00573805v1

HAL Id: hal-00573805

<https://hal.science/hal-00573805v1>

Preprint submitted on 4 Mar 2011 (v1), last revised 26 Oct 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A note on the equation $-\Delta u + e^u - 1 = 0$

Laurent Véron

Laboratoire de Mathématiques et Physique Théorique,
Université François Rabelais, Tours, FRANCE

Abstract If Ω is a bounded domain in \mathbb{R}^N , we study conditions on a Radon measure μ on $\partial\Omega$ for solving the equation $-\Delta u + e^u - 1 = 0$ in Ω with $u = \mu$ on $\partial\Omega$. The conditions are expressed in terms of nonlinear capacities.

2010 Mathematics Subject Classification. 35J60, 35J65, 28A12, 46E30.

Key words. Elliptic equations, Orlicz capacities, reduced measures, boundary trace

1 Introduction

Basic results

Consider the equation

$$-\Delta u + e^u - 1 = 0 \tag{1.1}$$

in Ω , a bounded domain in \mathbb{R}^N with smooth boundary $\partial\Omega = \Sigma$. Let $\rho(x) = \text{dist}(x, \Sigma)$. We denote by $\mathbb{P}^\Omega[\mu]$ the Poisson potential of a boundary measure μ , and by $\mathbb{G}^\Omega[\phi]$ the Green potential of $\phi \in \mathfrak{M}(\Omega)$. The study is initiated by M. Grillot and L. Véron¹

Boundary measures

$$\begin{aligned} -\Delta u + e^u - 1 &= 0 && \text{in } \Omega \\ u &= \mu && \text{on } \Sigma, \end{aligned} \tag{1.2}$$

with $\mu \in \mathfrak{M}_+(\Sigma)$. If the measure μ is such that (1.24) admits a solution, always unique, is it called **admissible**. The set of admissible positive boundary measures is denoted by $\mathfrak{M}_+^{exp}(\Sigma)$. Assume $\mu = \mu_S + \mu_R$ with $\mu_S \perp dH^{N-1}$, $\mu_R \ll dH^{N-1}$ then

¹Grillot M. et Véron L., *Boundary trace of solutions of the Prescribed Gaussian curvature equation*, Proc. Roy. Soc. Edinburgh **130 A**, 1-34 (2000)

Theorem 1 *Assume*

$$\exp(\mathbb{P}^\Omega[\mu_S]) \in L^1(\Omega; \rho dx) \quad (1.3)$$

then μ is admissible.

Proof. A weaker form is proved by Grillot and Véron. Moreover, the proof presented below is simpler. For $k > 0$, set $\mu_{R,k} = \inf\{k, \mu_R\}$ and denote by u_k the solution of

$$\begin{aligned} -\Delta u_k + e^{u_k} - 1 &= 0 & \text{in } \Omega \\ u_k &= \mu_S + \mu_{R,k} & \text{on } \Sigma. \end{aligned} \quad (1.4)$$

Such a solution exists because

$$\exp(\mathbb{P}^\Omega[\mu_S + \mu_{R,k}]) \leq e^k \exp(\mathbb{P}^\Omega[\mu_S])$$

by the maximum principle, and (1.3) implies that $\exp(\mathbb{P}^\Omega[\mu_S + \mu_{R,k}]) - 1 \in L^1(\Omega; \rho dx)$. The sequence u_k is nondecreasing. Since, for any $\zeta \in C_c^{1,1}(\bar{\Omega})$,

$$\int_{\Omega} (-u_k \Delta \zeta + (e^{u_k} - 1)\zeta) dx = \int_{\Sigma} \frac{\partial \zeta}{\partial \nu} d(\mu_S + \mu_{R,k}),$$

if we take in particular $\zeta = \zeta_0$ as being the solution of

$$\begin{aligned} -\Delta \zeta_0 &= 1 & \text{in } \Omega \\ \zeta_0 &= 0 & \text{on } \Sigma, \end{aligned} \quad (1.5)$$

we get

$$\int_{\Omega} (u_k + (e^{u_k} - 1)\zeta_0) dx = - \int_{\Sigma} \frac{\partial \zeta_0}{\partial \nu} d(\mu_S + \mu_{R,k}) \leq c \|\mu\|_{\mathfrak{M}}. \quad (1.6)$$

Thus $u = \lim_{k \rightarrow \infty} u_k$ is integrable,

$$\int_{\Omega} (u + (e^u - 1)\zeta_0) dx \leq c \|\mu\|_{\mathfrak{M}},$$

and the convergence of u_k and e^{u_k} to u and e^u hold respectively in $L^1(\Omega)$ and $L^1(\Omega; \rho dx)$. It follows that u satisfies (1.24). \square

The proof of the next result, inspired of Brezis, Marcus and Ponce ², is easy

Theorem 2 *The following properties hold:*

(i) *If $\mu \in \mathfrak{M}_+^{exp}(\Sigma)$ and $0 \leq \tilde{\mu} \leq \mu$, then $\tilde{\mu} \in \mathfrak{M}_+^{exp}(\Sigma)$.*

²Brezis H., Marcus M. and Ponce A., *Nonlinear elliptic equations with measures revisited*, preprint (2004).

(ii) Let $\{\mu_n\} \subset \mathfrak{M}_+^{exp}(\Sigma)$ be an increasing sequence which converges to μ in the weak sense of measures. Then $\mu \in \mathfrak{M}_+^{exp}(\Sigma)$.

(iii) $\mathfrak{M}_+^{exp}(\Sigma) + L_+^1(\Sigma) = \mathfrak{M}_+^{exp}(\Sigma)$.

Proof. I- Let $u = u_\mu$ be the solution of (1.24) and $w = \inf\{u, \mathbb{P}^\Omega[\tilde{\mu}]\}$. Since $\mathbb{P}^\Omega[\tilde{\mu}]$ is a supersolution for (1.1), w is a supersolution too. Furthermore w is nonnegative and $e^w - 1 \in L^1(\Omega; \rho dx)$. By Doob's theorem w admits a boundary trace $\mu^* \in \mathfrak{M}_+(\Sigma)$ and $\mu^* \leq \tilde{\mu} \leq \mu$. Let w^* be the solution of

$$\begin{aligned} -\Delta w^* + e^u - 1 &= 0 & \text{in } \Omega \\ w^* &= \tilde{\mu} & \text{on } \Sigma. \end{aligned}$$

then $u \geq w \geq w^*$ and ³,

$$\lim_t \int_{\Sigma_t} w^*(t, \cdot) \eta dS_t = \int_{\Sigma} \eta d\tilde{\mu} \quad \forall \eta \in C(\Sigma),$$

(here we denote by Σ_t the set of $x \in \Omega$ such that $\rho(x) = t > 0$). This implies that the boundary trace of w^* is $\tilde{\mu}$ and thus $\mu^* = \tilde{\mu}$. Set $\Omega_t = \{x \in \Omega : \rho(x) > t\}$ and let v_t be the solution of

$$\begin{aligned} -\Delta v_t + e^{v_t} - 1 &= 0 & \text{in } \Omega_t \\ v_t &= w & \text{on } \Sigma_t. \end{aligned}$$

Then $v_t \leq w$ in Ω_t . Furthermore

$$0 < t' < t \implies v_{t'} \leq v_t \quad \text{in } \Omega_{t'}.$$

Then $\tilde{u} = \lim_{t \rightarrow 0} v_t$ exists, the convergence holds in $L^1(\Omega)$ and $e^{v_t} \rightarrow e^{\tilde{u}}$ in $L^1(\Omega; \rho dx)$ (here we use the fact that $e^w \in L^1(\Omega; \rho dx)$). Because

$$\lim_{t \rightarrow 0} \int_{\Sigma_t} \tilde{w}(t, \cdot) \eta dS_t = \int_{\Sigma} \eta d\tilde{\mu} \quad \forall \eta \in C(\Sigma),$$

and $v_t = \tilde{w}$ on Σ_t , it follows that \tilde{u} admits $\tilde{\mu}$ for boundary trace and thus $\tilde{u} = u_{\tilde{\mu}}$.

II- Let $u_n = u_{\mu_n}$ be the solutions of (1.24) with boundary value μ_n . The sequence $\{u_n\}$ is increasing. Since

$$\int_{\Omega} (u_n + (e^{u_n} - 1)\zeta_0) dx = - \int_{\Sigma} \frac{\partial \zeta_0}{\partial \nu} d\mu_n \leq - \int_{\Sigma} \frac{\partial \zeta_0}{\partial \nu} d\mu, \quad (1.7)$$

we conclude as in the proof of Theorem 1, that u_n increases and converges to a solution $u = u_\mu$ of (1.24) with boundary value μ .

³Marcus M. and Véron L., *The boundary trace and generalized B.V.P. for semilinear elliptic equations with coercive absorption*, Comm. Pure Appl. Math. **56**, 689-731 (2003).

III- In the proof of I we have actually used the following result : *Let w be a non-negative supersolution of (1.1) such that $e^w \in L^1(\Omega; \rho dx)$ and let $\mu \in \mathfrak{M}_+(\Sigma)$ be the boundary trace of w . Then μ is admissible.* Let $f \in L^1_+(\Sigma)$ and μ be an admissible measure. We denote by $u = u_\mu$ the solution of (1.24). For $k > 0$, set $f_k = \min\{k, f\}$. The function $w_k = u_\mu + \mathbb{P}^\Omega[f_k]$ is a nonnegative supersolution, and, since $\mathbb{P}^\Omega[f_k] \leq k$, $e^{w_k} \in L^1(\Omega; \rho dx)$. Furthermore the boundary trace of w_k is $\mu + f_k$. Therefore $\mu + f_k$ is admissible. We conclude by II that $\mu + f$ is admissible \square

Remark. The assertions I and II in Theorem 1 are still valid if we replace $r \mapsto e^r - 1$ by any continuous nondecreasing function f vanishing at 0.

The Orlicz space framework

Let $M_{exp}(\Sigma)$ be the set of nonnegative boundary measures μ such that

$$\exp(\mathbb{P}^\Omega[\mu]) \in L^1(\Omega; \rho dx). \quad (1.8)$$

The set $M_{exp}(\Sigma)$ is not a linear space, but it is a convex subset of $\mathfrak{M}_+(\Sigma)$. Put

$$p(t) = \operatorname{sgn}(s)(e^s - 1), \quad P(t) = e^{|t|} - 1 - |t|,$$

and

$$\bar{p}(s) = \operatorname{sgn}(s) \ln(|s| + 1), \quad P^*(t) = (|t| + 1) \ln(|t| + 1) - |t|.$$

Then P and P^* are complementary functions in the sense that Young inequality holds

$$xy \leq P(x) + P^*(y) \quad \forall (x, y) \in \mathbb{R} \times \mathbb{R},$$

with equality if and only if $x = \bar{p}(y)$ or $y = p(x)$. It is classical to define

$$M_P(\Omega; \rho dx) = \{\phi \in L^1_{loc}(\Omega) : P(\phi) \in L^1(\Omega; \rho dx)\}, \quad (1.9)$$

$$M_{P^*}(\Omega; \rho dx) = \{\phi \in L^1_{loc}(\Omega) : P^*(\phi) \in L^1(\Omega; \rho dx)\}. \quad (1.10)$$

The Orlicz spaces $L_P(\Omega; \rho dx)$ and $L_{P^*}(\Omega; \rho dx)$ are the vector spaces spanned respectively by $M_P(\Omega; \rho dx)$ and $M_{P^*}(\Omega; \rho dx)$. They are endowed with the Luxemburg norms

$$\|\phi\|_{L_{P\rho}} = \inf \left\{ k > 0 : \int_{\Omega} P\left(\frac{f}{k}\right) \rho dx \leq 1 \right\}. \quad (1.11)$$

and

$$\|\phi\|_{L_{P^*\rho}} = \inf \left\{ k > 0 : \int_{\Omega} P^*\left(\frac{f}{k}\right) \rho dx \leq 1 \right\}. \quad (1.12)$$

Furthermore the Hölder-Young inequality holds ⁴

$$\left| \int_{\Omega} \phi \psi \rho dx \right| \leq \|\phi\|_{L_{P\rho}} \|\psi\|_{L_{P^*}} \quad \forall (\phi, \psi) \in L_P(\Omega; \rho dx) \times L_{P^*}(\Omega; \rho dx). \quad (1.13)$$

Since P^* satisfies the Δ_2 -condition, $M_{P^*}(\Omega; \rho dx) = L_{P^*}(\Omega; \rho dx)$. Furthermore, since

$$\frac{|a| \ln(1 + |a|)}{2} \leq P^*(a) \leq |a| \ln(1 + |a|) \quad \forall a \in \mathbb{R},$$

the space $L_{P^*}(\Omega; \rho dx)$ is associated with the class $L \ln L(\Omega; \rho dx)$ and to the Hardy-Littlewood maximal function. We recall its definition: we consider a cube Q_0 containing $\bar{\Omega}$, with sides parallel to the axes. If $f \in L^1(\Omega)$ we denote by \tilde{f} its extension by 0 in $Q_0 \setminus \Omega$ and put

$$M_{Q_0}[f](x) = \sup \left\{ \frac{1}{|Q|} \int_Q |f|(y) dy : Q \in \mathcal{Q}_x \right\}$$

where \mathcal{Q}_x denotes the set of all cubes containing x and contained in Q_0 , with sides parallel to the axes. Thus

$$\|f\|_{L \ln L_{\rho}} = \int_{Q_0} M_{Q_0}[f](x) \rho dx.$$

and

$$f \in L \ln L(\Omega; \rho dx) \iff \int_{\Omega} |f| \ln(1 + |f|) \rho dx$$

Furthermore, $L_P(\Omega; \rho dx)$ is the dual space of $L_{P^*}(\Omega; \rho dx)$, ⁵ if ℓ is a continuous linear functional on $L \ln L(\Omega; \rho dx)$, there exist a measurable function g_{ℓ} and some $\theta > 0$ such that

$$\begin{cases} \ell(f) = \int_{\Omega} g_{\ell} f dx \quad \forall f \in L \ln L(\Omega; \rho dx), \\ \int_{\Omega} e^{\theta |g_{\ell}|} \rho dx < \infty. \end{cases}$$

This can be seen as a consequence of Young's inequality.

Definition 1.1 *The space of all distributions on Σ such that*

$$\mathbb{P}^{\Omega}[\mu] \in L_P(\Omega; \rho dx)$$

⁴M.A. Krasnosel'skii and Y. B. Rutickii *Convex functions and Orlicz spaces* P. Noordhoff Ltd, Groningen (1961).

⁵M. Fuchs and G. Seregin *A regularity theory for variational integrals with $L \ln L$ -growth* **Calc. Var.** **6** (1998), 171-187.

is denoted by $B^{\text{exp}}(\Sigma)$, with norm

$$\|\mu\|_{B^{\text{exp}}} = \|\mathbb{P}^\Omega[\mu]\|_{L_{P_\rho}}. \quad (1.14)$$

The subset of distributions such that

$$\exp(\mathbb{P}^\Omega[\mu]) \in L^1(\Omega; \rho dx)$$

is denoted by $M^{\text{exp}}(\Sigma)$.

The analytic characterization of $B^{\text{exp}}(\Sigma)$ can be done in introducing the space of normal derivatives of Green potentials of $L \ln L$ functions:

$$N^{L \ln L}(\Sigma) = \{\eta : \rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta]) \in L \ln L(\Omega; \rho dx)\}. \quad (1.15)$$

where ρ^* is a smooth positive function with value ρ in a neighborhood of Σ . Then

$$\left| \int_\Sigma \eta d\mu \right| = \left| \int_\Omega \mathbb{P}^\Omega[\mu] \Delta(\rho^* \mathbb{P}^\Omega[\eta]) dx \right| \leq \|\mathbb{P}^\Omega[\mu]\|_{L_{P_\rho}} \|\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta])\|_{L_{P_\rho^*}}. \quad (1.16)$$

Notice that the actual regularity of the η function is not clear, although

$$\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta]) \in L \ln L(\Omega; \rho dx) \implies \Delta(\rho^* \mathbb{P}^\Omega[\eta]) \in L \ln L(\Omega).$$

If we take, as a norm on $N^{L \ln L}(\Sigma)$

$$\|\eta\|_{N^{L \ln L}} = \|\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta])\|_{L_{P_\rho^*}}, \quad (1.17)$$

and define the $C_{N^{L \ln L}}$ -capacity of a compact subset K of Σ by

$$C_{N^{L \ln L}}(K) = \inf\{\|\eta\|_{N^{L \ln L}} : \eta \in C^2(\Sigma), 0 \leq \eta \leq 1, \eta = 1 \text{ in a neighborhood of } K\}, \quad (1.18)$$

the following result follows from (1.16):

Proposition 1 *If $\mu \in B_+^{\text{exp}}(\Sigma)$, it does not charge Borel subsets with $C_{N^{L \ln L}}$ -capacity zero.*

Because P^* satisfies the Δ_2 -condition, $L_P(\Omega; \rho dx)$ is the dual space of $L_{P^*}(\Omega; \rho dx)$. Now we admit previsionally the following result of Feyel-de la Pradelle type ⁶

Theorem 3 *Let $\mu \in \mathfrak{M}_+(\Sigma)$, which does not charge Borel sets with $C_{N^{L \ln L}}$ -capacity zero. Then there exists an increasing sequence $\{\mu_n\} \subset B^{\text{exp}}(\Sigma)$ which converges*

⁶D. Feyel and A. de la Pradelle *Topologies fines et compactification associées à certains espaces de Dirichlet* **Ann. Inst. Fourier** **27** (1977), 121-146.

weakly to μ .

The proof could be a parallel copy of the original proof. It may have been proved for general capacities in duality (we shall see later on). A variant of this result is the Dal Maso form ⁷

Theorem 4 *Let $\mu \in \mathfrak{M}_+(\Sigma)$, which does not charge Borel sets with $C_{NL \ln L}$ -capacity zero. Then there exists $\nu \in N_+^{L \ln L}(\Sigma)$ and a nonnegative Borel function h such that*

$$\mu = h d\nu.$$

The role of Orlicz capacities: admissible measures and removable sets

As we have already seen it, a measure in $B_+^{exp}(\Sigma)$ is admissible, and does not charge Borel subsets of $C_{NL \ln L}$ -capacity zero. The following result is a slight extension of a result of Grillo-Véron, with a proof which inherits some observations of Brezis-Marcus-Ponce.

Theorem 5 *Let μ be an admissible measure. Then μ does not charge Borel subsets of $C_{NL \ln L}$ -capacity zero.*

Proof. Let K be a compact subset with $C_{NL \ln L}$ -capacity zero. There exist a sequence $\{\eta_n\} \subset C^2(\Sigma)$ such that $0 \leq \eta_n \leq 1$, $\eta_n = 1$ in a neighborhood of K and

$$\lim_{n \rightarrow \infty} \|\eta_n\|_{NL \ln L} = \|\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta_n])\|_{L_{P_\rho^*}} = 0. \quad (1.19)$$

Take $\rho^* \mathbb{P}^\Omega[\eta_n]$ as a test function, then

$$\int_{\Omega} (-u \Delta(\rho^* \mathbb{P}^\Omega[\eta_n]) + (e^u - 1) \rho^* \mathbb{P}^\Omega[\eta_n]) dx = - \int_{\Sigma} \frac{\partial(\rho^* \mathbb{P}^\Omega[\eta_n])}{\partial \nu} d\mu$$

Since $\frac{\partial(\rho^* \mathbb{P}^\Omega[\eta_n])}{\partial \nu} = \eta_n$ and $\mu > 0$, there holds $-\int_{\Sigma} \frac{\partial(\rho^* \mathbb{P}^\Omega[\eta_n])}{\partial \nu} d\mu \geq \mu(K)$.

Furthermore

$$\left| \int_{\Omega} u \Delta(\rho^* \mathbb{P}^\Omega[\eta_n]) dx \right| \leq \|u\|_{L_{P_\rho}} \|\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta_n])\|_{L_{P_\rho^*}}. \quad (1.20)$$

Then

$$\mu(E) \leq \int_{\Omega} (e^u - 1) \rho^* \mathbb{P}^\Omega[\eta_n] dx + \|u\|_{L_{P_\rho}} \|\rho^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta_n])\|_{L_{P_\rho^*}}.$$

⁷G. Dal Maso *On the integral representation of certain local functionals* **Recherche Math.** **32** (1983), 85-113.

By the same argument as in Brezis-Marcus-Ponce, $\lim_n \rightarrow \infty \rho^* \mathbb{P}^\Omega[\eta_n] = 0$, a.e. in Ω , and there exists a nonnegative L^1_ρ -function Φ such that $0 \leq \rho^* \mathbb{P}^\Omega[\eta_n] \leq \Phi$. By (1.19), (1.20) and Lebesgue's theorem, $\mu(E) = 0$. \square

Definition 1.2 *A subset $E \subset \Sigma$ is said removable, if and only if any positive solution u of (1.1) in Ω , which is continuous in $\bar{\Omega} \setminus E$ and vanishes on $\Sigma \setminus E$, is identically zero.*

Theorem 6 *A compact subset $K \subset \Sigma$ is removable if and only if $C_{NL \ln L}(K) = 0$.*

Proof. Let $u \in C(\bar{\Omega} \setminus K)$ be a solution of (1.1) which is zero on $\Sigma \setminus K$. Let $\{\eta_n\} \subset C^2(\Sigma)$ such that $0 \leq \eta_n \leq 1$, $\eta_n = 1$ in a neighborhood \mathcal{V} of K and (1.19) holds. Put $\theta_n = 1 - \eta_n$. Put $\rho_K(x) = \text{dist}(x, K)$. Then, as a consequence of Keller-Osserman estimate and the fact that u vanishes on K^c , there holds

$$u(x) \leq C \frac{\rho(x) \ln(2/\rho_K(x))}{\rho_K(x)} + D.$$

Thus the function $\zeta_n = \rho^* \mathbb{P}^\Omega[\theta_n]$ is an admissible test function for u , and

$$\int_{\Omega} (-u \Delta \zeta_n + (e^u - 1) \zeta_n) dx = 0.$$

Clearly $\mathbb{P}^\Omega[\theta_n] = 1 - \mathbb{P}^\Omega[\eta_n]$ and

$$\Delta \zeta_n = \Delta \rho^* - \Delta(\rho^* \mathbb{P}^\Omega[\eta_n])$$

Inasmuch we can modify ρ^* in order to have $-\Delta \rho^* \geq 0$, in which case $\rho^* = \rho$ near Σ is replaced by $\rho^* \approx \rho$, we derive

$$\begin{aligned} - \int_{\Omega} u \Delta \zeta_n dx &= - \int_{\Omega} \zeta_n^{-1} \Delta \zeta_n u \zeta_n dx \\ &\geq -2^{-1} \int_{\Omega} (e^u - 1 - u) \zeta_n dx - \int_{\Omega} Q(\zeta_n^{-1} \Delta(\rho^* \mathbb{P}^\Omega[\eta_n])) \zeta_n dx, \end{aligned}$$

where

$$Q(r) = (|r| + 2^{-1}) \ln(2|r| + 1) - |r| \leq C |r| \ln(|r| + 1) \quad \forall r \in \mathbb{R}.$$

Therefore

$$\int_{\Omega} (e^u - 1 - u) \zeta_n dx \leq 2C \int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_n])| \ln(1 + \rho^{-2} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_n])|) dx, \quad (1.21)$$

since $\zeta_n^{-1} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \leq \rho^{-2} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|$. Furthermore

$$\begin{aligned} \ln(1 + \rho^{-2} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) &= -\ln \rho + \ln(\rho + \rho^{-1} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) \\ &\leq -\ln \rho + \ln(1 + \rho^{-1} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) \end{aligned}$$

But (we can assume $\rho \leq 1$)

$$\begin{aligned} &\int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln(1 + \rho^{-2} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) dx \\ &\leq -\int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln \rho dx + \int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln(1 + \rho^{-1} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) dx, \end{aligned}$$

and

$$\begin{aligned} &\int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln \rho^{-1} dx \\ &= \int_{\{|\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \leq 1\}} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln \rho^{-1} dx + \int_{\{|\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| > 1\}} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln \rho^{-1} dx \\ &\leq \int_{\{|\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \leq 1\}} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln \rho^{-1} dx + \int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln(1 + \rho^{-1} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) dx \end{aligned}$$

But

$$\lim_{n \rightarrow \infty} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| = 0 \quad \text{a. e. in } \Omega,$$

at least up to some subsequence. Thus

$$\lim_{n \rightarrow \infty} \int_{\Omega} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])| \ln(1 + \rho^{-2} |\Delta(\rho^* \mathbb{P}^\Omega[\eta_m])|) dx = 0 \quad (1.22)$$

Using (1.30), we derive $u = 0$.

Conversely, assume that $C_{NL \ln L}(K) > 0$. Since the dual definition of $C_{NL \ln L}(K)$ appears to be⁸

$$C_{NL \ln L}(K) = \sup\{\mu(K) : \mu \in \mathfrak{M}_+(\Sigma), \mu(K^c) = 0, \|\mathbb{P}^\Omega[\mu]\|_{L_{P_\rho}} \leq 1\},$$

there exists a nonzero measure $\mu \in \mathfrak{M}_+(\Sigma)$ such that $\mu(K^c) = 0$ in the space $B_+^{exp}(\Sigma)$. This means that $\theta\mu \in M_+^{exp}(\Sigma)$ for some $\theta > 0$. Thus problem (1.24) admits a solution. \square

⁸S. E. Kuznetsov *Removable singularities for $Lu = \Psi(u)$ and Orlicz capacities* **J. Funct. Anal.** **170** (2000), 428-449.

By Theorems 1 and 3 we have a partial characterization of measures for which problems (1.24) admits a solution and K is not removable.

Theorem 7 *If a measure μ is admissible there exists an increasing sequence of $\{\mu_n\}$ satisfying*

$$\int_{\Omega} \exp(\theta_n \mathbb{P}^{\Omega}[\mu_n]) \rho dx < \infty \quad (1.23)$$

for some $\theta_n > 0$ which converges to μ .

Several questions can be adressed

1- If a singular measure μ is admissible does it exist an increasing sequence of $\{\mu_n\}$ converging to μ such that (1.23) holds with $\theta_n = 1$?

2- If a measure μ does not charge Borel sets with $C^{L \ln L}$ -capacity zero, does it exist $\theta > 0$ such that $\theta \mu \in M^{exp}(\Sigma)$?

3- If a singular measure μ is admissible, then $(1 - \delta)\mu \in M^{exp}(\Sigma)$ for any $\delta > 0$?

Further extensions

A part of the above construction could be extended to problems with more general nonlinearity such as

Boundary measures

$$\begin{aligned} -\Delta u + P(u) &= 0 \quad \text{in } \Omega \\ u &= \mu \quad \text{on } \Sigma, \end{aligned} \quad (1.24)$$

where P is a convex increasing function vanishing at 0 and such that $\lim_{r \rightarrow \infty} P(r)/r = \infty$: In Theorem 1- P , (1.3) should be replaced by

$$P(\mathbb{P}^{\Omega}[\mu_n]) \in L^1(\Omega; \rho dx). \quad (1.25)$$

In Theorem 2- P , (i) and (ii) still hold. For simplicity we assume that P is a N -function in the sense of Orlicz spaces

$$P(r) = \int_0^r p(s) ds$$

where p is increasing and vanishes at 0. Let P^* be the conjugate N -function, $L_P(\Omega; \rho dx)$ and $L_{P^*}(\Omega; \rho dx)$ the corresponding Orlicz spaces endowed with the

Luxemburg norms. Then Proposition 1- P is valid, provided the space $B^P(\Sigma)$ and $M^P(\Sigma)$ are accordingly defined with the following notations:

$$N^{P^*}(\Sigma) = \{\eta : \rho^{-1}\Delta(\rho^*\mathbb{P}^\Omega[\eta]) \in L_{P^*}(\Omega; \rho dx)\}$$

with corresponding norm

$$\|\eta\|_{N^{P^*}} = \|\rho^{-1}\Delta(\rho^*\mathbb{P}^\Omega[\eta])\|_{L_{P^*}}$$

and the corresponding capacity $C_{N^{P^*}}$. It is still likely that Theorem 3- P , 4- P hold. The proof of Theorem 5- P should be valid without any major modification. **However**, it appears that the characterization of removable sets cannot be adapted without further properties of the function P^* like the Δ_2 -condition. Such a condition holds usually when P has a power-like growth (> 1) and a logarithmic type growth.

Internal measures

Most of the above techniques can be extended to problem of the types

$$\begin{aligned} -\Delta u + e^u - 1 &= \mu & \text{in } \Omega \\ u &= 0 & \text{on } \Sigma, \end{aligned} \quad (1.26)$$

and

$$\begin{aligned} -\Delta u + P(u) &= \mu & \text{in } \Omega \\ u &= 0 & \text{on } \Sigma. \end{aligned} \quad (1.27)$$

where $\mu \in \mathfrak{M}_+^b(\Omega)$. If we denote by $\mathfrak{M}_+^{exp}(\Omega)$ the set of admissible measures for (1.26), then the internal outer part of Theorems 2 and 5 are already known (BMP). Inequality (1.16) becomes

$$\left| \int_\Omega \eta d\mu \right| = \left| \int_\Omega \eta \Delta \mathbb{G}^\Omega[\mu] dx \right| = \left| \int_\Omega \mathbb{G}^\Omega[\mu] \Delta \eta dx \right| \leq \|\mathbb{G}^\Omega[\mu]\|_{L_P} \|\Delta \eta\|_{L_{P^*}}. \quad (1.28)$$

for $\eta \in C_c^{1,1}(\bar{\Omega})$. Define the $C_{\Delta L \ln L}$ -capacity of a compact subset K of Σ by

$$C_{\Delta L \ln L}(K) = \inf\{\|\Delta \eta\|_{L_{P^*}} : \eta \in C_c^2(\Omega), 0 \leq \eta \leq 1, \eta = 1 \text{ in a neighborhood of } K\}, \quad (1.29)$$

Thus Proposition 1 is valid under the form

Proposition 1 bis *If $\mu \in B_+^{\text{exp}}(\Omega)$, it does not charge Borel subsets with $C_{\Delta L \ln L}$ -capacity zero.*

Theorem 6 becomes

Theorem 6 bis *A compact subset $K \subset \Omega$ is removable if and only if $C_{\Delta L \ln L}(K) = 0$.*

Proof. Let $u \in C(\Omega \setminus K)$ be a solution of (1.1) which is zero on Σ . Let $\{\eta_n\} \subset C^2(\Omega)$ such that $0 \leq \eta_n \leq 1$, $\eta_n = 1$ in a neighborhood \mathcal{V} of K and (1.19) holds. Put $\rho_K(x) = \text{dist}(x, K)$. Then, as a consequence of Keller-Osserman estimate, there holds

$$u(x) \leq C \ln(2/\rho_K(x)) + D.$$

Put $\theta_n = 1 - \eta_n$. Then the function $\zeta_n = \phi_1 \theta_n$ (ϕ_1 being the first eigenfunction of $-\Delta$) is an admissible test function for u , and

$$\int_{\Omega} (-u \Delta \zeta_n + (e^u - 1) \zeta_n) dx = 0.$$

We derive

$$\begin{aligned} - \int_{\Omega} u \Delta \zeta_n dx &= - \int_{\Omega} \zeta_n^{-1} \Delta \zeta_n u dx \\ &\geq -2^{-1} \int_{\Omega} (e^u - 1 - u) dx - \int_{\Omega} Q(\Delta(\zeta_n)) dx. \end{aligned}$$

Therefore

$$\int_{\Omega} (e^u - 1 - u) \zeta_n dx \leq 2C \int_{\Omega} |\Delta \zeta_n| \ln(1 + |\Delta \zeta_n|) dx, \quad (1.30)$$

The conclusion is as in BMP. \square

Reduced measures

What are the reduced measures both for the boundary and internal problems (1.24) (resp. 1.27)? A projection onto the closure of the sets $M^{\text{exp}}(\Sigma)$ (resp. $M^{\text{exp}}(\Omega)$) of positive measures in Σ (resp. Ω) satisfying

$$\int_{\Omega} \exp(\mathbb{P}^{\Omega}[\mu]) \rho dx < \infty$$

(resp.

$$\int_{\Omega} \exp(\mathbb{G}^{\Omega}[\mu]) dx < \infty.)$$

The definition of the projection is not clear, although an important fact is that it ensures uniqueness. This conjectures could be extended to problems involving operator $u \mapsto -\Delta u + P(u)$.