



**Validation of a confirmatory analytical method for the determination of aflatoxins B1, B2, G1 and G2 in foods and feed materials by HPLC with on-line photochemical derivatization and fluorescence detection**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-331.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	28-May-2009
Complete List of Authors:	Muscarella, Marilena; Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata Iammarino, Marco; Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata Nardiello, Donatella; University of Foggia, DiSACD & BIOAGROMED Lo Magro, Sonia; Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata Palermo, Carmen; University of Foggia, DiSACD & BIOAGROMED CENTONZE, DIEGO; University of Foggia, DiSACD & BIOAGROMED Palermo, Domenico; Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata
Methods/Techniques:	Clean-up - affinity columns, HPLC, Method validation
Additives/Contaminants:	Aflatoxins
Food Types:	Cereals

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7 2  
8  
9 **Validation of a confirmatory analytical method for the determination of**  
10 **aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> in foods and feed materials by HPLC with on-line**  
11 **photochemical derivatization and fluorescence detection**  
12  
13  
14  
15 6  
16  
17  
18 8 Marilena Muscarella<sup>a</sup>, Marco Iammarino<sup>a</sup>, Donatella Nardiello<sup>b</sup>, Sonia Lo Magro<sup>a</sup>, Carmen  
19 Palermo<sup>b</sup>, Diego Centonze<sup>b\*</sup>, Domenico Palermo<sup>a</sup>  
20  
21  
22 10  
23  
24  
25 12 <sup>a</sup> *Istituto Zooprofilattico Sperimentale della Puglia e della Basilicata, Via Manfredonia 20,*  
26 *71100 Foggia, Italy*  
27  
28 14 <sup>b</sup> *Dipartimento di Scienze Agro-Ambientali, Chimica e Difesa Vegetale and BIOAGROMED,*  
29 *Università degli Studi di Foggia, Via Napoli 25, 71100 Foggia, Italy*  
30  
31 16  
32  
33  
34  
35  
36 18  
37  
38  
39  
40 20  
41  
42  
43  
44  
45 22  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

\* Corresponding author. Tel.: +39-0881-589-104; fax: +39-0881-740211. *e-mail address*: centonze@unifg.it

## Abstract

A sensitive and selective analytical method for the simultaneous separation and quantitative determination of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> in foodstuffs and materials for feed has been validated. The method is based on high performance liquid chromatography with on-line post-column photochemical derivatization and fluorescence detection. The chromatographic separation of aflatoxins was accomplished by using a C<sub>18</sub> column eluted with an isocratic mobile phase consisting of water, methanol and acetonitrile. The sample preparation required a simple extraction of aflatoxins with MeOH/H<sub>2</sub>O (80:20, v/v) and a purification step by immunoaffinity column cleanup. The total analysis time, including sample preparation and chromatographic separation, did not exceed 40 min with a run time of 10 min. The on-line photochemical derivatization ensures better results in terms of simplicity, sensitivity and reproducibility with respect to chemical derivatization techniques, and provides an increase of the peak resolution and an extent of automation in comparison with the electrochemical ones. The procedure for the determination of aflatoxins in food samples and cereals for animal consumption was extensively validated following the Regulation (EC) No. 882/2004. Detection limits in wheat bran samples of 0.08 µgkg<sup>-1</sup> for AFB<sub>1</sub>, 0.02 µgkg<sup>-1</sup> for AFB<sub>2</sub>, 0.16 µgkg<sup>-1</sup> for AFG<sub>1</sub> and 0.04 µgkg<sup>-1</sup> for AFG<sub>2</sub> were attained. The method allows high recovery with mean values ranging from 72% to 94% and it satisfies the necessary requirements for sensitivity, linearity, selectivity, precision and ruggedness, demonstrating the conformity of the method with provisions of Regulation (EC) No. 401/2006.

**Keywords:** Aflatoxins, HPLC, photochemical derivatization, validation, Regulation (EC) No. 401/2006, Regulation (EC) No. 882/2004



1  
2  
3 aflatoxins have been developed, comprising immunological methods, such as enzyme linked  
4  
5  
6 74 immunosorbent assays (Trucksess and Stack, 1994, Park et al.,1989), and chromatographic  
7  
8 methods based on RP-HPLC and fluorescence detection (FLD) with chemical pre- or post-  
9  
10  
11 76 column (Wilson and Romer, 1991, Torres Espinosa et al., 1995, Roch et al., 1992, Saad et al.,  
12  
13 1995, Garcia-Villanova et al., 2004, Davis et al., 1981, Tuinstra and Haasnoot, 1983,  
14  
15 78 Niedwetzki and Lach, 1994, Ahmed and Robinson, 1998, Holcomb and Thompson, 1991,  
16  
17 Garner et al., 1993, Chan et al., 2004, Va'zquez et al., 1999, Cepeda et al., 1996) or on-line  
18  
19  
20 80 electrochemical (Kok et al., 1986, Kussak et al., 1995, Dorner et al., 1993, Scholten and Spanjer,  
21  
22 1996, Tavčar-Kalcher et al., 2007) derivatization of analytes.  
23  
24

25 82  
26  
27 Although ELISA methods are extensively used for rapid qualitative screenings of aflatoxins,  
28  
29  
30 84 they are not useful in providing a definitive confirmation of the toxins and an accurate  
31  
32 quantitative determination. The present trend is the use of HPLC for the determination of  
33  
34 86 aflatoxins because of characteristics of specificity, high sensitivity and simplicity of operation.  
35  
36 Additionally, the chromatographic determination allows quantifications of each toxin  
37  
38  
39 88 individually, obtaining more information than methods that give only a total quantification of  
40  
41 aflatoxins. This is particularly important for food analysis where the determination of AFB<sub>1</sub> is  
42  
43  
44 90 required.  
45  
46  
47  
48

49 92 The highly conjugated and rigid aflatoxin moieties give rise to native fluorescence  
50  
51 characteristics, therefore small structural modifications can have a strong influence on the  
52  
53  
54 94 fluorescence properties; for instance aflatoxins G<sub>2</sub> and B<sub>2</sub> are more fluorescent than B<sub>1</sub> and G<sub>1</sub>  
55  
56 that are required to be converted into more highly fluorescent derivatives. Several methods of  
57  
58  
59 96 derivatization are available, including pre-column treatment with trifluoroacetic acid (TFA)  
60  
(Garcia-Villanova et al, 2004) and post-column derivatization with iodine (Davis et al., 1981,

1  
2  
3 98 Tuinstra et al., 1983, Niedwetzki et al., 1994, Ahmed et al., 1998, Holcomb et al., 1991),  
4  
5 cyclodextrins (Va'zquez et al., 1999, Cepeda et al., 1996) and pyridinium hydrobromide  
6  
7  
8 100 perbromide (Garner et al., 1993, Chan et al., 2004). However, these methods are laborious and  
9  
10 present a number of disadvantages, such as handling of toxic reagents, instability of the  
11  
12 derivatives, a low day-to-day reproducibility. An alternative approach is based on post-column  
13 102 electrochemically generated bromine (Cepeda et al., 1996, Kok et al., 1986, Kussak et al., 1995,  
14  
15  
16  
17 104 Dorner et al., 1993, Scholten et al., 1996) that do not give a good chromatographic peak  
18  
19 resolution and allows a less extent of automation, since any accidental interruption of the mobile  
20  
21 phase flux determines a serious damage to the electrochemical cell. Nowadays, an on-line  
22 106 photochemical derivatization system is available (Joshua, 1993, Papadopoulou-Bouraoui et al.  
23  
24  
25  
26  
27 108 2002, Sobolev and Dorner, 2002, Sobolev, 2007, Waltking et al., 2006, Arranz et al., 2006,  
28  
29 Zang and Lim, 2005), but its potential has not fully investigated. It has been reported that the  
30  
31 on-line photochemical derivatization coupled to fluorescence detection guarantees an  
32 110 enhancement of efficiency and performances of the HPLC methods (Papadopoulou-Bouraoui et  
33  
34  
35  
36 112 al., 2002, Zang et al., 2005). Differently from chemical pre- and post-column derivatization  
37  
38 techniques, the photochemical derivatization does not involve any additional chemicals and,  
39  
40 consequently, no auxiliary pump is required, ensuring better results without any control of the  
41 114 derivatization reaction. Nevertheless, to the best of our knowledge chromatographic methods for  
42  
43  
44  
45  
46 116 the determination of aflatoxins, exploiting the photochemical derivatization and fluorescence  
47  
48  
49  
50  
51 118 detection, have been never validated according to the recent European Commission guidelines.

52  
53 In the present work, the procedure for the determination of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> in foods  
54  
55 and materials for feed, based on immunoaffinity column sample cleanup and liquid  
56 120 chromatography coupled to photochemical derivatization and fluorescence detection, has been  
57  
58  
59  
60 122 submitted to the validation procedure following the Regulation (EC) No. 882/2004. The

1  
2  
3 validation of the proposed analytical method is required in order to provide accurate and  
4  
5  
6 124 reproducible results within- and inter-official control laboratories, which are involved in the  
7  
8 monitoring and risk-assessment studies, as well as in official controls. Through validation  
9  
10  
11 126 procedure, linearity, selectivity, recovery, precision, detection and quantification limits (LODs,  
12  
13 LOQs), ruggedness of the method and measurement uncertainty have been evaluated.  
14  
15 128 Furthermore, laboratory performances were tested by taking part in a proficiency test round. The  
16  
17 results of the validation procedure demonstrate the conformity of the method of aflatoxins  
18  
19 analysis in foods and animal feeds with provisions of Regulation (EC) No. 401/2006, which, as  
20 130 underlined by European Commission Sanco 0895/2007, establishes more specific performance  
21  
22 criteria than Decision 657/2002/EC for methods of analysis to be used for official control of  
23  
24  
25 132 mycotoxins in foodstuffs.  
26  
27  
28  
29  
30 134

## 31 **Materials and Methods**

32  
33  
34 136 **Chemicals.** A standard solution of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> with a total certified  
35  
36 concentration of 5 µg mL<sup>-1</sup> corresponding to a concentration of 2.0 µg mL<sup>-1</sup> each for B<sub>1</sub> and G<sub>1</sub>  
37  
38 and of 0.5 µg mL<sup>-1</sup> each for B<sub>2</sub> and G<sub>2</sub>, was supplied by Riedel-de Haën (Sigma-Aldrich  
39 138 Laborchemikalien, Seelze, Germany). Water, methanol and acetonitrile of HPLC grade were  
40  
41 purchased from Baker (Deventer, Holland). Sodium chloride (≥99.0%), Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O  
42 140 (≥99.0%), NaH<sub>2</sub>PO<sub>4</sub>·H<sub>2</sub>O (≥99.0%) and TWEEN 20® were purchased from Sigma-Aldrich. All  
43  
44  
45  
46  
47  
48  
49 142 chemicals were of ACS grade. A 500 µg L<sup>-1</sup> stock solution of aflatoxins standard, containing 200  
50  
51 µg L<sup>-1</sup> of AFB<sub>1</sub>, 50 µg L<sup>-1</sup> of AFB<sub>2</sub>, 200 µg L<sup>-1</sup> of AFG<sub>1</sub> and 50 µg L<sup>-1</sup> of AFG<sub>2</sub>, were prepared in  
52  
53 acetonitrile and stored in amber flask at -20°C up to 6 months. Working standard solutions were  
54 144 prepared by dilution in mobile phase and stored at -20°C when not in use; these solutions were  
55  
56  
57  
58  
59 146 stable for at least 4 weeks. Phosphate buffer salinum (PBS) was prepared by dissolving 9.0 g of  
60  
sodium chloride, 2.85 g of Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O, 0.55 g of NaH<sub>2</sub>PO<sub>4</sub>·H<sub>2</sub>O and 1.0 mL of TWEEN

1  
2  
3  
4 148 20 $\mu$ L in 990 mL of water. The pH value was adjusted to 7.2 with concentrated hydrochloric acid  
5  
6 or NaOH 1M and the volume made to 1L with water.  
7

8  
9 150 **Instrumentation.** Chromatographic separations were performed on a HPLC system, Agilent  
10  
11 Technologies SL 1200 Series (Waldbronn, Germany) composed of a binary pump equipped with  
12  
13 152 micro vacuum degasser, thermostated autosampler, column compartment and fluorescence  
14  
15 detector (model G1321A). On-line photochemical derivatization was performed using a  
16  
17 commercially available system UVE<sup>TM</sup> LCTech GmbH (Dorfen, Germany) placed between the  
18 154 separation column and the fluorescence detector, which consisted of a 254 nm low-pressure  
19  
20 mercury lamp and a 1 mL knitted reaction coil, fitted around the UV lamp. All the separations  
21  
22 156 were performed by using a ZORBAX Eclipse XDB C<sub>18</sub> column (150  $\times$  4.6 mm i.d., particle size  
23  
24 5  $\mu$ m, Agilent Technologies) operating at a flow-rate of 1.0 mLmin<sup>-1</sup> in isocratic elution with a  
25  
26 mixture of water, methanol and acetonitrile (55:15:30, v/v/v). The injection volume was 20  $\mu$ L  
27  
28 158 and the column temperature was set at 40°C. The fluorescence detection was carried out at a  $\lambda_{exc}$   
29  
30 and  $\lambda_{em}$  of 365 and 435 nm, respectively. The system was interfaced, via network  
31  
32 160 chromatographic software (Agilent ChemStation), to a personal computer for instrumentation  
33  
34 control, data acquisition and processing.  
35  
36  
37  
38  
39  
40  
41

42 164 **Sample preparation.** Samples of wheat bran, almonds and pistachios were bought in local  
43  
44 stores; durum wheat, maize and oats were supplied by farms chosen among the major cereal  
45  
46 166 producers in the province of Foggia (Italy). Both sample extraction and cleanup procedure were  
47  
48 those recommended by LCTech and slightly modified in order to simplify the analytical  
49  
50 protocol and to reduce reagents and chemical wastes. The method did not require vacuum or  
51  
52 168 pumping devices ensuring better recoveries and reproducibility. For each matrix, from a sample  
53  
54 weight of 1 kg, an amount of 15 g of ground sample was suspended in 30 mL of 80 % methanol  
55  
56 170 and vortexed for few min. For the validation measurements, samples were spiked, prior to the  
57  
58 extraction, with the proper amounts of mixed aflatoxin standard solutions to give required  
59  
60



1  
2  
3 concentrations. After filtration with paper (Whatman No. 40), 2 mL of the mixture were added  
4  
5  
6 174 to 8 mL of PBS solution and passed through the immunoaffinity columns Aflaclean™ (LCTech  
7  
8 GmbH, Dorfen, Germany) for sample clean-up. After washing with 10 mL of water, the fraction  
9  
10 containing the aflatoxins was slowly eluted with 2 mL of methanol by gravity at a flow rate  
11 176 lower than of 0.5 mL min<sup>-1</sup>. The eluate was evaporated to dryness at 40 °C under nitrogen  
12  
13 stream (Dubnof Bath BSD/D). Finally, the residue was solubilized in 2 mL of the mobile phase,  
14  
15 178 filtered by an Anotop 10 LC (0.2 µm, 10 mm, Whatman) and then injected. The analytical  
16  
17 procedure resulted in a 2-fold dilution of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub>, evaluated against a  
18  
19 known concentration added to a blank real sample. The stability of purified extracts was high  
20 180 enough to allow autosampler overnight injections. No significant change in concentration or  
21  
22 purity of aflatoxins within 8 hours of analysis was observed.  
23  
24  
25 182  
26  
27  
28

29 184 **Validation procedure.** In consequence of different legal limits for aflatoxins in foodstuffs and  
30  
31 cereals for animal consumption, different sets of standard solutions and spiked samples were  
32  
33 used for foodstuffs (wheat bran, almonds and pistachios) and materials for feed (durum wheat,  
34 186 maize and oats). The linearity test in foodstuffs was performed by three series of analyses on  
35  
36 three different days, by injecting four standard solutions of aflatoxins B<sub>1</sub> and G<sub>1</sub> each at  
37  
38 concentrations of 0.25, 0.5, 1.0 and 2.0 µgL<sup>-1</sup> and aflatoxins B<sub>2</sub> and G<sub>2</sub> each at concentrations of  
39 188 0.0625, 0.125, 0.25 and 0.5 µgL<sup>-1</sup>. Likewise, method linearity in feed materials was determined  
40  
41 by using working standard solutions of AFB<sub>1</sub> at concentration of 2.5, 5.0, 10 and 20 µgL<sup>-1</sup>. The  
42  
43 method selectivity was tested by the analysis of 20 independent blank samples of foods (7 of  
44 190 wheat bran, 7 of almonds and 6 of pistachios) and 20 samples of cereals (7 of durum wheat, 7 of  
45  
46 maize and 6 of oats), found negative by ELISA screening. The absence of interfering peaks in  
47  
48 the retention time-window of interest was checked for each analyte within the ± 2.5% retention  
49 192 time range. Precision, recovery and both detection and quantification limits of the method were  
50  
51 determined by performing tests on three sets of blank wheat bran samples (six replicates each)  
52  
53  
54 194  
55  
56  
57  
58  
59 196  
60

1  
2  
3  
4 198 fortified with AFB<sub>1</sub> at concentration of 0.5, 1.0 and 1.5 times the ML in foodstuffs (i.e., 1.0, 2.0  
5  
6 and 3.0 µgkg<sup>-1</sup>) and aflatoxins B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> at concentration of 0.625, 1.25 e 1.875 times the  
7  
8  
9 200 ML of the sum of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> (i.e., 1.0, 2.0 and 3.0 µgkg<sup>-1</sup> for B<sub>1</sub> and G<sub>1</sub>, and  
10  
11 0.25, 0.5 and 0.75 µgkg<sup>-1</sup> for B<sub>2</sub> and G<sub>2</sub>). As far as the analysis of feed materials is concerned,  
12  
13  
14 202 only two sets of blank durum wheat samples (six replicates each) fortified with AFB<sub>1</sub> at  
15  
16 concentration of 0.5, 1.0 and 1.5 times the ML in cereals for animals (i.e., 10, 20 and 30 µgkg<sup>-1</sup>)  
17  
18 204 were analysed. The experiments, corresponding to a total number of 54 and 36 for food samples  
19  
20 and animal feed materials, respectively, were performed in different days with the same  
21  
22  
23 206 instruments but different operators.

## 27 208 **Results and Discussion**

29  
30 **Method Validation.** The proposed method has been validated according to Regulation (EC) No.  
31  
32 210 882/2004, a document about official controls performed to ensure the verification of compliance  
33  
34 with feed and food laws, animal health and animal welfare rules, which demonstrates the  
35  
36 conformity of the analytical performances with criteria established in Regulation (EC) No.  
37 212 401/2006. Therefore, through validation procedure, linearity, selectivity, recovery, precision,  
38  
39  
40  
41 214 detection and quantification limits, ruggedness of the method and measurement uncertainty have  
42  
43  
44 been evaluated.

45  
46  
47 216 **Selectivity towards interferences.** Figure 1 shows, as an example, the fluorescence enhancement  
48  
49 attained by on-line photochemical derivatization for a wheat bran sample fortified with AFB<sub>1</sub>  
50  
51 and AFG<sub>1</sub> at 3.0 µgkg<sup>-1</sup> and AFB<sub>2</sub> and AFG<sub>2</sub> at 0.75 µgkg<sup>-1</sup>. Through the post-column UV  
52  
53 irradiation, AFB<sub>1</sub> and AFG<sub>1</sub> are converted by the reaction with water contained in the mobile  
54  
55  
56 220 phase to fluorescent hemiacetals, which are similar compounds to those obtained by  
57  
58 derivatization with TFA (Garcia-Villanova et al., 2004). This photochemical modification  
59  
60  
222 determines an increase of the response sensitivity of AFB<sub>1</sub> and AFG<sub>1</sub>, while a very slight

1  
2  
3  
4 modification of signal intensities has been observed for AFB<sub>2</sub> and AFG<sub>2</sub>. Use of both the  
5  
6 224 immunoaffinity sample cleanup and post-column photochemical derivatization permitted an  
7  
8 efficient elimination of possible endogenous matrix interfering substances. In pre-validation  
9  
10 experiments the probable interfering compounds, ochratoxin A and fumonisins B<sub>1</sub> and B<sub>2</sub>,  
11 226  
12 common contaminants of cereals, nuts and dried fruit, have been tested. These compounds  
13  
14 showed retention times significantly different from those of aflatoxins or undetectable signals  
15 228  
16 under the adopted experimental conditions. Chromatograms of Figure 2, obtained for blank and  
17  
18 spiked samples with aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub>, demonstrate the selectivity of the method  
19  
20 230  
21 toward other naturally present compounds, since at the retention times of the aflatoxins no  
22  
23 interfering peaks were observed.  
24  
25 232

26  
27 **Calibration curves and limits of detection and quantification.** The goodness-of-fit of the data  
28  
29 to the calibration curve is obtained in terms of response factor distribution (signal-to-  
30 234  
31 concentration ratio,  $y_i/x_i$ ) whose reference range is  $y/x_{\text{mean}} \pm 10\%$ . Furthermore, any systematic  
32  
33 instrumental bias can be ruled out since the intercept includes the zero value at 95% confidence  
34 236  
35 level. The calibration parameters evaluated for each aflatoxin by using standard solutions are  
36  
37 reported in Table 1. Limits of detection and quantification were calculated according to the  
38  
39 238 following equations (Miller and Miller, 1993):  $\text{LOD} = 3.3 s_a/b$  and  $\text{LOQ} = 10 s_a/b$ , where  $s_a$  is the  
40  
41 standard deviation of the intercept and  $b$  is the slope of the regression line, obtained from the  
42  
43 calibration curve. The limit of quantification for AFB<sub>1</sub> of  $0.11 \mu\text{gL}^{-1}$  corresponding to a value of  
44 240  
45  $0.22 \mu\text{gkg}^{-1}$  in matrix is satisfactory with respect to the allowed maximum content of  $2 \mu\text{gkg}^{-1}$   
46  
47 (Regulation (EC) No. 1881/2006). Fit of the data to the calibration curve of AFB<sub>1</sub> in cereals for  
48  
49 242 feeds gave the linear regression equation  $y = (1.002 \pm 0.006) x + (0.09 \pm 0.06)$  with a correlation  
50  
51 coefficient of 0.9999, where  $y$  is the peak area expressed in luminescence unit (LU) and  $x$  is the  
52  
53 244 concentration of AFB<sub>1</sub> in  $\mu\text{gL}^{-1}$ . The obtained limit of quantification of  $1.2 \mu\text{gkg}^{-1}$  is noticeably  
54  
55  
56  
57  
58 246  
59  
60

1  
2  
3  
4 lower than the maximum content of  $20 \mu\text{gkg}^{-1}$  set for feed materials (Italian Ministerial Decree  
5  
6 248 149/2004, Italian Official Journal n.139, June 2004).

7  
8 **Precision and recovery.** Precision results have been previously processed by the Shapiro–Wilk  
9  
10 test to verify the normal distribution. Afterwards, ANOVA one way test was performed in order  
11 250 to verify the homogeneity of the concentration mean values evaluated on three different days. In  
12  
13 Table 2 and 3 the results of repeatability ( $\text{RSD}_r$ ) and within-laboratory reproducibility ( $\text{RSD}_R$ )  
14  
15 252 are given for each aflatoxin at the three contamination levels in spiked wheat bran samples.  
16  
17 Furthermore, Table 4 shows the results of repeatability and within-laboratory reproducibility  
18  
19 254 determined for  $\text{AFB}_1$  in spiked durum wheat (cereal for feeds). Regulation (EC) No. 401/2006  
20  
21 issued that the recommended values of experimental RSD for each concentration level must be  
22  
23 256 lower or equal than value derived by Horwitz equation, which provides the expected  $\text{RSD}\%$   
24  
25 lower or equal than value derived by Horwitz equation, which provides the expected  $\text{RSD}\%$   
26  
27 258 only on the basis of the concentration, independently of the matrix and analytical method used.  
28  
29 Nevertheless, for mass fractions lower than  $120 \mu\text{gkg}^{-1}$ , the  $\text{RSD}\%$  can be predicted in a more  
30  
31 260 restrictive manner by the Thompson equation (Thompson, 2000), which provides a  
32  
33 recommended value of  $\text{RSD}_r$  of 15 % and a reference inter-laboratory reproducibility of 23%.  
34  
35 As can be seen in Table 2 and 4,  $\text{RSD}_r$  % values are in agreement with the calculated  
36  
37 262 repeatability by Thompson equation. In absence of within-laboratory reproducibility reference  
38  
39 values, our experimental data have been compared with the repeatability maximum permitted  
40  
41 264 limit of 30%, which has been calculated as 2 fold the  $\text{RSD}_r$  recommended value, by using a  
42  
43 similar approach proposed for the calculation of the maximum permitted value of Horwitz  $\text{RSD}_R$   
44  
45 266 (Regulation (EC) No. 401/2006). As shown in Table 3, all  $\text{RSD}_R$  values resulted well below of  
46  
47 268 30%, and even in agreement with the repeatability recommended values of 15%, apart from  
48  
49  $\text{AFG}_2$ .  
50  
51 In Table 5 mean recoveries of aflatoxins evaluated at each fortification level in spiked wheat  
52  
53 270 bran and durum wheat samples are reported. As can be noticed, the recovery values obtained for  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 272 food products ranged from 76.4 to 94.6% for aflatoxins B<sub>1</sub> and G<sub>1</sub> and from 79.5 to 92.8% for  
5  
6 aflatoxins B<sub>2</sub> and G<sub>2</sub>. These results are in agreement with Regulation (EC) No. 401/2006 that  
7  
8 274 establishes recoveries in the range 70-110% and 50-120% for concentrations in the range 1-10  
9  
10  $\mu\text{gkg}^{-1}$  and less than 1  $\mu\text{gkg}^{-1}$ , respectively. In the analysis of cereals for animal feeds, mean  
11  
12  
13 276 recovery data higher than 72.6 % were obtained. It was not possible to compare experimental  
14  
15 mean recoveries with reference values since so far no legal criteria have been set for these  
16  
17  
18 278 matrices.

19  
20 Assessment of LODs and LOQs in real samples has been calculated for each analyte by mean  
21  
22  
23 280 the fit of the data to calibration curves obtained by plotting the fluorescence signals of 18 spiked  
24  
25 wheat bran samples (six replicates evaluated in the three different working sessions at each  
26  
27 282 validation level) as a function of the nominal fortification level. LOD and LOQ values, which  
28  
29 have been calculated by using the same equations previously described, ranged from 0.06 to 0.3  
30  
31  
32 284  $\mu\text{gkg}^{-1}$  and from 0.2 to 0.8  $\mu\text{gkg}^{-1}$ , respectively, with LOD and LOQ for AFB<sub>1</sub> of 0.2  $\mu\text{gkg}^{-1}$  and  
33  
34 0.6  $\mu\text{gkg}^{-1}$ , respectively. As expected, taking into account the loss of analyte during extraction  
35  
36  
37 286 and cleanup steps, LODs and LOQs are higher than those determined by calibration curves of  
38  
39 standard solutions, but still considerably lower than legal limits.

40  
41  
42 288 **Ruggedness (major changes).** The method ruggedness under conditions of major changes was  
43  
44 evaluated by using the Youden experimental design as reported elsewhere (Muscarella et al.,  
45  
46 290 2007). Two independent Youden tests were performed for different sets of real samples:  
47  
48 almonds and pistachios (foodstuffs) and maize and oats (feed materials) spiked at the AFB<sub>1</sub>  
49  
50  
51 292 maximum level. The seven factors chosen as variables for the Youden test were the matrix and  
52  
53 six fictitious factors; the use of a fictitious variable means no variation of analysis conditions. In  
54  
55  
56 294 this case the Youden experimental design requires eight independent experiments: four with the  
57  
58 matrix of validation (e.g. wheat bran and durum wheat) and four with each testing matrices  
59  
60  
296 (almonds, pistachios, maize and oats). Analysis of foods for human consumption gave a

1  
2  
3  
4 calculated standard deviation of difference  $S_{D_i}$  ( $0.09 \mu\text{gkg}^{-1}$  for almonds and pistachios) lower  
5  
6 298 than the estimated method precision at the ML ( $0.21 \mu\text{gkg}^{-1}$ ). Similar considerations are valid  
7  
8 for maize sample whose standard deviation of difference  $S_{D_i}$  ( $1.40 \mu\text{gkg}^{-1}$ ) resulted slightly  
9  
10 300 lower than method precision ( $1.45 \mu\text{gkg}^{-1}$ ). In the case of oats matrix, test F (performed at n-1  
11  
12 degrees of freedom and 95% confidence level) demonstrated that  $S_{D_i}$  ( $2.36 \mu\text{gkg}^{-1}$ ) was not  
13  
14 statistically different from the reproducibility standard deviation ( $1.45 \mu\text{gkg}^{-1}$ ) at the ML,  
15 302 therefore the variation of the matrix has no effect on the method performances.  
16  
17  
18

19  
20 304 **Measurement uncertainty.** The evaluation of uncertainty of analytical results is compulsory for  
21  
22 laboratories accredited according to ISO 17025 (UNI CEI EN ISO/IEC 17025/2000), and  
23  
24 several methods for the determination of this parameter have been proposed (EURACHEM  
25 306 Guide, 1995; EURACHEM/CITAC GUIDE CG 4, 2000; Nordic Committee on Food Analysis,  
26  
27 Guide, 1995; EURACHEM/CITAC GUIDE CG 4, 2000; Nordic Committee on Food Analysis,  
28  
29 NMKL, 2003). In this work we have used the bottom-up method together with validation data  
30 308 obtained from each step of the analytical procedure (Hund et al., 2001). On the basis of  
31  
32 uncertainties propagation law, for each aflatoxin the concentration relative uncertainty has been  
33  
34 310 calculated by the equation:  
35  
36  
37

$$38 \quad \bar{u} = \sqrt{(\bar{u}(C))^2 + (\bar{u}(V_f))^2 + (\bar{u}(w))^2}$$

39 312  
40  
41 where  $\bar{u}$  indicates the relative uncertainty, C is the analyte concentration in the sample,  $V_f$  is the  
42  
43 volume of the final extract, and w is the sample weight. The determination of  $\bar{u}$  has been  
44 314 performed by considering four sources of uncertainty: (a) preparation of the standard; (b)  
45  
46 method reproducibility; (c) method recovery; (d) calibration curve.  
47  
48  
49 316

50  
51 A relative expanded measurement uncertainty was calculated by using a coverage factor  $k$  of 2,  
52  
53 corresponding approximately to a 95% confidence level (Nordic Committee on Food Analysis,  
54 318 NMKL, 2003) obtaining values of 8.4%, 12.9 %, 9.9 % and 14.2 % for aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and  
55  
56 G<sub>2</sub>, in foodstuffs. In the validation on cereals for animal feeds, the relative expanded  
57  
58 320 measurement uncertainty calculated for AFB<sub>1</sub> was 7.6 %. These results were compared with the  
59  
60

1  
2  
3  
4 322 uncertainty function (Uf), reported in the Regulation (EC) No. 401/2006, to evaluate if the  
5  
6 method of analysis is “fit-for-purpose”. This function, which gives the acceptable maximum  
7  
8 324 uncertainty level of the method, is calculated by the following equation:

$$Uf = \sqrt{\left[\left(\frac{LOD}{2}\right)^2 + (\alpha C)^2\right]}$$

10  
11  
12  
13  
14  
15 326 where C is the concentration of interest and  $\alpha$  is a C dependent constant factor. Measurement  
16  
17 uncertainties evaluated for AFB<sub>1</sub> in food samples and feed materials (8.4% and 7.6%,  
18  
19 respectively) are considerably lower than corresponding Uf values (20% at C = 2  $\mu\text{gkg}^{-1}$ ,  $\alpha =$   
20 328 0.2 and 21 % at C = 20  $\mu\text{gkg}^{-1}$ ,  $\alpha = 0.2$ , respectively). The above results confirm that method is  
21  
22  
23  
24  
25 330 “fit-for-purpose”.

26  
27  
28 **Proficiency test round.** The proficiency test round was organised by Milk Standard Laboratory  
29  
30 332 of A.I.A. (Italian Association of Breeders) in April 2007. Three unknown maize flour samples  
31  
32 were analyzed in duplicate and the results were assessed in terms of Z-score, as calculated by the  
33  
34 proficiency test organizer. The Z-scores obtained in the A.I.A. ring test performed on unknown  
35 334 maize flour samples ( $Z_1=-0.849$ ,  $Z_2=-0.043$ ,  $Z_3=-0.040$ ) and the value of the laboratory Z-score  
36  
37 ( $Z_{\text{LAB}}=-0.322$ ) were all below the reference value of 1, proving the technical competence of the  
38  
39 336 laboratory and the method reliability.  
40  
41  
42  
43

44 338

## 46 47 48 49 340 50 51 52 53 54 342 55 56 57 58 59 344 60

51  
52  
53  
54 342  
55  
56  
57  
58  
59 344  
60

A fast, reliable and automated method for the simultaneous determination of aflatoxins in a variety of foods and animal feed materials, based on HPLC-FLD with post-column photochemical derivatization, has been validated. Photochemical derivatization results reagentless, no time consuming and easy to use, ensuring high sensitivity and reproducibility without any control of the derivatization step with respect to chemical derivatization systems, and it provides a better peak resolution and an extent of automation in comparison with the

1  
2  
3  
4 346 electrochemical ones. The accurate and highly sensitive proposed method offers several  
5  
6 advantages in terms of simplicity, rapidity and efficiency. In fact, the results of the method  
7  
8 348 validation, performed according to Regulation (EC) No. 882/2004, demonstrate the method  
9  
10 conformity with provisions of Regulation (EC) No. 401/2006 and indicate that the procedure  
11  
12 can be successfully applied for accurate confirmation analyses. The expanded measurement  
13  
14 uncertainties and excellent results achieved in a proficiency test round confirmed the laboratory  
15  
16 technical competence and the method reliability in the determination of aflatoxins in real  
17  
18 352 samples, with the accuracy level corresponding to the requirements of the recent European  
19  
20 regulation. Therefore, the proposed method is well suited to satisfy the demands for accurate  
21  
22 354 and sensitive detection of aflatoxins with minimal sample preparation and cleanup steps.  
23  
24  
25  
26  
27 356

#### 28 29 **ACKNOWLEDGMENT**

30  
31  
32 358 Ministero della Salute (Rome, Italy) is gratefully acknowledged for providing the financial  
33  
34 support. Mr. Pasquale D'Antini and Mr G. Battafarano, from Istituto Zooprofilattico  
35  
36 360 Sperimentale di Puglia e Basilicata-Foggia, are gratefully acknowledged for technical  
37  
38 assistance.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4 362 **REFERENCES**

- 5  
6 Ahmed YA, Robinson RK. 1998. Selection of a Suitable Method for Analysis of Aflatoxins in  
7  
8 364 Date Fruits. *J. Agric. Food Chem.* 46(2):580-584.
- 9  
10  
11 Arranz I, Sizoo E, Van Egmond H, Kroeger K, Legarda TM, Burdaspal P, Reif K, Stroka J.  
12  
13 366 2006. Determination of Aflatoxin B<sub>1</sub> in Medical Herbs: Interlaboratory Study. *J. AOAC Int.*  
14  
15 89(3):595-605.
- 16  
17  
18 368 Bullerman LB. 1979. Significance of Mycotoxins to Food Safety and Human Health. *J. Food*  
19  
20 *Protect.* 42(1):65-86.
- 21  
22  
23 370 Cepeda A, Franco CM, Fente CA, Vázquez BI, Rodríguez JR, Prognon P, Mahuzier G. 1996.  
24  
25 Postcolumn excitation of aflatoxins using cyclodextrins in liquid chromatography for food  
26  
27 analysis. *J. Chromatogr. A*, 721(1):69–74.
- 28 372  
29  
30  
31 Chan D, MacDonald SJ, Boughtflower V, Brereton P. 2004. Simultaneous determination of  
32  
33 374 aflatoxins and ochratoxin A in food using a fully automated immunoaffinity column clean-up  
34  
35 and liquid chromatography–fluorescence detection. *J. Chromatogr. A*. 1059:13-16.
- 36  
37  
38 376 Davis ND, Guy ML, Diener U.L. 1981. Improved fluorometric-iodine method for determination  
39  
40 of aflatoxin in corn. *J. AOAC Int.* 64(5):1974-1976.
- 41  
42  
43 378 Dickens F, Jones HE. 1963. The carcinogenic action of aflatoxin after its subcutaneous injection  
44  
45 in the rat. *Brit. J. Cancer.* 17:691–698.
- 46  
47  
48 380 Dorner JW, Blankenship PD, Cole RJ. 1993. Performance of two immunochemical assays in the  
49  
50 analysis of peanuts for aflatoxin at 37 field laboratories. *J. AOAC Int.* 76:637-643.
- 51  
52  
53 382 Eaton DL, Groopman JD. 1994. *The Toxicology of Aflatoxins*. San Diego: Academic Press.
- 54  
55  
56 EURACHEM Guide. 1995. *Quantifying Uncertainty in Analytical Measurement*. 1<sup>st</sup> ed. Ellison  
57  
58 SLR, Rosslein M, Williams A. Eds.
- 59 384  
60

- 1  
2  
3 EURACHEM/CITAC GUIDE CG 4. 2000. Quantifying Uncertainty in Analytical  
4  
5  
6 386 Measurement. 2<sup>nd</sup> ed. Ellison SLR, Rosslein M, Williams A. Eds.  
7  
8  
9 European Commission. Regulation (EC) No. 1881/2006. 19 December 2006. Off. J. European  
10  
11 388 Union. L364:5-24.  
12  
13  
14 European Commission. Regulation (EC) No. 683/2004. 13 April 2004. Off. J. European Union.  
15  
16 390 L106:3-5.  
17  
18  
19 European Commission. Regulation (EC) No. 882/2004. 29 April 2004. Off. J. European Union.  
20  
21 392 L165:1–141.  
22  
23  
24 European Commission. Regulation (EC) No. 401/2006. 23 February 2006, Off. J. European  
25  
26 394 Union. L70:12–34.  
27  
28  
29 European Commission. Health & Consumer Protection Directorate-General,  
30  
31 396 SANCO/0895/2007. Guidelines for the implementation of Decision 2002/657/ec regarding some  
32  
33 contaminants (mycotoxins, dioxins and dioxin-like PCBs and heavy metals).  
34  
35  
36 398 European Commission. Decision (EC) No. 657/2002. 12 August 2002. Off. J. European Union  
37  
38 L221:8–36.  
39  
40  
41 400 European Commission. Directive (EC) No. 100/2003. 31 October 2003. Off. J. European Union.  
42  
43 L285:33-37.  
44  
45  
46 402 Garner RC, Kingfisher CN. 1979. Chemical Carcinogens and DNA. (Vol. 1). Grover PL Ed.  
47  
48 Boca Raton: CRC Press.  
49  
50  
51 404 Garner RC, Whattam MW, Taylor PJJ, Stow MW. 1993. Analysis of United Kingdom  
52  
53 purchased spices for aflatoxins using an immunoaffinity column clean-up procedure followed by  
54  
55 high-performance liquid chromatographic analysis and post-column derivatization with  
56 406 pyridinium bromide perbromide. J. Chromatogr. 648:485-490.  
57  
58  
59  
60

- 1  
2  
3  
4 408 Garcia-Villanova RJ, Cordon C, Gonzales Paramas AM, Aparicio P, Garcia Rosales ME. 2004.  
5  
6 Simultaneous Immunoaffinity Column Cleanup and HPLC Analysis of Aflatoxins and  
7  
8 410 Ochratoxin A in Spanish Bee Pollen. *J. Agric. Food Chem.* 52:7235-7239.  
9  
10  
11 Harwig J, Munro IC. 1975. Mycotoxins of possible importance in diseases of Canadian farm  
12  
13 412 animals. *Can. Vet. J.* 16(5):125-141.  
14  
15  
16 Holcomb M, Thompson HC. 1991. Analysis of Aflatoxins (B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>, and G<sub>2</sub>) in Rodent Feed  
17  
18 414 by HPLC Using Postcolumn Derivatization and Fluorescence Detection. *J. Agric. Food Chem.*  
19  
20 39:137-140.  
21  
22  
23 416 Hund E, Massart DL, Smeyers-Verbeke J. 2001. Operational definitions of uncertainty. *TrAC.*  
24  
25 20(8):394-406.  
26  
27  
28 418 International Agency for Research on Cancer (IARC). Lyon 1993. IARC Monographs on the  
29  
30 Evaluation of Carcinogenic Risks to Humans. 56:245-362.  
31  
32  
33 420 Italian Ministerial Decree 149/2004. June 2004. *Ital. Off. J.* 139:14-18.  
34  
35  
36 Joshua H. 1993. Determination of aflatoxins by reversed-phase high-performance liquid  
37  
38 422 chromatography with post-column in-line photochemical derivatization and fluorescence  
39  
40 detection. *J. Chromatogr. A.* 654(2):247-254.  
41  
42  
43 424 Kok WT, Van Neer TCH, Traag WA, Tuinstra LGMT. 1986. Determination of aflatoxins in  
44  
45 cattle feed by liquid chromatography and post-column derivatization with electrochemically  
46  
47 generated bromine. *J. Chromatogr.* 367(1):231-236.  
48 426  
49  
50  
51 Kussak A, Andersson B, Andersson K. 1995. Determination of Aflatoxins in Hot Chilli  
52  
53 428 Products by Matrix Solid-Phase Dispersion and Liquid Chromatography. *J. Chromatogr. A.*  
54  
55 708(1):55-60.  
56  
57  
58 430 Miller JC, Miller JN. 1993. *Statistics for Analytical Chemistry*. 3<sup>rd</sup> ed. New York:Ellis Horwood  
59  
60 PTR Prentice-Hall.

- 1  
2  
3  
4 432 Muscarella M, Lo Magro S, Palermo C, Centonze D. 2007. Validation according to European  
5  
6 Commission Decision 2002/657/EC of a confirmatory method for aflatoxin M<sub>1</sub> in milk based on  
7  
8  
9 434 immunoaffinity columns and high performance liquid chromatography with fluorescence  
10  
11 detection. *Anal. Chim. Acta.* 594:257-264.  
12  
13 436 Niedwetzki G, Lach G. 1994. Determination of aflatoxins in food by use of an automatic work  
14  
15 station. *J. Chromatogr. A.* 661:175-180.  
16  
17  
18 438 Nordic Committee on Food Analysis (NMKL). 2003. Estimation and expression of  
19  
20 measurement uncertainty in chemical analysis, 2<sup>nd</sup> ed. NMKL Secretariat. Finland procedure  
21  
22  
23 440 No. 5.  
24  
25  
26 Papadopoulou-Bouraoui A, Stroka J, Anklam E. 2002. Comparison of Two Post-Column  
27  
28 442 Derivatization Systems, Ultraviolet Irradiation and Electrochemical Determination. *J. AOAC*  
29  
30 *Int.* 85:411-416.  
31  
32  
33 444 Park DL, Miller BM, Hart LP, Yang G, McVey J, Page SW, Pestka J, Brown LH. 1989.  
34  
35 Enzyme-linked immunosorbent assay for screening aflatoxin B<sub>1</sub> in cottonseed products and  
36  
37 446 mixed feed: collaborative study. *J. AOAC.* 72(2):326-332.  
38  
39  
40 Peraica M, Radić B, Lucić A, Pavlović M. 1999. Toxic effects of mycotoxins in humans.  
41  
42 448 *Bulletin of World Health Organization.* 77:754–766.  
43  
44  
45 Roch OG, Blunden G, Coker RD, Nawaz S. 1992. The development and validation of a solid  
46  
47 450 phase extraction/HPLC method for the determination of aflatoxins in groundnut meal  
48  
49 *Chromatographia.* 33:208-212.  
50  
51  
52  
53 452 Saad AM, Abdelgadir M, Moss M.O. 1995. Exposure of infants to aflatoxin M<sub>1</sub> from mothers'  
54  
55 breast milk in Abu Dhabi, UAE. *Food Addit. Contam.* 12(2):225-261.  
56  
57  
58 454 Scholten JM, Spanjer MC. 1996. Determination of Aflatoxin B<sub>1</sub> in Pistachio Kernels and Shells.  
59  
60 *J. AOAC Int.* 79(6):1360-1364.

- 1  
2  
3  
4 456 Sobolev VS, Dorner JW. 2002. Cleanup Procedure for Determination of Aflatoxins in Major  
5  
6 Agricultural Commodities by Liquid Chromatography. *J. AOAC Int.* 85(3):642-645.  
7  
8  
9 458 Sobolev VS. 2007. Simple, Rapid, and Inexpensive Cleanup Method for Quantitation of  
10  
11 Aflatoxins in Important Agricultural Products by HPLC. *J. Agric. Food Chem.* 55:2136-2141.  
12  
13  
14 460 Tavčar-Kalcher G, Vrtač K, Pestešek U, Vengušt A. 2007. Validation of the procedure for the  
15  
16 determination of aflatoxin B<sub>1</sub> in animal liver using immunoaffinity columns and liquid  
17  
18 462 chromatography with postcolumn derivatization and fluorescence detection. *Food Control.*  
19  
20 18(4):333-337.  
21  
22  
23 464 Thompson M. 2000. Recent trends in inter-laboratory precision at ppb and sub-ppb  
24  
25 concentrations in relation to fitness for purpose criteria in proficiency testing. *Analyst.* 125:385–  
26  
27 386.  
28 466  
29  
30 Torres Espinosa E, Acuna Askar K, Naccha Torres LR, Montoya Olvera R, Castellon Santa  
31  
32 Anna JP. 1995. Quantification of aflatoxins in corn distributed in the city of Monterrey, Mexico.  
33 468 *Food Addit. Contam.* 12(3):383-386.  
34  
35  
36  
37  
38 470 Trucksess MW, Stack ME. 1994. Enzyme-linked immunosorbent assay of total aflatoxins B<sub>1</sub>,  
39  
40 B<sub>2</sub>, and G<sub>1</sub> in corn: follow-up collaborative study. *J. AOAC Int.* 77(3):655–658.  
41  
42  
43 472 Tuinstra LGMT, Haasnoot W. 1983. Rapid determination of aflatoxin B<sub>1</sub> in dutch feeding stuffs  
44  
45 by high-performance liquid chromatography and post-column derivatization. *J. Chromatogr. A.*  
46  
47 282:457-462.  
48 474  
49  
50  
51  
52  
53 476 Vázquez BI, Fente CA, Franco CM, Cepeda A, Mahuzier G, Prognon P. 1999. Preliminary  
54  
55 study on fluorimetric detection of aflatoxins Q<sub>1</sub>, P<sub>1</sub> and B<sub>1</sub> using heptakis-di-O-methyl-β-  
56  
57 cyclodextrin as post-column HPLC reagent. *Anal. Commun.* 36:5-7.  
58 478  
59  
60

- 1  
2  
3  
4       Waltking AE, Wilson D, Chan D, Dunn E, Humphries J, Kandler H, Sizoo E, Wilson D. 2006.  
5  
6       480   Liquid Chromatographic Analysis of Aflatoxin Using Post-Column Photochemical  
7  
8       Derivatization: Collaborative Study. J. AOAC Int. 89(3):678-692.  
9  
10  
11       482   Wilson TJ, Romer TR. 1991. Use of MycoSep multifunctional cleanup column for liquid  
12  
13       chromatographic determination of aflatoxin in agricultural product J. AOAC. 74(6):951-956.  
14  
15  
16       484   Zang LH, Lim JK. 2005. Hitachi HPLC aflatoxin photochemical post-column derivatization.  
17  
18       LC-GC North America. 23(suppl. 1):47.  
19

20  
21       486  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**FIGURE CAPTIONS**

1  
2  
3  
4  
5  
6 488 **Figure 1.** Chromatograms of a wheat bran sample spiked with AFB<sub>1</sub> and AFG<sub>1</sub> at 3.0 µgkg<sup>-1</sup>  
7  
8 and AFB<sub>2</sub> and AFG<sub>2</sub> at 0.75 µgkg<sup>-1</sup>. A) with and B) without on-line photochemical derivation.  
9  
10 Separation carried out by a ZORBAX Eclipse XDB C<sub>18</sub> column eluted with a mixture of water,  
11 490 methanol and acetonitrile 55:15:30 (v/v) at a flow rate of 1.0 mLmin<sup>-1</sup>; column temperature  
12  
13 40°C; injected volume: 20 µL; λ<sub>exc</sub> 365 nm and λ<sub>em</sub> 435 nm.  
14  
15  
16 492  
17  
18  
19  
20

21 494 **Figure 2.** Chromatograms of a) blank samples and b) spiked samples of almonds (A), pistachios  
22  
23 (B), maize (C) and oats (D). Almonds and pistachios samples have been fortified with AFB<sub>1</sub> and  
24  
25 AFG<sub>1</sub> at 2.0 µgkg<sup>-1</sup>, and AFB<sub>2</sub> and AFG<sub>2</sub> at 0.5 µgkg<sup>-1</sup>. Maize and oats samples have been  
26 496 fortified with AFB<sub>1</sub> and AFG<sub>1</sub> at 20 µgkg<sup>-1</sup>, and AFB<sub>2</sub> and AFG<sub>2</sub> at 5 µgkg<sup>-1</sup>. Experimental  
27  
28 conditions as reported in Figure. 1.  
29  
30  
31 498  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Validation of a confirmatory analytical method for the determination of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> in foods and feed ...

**Table 1.** Performance parameters for aflatoxins in foodstuffs.

$y = a + bx^a$					
Aflatoxin	$b \pm SD$	$a \pm SD$	$r^b$	LOD <sup>c</sup>	LOQ <sup>c</sup>
<b>B<sub>1</sub></b>	0.87 ± 0.01	0.02 ± 0.01	0.9998	0.04	0.11
<b>B<sub>2</sub></b>	2.31 ± 0.04	0.02 ± 0.01	0.9996	0.01	0.04
<b>G<sub>1</sub></b>	0.368 ± 0.009	0.014 ± 0.009	0.9991	0.08	0.24
<b>G<sub>2</sub></b>	0.87 ± 0.01	0.005 ± 0.004	0.9996	0.02	0.05

<sup>a</sup> y is the signal in luminescence unit (LU); x is the value of concentration in µg L<sup>-1</sup>. <sup>b</sup> Correlation coefficient. <sup>c</sup> Values expressed in µg L<sup>-1</sup>.



50 **Table 2.** Repeatability for the determination of aflatoxins in spiked wheat bran samples.

Aflatoxin	Operator	Contamination Level ( $\mu\text{g kg}^{-1}$ )	Determined Concentration Mean $\pm$ SD (n = 6)	RSD <sub>r</sub> <sup>a</sup> (%)
<b>B<sub>1</sub></b>	1	1	1.05 $\pm$ 0.04	3.5
	2		1.0 $\pm$ 0.1	9.8
	3		1.05 $\pm$ 0.06	5.5
	1	2	1.9 $\pm$ 0.1	7.1
	2		2.0 $\pm$ 0.2	10
	3		1.9 $\pm$ 0.1	4.9
	1	3	3.05 $\pm$ 0.14	4.8
	2		3.00 $\pm$ 0.08	2.7
	3		3.1 $\pm$ 0.3	9.5
<b>B<sub>2</sub></b>	1	0.25	0.27 $\pm$ 0.06	11
	2		0.25 $\pm$ 0.02	9.1
	3		0.26 $\pm$ 0.03	10
	1	0.50	0.46 $\pm$ 0.03	7.0
	2		0.49 $\pm$ 0.05	9.8
	3		0.47 $\pm$ 0.03	5.9
	1	0.75	0.77 $\pm$ 0.06	7.4
	2		0.75 $\pm$ 0.02	3.0
	3		0.76 $\pm$ 0.07	9.6
<b>G<sub>1</sub></b>	1	1	1.10 $\pm$ 0.08	7.7
	2		1.0 $\pm$ 0.1	10
	3		1.0 $\pm$ 0.1	9.2
	1	2	1.8 $\pm$ 0.2	11
	2		2.0 $\pm$ 0.2	9.0
	3		1.9 $\pm$ 0.1	6.2
	1	3	3.1 $\pm$ 0.2	7.0
	2		3.0 $\pm$ 0.1	4.4
	3		3.0 $\pm$ 0.3	9.2
<b>G<sub>2</sub></b>	1	0.25	0.27 $\pm$ 0.02	8.8
	2		0.24 $\pm$ 0.04	15
	3		0.27 $\pm$ 0.04	15
	1	0.50	0.46 $\pm$ 0.06	13
	2		0.51 $\pm$ 0.07	14
	3		0.47 $\pm$ 0.05	11
	1	0.75	0.77 $\pm$ 0.07	9.0
	2		0.74 $\pm$ 0.05	6.6
	3		0.77 $\pm$ 0.08	11

<sup>a</sup> Relative standard deviation under repeatability conditions.

Validation of a confirmatory analytical method for the determination of aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> in foods and feed ...

58 **Table 3.** Within-laboratory reproducibility for the determination of  
 59 aflatoxins in spiked wheat bran samples.

Aflatoxin	Contamination Level ( $\mu\text{g kg}^{-1}$ )	Determined Concentration RSD <sub>R</sub> <sup>a</sup>	
		Mean $\pm$ SD (n = 18)	(%)
B <sub>1</sub>	1	1.0 $\pm$ 0.1	9.9
	2	1.9 $\pm$ 0.2	11
	3	3.0 $\pm$ 0.2	6.8
B <sub>2</sub>	0.25	0.26 $\pm$ 0.03	12
	0.50	0.47 $\pm$ 0.05	11
	0.75	0.76 $\pm$ 0.06	7.6
G <sub>1</sub>	1	1.05 $\pm$ 0.15	14
	2	1.9 $\pm$ 0.3	15
	3	3.0 $\pm$ 0.2	8.1
G <sub>2</sub>	0.25	0.26 $\pm$ 0.05	20
	0.50	0.48 $\pm$ 0.09	19
	0.75	0.76 $\pm$ 0.08	10

62 <sup>a</sup> RSD<sub>R</sub>, relative standard deviation under within-laboratory reproducibility conditions.

64 **Table 4.** Repeatability and within-laboratory reproducibility for the determination of  
 65 aflatoxin B<sub>1</sub> in spiked durum wheat samples.

Aflatoxin	Contamination Level ( $\mu\text{g kg}^{-1}$ )	Operator	Determined Concentration RSD <sup>a</sup>	
			Mean $\pm$ SD	(%)
<i>Repeatability (n = 6)</i>				
B <sub>1</sub>	10	1	10 $\pm$ 1	10
		2	10.0 $\pm$ 0.8	7.9
	20	1	20 $\pm$ 1	6.1
		2	20.0 $\pm$ 1.5	7.4
	30	1	29.8 $\pm$ 1.6	5.2
		2	30 $\pm$ 2	5.7
<i>Within-laboratory Reproducibility (n = 12)</i>				
B <sub>1</sub>	10		9.9 $\pm$ 0.9	9.6
	20		20.2 $\pm$ 1.5	7.2
	30		29.9 $\pm$ 1.6	5.5

68 <sup>a</sup> RSD, relative standard deviation under repeatability (RSD<sub>r</sub>) and within-laboratory reproducibility  
 69 (RSD<sub>R</sub>) conditions.

**Table 5.** Recovery of aflatoxins in spiked wheat bran (foodstuffs) and durum wheat samples (feed materials).

Aflatoxin	Contamination Level ( $\mu\text{g kg}^{-1}$ )	Recovery $\pm$ SD (%)	Recommended Range (%)
<i>Foodstuffs (n = 18)</i>			
<b>B<sub>1</sub></b>	1	76.4 $\pm$ 6.5	70-110
	2	83.0 $\pm$ 9.8	
	3	92.2 $\pm$ 8.3	
<b>B<sub>2</sub></b>	0.25	79.5 $\pm$ 11.6	50-120
	0.50	82.1 $\pm$ 9.9	
	0.75	92.8 $\pm$ 8.0	
<b>G<sub>1</sub></b>	1	80.4 $\pm$ 9.6	70-110
	2	83.4 $\pm$ 11.1	
	3	94.6 $\pm$ 8.2	
<b>G<sub>2</sub></b>	0.25	89.2 $\pm$ 16.9	50-120
	0.50	82.0 $\pm$ 10.7	
	0.75	87.7 $\pm$ 10.1	
<i>Feed materials (n = 12)</i>			
<b>B<sub>1</sub></b>	10	75.7 $\pm$ 6.3	N.A. <sup>a</sup>
	20	74.3 $\pm$ 4.7	
	30	72.6 $\pm$ 3.9	

<sup>a</sup> N.A., Not Available; a legal value has not been established.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1

*M. Muscarella et al.*

Figure 1  
839x1187mm (150 x 150 DPI)




Figure 2

*M. Muscarella et al.*

Figure 2  
839x1187mm (150 x 150 DPI)