



Impact of food consumption habits on the pesticide dietary intake: Comparison between a French vegetarian population and the general one.

Marieke van Audenhaege, Fanny Héraud, Céline Menard, Juliette Bouyrie,
Sophie Morois, Gloria Calamassi-Tran, Sébastien Lesterle, Jean-Luc J.L.
Volatier, Jean-Charles Leblanc

► To cite this version:

Marieke van Audenhaege, Fanny Héraud, Céline Menard, Juliette Bouyrie, Sophie Morois, et al.. Impact of food consumption habits on the pesticide dietary intake: Comparison between a French vegetarian population and the general one.. Food Additives and Contaminants, 2009, 26 (10), pp.1372-1388. 10.1080/02652030903031171 . hal-00573899


HAL Id: hal-00573899

<https://hal.science/hal-00573899>

Submitted on 5 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Impact of food consumption habits on the pesticide dietary intake: Comparison between a French vegetarian population and the general one.**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2009-014.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	07-May-2009
Complete List of Authors:	Van Audenhaege, Marieke; Agroparistech Héraud, Fanny; French Food Safety Agency, Office of Scientific Support to the Risk Assessment MENARD, Céline; French Food Safety Agency, Office of Scientific Support to the Risk Assessment Bouyrie, Juliette; Agroparistech Morois, Sophie; Agroparistech Calamassi-Tran, Gloria; French Food Safety Agency, Office of Scientific Support to the Risk Assessment Lesterle, Sébastien; French Food Safety Agency, Crops and Environment Department Volatier, Jean-Luc; French Food Safety Agency, Office of Scientific Support to the Risk Assessment Leblanc, Jean-Charles; French Food Safety Agency, Office of Scientific Support to the Risk Assessment
Methods/Techniques:	Exposure assessment
Additives/Contaminants:	Pesticide residues
Food Types:	

SCHOLARONE™
Manuscripts

For Peer Review Only

1 2 Introduction 3

4 Pesticide residues are regularly found in foodstuffs, as revealed by the European Commission
5 which recently acknowledged that they were detected in 45.7% of the 62,569 samples
6 analysed throughout the European Union in 2005 (European Commission, 2007). Pesticide
7 residues are mainly present in cereals, vegetables and fruit. Thus, vegetables and fruit are
8 found to contribute respectively to 75.8% and 20.5% of the estimated daily intakes of the
9 Korean population (Chun and Kang, 2003). Some pesticides, mostly Persistent Organic
10 Pollutants (POPs), also concentrate all along the food chain and contaminate foodstuffs of
11 animal origin (IPCS, 1996). These commodities are found to be the main sources of
12 organochlorine dietary exposure for the Australian population (Kannan *et al.*, 1997) as well as
13 for the Spanish one (Herrera *et al.*, 1996).

14 Considering pesticide residues in foodstuffs as a public health issue, it is important to
15 characterise which population is at risk. The pesticide dietary intake is a function of both
16 residue levels in food and consumption habits. Vegetarians typically consume more fruit,
17 vegetables and cereal products compared to individuals of the general population whose diet
18 also includes animal-derived products. The aim of this study is to compare the impact of
19 consumption habits of a French vegetarian population and the general one on their pesticide
20 dietary intake.

21 22 Material and methods 23

24 Consumption data 25

26 French general population. Consumption data for the general population came from the
27 French individual consumption survey “*enquête Individuelle et Nationale de Consommation
28 Alimentaire*” (INCA), conducted in 1999. The original sample of 1985 healthy adults (aged 15
29 and over) was representative of the French population through the stratification (region of
30

Deleted: are privileged

Deleted: rs

Deleted: of

Deleted: Are they consequently more exposed to pesticides than the general population?

Formatted

Formatted

Deleted: INCA

residence, agglomeration size) and the use of the quota method (age, gender, household size, individual occupation and socio-professional status) (Volatier, 2000). Subjects were asked to complete a seven-day food record diary (consecutive days) as well as other questionnaires on anthropometrical and socio-economic factors. Participants could estimate portion sizes through comparison of their actual consumption with photographs compiled in a manual adapted from the SuViMax picture booklet (Hercberg *et al.*, 1994). Adult people identified as under-reporters by comparing their reported energy intake to their basal metabolic rate as estimated from the Schofield equations (Black, 2000) were excluded from the study. Thus, the final sample contained 1474 individuals.

The vegetarian population. Vegetarian subjects were recruited among adherents of the main French vegetarian or vegan societies between March and July 1997, through questionnaires published in the society's journal or directly distributed to the adherents (Leblanc *et al.*, 2000). The total participation rate was 19.5% with 145 respondents. After deletion of incomplete questionnaires, consumption data of 137 individuals were available for analysis.

Deleted: answers

Deleted:

For each participant, consumption data were collected in a 5-day food diary. Age, gender, weight and height were also listed. A classification of the vegetarian subjects was adopted on the basis of their effective food intakes (Table 1): 26 subjects considering themselves as vegetarians but having an omnivorous diet were gathered in the pseudo-vegetarian (OMN) group. The lacto-vegetarian (LV) group was composed of 41 subjects that only stand for dairy products in their diet. On the contrary, in the ovo-lacto-vegetarian (OLV) and pesco-lacto-vegetarian (PLV) groups – 43 and 16 adults – subjects get their proteins via both dairy products and respectively eggs or fish consumptions. At last, strict vegetarians, for whom all commodities of animal origin are excluded from the diet, were gathered in the vegan (VG) group (11 subjects). Although the number of subjects in each subgroup is rather low, the homogeneity of eating habits within each subgroup allows to consider it sufficient for further

1
2 analysis (Leblanc, 2001). No adjustment was done according to the potential food
3 consumption underreporting.
4
5
6
7
8

Deleted: ¶

9 Residue data

10
11 Maximum Residue Limits (MRLs) and Acceptable Daily Intake (ADI). To assess the impact of
12 consumption habits on the pesticide dietary intake, a fixed contamination level, set at the
13 MRL, was considered whatever the population. Priority was given to MRLs in force within
14 the European Union in July 2007, and next to French MRLs. Only substances, with an ADI
15 and at least one MRL, were included in the study (N = 421) (Table 2).
16
17

18 Sensitivity analysis on fish consumption habits. For a given pesticide, commodities with no
19 existing MRL were not taken into account in the intake assessment. In particular, no MRL are
20 actually defined at the European level for fish or fish-based products. However, several
21 studies have already shown that they are potentially contaminated by pesticides, especially
22 organochlorine compounds (Hites *et al.*, 2004), and could even explain the presence of
23 pesticides in human samples (Rivas *et al.*, 2007). Moreover, PLV eat large amounts of fish as
24 a source of proteins. A sensitivity analysis on fish consumption habits was consequently
25 conducted on the Aldrin case.

26 The total Aldrin intake was first calculated based on the contribution of all food items for which an MRL is defined. Then, it was compared to a total intake calculated considering a fictive maximum limit for the fish and fish-based products.

27 This fictive value is assumed at the MRL defined for an other product of animal origin: meat.

28 As Aldrin is a lipophilic compound, its MRL is defined on the fat. Thus, an average fat
29 content of 10% was considered to define the fictive maximum limit for the fish and fish-based
30 products.

Deleted: of meat products

Deleted: express

Deleted: MRL

Deleted: on a total weight basis

Deleted: The fish residue level was set at the MRL defined for meat, corrected by an average meat fat content of 10% in order to get a residue level for the whole product.

51 Recipes

In order to match consumption data to MRLs, which are established on Raw Agricultural Commodities (RAC), the 895 food items defined in the INCA Survey and the 545 food items defined in the Vegetarian Consumption Survey were respectively broken down into 153 and 137 RAC. 402 standardized recipes, which have been defined taking account of industrial processes, home cooking habits and edible portions for the INCA Survey (Volatier, 2000), have been used. For the few vegetarian food items without any standardized recipe available, a recipe has been specifically defined according to the same approach than the one used to create the standardized recipes.

Deleted: and 254 recipes were respectively defined for the general and vegetarian populations in order to take account of industrial processes, home cooking habits and edible portions.

Intake assessment

Formatted

The Theoretical Maximum Daily Intake (TMDI) (in mg. kg⁻¹ bw. day⁻¹) as defined by WHO (WHO, 1997) was estimated for each pesticide at the individual level, according to equation 1. The average was then derived for each population and expressed as a percentage of the ADI.

$$(1) \quad TMDI_{p,i} = \frac{\sum_{a \in A_p}^{\sum_{k=1}^{N_i}} C_{k,a,i} \times MRL_{p,a}}{N_i \times bw_i}$$

Where:

TMDI_{p,i}: Theoretical Maximum Daily Intake of the individual *i* for the pesticide *p*,

C_{k,a,i}: quantity of foodstuff *a* consumed by individual *i* on day *k*,

A_p: list of foodstuffs for which a MRL is defined for the pesticide *p*,

MRL_{p,a}: Maximum Residue Limit for the pesticide *p* in the foodstuff *a*,

N_i: number of days of consumption for the individual *i*,

bw_i: body weight of individual *i*.

1
2 The contribution of ten food groups (fruit, vegetables, cereals, potatoes, other products of plant
3
4 origin (mainly oils and oilseeds), milk and dairy products, eggs, meat and other products of
5 animal origin (honey), water and wine) was characterized as a ratio between the food group
6 intake and the total one.
7
8

Deleted: vegetable

12 Statistical analysis

13
14 The statistical analysis was conducted using the SAS® software (SAS, Cary, NC 27513)
15
16 running on the UNIX® system in order to characterize the determining factors of the total
17
18 dietary intake and its differences among the studied populations.

Deleted: S

19 First, the physiological characteristics (age and body mass index) were analysed in order to
20 ensure the two populations to be comparable. Consumption habits were also compared
21 between the general population and the vegetarian cohort based on individual average intakes.
22
23 As variables were not shown to have a Gaussian distribution according to the Shapiro-Wilk
24 test, all comparisons between the general population and the vegetarian cohort were based on
25 Kruskal-Wallis.
26
27

Deleted: using Student tests

Deleted: in terms of risk of exposure

Deleted: five

Deleted: subgroups using tests of Bonferroni (Dunn)

Deleted: not

28 Into the vegetarian cohort, the five subgroups were considered to be independent due to the
29 fact one individual can only belong to only one subgroup. As a Gaussian distribution was not
30 assured within the studied subgroups, physiological comparisons were based on Kruskal-
31 Wallis tests. Similar tests were conducted to assess whether mean consumption levels were
32 significantly different between two groups – for example the PLV group *vs.* the four other
33 vegetarian subgroups.
34
35

Deleted: population

Deleted: repartition

Deleted: the

Deleted: Tests of Bonferroni (Dunn) were chosen

36 Then, a specific analysis was done on few pesticides chosen according to their main
37 contributors to the total dietary intake previously described by Menard *et al.* (2008):
38

- 39 - products of animal origin for Aldrin,
40 - fruit and vegetables for Diazinon,
41
42

1 - cereals for Chlorpyrifos-methyl.

2 Carbamates (Aldicarb, Tri-allate, Propineb and Thiram) were also chosen for a rapid analysis
3 (linear regression between subgroup average consumptions and TMDIs) because of total
4 intakes differences between the population subgroups.

Deleted: D

5 For Aldrin, Diazinon and Chlorpyrifos-methyl, the comparison was completed with multi
6 linear regressions (ANOVA procedure) in order to explain the total dietary intake given by
7 the TMDI/ADI ratio. The determining factors were consumptions levels of the different food
8 groups, age, gender, Body Mass Index and pertinent interactions between those variables (for
9 example BMI*gender).

Deleted: these pesticides

Deleted: and

Results

Description of the populations

Physiological characteristics of the studied populations. Characteristics of both populations are included in Table 1 according to diet group. The sex ratio (M/F) is similar between the studied populations (0.79 and 0.83 in the vegetarian and the general population respectively).

Age distributions are relatively close in both populations with mean age of 49 ± 15 years for vegetarian population and 43 ± 18 for the general one (not significantly different).

The vegetarian population shows a lower mean BMI compared to the general population's one (21.3 ± 2.7 kg. m⁻² vs. 23.5 ± 3.8 , not significantly different). Indeed, respectively 74.5 and 64% of the vegetarian and the general populations have a BMI between 18.5 and 24.9.

The vegetarian group includes less overweight or obese subjects (BMI higher than 25) than the general one (10.2% vs. 30.6%) and more thin subjects with BMI below 18.5 (15.3% vs. 5.4%).

Deleted: population

Some differences can also be observed among the subgroups. The mean age of the LV subgroup (42.2 ± 11.9 years) is significantly the lowest of the vegetarian subgroups ($p=0.003$) and especially when compared to the OMN subgroup (56.6 ± 19.4 years, $p=0.004$). Women

Deleted: 004

Deleted: $<10^{-3}$

1
2 are dominant in all subgroups except in the LV (M/F ratio=1.21) and OLV (M/F ratio=1.05)
3 ones. The BMI of the PLV subgroup is shown to be significantly the lowest of the vegetarian
4 subgroups ($p=0.002$), below the LV ($p=0.004$), OMN ($p=0.012$) and OLV ($p=0.005$) ones.

Deleted: 015
Deleted: 012
Deleted: 049

5
6 *Consumptions habits of the studied populations.* As shown in Table 1, the total consumption,
7 beverages excepted, is higher, but not significantly different, in the general population than in
8 the vegetarian subgroups (1241.4 g. day⁻¹ for the general population vs. 782.6 – 1174.2 g.day⁻¹
9 for the vegetarian subgroups), especially when it is compared to the PLV group in which
10 subjects eat the least.

Deleted: quasi non-eaten

11
12 The general population consumes some products, which are hardly even eaten by vegetarians
13 such as eggs ($p<10^{-4}$, 29.0 g. day⁻¹ vs. 0.1 – 4.0 g. day⁻¹), dairy products ($p<10^{-4}$, 299.6 g. day⁻¹
14 vs. 2.9 – 119.4 g. day⁻¹) and other products of animal origin ($p<10^{-4}$, 161.2 g. day⁻¹ for the
15 general population vs. 1.5 – 9.0 g. day⁻¹ for the vegetarian subgroups). Vegetarians would also
16 be more abstemious than subjects of the general population ($p<10^{-4}$): their wine consumption
17 goes from 42.6 to 105.2 g. day⁻¹ while its reaches 160.3 g. day⁻¹ for the non vegetarian
18 population.

19
20 On the contrary, compared to the general population, vegetarians eat more fruit (247.7 - 317.6
21 g. day⁻¹ for OLV, VG, LV and OMV groups, 124.7 g. day⁻¹ for the PLV group vs. 173.4 g.
22 day⁻¹ for the general population), vegetables (286.1 – 381.8 g. day⁻¹ vs. 170.1 g. day⁻¹), cereals
23 (189.9 and 206.3g. day⁻¹ for PLV and LV groups, 128.7 - 157.3g. day⁻¹ for VG, OLV and
24 OMV groups vs. 173.0 g. day⁻¹ for the general population) and potatoes (53.3-86.9 g. day⁻¹ for
25 VG, LV, OLV, 41.0 and 22.3 g. day⁻¹ for OMV and PLV groups vs. 52.6 g. day⁻¹ for the
26 general population) (figures not statistically different). Finally, consumption of other products
27
28 **Deleted:** oilseeds
29 of plant origin would be similar between the general population (22.3 g. day⁻¹) and
30 vegetarians groups (18.8 – 36.3 g. day⁻¹).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The LV diet differs from the other vegetarian ones: fruit, vegetables and cereals are consumed in the same amount while one or several foodstuffs are predominant in other diets. The OLV diet and the VG one (to a lower extent) distinguish themselves by a low cereal consumption (respectively 137.6 and 157.3 g. day⁻¹ vs. 173.0 – 206.3 g. day⁻¹ for the other vegetarian subgroups): meals are essentially composed of fruit and vegetables (respectively 62% and 64% of the total diet). The PLV diet is characterized by the opposite tendency: individuals of this subgroup have the highest cereal consumption (206.3 g. day⁻¹ vs. 128.7 – 189.9 g. day⁻¹ for the other vegetarian subgroups) and the lowest fruit consumption among vegetarians (124.7 g. day⁻¹ vs. 247.7 – 317.6 g. day⁻¹ for the other vegetarian subgroups).

Deleted: quasi-

Deleted: amount

Comparison of pesticide dietary intakes. Table 2 presents detailed results for pesticides with at least one TMDI above the ADI (detailed results for the other pesticides are provided in a supplemental material table).

Deleted: the

Deleted: expressed in percentage of the ADI for each pesticide in the general population and in the five vegetarian subgroups

Among the 421 pesticides taken into account, only 48 molecules (11.4%) have a TMDI value above the ADI for at least one of the studied populations, half of them (23 molecules) representing a potential health risk in all subgroups. However, some differences exist between the two populations. The number of pesticides with TMDI above ADI is always higher in vegetarian subgroups (30 molecules for PLV, 41 for LV, 42 for OMN, 43 for VG and 44 for OLV) than in the general population (29 molecules). It should however be noticed that in terms of number of pesticides exceeding the ADI, the PLV group is extremely close to the general population (30 vs. 29) and that other vegetarians groups are quite similar.

All the pesticides with TMDI higher than ADI for the general population have also at least one TMDI above ADI in one of the vegetarian subgroups. According to Table 2, most of the pesticides with TMDI above ADI are detected in all the vegetarian subgroups (25/48). Few pesticides have a TMDI higher than the ADI only for one of the vegetarian subgroups (5/48):

1 Imazalil (VG), Formetanate hydrochlorid (OLV), Deltamethrin (PLV), Diquat (PLV) and
2 Fenpropimorph (OLV). When pesticides are detected in three vegetarian subgroups (5/48), it
3 corresponds mostly to the [OMN, OLV, VG] subgroups (3/48) but also to the [LV, PLV, VG]
4 and the [OMN, LV, OLV] ones (2/48). Finally, when pesticides are detected in four out of the
5 five vegetarian subgroups (13/48), it always corresponds to the OMN, LV, OLV and VG
6 subgroups.

7 ADI overruns are different from one pesticide class to another or even in the same class. The
8 pesticides with TMDI above ADI are mainly organophosphates, organochlorine compounds
9 and carbamates. All populations confounded, respectively 44%, 36% and 17% of the
10 organophosphates, organochlorines and carbamates have a TMDI above ADI vs. 5% for the
11 other pesticides. The organochlorine and organophosphate classes appear with TMDI levels
12 significantly higher in both the vegetarian ($p=0.048$ and 10^{-4} respectively) and the general
13 population ($p<10^{-4}$ and 0.001 respectively) than the other pesticides classes. Carbamates also
14 appear with TMDI significantly higher than the other pesticides but only for the vegetarian
15 population ($p=0.007$). In another hand, some pesticide classes would not present any long-
16 term risk through dietary intake in both studied populations. For instance, the triazine and
17 urea classes have no ADI exceedance. In the class of pyrethroïds, only Deltamethrin intakes
18 are above ADI for the PLV subgroup (TMDI = 101.9% ADI). Finally, only two molecules are
19 potentially representing a risk in the azole class: Imazalil for the VG subgroup (TMDI =
20 102.5% ADI) and Prochloraz for all vegetarian subgroups except the PLV one (TMDI = 107.5
21 to 212.3% ADI).

45 *Determining factors of total pesticide intake*

46 *Impact of the physiological characteristics.* The impact of physiological characteristics of the
47 populations has been analysed through multi-linear regression on few pesticides (Aldrin,
48

49 Formatted

Diazinon, Chlorpyrifos-methyl and several carbamates). The structure of the population in term of age or sex ratio is similar between the populations and did not have any significant effect on the total intake. Nevertheless, the interaction BMI*diet was found to explain the Chlorpyrifos-methyl intake variability ($p < 0.07$) as well as the Aldrin intake one ($p < 0.07$).

Deleted: and its interaction with the

Deleted: the interaction between BMI and the diet for

A strong impact of fruit, vegetables and cereal consumptions on pesticide dietary intake.

Results of the specific food contributions analyses reveal a strong correlation between the total dietary intake of carbamates and the consumption of plant origin foodstuffs for all vegetarian subgroups (Figures 1a & 1b). By instance, TDMI is linear linked to the total average consumption of vegetables, fruit and wine for Tri-allate ($R^2=0.99$), Thiram ($R^2=0.97$) and Propineb ($R^2=0.85$) and to potato, vegetables and fruit consumption levels for Aldicarb ($R^2=0.90$). But food contributions are not the same between the different vegetarian groups.

Deleted: vegetal

Deleted: -

For Carbaryl, the intake of the PLV subgroup, whose TDMI is lower, mainly comes from cereals. On the contrary, individuals of the other groups, whose intake mainly results from the contribution of vegetables, are more exposed (Figure 1c). The same conclusion rises for Aldicarb: individuals of the PLV subgroup have lower intakes while their cereal consumption is the highest among vegetarian subgroups (TDMI = 97.6% ADI vs. 123.2% to 174.5% ADI). Organophosphate intakes are also significantly different between the five vegetarian subgroups (respectively $p=0.003$ and $p=0.004$ for Chlorpyrifos-methyl and Diazinon). For Chlorpyrifos-methyl, cereals are major contributors to the dietary exposure (Figure 2) and PLV subjects are more exposed than the other groups. The classification order of vegetarian subgroups to Chlorpyrifos-methyl exposure is the same than for cereal consumption levels: PLV > LV > VG > OLV > OMN (Table 1). For Diazinon, for which vegetarians are more exposed than the general population (except PLV group), fruit are major contributors with respectively 54.2%, 58.7%, 63.7%, 67.9% and 69.7% of the Diazinon TMDI for the PLV, LV, OMN, OLV and VG subgroups (Figure 2). Therefore the OLV subgroup for which fruit

1 consumption level is the highest (Table 1) has the highest intake (TMDI = 130.4% ADI). On
2 the contrary, subjects of the PLV subgroup for which fruit consumption level is the lowest
3 (Table 1), are less exposed than subjects of the OMN ($p=0.002$), LV ($p<10^{-4}$) and OLV
4 ($p=0.001$) subgroups (TMDI = 55.7% ADI vs. TMDI = 103.7%, 115.1% and 130.4% ADI
5 respectively).

6
7
8
9
10
11
12 *The PLV diet: an apparent protection against organochlorines?* The Aldrin dietary intake is
13 higher in the general population than in the vegetarian cohort (TMDI = 348.7% ADI vs.
14 TMDI = 146.2 to 183.9% ADI), mainly explained by the contribution of animal-origin
15 commodities (respectively 59.5% and 9.3% TDMI for meat and dairy products) (Figure 2).
16 PLV subjects were previously shown to be the least exposed subjects to Aldrin (Figure 2) but
17 fish-derived products which represent a significant part of their diet were not taken into
18 account in the assessment. When a fictive contamination is attributed to these foodstuffs, the
19 order is reversed: subjects from the OMN, LV, OLV and VG groups become less exposed to
20 Aldrin than PLV subjects (TMDI > 300% ADI) (Figure 3).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Deleted: fictive

Deleted: one

Formatted

Discussion

48 out of the 421 pesticides included in the study may theoretically represent a long-term risk
49 for the French general population or any of the 5 vegetarian subgroups due to the dietary
50 intake. These molecules correspond mainly to the carbamate, organochlorine and
51 organophosphate classes. On the contrary, most of pesticides belonging to the azole,
52 pyrethroïd, triazine and urea classes have TMDIs below ADI. Vegetarians are critically
53 exposed to a higher number of pesticides ($N = 48$) than subjects of the general population (N
54 = 29). All the pesticides with TMDI above ADI for the general population would be detected
55 as such in at least one vegetarian subgroup.

1
2 Specific analysis about food contributions shows that fruit and vegetables consumption levels
3 explain vegetarians overexposure to carbamates and Diazinon. The PLV subgroup, identified
4 with a high cereal consumption level, has consequently higher TDMI for pesticides with
5 cereals as main contributors, like Chlorpyrifos-methyl. On the contrary, vegetarians are less
6 exposed to organochlorine compound like Aldrin, due to a lower consumption of animal
7 origin commodities.
8
9

10 In order to eliminate any methodological bias in the comparison analysis, physiological
11 characteristics of the two populations have first been compared.

12 No significantly differences could be shown for sex ratio, age distributions and BMI. Moreover, consumption surveys used for this analysis are quite comparable. First, they have been conducted within the same timeframe, in the late 1990s. A new individual survey conducted among the French general population in 2006 – 2007 tends to show that consumption levels of fruit and vegetables haven't significantly changed since 1999 (Lafay, 2009). Thus the trends observed in this study should still be relevant. Secondly, both surveys are based on self-administered food diaries.

13 Food diary allows to estimate a wide range of foods intake (Willet, 1998), which is adapted to
14 this issue, as many food items may contain pesticides residues. Nevertheless, they can be
15 associated with an alteration of the eating behaviour, conducting to a potential
16 underestimation of real consumption level. They also require a high motivation from the
17 subject which can lead to a poor response rate or a lowering quality of answers according to
18 the days of inquiry (Willet, 1998). The food underreporting has been adjusted for subjects of
19 the general population. It has not been the case for vegetarian ones, because of the low
20 number of vegetarian subjects available for this analysis. This methodological bias may have
21 minimized the differences of exposure between the two groups.

22 Nevertheless, dietary habits appear to be the main explanation to the observed differences. If
23 amount levels of fruit, vegetables and cereals consumed are about equivalent for both subjects

Deleted: The BMI is found to explain the Aldrin and Chlorpyrifos-methyl intake variability. In fact, the ratio "amount of food consumed / body weight" is part of the TMDI equation. According to Schutz *et al.* (1989), if the basal metabolic rate (BMR) increases with weight, when expressed on a body weight basis, it is less important for obese individuals than for individuals with normal weight (Schutz *et al.*, 1989). This could indirectly lead to an average ratio "amount of food consumed / body weight" less important in the general population than in the vegetarian one, and could contribute to a higher total intake in the vegetarian population

Deleted: ¶

Deleted: N

of the LV subgroup and the general population, one or several of these foodstuffs are predominant in the other vegetarian diets, in which foods of animal origin are limited or evicted. This "substitution" of animal commodities by plant origin ones impacts both the consumption and contamination parameters of the TDMI equation.

First, it increases the consumption level of plant origin products ($591.4 \text{ g. day}^{-1}$ for the general population vs. between 712.1 and $1037.1 \text{ g. day}^{-1}$ for the vegetarian subgroups), all the more that they are less energy-dense than animal origin foodstuffs so larger amounts are consequently needed to reach the same energy level.

Secondly, it impacts the contamination level as the MRL set for an animal origin product is replaced by a MRL set for a product of plant origin. But MRLs are not systematically defined for animal commodities. In fact, they are set only when feed may contain residues at a sufficiently high level ($> 0.1 \text{ mg. kg}^{-1}$) and when metabolism studies show they may be found at levels higher than 0.1 mg. kg^{-1} in products of animal origin (European Commission, 2002).

Moreover, except for pesticides also used as veterinary drugs or biocides, residues in food of animal origin result from an indirect usage via feed and water. Levels are consequently generally less important than for plant origin food on which the farmer may directly apply the pesticide. Exception can be underlined for lipophilic compounds such as for organochlorine and some organophosphate compounds, which tend to concentrate in animal commodities (IPCS, 1996). This explains why dietary intake is higher for the vegetarian population than for the general one, excepted for few compounds such as organochlorine ones.

The impact of the substitution is also observable between the different classes of plant origin commodities. For some pesticides, such as carbamates and Diazinon (organophosphate), intakes are mainly indicated by fruit and/or vegetables contributions. This explains why OLV and VG are more exposed than the other population subgroups. On the contrary, subjects of the PLV subgroup are less exposed to these compounds but more exposed to pesticides for

Deleted: as

Deleted: , 1

1 which cereals are the main contributors (Pirimiphos-methyl, Malathion and Chlorpyrifos-
2 methyl).

3 This substitution seems to have little impact on the total intake of some pesticides, such as
4 Carbaryl. In fact, Carbaryl MRLs are set at 0.05 mg. kg⁻¹ in meat, eggs, dairy products and
5 most of vegetables and oil seeds, whereas it is 10 to 100 times higher than MRL set for
6 cereals, wine, tomatoes and olives (0.5-1, 3, 5 and 5 mg. kg⁻¹ respectively). Some differences
7

8 among the ~~subgroups~~ exposure are consequently expected. In fact, no clear effect of
9 consumption habits on the residue intake can be shown. This could be explained by the fact
10 that TMDI as expressed in this study depends not only on the residue intake but also on the
11 ADI level. Indeed, it has previously been shown that when the ADI is very low, due to high
12 toxicity of the pesticide, residue and consumption levels have a very limited impact on the
13 TMDI (Menard *et al.*, 2008). According to Table 2, it can be determined that the ADI for
14 pesticides with TMDI > ADI for all population groups is of 0.004 mg. kg bw⁻¹. day⁻¹ on
15 average, 25 times lower than the average ADI (0.09 mg. kg bw⁻¹. day⁻¹) based on pesticides
16 with no ADI overrun.

Deleted: population

17 Nevertheless, this study shows the importance in terms of risk assessment to take account of
18 specific consumption habits among the population. Vegetarians actually represent 2% of the
19 French population (Leblanc *et al.*, 2000). Considering only the general population would
20 potentially put vegetarians in situation of risk. This is all the more important that pesticide
21 uses are sometimes limited on some specific food crops with high MRLs, such as cereals
22 (Delthamethrin 3 mg. kg⁻¹) or fruit (Imazalil 2 mg. kg⁻¹ for citrus crop). Although the general
23 population is protected from these two pesticides, the everyday high cereal consumers such as
24 in the PLV group are theoretically overexposed to Delthametrin as well as the everyday high
25 fruit consumers such as in the VG group for Imazalil. Considering that some ADI overruns
26 would not be identified with a prediction model based only on the general population, it
27

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

would be pertinent to develop new pesticide dietary intake model including vegetarian diets.

In that way, European Food Safety Authority has recently built a model for chronic and acute risk assessment at the European level, which includes a vegetarian population (from UK) (EFSA, 2007).

These conclusions need to be further investigated. In fact, it should be underlined that some foodstuffs, such as fish, are not taken into account in the risk assessment. This study aims to characterise this bias on the Aldrin case. The PLV group appears first to be more protected from Aldrin exposure than the other vegetarian subgroups. This is not any more the case when a theoretical fish contribution is taken into account. Dietary intakes of organochlorine compound and other lipophilic substances may have been consequently underestimated.

Secondly, the number of respondents standing between 11 and 43 does not allow robust comparisons between the vegetarian subgroups. Conclusions would need to be confirmed on more individuals.

Moreover, this analysis is based on the hypothesis that the food contamination level consumed by the general population and the vegetarian one are on average the same. However, some studies show that individuals with vegetarian habits would preferably buy organic or non-sprayed products (Leitzmann, 2003, Santucci, 2002, Schryver *et al.*, 2007). These foods are supposed to be less contaminated than conventional products, even if no study has really proven it yet. Consequently, the higher consumption level of products potentially representing a risk could be offset by lower contamination levels. Further investigations based on the food supply habits of the population and the food contamination levels according to farming practices could be conducted in order to better characterise the sensitive population to be considered during pesticides residues risk assessment.

Finally, even if this study tends to show a higher risk of pesticides exposure for vegetarians, the vegetarian diet is generally associated with some beneficial effects. This diet would imply

Deleted: FSA

lower risk of ischemic diseases through its low rate in saturated fatty acids (Rajaram, 2003). A lower mortality rate for the vegetarian population has been also shown when compared to the general population (Key *et al.*, 1999). Thus, further benefit-risk analysis of fruit and vegetables consumption aimed to balance the risk of pesticides exposure and nutritional beneficial effects could be conducted to handle the present issue in term of public health.

Conclusion

The goal of the present study was to understand how consumption habits, especially the vegetarian ones, can influence pesticide dietary intakes. The analysis of pesticide residue exposure was carried out via a maximalist approach, using MRL values as contamination levels of foodstuffs. Vegetarians were shown to be globally more exposed than the general population even if differences exist from one population subgroup to another and depending on the pesticide considered. The general population is exposed to higher levels than the vegetarian one for pesticides with animal products as main contributors (mainly organochlorines). On the contrary, pesticide dietary intakes coming from fruit, vegetables and cereals are higher in the vegetarian population. Finally, this analysis demonstrates that among the vegetarians included in the study, dietary intakes from plant origin foodstuffs (eaten in

great quantities to comfort the absence of animal commodities such as meat, eggs and dairy products) largely offset the limited dietary intake associated to a negligible consumption of origin-animal commodities. The PLV diet is the only one that seems to be advantageous in terms of number of pesticides exceeding the ADI but the lack of MRL for fish certainly distorts the evaluation of pesticide intakes, particularly for residues that are persistent in the environment. Such results should however be considered cautiously because the approach using TMDI values is a very conservative one. Afterwards, it should be interesting to adopt a similar procedure using real food contamination levels instead of MRLs that represent

Deleted: study

Deleted: tends to prove

Deleted: that for vegetarians

Deleted: vegetal

Deleted: dietary

1 theoretical estimations. Even if this study tends to show a higher risk for vegetarians
2 concerning some pesticides, such a diet (even a lifestyle) could also presents great advantages
3 for health. Thus it would be important to develop methodology in order to consider
4 simultaneously benefits and risks associated with this particular diet.
5
6
7
8
9

10
11
12 **References**
13

- 14 Black AE. 2000. The sensitivity and specificity of the Goldberg cut-off for EI:BMR for
15 identifying diet reports of poor validity. *Eur J Clin Nutr.* 54(5): 395-404.
16
17 Chun OK, Kang HG. 2003. Estimation of risks of pesticide exposure, by food intake, to
18 Koreans. *Food Chem Toxicol.* 41(8): 1063-76.
19
20 EFSA. 2007. Reasoned Opinion on the potential chronic and acute risk to consumers' health
21 arising from the proposed temporary EU MRLs according to Regulation EC
22 No396/2005 on Maximum Residue Levels of Pesticides in Food and Feed of Plant and
23 Animal Origin. Parma (IT): EFSA. Available from:
24
25 http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753816_1178620776373.htm.
26
27 European Commission. 2002. Guidelines Related to Setting Maximum Residue Limits.
28 Appendix A, F, G, I and K. Brussels (BE): European Commission. Available from:
29
30 http://ec.europa.eu/food/plant/protection/resources/publications_en.htm.
31
32 European Commission. 2007. Monitoring of pesticide residues in products of plant origin in
33 the European Union, Norway, Iceland and Liechtenstein. Brussels: Available from:
34
35 Hercberg S, Deheeger M, Preziosi P. 1994. SU-VI-MAX. Portions alimentaires. Manuel
36 photos pour l'estimation des quantités. Paris (FR): Editions Poly Technica.
37
38 Herrera A, Arino A, Conchello P, Lazaro R, Bayarri S, Perez-Arquillue C, Garrido MD,
39 Jodral M, Pozo R. 1996. Estimates of mean daily intakes of persistent organochlorine
40 pesticides from Spanish fatty foodstuffs. *Bull Environ Contam Toxicol.* 56(2): 173-7.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 Hites RA, Foran JA, Carpenter DO, Hamilton MC, Knuth BA, Schwager SJ. 2004. Global
3
4 assessment of organic contaminants in farmed salmon. Science. 303(5655): 226-9.
5

6 IPCS. 1996. Persistent organic pollutants: an assessment report on DDT, aldrin, dieldrin,
7
8 endrin, chlordane, heptachlor, hexachlorobenzene, mirex, toxaphene, polychlorinated
9
10 biphenyls, dioxins and furans. Geneva: UNEP. Available from:
11
12 <http://www.chem.unep.ch/pops/ritter/en/ritteren.pdf>.
13

14 Kannan K, Tanabe S, Giesy JP, Tatsukawa R. 1997. Organochlorine pesticides and
15
16 polychlorinated biphenyls in foodstuffs from Asian and oceanic countries. Rev
17
18 Environ Contam Toxicol. 152 :1-55.
19

20 Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G, Burr ML, Chang-Claude
21
22 J, Frentzel-Beyme R, Kuzma JW, Mann J, McPherson K. 1999. Mortality in
23
24 vegetarians and nonvegetarians: detailed findings from a collaborative analysis of 5
25
26 prospective studies. Am J Clin Nutr. 70(3 Suppl): 516S-524S.
27

28 Lafay L. 2009. Rapport de l'étude Individuelle Nationale des Consommations Alimentaires 2
29
30 (INCA 2) 2006/2007. Paris (FR) : Les rapports bleus de l'AFSSA.
31

32 Leblanc JC, Yoon H, Kombadjian A, Verger P. 2000. Nutritional intakes of vegetarian
33
34 populations in France. Eur J Clin Nutr. 54(5): 443-9.
35

36 Leblanc JC. 2001. Influence de l'environnement sur le comportement alimentaire de groupes
37
38 particuliers de consommateurs : Etude de l'équilibre nutritionnel et sanitaire des
39
40 ingesta. Thèse pour l'obtention du diplôme de docteur de l'université Paris 7,
41
42 spécialité : « Physiologie et Physiopathologie de la Nutrition ». Soutenue le 1 juin
43
44 2001.

45 Leitzmann C. 2003. Nutrition ecology: the contribution of vegetarian diets. Am J Clin Nutr.
46
47 78(3 Suppl): 657S-659S.
48

1
2 Menard C, Heraud F, Nougadere A, Volatier JL, Leblanc JC. 2008. Relevance of integrating
3 agricultural practices in pesticide dietary intake indicator. Food Chem Toxicol. 46(10):
4 3240-53.
5
6

7 Rajaram S. 2003. The effect of vegetarian diet, plant foods, and phytochemicals on
8 hemostasis and thrombosis. Am J Clin Nutr. 78(3 Suppl): 552S-558S.
9

10 Rivas A, Cerrillo I, Granada A, Mariscal-Arcas M, Olea-Serrano F. 2007. Pesticide exposure
11 of two age groups of women and its relationship with their diet. Sci Total Environ.
12 382(1): 14-21.
13
14

15 Santucci F. 2002. Market issues in organic meat and dairy markets. Paper presented at:
16 Symposium on organic markets for meat and dairy products: trade opportunities for
17 developing countries. Nineteenth Session of the Intergovernmental group on meat and
18 dairy products; 27 - 29 August 2002; Rome (IT).
19
20

21 Schryver T, Smith C, Wall M. 2007. Self-identities and BMI of Minnesotan soy consumers
22 and non-consumers. Obesity (Silver Spring). 15(5): 1101-6.
23
24

25 Schutz Y, Flatt JP, Jequier E. 1989. Failure of dietary fat intake to promote fat oxidation: a
26 factor favoring the development of obesity. Am J Clin Nutr. 50(2): 307-14.
27
28

29 Volatier J-L. 2000. Enquête INCA individuelle et nationale sur les consommations
30 alimentaires. Paris (FR): Tech & Doc.
31
32

33 WHO. 1997. Guidelines for predicting dietary intake of pesticides residues (revised).
34 Prepared by the Global Environment Monitoring System - Food Contamination
35 Monitoring and Assessment Programme (GEMS/Food) in collaboration with the
36 Codex Committee on Pesticide Residues. Geneva (CH): WHO. Available from:
37
38 <http://www.who.int/foodsafety/chem/gems/en/index.html>.
39
40

41 Willet W. 1998. Nutritional epidemiology. Monographs in Epidemiology and Biostatistics
42
43 Volume 30. Oxford (UK): Oxford University Press.
44
45

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 1. General characteristics and food intakes of the study population.

	Pseudo-vegetarians (OMN)	Lacto-vegetarians (LV)	Ovo-lacto-vegetarians (OLV)	Pesco-vegetarians (PLV)	Vegans (VG)	General population
Number of subjects	26	41	43	16	11	1474
Average age (years)	56.6 (± 19.4)	42.2 (± 11.9) [*, p= $3 \cdot 10^{-3}$]	51.1 (± 15.9)	47.6 (± 14.1)	48.0 (± 13.9)	43.0 (± 18.0)
Sex ratio M/F	0.44	1.41	1.05	0.33	0.57	0.83
Average BMI (kg/m²)	21.3 (± 2.6)	21.6 (± 2.5)	21.8 (± 3.0)	19.6 (± 2.2) [*, p= $2 \cdot 10^{-2}$]	20.8 (± 2.2)	23.5 (± 3.8)
Average daily intakes (g/capita/day)	956.6	992.6	1121.2	782.6	1174.2	1241.4
<i>Fruits</i>	247.7 (± 128.4)	265.0 (± 144.9)	317.6 (± 228.3)	124.7 (± 73.3) [*; p= $2 \cdot 10^{-4}$]	297.0 (± 292.4)	173.4 (± 135.0)
<i>Vegetables</i>	311.3 (± 134.3)	286.1 (± 112.9)	381.8 (± 144.8)	339.1 (± 169.8)	459.6 (± 300.2)	170.1 (± 79.9)
<i>Cereals</i>	128.7 (± 79.5)	189.9 (± 84.8)	137.6 (± 121.2)	206.3 (± 110.4)	157.3 (± 93.5)	173.0 (± 124.4)
<i>Potatoes</i>	41.0 (± 35.2)	68.9 (± 47.3)	53.3 (± 43.2)	22.3 (± 19.2)	86.9 (± 120.4)	52.6 (± 38.1)
<i>Other products of plant origin</i>	19.3 (± 14.5)	18.8 (± 10.7)	23.4 (± 15.7)	19.7 (± 12.2) [*, p= $2 \cdot 10^{-3}$]	36.3 (± 37.5)	22.3 (± 10.2)
<i>Milk and dairy products</i>	119.4 (± 101.6)	111.5 (± 136.3)	89.3 (± 96.1)	23.6 (± 46.2)	2.9 (± 3.3)	299.6 (± 160.0) [**, p< 10^{-4}]
<i>Eggs</i>	3.1 (± 1.8)	0.9 (± 0.7)	4.0 (± 2.7)	1.0 (± 1.1)	0.1 (± 0.0)	29.0 (± 22.1) [**, p< 10^{-4}]
<i>Meat and other products of animal origin</i>	5.8 (± 9.4)	9.0 (± 13.5)	9.0 (± 12.4)	1.5 (± 1.8)	7.1 (± 12.7)	161.2 (± 72.0) [**, p< 10^{-4}]
Beverages (g/day)	634.3	783.4	1014.1	493.7	720.3	922.7
<i>Water</i>	554.0 (± 358.0)	740.8 (± 602.4)	908.9 (± 946.6)	449.3 (± 185.0)	623.0 (± 265.1)	762.4 (± 409.3)
<i>Wine</i>	80.3 (± 104.1)	42.6 (± 93.8)	105.2 (± 188.9)	44.4 (± 47.7)	97.3 (± 110.4)	160.3 (± 249.3) [**, p< 10^{-4}]

[*, p= $2 \cdot 10^{-3}$] represents a comparison inside the vegetarian population: the value given for this group is significantly different with a probability of $2 \cdot 10^{-3}$ from those of the four other groups.

[**, p= $2 \cdot 10^{-3}$] represents a comparison between the two populations: the value given for the general population is significantly different from the one of the global vegetarian population with a probability of $2 \cdot 10^{-3}$.

All probabilities are derived from non-parametric Kruskal-Wallis tests.

Deleted: 4

Deleted: 1,5

Deleted: 7

Deleted: vegetables

Deleted: D

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

Table 2. Detailed results concerning the 48 pesticides with at least one TMDI over ADI

	Active substance	Class	ADI (source, year)	MRL reference	TMDI (%ADI)					
					GP	OMN	LV	OLV	PLV	VG
Imazalil	AZOL.	0.025 (a, 1997)	EU/N	52,4	71,4	79,8	85,9	39,0	102,5	
Prochloraz	AZOL.	0.01 (a, 1988)	EU/N	107,5	139,1	162,3	212,4	95,5	207,4	
Aldicarb	CARB.	0.001 (a, 1994)	EU/N	105,9	123,2	145,1	148,3	97,6	174,5	
Carbaryl	CARB.	0.0008 (a, 2006)	EU	876,9	858,1	1158,2	984,6	679,4	844,1	
Formetanate hydrochlorid	CARB.	0.004 (a, 2005)	N	33,8	61,6	28,5	100,9	38,0	23,6	
Propineb	CARB.	0.007 (a, 2003)	EU/N	217,6	298,3	305,9	366,2	202,2	365,1	
Thiram	CARB.	0.01 (a, 2003)	EU	177,3	290,5	258,1	347,4	201,1	362,7	
Tri-allate	CARB.	0.001 (a, 2003)	EU/N	74,5	114,4	110,2	133,4	102,3	143,2	
Aldrin	OC	0.0001 (a, -)	EU/N	348,8	163,5	159,4	175,9	146,4	183,9	
Dicofol	OC	0.0025 (a, -)	EU/N	165,1	146,4	172,4	187,0	60,0	139,8	
Dieldrin	OC	0.0001 (a, -)	EU/N	348,8	163,5	159,4	175,9	146,4	183,9	
Heptachlor	OC	0.0001 (a, -)	EU/N	331,0	152,6	147,1	163,5	135,7	175,7	
Azinphos-methyl	OP	0.005 (a, 1991)	EU/N	83,6	122,5	118,8	146,9	90,8	155,3	
Chlorfenvinphos	OP	0.0005 (a, -)	EU	95,0	192,0	140,8	176,9	198,8	259,2	
Chlorpyrifos-methyl	OP	0.01 (a, 1976)	EU/N	96,0	89,7	116,7	94,4	129,4	105,7	
Diazinon	OP	0.002 (a, 1999)	EU/N	75,4	103,7	115,1	130,4	55,7	120,0	
Dichlorvos	OP	8E-05 (a, 2005)	EU/N	7355,9	6858,0	8901,7	7151,0	10729,6	8312,1	
Dimethoate	OP	0.001 (a, 2006)	EU	92,9	100,5	114,7	120,6	87,2	143,0	
Disulfoton	OP	0.0003 (a, -)	EU/N	168,8	151,0	156,3	155,0	134,1	152,6	
Ethion	OP	0.002 (a, -)	EU/N	165,2	203,0	220,3	253,0	135,2	237,3	
Malathion	OP	0.03 (a, 2005)	EU/N	102,0	111,1	129,5	123,1	141,4	130,6	
Methidathion	OP	0.001 (a, -)	EU/N	187,3	200,1	252,2	303,2	89,2	258,8	
Mevinphos	OP	0.0008 (a, -)	EU/N	134,0	233,4	211,7	279,2	182,5	303,3	
Monocrotophos	OP	0.0006 (a, -)	EU/N	87,5	109,7	100,9	121,2	56,5	131,1	
Naled	OP	0.002 (a, 2003)	EU/N	60,0	101,8	92,9	120,3	81,6	127,7	
Omethoate	OP	0.0004 (d, -)	EU/N	271,3	479,7	438,2	567,4	364,2	582,4	
Oxydemeton-methyl	OP	0.0003 (a, 2006)	EU/N	114,6	114,6	115,9	132,1	112,1	141,4	
Phorate	OP	0.0005 (a, 1999)	EU/N	131,2	144,5	146,4	162,5	134,6	175,2	
Phosalone	OP	0.01 (a, 1999)	EU/N	66,8	110,1	100,9	127,6	79,2	135,5	
Phosmet	OP	0.003 (a, 2006)	EU/N	98,9	189,9	166,6	242,4	89,2	199,6	
Phosphamidon	OP	0.0005 (a, -)	EU/N	206,5	327,8	309,7	383,5	283,7	409,9	
Pirimiphos-methyl	OP	0.004 (a, 2005)	EU/N	429,6	397,7	499,8	426,3	556,5	454,6	
Quinalphos	OP	0.0005 (c, 1992)	EU/N	70,3	112,9	107,9	133,7	93,7	147,8	
Sulfotep	OP	0.0005 (c, 1995)	N	231,1	393,5	350,3	466,0	281,9	488,9	
Thiometon	OP	0.003 (a, -)	N	97,0	154,0	145,3	189,1	123,2	194,8	
Deltamethrin	PYR.	0.01 (a, 2002)	EU	73,4	67,0	85,4	72,0	101,9	83,6	
Biphenyl	OTH.	0.05 (c, 1989)	N	138,4	134,7	172,6	234,0	56,6	157,0	
Cyhexatin	OTH.	0.0005 (a, 1987)	EU/N	268,8	276,3	263,5	296,7	202,8	299,5	
Diquat	OTH.	0.002 (a, 2001)	EU/N	54,1	53,3	62,0	65,7	109,0	75,7	
Ethoxyquin	OTH.	0.005 (a, 1998)	N	71,5	119,7	97,3	109,2	71,8	152,2	
Fenpropimorph	OTH.	0.003 (a, 1999)	EU/N	70,4	83,1	91,3	108,6	61,6	99,8	
Fentin acetate	OTH.	0.0004 (a, -)	EU/N	208,3	200,6	200,8	218,9	181,3	227,1	

1	Fentin								
2	hydroxide	OTH.	0.0004 (a, -)	EU/N	208,3	200,6	200,8	218,9	181,3
3	Hydrogen								
4	cyanide	OTH.	0.02 (c, 1993)	EU/N	198,2	167,7	232,3	169,4	291,3
5	Metaldehyde	OTH.	0.025 (a, 2001)	N	65,6	122,3	99,5	116,3	91,2
6	Ofurace	OTH.	0.0007 (d, -)	N	178,6	133,4	141,2	109,1	39,7
7	Rotenone	OTH.	0.0001 (a, 2004)	EU/N	240,7	407,8	372,4	482,0	326,9
8	Vinclozolin	OTH.	0.005 (a, 2004)	EU/N	184,2	266,4	216,9	358,9	171,0
9									
10									
11									
12									
13									
14									

The abbreviations AZOL., CARB., OC, OP, PYR., TRIAZ., UREA and OTH. refer respectively to the following classes: azoles, carbamates, organochlorines, organophosphates, pyrethroïds, urea and other pesticides.

ADI values are from Agritox database (<http://www.dive.afssa.fr/agritox/php/donnees-essentielles.php?or>), EFSA opinions (http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753816_home.htm) or European directives


(<http://eur-lex.europa.eu/JOIndex.do?ihmlang=fr>) (a), JMPR

(http://www.codexalimentarius.net/mrls/pestdes/pest_ref/plst-f.htm) (b), Iris

(<http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showSubstanceList>) (c) and Australie

(<http://www.tga.gov.au/docs/pdf/adi.pdf>) (d). No year is given for the ADI value if the pesticide is now forbidden. MRL values in force are national values (N) or European ones (EU). Pesticides potentially at risk for one group are highlighted in grey; TDMI values above ADI are in bold type.

Figure 1. Determining factors of the Theoretical Maximum Daily Intakes of carbamates.



Estimations of TMDIs can be obtained from the average consumption of vegetal-origin products. (a) For Tri-allate, Thiram and Propineb, TMDI is a function of the average consumption of vegetables, fruit and wine in each vegetarian subgroup. (b) For Aldicarb, TMDI is a function of the average consumption of potatoes, vegetables and fruit in each vegetarian subgroup. (c) Contributions of vegetal-origin products to the Carbaryl dietary intake.

Deleted: ¶

Figure 2. Contribution of foodstuffs to Aldrin (organochlorine), Chlорpyrifos-methyl and Diazinon (organophosphates) dietary intakes.





Figure 3. Impact of fish or fish-derived products consumption on the Adrin dietary intake.



Supplemental Table: Detailed results concerning the other pesticides studied

Active substance	Class	ADI (source, year)	MRL reference	GP	OMN	TMDI (%ADI)			
						LV	OLV	PLV	VG
Amitrole (Aminotriazole)	AZOL.	0.001 (a, 2001)	EU	10,0	14,1	14,6	16,4	14,2	18,7
Azaconazole	AZOL.	0.025 (d, -)	N	0,0	0,0	0,0	0,0	0,0	0,0
Benomyl	AZOL.	0.1 (a, -)	EU/N	2,3	2,3	2,5	2,6	2,0	2,5
Bitertanol	AZOL.	0.01 (a, 2004)	EU/N	47,0	76,0	86,5	89,0	40,6	88,1
Bromuconazole	AZOL.	0.025 (a, 1999)	N	0,7	0,8	0,9	0,9	0,7	1,0
Carfentrazone-ethyl	AZOL.	0.03 (a, 2003)	EU	0,7	0,8	0,9	0,8	1,0	1,0
Chlorfenapyr	AZOL.	0.02 (d, -)	EU	2,5	3,5	3,6	4,0	3,5	4,5
Cyazofamid	AZOL.	0.17 (a, 2003)	EU	0,4	0,3	0,4	0,4	0,2	0,3
Cyproconazole	AZOL.	0.01 (a, 1989)	N	3,2	3,5	3,3	3,2	2,5	3,6
Difenconazole	AZOL.	0.01 (a, 1990)	N	9,8	13,1	13,1	13,1	7,3	12,8
Diniconazole	AZOL.	0.005 (a, 1987)	N	6,6	5,3	5,8	5,0	5,0	5,3
Epoxiconazole	AZOL.	0.005 (a, 1992)	N	2,3	1,8	2,1	1,7	1,7	1,6
Fenbuconazole	AZOL.	0.036 (a, 1999)	N	1,8	2,0	2,5	2,1	1,0	1,6
Fipronil	AZOL.	0.0002 (a, 2003)	N	16,5	16,5	21,3	18,2	21,8	20,4
Fluquinconazole	AZOL.	0.003 (a, 1995)	N	10,9	11,8	11,6	10,0	8,7	11,8
Flusilazole	AZOL.	0.002 (a, 2004)	N	27,8	33,1	31,2	33,0	25,5	39,8
Flutriafol	AZOL.	0.01 (a, 1987)	N	2,8	2,5	3,3	2,7	4,5	3,2
Hexaconazole	AZOL.	0.005 (a, 1989)	EU	11,8	14,7	14,7	16,0	11,2	16,1
Metconazole	AZOL.	0.01 (a, 2006)	N	0,6	0,6	0,6	0,7	0,6	0,4
Myclobutanil	AZOL.	0.02 (a, 1989)	EU/N	23,3	29,2	32,2	40,4	12,7	37,4
Paclobutrazol	AZOL.	0.1 (a, 1998)	EU/N	1,3	2,3	2,2	2,8	1,2	3,0
Penconazole	AZOL.	0.03 (a, 1992)	EU/N	3,0	3,8	3,4	3,9	3,0	4,5
Propiconazole	AZOL.	0.04 (a, 2003)	EU	1,5	2,0	2,0	2,3	2,0	2,5
Tebuconazol	AZOL.	0.03 (a, 1989)	N	2,8	3,1	2,7	2,9	2,1	3,0
Tebufenpyrad	AZOL.	0.01 (a, 1998)	N	7,6	11,1	9,0	10,5	6,5	13,7
Tetraconazole	AZOL.	0.008 (a, 2006)	N	7,0	9,4	7,5	8,6	5,5	11,2
Triadimefon	AZOL.	0.03 (a, 1985)	EU/N	12,9	13,1	12,2	13,4	9,1	13,0
Triadimenol	AZOL.	0.05 (a, 1989)	EU/N	7,7	7,9	7,3	8,0	5,5	7,8
Triazamate	AZOL.	0.002 (a, 1993)	N	1,7	2,1	2,4	2,3	1,6	2,9
Triticonazole	AZOL.	0.025 (a, 2005)	N	0,2	0,2	0,3	0,2	0,3	0,2
Asulam	CARB.	0.05 (c, 1995)	N	0,0	0,1	0,0	0,1	0,1	0,0
Benfuracarb	CARB.	0.015 (a, 2002)	EU/N	8,5	5,3	5,2	5,7	4,6	5,8
Carbendazim	CARB.	0.02 (a, 2005)	EU	9,9	11,4	12,7	13,1	10,9	13,2
Carbofuran	CARB.	0.002 (a, 1999)	EU/N	54,5	36,6	38,8	42,1	22,2	32,9
Carbosulfan	CARB.	0.01 (a, 1986)	EU/N	12,9	8,5	8,2	8,9	7,3	9,4
Chlorpropham	CARB.	0.05 (a, 2003)	EU/N	10,9	9,2	13,0	10,7	5,0	15,0
Desmedipharm	CARB.	0.03 (a, 2004)	EU	1,7	2,3	2,4	2,7	2,4	3,1
Diethofencarb	CARB.	0.1 (a, 1991)	N	0,9	1,1	1,4	1,7	0,6	1,1
Ethiofencarb	CARB.	0.1 (a, -)	N	3,8	7,4	5,5	7,8	6,2	9,1
Fenoxy carb	CARB.	0.05 (a, 1999)	N	1,5	2,3	2,0	2,2	1,4	2,7
Furathiocarb	CARB.	0.03 (d, -)	EU/N	4,3	2,7	2,7	2,9	2,4	3,0
Iprovalicarb	CARB.	0.015 (a, 2002)	EU	18,1	19,4	19,6	22,7	12,7	21,9
Mancozeb	CARB.	0.05 (a, 1993)	EU/N	30,5	41,8	42,8	51,3	28,3	51,1
Maneb	CARB.	0.05 (a, 2005)	EU/N	30,5	41,8	42,8	51,3	28,3	51,1
Methiocarb	CARB.	0.013 (a, 2006)	N	3,1	5,0	4,5	5,9	4,7	5,7
Methomyl	CARB.	0.0025 (a, 2006)	EU/N	76,8	81,7	81,9	99,9	56,5	93,5
Metiram	CARB.	0.03 (a, 1993)	EU/N	50,8	69,6	71,4	85,5	47,2	85,2

3	Molinate	CARB.	0.008 (a, 2003)	EU	6,1	8,5	8,7	9,9	8,3	11,2
4	Oxamyl	CARB.	0.001 (a, 2005)	EU	10,7	14,7	15,9	17,4	14,3	18,5
5	Phenmedipham	CARB.	0.03 (a, 2004)	EU	3,0	3,3	3,0	3,4	2,7	3,5
6	Pirimicarbe	CARB.	0.035 (a, 2003)	EU/N	8,9	14,8	13,6	17,5	11,9	18,5
7	Propamocarbe	CARB.	0.29 (a, 2006)	N	0,8	1,4	1,3	2,4	1,3	2,4
8	chlorhydrate									
9	Propham	CARB.	0.02 (c, 1987)	EU/N	4,1	3,9	3,9	4,3	3,4	4,4
10	Propoxur	CARB.	0.02 (a, -)	EU/N	5,2	5,7	5,4	6,2	5,8	6,3
11	Prosulfocarbe	CARB.	0.005 (a, 2005)	N	1,2	1,0	1,3	1,0	0,9	1,2
12	Thiabendazole	CARB.	0.1 (a, 2001)	EU/N	22,5	30,3	33,7	37,2	16,1	49,7
13	Thiodicarb	CARB.	0.01 (a, 2005)	EU	19,2	20,4	20,5	25,0	14,1	23,4
14	Thiophanate-	CARB.	0.08 (a, 1998)	EU	6,1	6,4	6,6	7,1	4,0	6,2
15	methyl									
16	Zineb	CARB.	0.03 (a, -)	EU/N	1,4	69,6	71,4	85,5	47,2	85,2
17	No ADI available	CARB.								
18										
19										
20										
21	Chlordane	OC	0.0005 (a, -)	EU/N	38,5	33,8	35,7	37,5	36,5	41,9
22	Chlorothalonil	OC	0.015 (a, 2003)	EU/N	52,4	74,0	61,5	77,1	55,2	71,7
23	DDT	OC	0.01 (a, -)	EU/N	17,5	8,0	7,7	8,4	6,8	8,8
24	Endosulfan	OC	0.006 (a, 1998)	EU/N	27,1	28,6	32,8	36,1	17,3	31,7
25	Endrin	OC	0.0002 (a, -)	EU/N	77,7	70,0	70,8	79,5	67,0	87,9
26	Lindane	OC	0.005 (a, -)	EU/N	4,2	3,0	2,9	3,3	2,7	3,5
27	Methoxychlor	OC	0.1 (a, -)	EU/N	0,1	0,1	0,1	0,2	0,1	0,2
28	No ADI available	OC								
29										
30										
31										
32	Acephate	OP	0.03 (a, -)	EU	1,1	1,1	1,1	1,2	1,0	1,2
33	Azinphos-ethyl	OP	0.002 (d, -)	EU	17,1	18,1	19,6	19,9	19,4	21,9
34	Bromophos	OP	0.04 (a, -)	N	15,0	25,4	23,2	30,1	20,4	31,9
35	Bromophos-ethyl	OP	0.003 (a, -)	EU/N	11,7	18,8	18,0	22,3	15,6	24,6
36	Chlorpyrifos-ethyl	OP	0.01 (a, 2005)	EU	25,7	39,7	44,5	57,8	21,1	52,9
37	Dioxathion	OP	0.0015 (a, -)	EU/N	23,4	37,6	36,0	44,6	31,2	49,3
38	Ethephon	OP	0.03 (a, 2005)	EU/N	9,9	11,7	13,4	12,2	7,2	12,0
39	Ethoprophos	OP	0.0004 (a, 1999)	N	9,6	14,2	14,2	17,5	11,5	18,8
40	Etrimesfos	OP	0.003 (a, -)	N	4,5	3,9	3,5	3,2	2,5	3,9
41	Fenamiphos	OP	0.0008 (a, 1999)	EU/N	34,7	44,5	47,3	51,7	41,0	54,4
42	Fenitrothion	OP	0.005 (a, 2006)	EU	8,3	6,3	4,3	7,6	4,2	7,6
43	Fenthion	OP	0.007 (a, 1997)	EU/N	35,9	39,7	49,8	58,8	14,1	47,1
44	Fonofos	OP	0.002 (c, 1991)	N	5,9	28,8	16,8	22,4	24,9	31,3
45	Fosthiazate	OP	0.004 (a, 2003)	EU	5,2	7,3	7,6	8,7	7,4	10,0
46	Isofenphos	OP	0.001 (a, -)	N	8,6	13,6	13,7	17,1	9,6	17,7
47	Mecarbam	OP	0.002 (a, -)	EU/N	24,2	33,8	34,7	39,1	33,1	43,7
48	Methacrifos	OP	0.006 (a, -)	EU/N	9,2	11,6	11,9	13,3	11,1	14,6
49	Methamidophos	OP	0.001 (a, 2005)	EU/N	30,4	35,6	29,2	30,3	32,4	28,5
50	Parathion	OP	0.004 (a, -)	EU/N	20,5	19,6	19,6	21,3	17,1	21,9
51	Parathion-methyl	OP	0.003 (a, -)	EU/N	46,4	70,7	64,6	82,9	54,4	87,2
52	Phoxim	OP	0.004 (a, -)	EU/N	6,6	5,6	7,8	6,1	9,7	6,7
53	Pirimiphos ethyl	OP	0.0002 (d, -)	N	30,6	51,4	47,0	60,8	41,2	64,3
54	Profenofos	OP	0.01 (a, -)	EU/N	6,0	7,0	7,1	8,0	6,7	8,7
55	Pyrazophos	OP	0.004 (a, -)	EU/N	15,9	18,1	18,3	20,4	16,8	21,9
56	Terbufos	OP	0.0002 (a, 1999)	N	9,2	19,1	18,0	22,4	22,9	27,3

1	Tetrachlor-								
2	vinphos	OP	0.05 (d, -)	N	15,5	24,6	22,7	30,0	14,5
3	Triazophos	OP	0.001 (a, -)	EU	19,2	18,1	19,0	19,6	17,7
4	Trichlorfon	OP	0.01 (a, 1978)	EU	31,0	49,6	45,4	59,1	37,0
5	Vamidothion	OP	0.008 (a, -)	EU/N	8,9	16,7	14,3	17,5	11,8
6	No ADI available		OP	Chlormephos, Ditalimfos, Fenchlorphos, Formothion, Heptenophos, Pyridiphenthion, TEPP					
7	Acrinathrin	PYR.	0.02 (a, 1990)	N	1,4	1,9	2,3	2,1	0,7
8	Alphamethrin	PYR.	0.015 (a, 2004)	EU/N	28,2	48,1	43,7	55,4	29,9
9	Beta-Cyfluthrin	PYR.	0.003 (a, 2002)	EU/N	26,9	30,2	27,6	38,0	24,5
10	Bifenthrin	PYR.	0.015 (a, 2005)	EU/N	18,0	24,0	22,8	30,4	21,2
11	Bioresmethrin	PYR.	0.03 (a, -)	N	0,6	0,9	1,0	0,9	0,3
12	Cyfluthrin	PYR.	0.003 (a, 2002)	EU/N	31,6	40,5	36,3	48,3	31,0
13	Cypermethrin	PYR.	0.05 (a, 1994)	EU/N	9,9	15,1	13,7	17,4	9,4
14	Esfenvalerate	PYR.	0.02 (a, 2000)	EU/N	1,6	2,0	2,1	2,3	2,0
15	Fenpropathrin	PYR.	0.03 (a, -)	N	5,8	8,7	7,1	8,4	4,9
16	Fenvalerate	PYR.	0.02 (a, -)	N	3,8	0,6	0,5	0,4	0,1
17	Flucythrinate	PYR.	0.02 (a, -)	EU/N	4,1	3,9	3,9	4,3	3,4
18	Fluvalinate	PYR.	0.01 (c, 1991)		0,0	0,0	0,0	0,0	0,0
19	Lambda-								
20	Cyhalothrin	PYR.	0.005 (a, 2000)	EU/N	33,4	33,7	30,2	43,0	26,5
21	Permethrin	PYR.	0.05 (a, -)	EU/N	2,6	1,6	1,6	1,7	1,4
22	Resmethrin	PYR.	0.03 (c, 1988)	EU/N	5,1	5,0	4,9	5,5	4,2
23	Tau-Fluvalinate	PYR.	0.01 (a, 1984)	N	9,7	14,2	11,8	17,5	11,1
24	Tefluthrin	PYR.	0.013 (a, 2002)	N	0,4	0,3	0,3	0,3	0,3
25	No ADI available		PYR.	Tralomethrin					
26	Ametryne	TRIAZ.	0.009 (c, 1989)	N	0,8	1,8	2,4	2,6	1,0
27	Anilazine	TRIAZ.	0.1 (a, -)	N	2,2	1,8	2,0	1,7	1,7
28	Atrazine	TRIAZ.	0.035 (c, 1993)	EU	1,0	1,6	1,6	1,9	1,4
29	Cyromazine	TRIAZ.	0.17 (a, 1984)	EU/N	3,0	6,8	5,7	10,2	5,9
30	Prometryne	TRIAZ.	0.004 (c, 1992)	N	16,5	30,6	25,0	28,6	23,0
31	Pymetrozine	TRIAZ.	0.03 (a, 2001)	EU/N	5,3	8,3	8,4	11,5	6,1
32	Simazine	TRIAZ.	0.005 (a, 2003)	N	5,6	9,8	5,0	8,3	3,2
33	Terbutylazine	TRIAZ.	0.003 (d, -)	N	6,4	8,1	6,7	6,9	5,1
34	Terbutryn	TRIAZ.	0.001 (c, 1988)	N	0,5	0,8	0,9	1,2	0,7
35	No ADI available		TRIAZ.	Cyanazine, Desmetyne, Terbumeton					
36	Amidosulfuron	UREA	0.2 (a, 1995)	N	0,1	0,1	0,1	0,1	0,1
37	Azimsulfuron	UREA	0.1 (a, 1999)	EU	0,2	0,3	0,3	0,3	0,4
38	Bensulfuron-methyl	UREA	0.2 (a, 1990)	N	0,0	0,0	0,0	0,0	0,1
39	Chloroxuron	UREA	0.05 (c, 1990)	EU/N	1,0	1,4	1,4	1,6	1,3
40	Chlortoluron	UREA	0.04 (a, 2005)	N	0,3	0,2	0,3	0,2	0,2
41	Cinosulfuron	UREA	0.09 (a, 1993)	N	0,0	0,0	0,0	0,0	0,1
42	Diflubenzuron	UREA	0.02 (a, 1986)	N	5,5	12,3	11,7	13,2	6,9
43	Diuron	UREA	0.007 (a, 2005)	N	1,7	2,8	2,3	2,3	1,7
44	Ethoxysulfuron	UREA	0.04 (a, 2003)	EU/N	1,2	1,7	1,7	2,0	1,7
45	Flazasulfuron	UREA	0.013 (a, 2004)	N	0,1	0,1	0,1	0,1	0,1
46	Flufenoxuron	UREA	0.01 (a, 1999)	N	9,2	11,9	9,8	10,0	7,0
47									

1	Flupyrusulfuron-									
2	methyl	UREA	0.035 (a, 2001)	EU	0,6	0,8	0,8	0,9	0,8	1,1
3	Foramsulfuron	UREA	0,5 (a, 2003)	EU/N	0,0	0,0	0,0	0,0	0,0	0,0
4	Hexaflumuron	UREA	0,005 (a, 1991)	N	12,2	20,3	16,7	18,6	12,1	25,9
5	Iodosulfuron-									
6	methyl-sodium	UREA	0,03 (a, 2003)	EU	0,6	0,9	0,9	1,0	0,8	1,2
7	Isoproturon	UREA	0,015 (a, 2002)	EU/N	5,5	5,3	5,4	5,8	4,8	6,1
8	Linuron	UREA	0,003 (a, 2003)	EU/N	18,3	29,9	27,8	31,4	28,5	40,2
9	Lufenuron	UREA	0,01 (a, 1993)	N	14,2	15,7	12,3	14,5	8,5	17,7
10	Methabenz-									
11	thiazuron	UREA	0,004 (d, -)	N	5,6	4,5	5,1	4,2	4,3	3,9
12	Metsulfuron-									
13	methyl	UREA	0,22 (a, 2000)	EU	0,2	0,3	0,3	0,4	0,3	0,4
14	Monolinuron	UREA	0,003 (a, -)	EU/N	27,3	26,1	26,2	28,5	22,8	29,2
15	Nicosulfuron	UREA	0,4 (a, 1994)	N	0,0	0,0	0,0	0,0	0,0	0,0
16	Oxasulfuron	UREA	0,013 (a, 2003)	EU/N	4,1	5,2	5,3	6,0	5,1	6,7
17	Pencycuron	UREA	0,025 (a, 1995)	N	1,8	4,8	4,4	7,7	5,1	12,2
18	Prosulfuron	UREA	0,02 (a, 2002)	EU	1,1	1,5	1,6	1,8	1,8	2,1
19	Rimsulfuron	UREA	0,1 (a, 2005)	N	0,0	0,0	0,0	0,0	0,0	0,0
20	Sulfosulfuron	UREA	0,24 (a, 2002)	EU/N	0,3	0,3	0,3	0,4	0,3	0,4
21	Teflubenzuron	UREA	0,035 (a, 1995)	N	1,7	2,9	2,3	2,6	1,7	3,6
22	Thifensulfuron-									
23	methyl	UREA	0,01 (a, 2001)	EU	4,8	6,8	6,9	7,8	6,6	8,7
24	Triasulfuron	UREA	0,01 (a, 2000)	EU	4,8	6,8	6,9	7,8	6,6	8,7
25	Triflumuron	UREA	0,005 (a, 1985)	N	23,8	39,9	32,5	36,4	23,9	50,7
26	Triflusulfuron-									
27	methyl	UREA	0,04 (a, 1996)	N	0,0	0,0	0,0	0,0	0,0	0,0

34	No ADI available	UREA	Dimefuron, Neburon							
35	1,3dichloro-propene	OTH.	0,0125 (a, 2005)	N	0,5	1,5	0,9	1,4	1,6	2,0
36	2,4,5-T	OTH.	0,03 (a, -)	EU/N	1,2	1,9	1,8	2,2	1,6	2,5
37	2,4-D	OTH.	0,05 (a, 2001)	EU/N	2,6	2,9	3,4	4,0	1,8	3,5
38	2,4-DB	OTH.	0,02 (a, 2003)	EU/N	3,0	3,5	3,6	4,0	3,3	4,4
39	Abamectine	OTH.	0,002 (a, 1997)	EU/N	8,8	11,9	11,0	15,1	10,3	15,1
40	Acetochlor	OTH.	0,02 (a, 1996)	N	0,0	0,0	0,0	0,0	0,0	0,0
41	Acibenzolar-S-methyl	OTH.	0,1 (a, 2001)	EU/N	0,9	1,0	1,4	1,2	0,6	0,8
42	Acifluorfen, sodium salt	OTH.	0,013 (c, 1988)	N	0,0	0,1	0,0	0,1	0,1	0,1
43	Aclonifen	OTH.	0,02 (a, 1994)	N	0,2	0,2	0,2	0,2	0,3	0,3
44	Alachlor	OTH.	0,0005 (a, 1984)	N	1,3	1,8	1,7	1,9	4,2	3,4
45	Amitraz	OTH.	0,01 (a, 1998)	EU	4,9	6,8	7,0	7,9	6,7	8,9
46	Azocyclotin	OTH.	0,01 (a, 1989)	N	12,2	12,4	12,3	14,0	9,5	14,6
47	Azoxystrobin	OTH.	0,1 (a, 1998)	EU	8,0	10,5	13,8	14,0	15,4	15,2
48	Benalaxyl	OTH.	0,04 (a, 2004)	EU/N	3,8	2,7	2,8	3,0	2,1	2,9
49	Benoxacor	OTH.	0,004 (a, 1996)	N	0,1	0,0	0,1	0,1	0,1	0,1
50	Bentazone	OTH.	0,1 (a, 2000)	EU/N	1,1	1,4	1,4	1,6	1,3	1,7
51	Bifenox	OTH.	0,3 (a, 2000)	N	0,0	0,0	0,0	0,0	0,0	0,0
52	Binapacryl	OTH.	0,002 (d, -)	EU/N	17,6	28,2	27,0	33,4	23,4	36,9
53	Bromopropylate	OTH.	0,03 (a, -)	EU/N	2,8	2,7	2,7	2,9	2,4	3,0
54	Bromoxynil phenol	OTH.	0,01 (a, 2004)	N	2,7	1,5	1,9	1,4	2,1	1,5
55	Bupirimate	OTH.	0,05 (a, 1982)	N	1,9	2,6	2,1	2,5	1,4	2,6

1										
2										
3	Buprofezin	OTH.	0.01 (a, 1990)	N	1,2	1,6	2,9	2,3	0,7	1,6
4	Butralin	OTH.	0.003 (a, 2005)	N	0,8	0,6	0,7	0,5	0,2	0,5
5	Captan	OTH.	0.1 (a, 2006)	EU/N	13,6	15,7	13,0	16,4	8,5	15,2
6	Captofol	OTH.	0.002 (c, 1987)	EU/N	13,7	16,9	18,6	19,1	19,1	21,5
7	Carbetamide	OTH.	0.021 (a, 2005)	N	0,1	0,2	0,1	0,1	0,0	0,0
8	Carbon disulfide	OTH.	0.1 (c, 1990)	EU	0,3	0,2	0,3	0,2	0,4	0,3
9	Carbon tetrachloride	OTH.	0.0007 (c, 1991)	EU	37,9	32,1	44,4	32,5	55,6	38,5
10	Chinomethionat (oxythioquinox)	OTH.	0.006 (a, -)	EU	28,3	47,3	42,4	56,8	33,2	59,7
11	Chlorbenside	OTH.	0.01 (b, 1965)	EU/N	4,3	2,4	2,3	2,3	1,5	1,8
12	Chlorfenson	OTH.	0.01 (a, -)	EU/N	1,0	1,3	1,4	1,5	1,3	1,7
13	Chlormequat	OTH.	0.05 (a, 1997)	EU/N	11,7	11,2	11,0	12,0	9,6	9,9
14	Chlorobenzilate	OTH.	0.02 (a, -)	EU/N	1,0	1,4	1,4	1,6	1,3	1,7
15	Chlozolinate	OTH.	0.1 (a, -)	EU	0,5	0,7	0,7	0,8	0,7	0,9
16	Cinidon-ethyl	OTH.	0.01 (a, 2002)	EU	6,3	8,1	8,7	9,2	9,0	10,4
17	Clethodim	OTH.	0.01 (a, 1995)	N	1,6	2,5	3,2	3,0	3,6	4,6
18	Clodinafop-propargyl	OTH.	0.003 (a, 2000)	N	3,8	2,9	3,2	2,6	2,5	2,4
19	Clofentezine	OTH.	0.02 (a, 1986)	EU/N	12,1	14,3	11,6	16,3	6,9	13,4
20	Clomazone	OTH.	0.043 (a, 2001)	N	0,0	0,0	0,0	0,0	0,0	0,0
21	Clopyralid	OTH.	0.15 (a, 2005)	N	1,5	1,3	1,4	1,2	1,2	1,1
22	Cloquintocet-mexyl	OTH.	0.04 (a, 1993)	N	0,3	0,2	0,2	0,2	0,2	0,2
23	Copper	OTH.	0.5 (a, -)	N	3,9	3,3	2,9	2,4	1,2	2,2
24	Cyclanilide	OTH.	0.0075 (a, 2001)	EU/N	7,3	9,3	9,5	10,6	8,9	11,6
25	Cycloxydim	OTH.	0.07 (a, 2004)	N	2,8	4,2	4,1	3,8	3,1	5,0
26	Cyhalofop-butyl	OTH.	0.003 (a, 2002)	EU	6,7	9,4	9,6	10,9	9,7	12,3
27	Cymoxanil	OTH.	0.016 (a, 1999)	N	1,2	1,2	1,4	1,2	0,5	1,2
28	Cyprodinyl	OTH.	0.03 (a, 2003)	N	15,5	27,2	21,8	37,9	24,8	46,1
29	Daminozide	OTH.	0.45 (a, 2005)	EU/N	0,1	0,1	0,1	0,1	0,1	0,1
30	Dicamba	OTH.	0.02 (a, 1995)	N	0,0	0,0	0,0	0,0	0,0	0,0
31	Dichlobenil	OTH.	0.01 (a, 1997)	N	0,6	0,5	0,5	0,4	0,1	0,3
32	Dichlofluanid	OTH.	0.3 (a, -)	EU/N	11,9	19,3	17,0	23,0	15,0	24,0
33	Dichlorprop	OTH.	0.03 (d, -)	EU/N	1,6	2,3	2,3	2,6	2,2	2,9
34	Dichlorprop-P	OTH.	0.06 (a, 2006)	EU	0,6	0,9	0,9	1,1	0,8	1,2
35	Diclobutrazol	OTH.	0.03 (d, -)	N	2,4	1,8	1,9	1,5	0,7	1,3
36	Diclofop-methyl	OTH.	0.001 (a, 2005)	N	27,8	39,5	35,1	37,5	31,5	37,4
37	Diflufenican	OTH.	0.25 (a, 2000)	N	0,0	0,0	0,0	0,0	0,0	0,0
38	Dimethachlore	OTH.	0.11 (a, 1999)	N	0,0	0,0	0,0	0,0	0,0	0,0
39	Dimethenamid	OTH.	0.02 (a, 2005)	N	0,0	0,0	0,0	0,0	0,0	0,0
40	Dimethenamid-P	OTH.	0.02 (a, 2003)	EU	0,0	0,0	0,0	0,0	0,0	0,0
41	Dimethipin	OTH.	0.02 (a, 1999)	N	0,4	0,4	0,7	0,5	0,3	0,4
42	Dimethomorph	OTH.	0.05 (a, 2006)	N	4,9	3,6	3,8	2,9	1,0	2,6
43	Dinocap	OTH.	0.004 (a, 2004)	EU/N	9,0	15,5	14,0	18,8	9,0	19,8
44	Dinoseb	OTH.	0.003 (c, 1989)	EU/N	11,7	18,8	18,0	22,3	15,6	24,6
45	Diphenylamine	OTH.	0.08 (a, 1998)	EU	7,0	13,6	10,3	14,0	8,2	18,4
46	Dithianon	OTH.	0.01 (a, 1992)	N	10,4	21,3	17,3	18,8	12,2	25,4
47	Dodine	OTH.	0.1 (a, 2000)	EU/N	2,1	3,8	3,3	4,1	2,7	4,6
48	Ethofumesate	OTH.	0.07 (a, 2002)	EU/N	1,7	1,3	1,3	1,4	1,1	1,3

1	Ethylene dibromide	OTH.	0.009 (c, 2004)	EU	1,1	1,5	1,6	1,7	1,5	1,9
2	Etofenprox	OTH.	0.03 (a, 1995)	N	0,7	0,5	1,5	2,5	0,8	2,9
3	Famoxadone	OTH.	0.012 (a, 1995)	EU/N	20,6	15,0	14,0	15,3	8,3	13,1
4	Fenamidone	OTH.	0.03 (a, 2003)	EU/N	3,9	6,3	6,1	9,0	5,3	9,9
5	Fenarimol	OTH.	0.01 (a, 1995)	EU/N	12,6	17,2	17,8	18,6	9,2	16,7
6	Fenazaquin	OTH.	0.005 (a, 1993)	N	11,9	16,0	18,8	16,3	6,3	11,7
7	Fenbutatin oxide	OTH.	0.025 (a, 1973)	EU/N	35,9	45,3	49,6	59,8	21,2	54,7
8	Fenchlorim	OTH.	0.05 (a, 1988)	N	0,1	0,1	0,1	0,1	0,2	0,1
9	Fenhexamid	OTH.	0.2 (a, 2000)	EU/N	6,6	12,9	10,4	19,3	10,2	20,7
10	Fenoxyprop ethyl	OTH.	0.01 (a, 1988)	N	1,1	0,9	1,0	0,8	0,9	0,8
11	Fenpropidine	OTH.	0.005 (a, 1995)	N	4,5	3,6	4,1	3,3	3,5	3,1
12	Fenpyroximate	OTH.	0.01 (a, 1999)	N	2,3	3,8	3,3	3,0	2,3	4,5
13	Florasulam	OTH.	0.05 (a, 2002)	EU	0,3	0,4	0,4	0,4	0,4	0,5
14	Fluazifop-P-butyl	OTH.	0.01 (a, 2001)	N	6,3	21,6	22,0	28,4	61,1	47,1
15	Fluazinam	OTH.	0.01 (a, 1997)	N	36,5	27,1	28,2	21,8	7,9	19,2
16	Fludioxonil	OTH.	0.03 (a, 2003)	N	10,3	22,9	20,6	34,9	22,3	52,4
17	Flufenacet	OTH.	0.005 (a, 2003)	EU	10,6	14,4	15,1	16,7	13,8	19,3
18	Flumioxazine	OTH.	0.009 (a, 2002)	EU	5,5	7,7	7,9	8,9	7,8	10,1
19	Fluroxypyr	OTH.	0.8 (a, 2000)	EU	0,1	0,1	0,1	0,1	0,1	0,1
20	Flutolanil	OTH.	0.01 (a, 1992)	N	0,8	0,7	1,1	0,9	0,4	1,4
21	Folpet	OTH.	0.1 (a, 2006)	EU/N	14,0	16,2	13,6	17,0	9,9	15,8
22	Fomesafen	OTH.	0.01 (a, -)	N	0,3	0,4	0,3	0,3	0,4	0,1
23	Fosetyl-Al	OTH.	3 (a, 2005)	N	0,4	0,5	0,4	0,6	0,3	0,5
24	Glufosinate	OTH.	0.02 (a, 2005)	N	12,2	18,9	17,2	20,8	14,3	24,4
25	Glyphosate	OTH.	0.3 (a, 2001)	EU/N	5,6	5,7	6,8	7,0	10,3	7,5
26	Haloxyfop-Ethoxyethyl	OTH.	0.0007 (a, -)	N	3,8	2,9	2,7	3,2	1,8	1,8
27	Haloxyfop-R-methyl	OTH.	0.0007 (a, 1991)	N	4,6	7,6	6,4	7,8	12,5	10,0
28	Hexythiazox	OTH.	0.03 (a, 1988)	N	5,8	8,6	7,5	9,2	6,6	10,1
29	Hydrogen phosphide	OTH.	0.0003 (c, 1993)	EU	88,5	74,8	103,6	75,8	129,7	89,9
30	Imazamethabenz	OTH.	0.06 (a, 2004)	N	0,2	0,2	0,2	0,2	0,2	0,1
31	Imazamox	OTH.	9 (a, 2002)	EU/N		0,0	0,0	0,0	0,0	0,0
32	Imazaquin	OTH.	0.25 (c, 1990)	N	0,0	0,0	0,0	0,0	0,0	0,0
33	Imidacloprid	OTH.	0.05 (a, 1991)	N	1,3	1,7	1,5	1,6	1,2	1,9
34	Ioxynil	OTH.	0.005 (a, 2004)	N	5,2	2,7	3,5	2,6	4,0	2,7
35	Iprodione	OTH.	0.06 (a, 2003)	EU/N	41,1	59,0	54,8	70,2	42,3	67,8
36	Isoxaben	OTH.	0.05 (a, 1995)	N	0,1	0,2	0,1	0,2	0,1	0,2
37	Isoxaflutole	OTH.	0.02 (a, 2003)	EU	2,5	3,5	3,5	4,0	3,5	4,6
38	Kresoxim-methyl	OTH.	0.4 (a, 1999)	EU	0,5	0,6	0,5	0,6	0,3	0,5
39	Maleic hydrazide	OTH.	0.25 (a, -)	EU	17,5	17,1	24,2	20,3	12,0	31,7
40	Mecoprop	OTH.	0.01 (a, 2003)	EU	4,9	6,8	7,0	7,9	6,7	9,0
41	Mecoprop-p	OTH.	0.01 (a, 2003)	EU	4,9	6,8	7,0	7,9	6,7	9,0
42	Mepiquat	OTH.	0.58 (a, 2004)	N	0,6	0,5	0,5	0,5	0,4	0,3
43	Mesotriione	OTH.	0.01 (a, 2003)	EU	4,9	6,8	7,0	7,9	6,7	9,0
44	Metalaxyl	OTH.	0.03 (a, -)	EU/N	8,9	11,8	10,9	12,4	8,0	14,6
45	Metalaxyl-M	OTH.	0.08 (a, 2002)	EU/N	3,6	5,2	4,8	5,9	3,3	6,5

1	Metamitron	OTH.	0.025 (a, 2003)	N	0,0	0,1	0,1	0,1	0,0	0,0
2	Metazachlor	OTH.	0.036 (a, -)	N	0,0	0,0	0,1	0,1	0,1	0,1
3	Methoprene (Methrprene)	OTH.	0.4 (d, -)	N	0,0	0,0	0,1	0,1	0,0	0,0
4	Methyl bromide	OTH.	0.1 (a, 1966)	EU/N	0,6	0,8	0,9	0,9	0,9	1,0
5	Metolachlor	OTH.	0.15 (c, 1994)	N	0,1	0,1	0,1	0,1	0,0	0,1
6	Metosulam	OTH.	0.005 (a, 2006)	N	0,9	0,7	0,9	0,7	0,8	0,7
7	Metribuzine	OTH.	0.013 (a, 1999)	N	1,9	2,0	2,6	2,1	1,5	2,4
8	Napropamide	OTH.	0.125 (a, 1999)	N	0,1	0,2	0,2	0,2	0,1	0,2
9	Nitrothal- isopropyl	OTH.	0.05 (d, -)	N	2,4	4,0	3,2	3,6	2,4	5,1
10	Norflurazon Ortho- phenylphenol	OTH.	0.04 (c, 1991)	N	0,2	0,4	0,4	0,4	0,5	0,6
11	Oryzalin	OTH.	0.05 (c, 1991)	N	0,1	0,1	0,1	0,2	0,1	0,2
12	Oxadiargyl	OTH.	0.008 (a, 2003)	EU/N	1,2	1,7	1,7	2,0	1,7	2,2
13	Oxadiazon	OTH.	0.004 (a, 2004)	EU/N	4,2	6,6	6,2	8,0	4,0	8,5
14	Oxadixyl	OTH.	0.01 (d, -)	N	16,4	16,9	18,2	17,9	6,7	13,1
15	Oxyfluorfen	OTH.	0.004 (a, 1997)	N	0,7	0,6	0,6	0,5	0,2	0,3
16	Paraquat	OTH.	0.004 (a, 2003)	EU	3,6	5,7	5,5	6,8	4,7	7,6
17	Pendimethalin	OTH.	0.125 (a, 2003)	EU/N	0,7	0,8	0,8	0,8	0,7	0,9
18	Picloram	OTH.	0.2 (a, 2003)	N	0,0	0,0	0,0	0,0	0,0	0,0
19	Picolinafen	OTH.	0.014 (a, 2002)	EU	3,5	5,0	5,1	5,7	5,0	6,5
20	Picoxystrobin	OTH.	0.043 (a, 2003)	EU	1,5	1,7	1,7	1,9	1,7	2,1
21	Piperonyl butoxide	OTH.	0.2 (a, 2001)	N	13,2	11,2	15,5	11,3	19,4	13,4
22	Pretilachlor	OTH.	0.018 (a, 1988)	N	0,1	0,2	0,3	0,2	0,6	0,3
23	Procymidone	OTH.	0.025 (a, 2003)	EU/N	35,6	46,6	40,1	61,6	27,4	45,8
24	Prohexadione calcium	OTH.	0.2 (a, 2000)	EU/N	0,5	0,5	0,5	0,5	0,5	0,6
25	Propachlor	OTH.	0.013 (c, 1992)	N	0,0	0,0	0,0	0,0	0,0	0,0
26	Propaqvizafop	OTH.	0.003 (a, 1995)	N	1,6	1,8	2,5	2,5	1,5	2,7
27	Propargite	OTH.	0.02 (a, 1995)	EU/N	19,9	24,8	21,9	22,1	14,3	27,7
28	Propyzamide	OTH.	0.085 (a, 2003)	EU/N	0,6	1,0	0,9	1,5	1,0	1,8
29	Pyraclostrobin	OTH.	0.03 (a, 2004)	EU/N	11,6	14,2	12,7	18,7	10,3	19,6
30	Pyraflufen-ethyl	OTH.	0.2 (a, 2001)	EU	0,1	0,1	0,1	0,2	0,2	0,2
31	Pyrethrins	OTH.	0.04 (a, -)	EU/N	34,8	42,2	46,4	47,0	49,5	52,1
32	Pyridabene	OTH.	0.008 (a, 1990)	N	2,5	3,2	2,7	2,7	1,9	3,5
33	Pyridate	OTH.	0.036 (a, 2001)	EU/N	2,6	2,7	2,6	2,8	2,5	2,8
34	Pyrifenoxy	OTH.	0.007 (d, -)	N	2,1	3,0	2,7	2,7	1,9	3,6
35	Pyrimethanil	OTH.	0.17 (a, 2006)	N	1,7	2,2	2,0	2,6	1,4	2,6
36	Pyriproxyfen	OTH.	0.07 (a, 1993)	N	0,1	0,1	0,2	0,1	0,0	0,1
37	Quinclorac	OTH.	0.3 (d, -)	N	0,0	0,0	0,0	0,0	0,0	0,0
38	Quinmerac	OTH.	0.08 (a, 1993)	N	0,1	0,1	0,1	0,1	0,1	0,1
39	Quinoxifen	OTH.	0.2 (a, 2004)	N	0,8	0,5	0,5	0,4	0,2	0,3
40	Quintozene	OTH.	0.1 (a, -)	EU/N	0,3	0,3	0,3	0,3	0,3	0,4
41	Quizalofop- Ethyl	OTH.	0.009 (a, -)	N	2,8	4,5	4,1	4,8	3,6	5,7
42	Sethoxydim	OTH.	0.09 (c, 1989)	N	3,8	6,1	5,8	7,2	4,8	7,9
43	Silthiopham	OTH.	0.064 (a, 2003)	EU	0,8	1,1	1,1	1,2	1,0	1,4
44	Spiroxamine	OTH.	0.025 (a, 1999)	EU/N	5,7	5,0	4,6	5,4	3,8	5,5
45	Sulfosate	OTH.	0.2 (a, 2001)	N	5,9	4,8	6,0	5,0	6,2	5,0
46	Tebufenozide	OTH.	0.02 (a, 1993)	N	7,6	8,4	7,3	8,0	6,5	10,0

1	Tecnazene	OTH.	0.02 (a, -)	EU/N	4,1	3,9	3,9	4,3	3,4	4,4	
2	Tetradifon	OTH.	0.0075 (a, -)	N	58,2	74,5	62,4	70,8	42,0	81,9	
3	Tolylfluanide	OTH.	0.1 (a, 2005)	N	2,4	4,0	3,2	3,6	2,4	5,1	
4	Tralkoxydime	OTH.	0.005 (a, 1991)	N	0,9	0,7	0,8	0,7	0,6	0,6	
5	Triclopyr	OTH.	0.03 (a, 2005)	N	1,7	1,9	4,0	2,5	7,6	4,0	
6	Tridemorph	OTH.	0.01 (a, 2000)	EU/N	8,4	8,1	8,2	9,0	7,7	9,3	
7	Trifloxystrobin	OTH.	0.1 (a, 2003)	EU	5,6	5,2	4,5	5,1	2,5	4,7	
8	Trifluralin	OTH.	0.015 (a, 2005)	N	0,4	0,9	0,7	0,8	0,8	1,0	
9	Triforine	OTH.	0.02 (a, -)	EU/N	15,7	26,3	22,5	25,4	16,8	30,0	
10	Trinexapac - ethyl	OTH.	0.32 (a, 2005)	N	0,1	0,1	0,1	0,1	0,1	0,1	
11	No ADI available				Aramite, Benazolin ethyl, Bensultap, Benzoxamate, Butylhydroxyanisol (BHA), Cartap, Chlome-thoxyfen, Difenzoquat, Diflufenicanil, Dinoterb, DNOC, Fentin compounds, Flubenzimine, Flumequine, Fluorochloridone, Fluoroglycofen, Flupoxam, Hymexazol, Mefenoxam, Mepronil, Naphthalene acetic acid (NAA), Nitrofen, Nuarimol, Ortho-phenylphenol sodium salt, Phosmetine, Sulcotrione, Sulfur, Tetrasul, Thiazopyr, Trimethylsulfonium.						
12		OTH.									