



**HAL**  
open science

# The chemistry and analysis of annatto food colouring: a review

Michael Scotter

► **To cite this version:**

Michael Scotter. The chemistry and analysis of annatto food colouring: a review. *Food Additives and Contaminants*, 2009, 26 (08), pp.1123-1145. 10.1080/02652030902942873 . hal-00573911

**HAL Id: hal-00573911**

**<https://hal.science/hal-00573911>**

Submitted on 5 Mar 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**The chemistry and analysis of annatto food colouring: a review**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2009-079.R1
Manuscript Type:	Review
Date Submitted by the Author:	31-Mar-2009
Complete List of Authors:	Scotter, Michael; DEFRA Central Science Laboratory, Food Safety and Quality
Methods/Techniques:	Analysis - NMR, Chromatography - GC/MS, Chromatography - HPLC, LC/MS
Additives/Contaminants:	Colours, Process contaminants, Volatiles
Food Types:	Ingredients, Processed foods

SCHOLARONE™  
Manuscripts

## The chemistry and analysis of annatto food colouring: a review

### Abstract

Annatto food colouring (E160b) has a long history of use in the food industry for the colouring of a wide range of food commodities. The principle colouring components of annatto is the oil-soluble diapo carotenoid bixin, which is the methyl ester of the dicarboxylic acid norbixin, which is soluble in aqueous alkali. Bixin and norbixin therefore exhibit not only physicochemical properties normally associated with carotenoids but also certain anomalous properties that have an impact on the stability, food colouring applications and importantly the analysis of annatto. This review summarizes the key aspects of the structural determination of bixin (and norbixin) with special attention to *cis-trans* isomerization and how this links with its chemical structure, spectroscopic properties and stability. The oxidative, thermal and photo stability of annatto and the subsequent implications for its use in the colouring of foods, food processing, and the analysis of foods and beverages are discussed along with important mechanistic, thermodynamic and kinetic aspects. The main analytical techniques used for the chemical characterization of annatto i.e. spectrophotometry, NMR, chromatography (particularly HPLC) and mass spectrometry are reviewed in detail and other methods discussed. This links in with a review of the methods available for the detection and measurement of annatto in colour formulations and foods and beverages, which highlights the importance of the need for a good understanding and knowledge of the chemistry of bixin and norbixin in order to meet new analytical challenges.

**Keywords:** annatto, bixin, norbixin, food additives, chemistry, stability,

26 analysis, E160b

For Peer Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 27 **Overview**

28 Annatto is a natural colouring agent obtained from the outer coats of the seeds of the  
29 tropical shrub *Bixa orellana*. Annatto and its extracts are designated collectively as  
30 E160b and permitted as a food additive in the European Union and elsewhere, and  
31 have widespread use in the food industry for the colouring of many commodities  
32 including flour and sugar confectionery, dairy and savoury products, soft drinks and  
33 fish. The major colour principles of annatto are the carotenoids bixin and norbixin.  
34 Though chemically very similar, differences in their chemical properties present  
35 several challenges to the analytical chemist with respect to stereochemistry, solubility,  
36 chromatographic behaviour and stability. While current legislation on the extraction  
37 and use of annatto colours and their applications in food are addressed briefly, this  
38 review focuses on the chemistry, stability and analysis of annatto pertaining to its use  
39 as a permitted food colouring.

## 41 **Annatto in foods**

### 42 *Legislative aspects*

43 The use of food colours in the European Union is controlled by European Community  
44 Directive 94/36/EC (EC, 1994 as amended) which contains a list of permitted colours,  
45 a list of foodstuffs to which these colours may be added, and where appropriate,  
46 maximum limits on the level of addition. The permitted uses of annatto and the  
47 maximum levels of addition are given in Table 1. Annatto extracts are listed amongst  
48 those colours that may be used singly or in combination in certain foods up to the  
49 maximum levels specified (on a ready-for-consumption basis). Comprehensive on-line  
50 sources of information on permitted food colour regulations and specifications may be

1  
2  
3 51 found at the Nordic Food Additives Database (NNT, 2008) and the Food Law site of  
4  
5  
6 52 the Department of Food Biosciences, University of Reading (Jukes, 2008).  
7  
8  
9 53

10 54 [Insert Table 1 about here]  
11  
12  
13 55

14  
15 56 In July 2006, the Commission published a set of four proposed Regulations that are  
16  
17 57 intended to replace the current system and provide a common basis for controls on  
18  
19  
20 58 food additives, food flavourings and food enzymes. The proposals were published as  
21  
22 59 separate Commission Documents on additives, flavourings, enzymes and a common  
23  
24 60 authorization procedure (EC, 2006). The proposal brings together all of the existing  
25  
26  
27 61 food additive regulations and plans to introduce comitology for additive approvals in  
28  
29 62 place of the cumbersome co-decision procedure.  
30  
31  
32 63

33  
34 64 The specifications for food colours are laid down in Commission Directive 95/45/EC  
35  
36 65 (EC, 1995) in which separate definitions and purity criteria are prescribed for (i)  
37  
38 66 solvent-extracted bixin and norbixin, (ii) alkali-extracted annatto and (iii) oil-  
39  
40  
41 67 extracted annatto. Solvent-extracted bixin and norbixin formulations are often referred  
42  
43 68 to as indirectly-extracted annatto formulations, whereas alkali- and oil-extracted  
44  
45 69 annatto are termed directly-extracted. The purity specifications include definition of  
46  
47 70 the source material(s) and the solvents permitted for extraction, the identification and  
48  
49 71 the minimum content of the colouring material (measured by spectrophotometry), and  
50  
51 72 the limits for residual solvents and heavy metals.  
52  
53  
54  
55 73

74 *Use of annatto in foods*

75 Annatto was reported to be the most commonly consumed natural colour additive in  
76 the UK (MAFF, 1987 and 1993) where the *per capita* consumption was estimated to  
77 be 0.065 mg/kg bw/day based on pure colouring component, representing some  
78 12.5% of the Acceptable Daily Intake (ADI). The chemistry and applications of oil-  
79 and water-soluble annatto colours in terms of their modes of applications to a wide  
80 range of food products and the usage levels required to obtain the desired colour  
81 shades has been reviewed (Collins, 1992; Levy and Rivadeneira, 2000). Crystalline  
82 bixin products of 80-97% purity may be obtained by extraction of annatto seed with  
83 certain permitted organic solvents and subsequent production of a solvent-free  
84 product, which is then processed to give a range of high purity oil- and water-soluble  
85 annatto formulations. Oil-soluble bixin is generally used in fatty food applications,  
86 whereas norbixin, because of its ability to bind strongly with protein, is especially  
87 suited for the colouring of high protein content foods. Annatto colours are often  
88 formulated with other additives such as emulsifiers to produce forms of water-soluble  
89 annatto that are stable to the effects of e.g. acids, metal ions and salts. The  
90 applications and stability of spray-dried annatto formulations in fruit and vegetable  
91 products have been studied (Satyanarayana *et al.*, 2006).

92  
93 *Annatto intake*

94 Bixin is reported to be rapidly absorbed in the bloodstream, comparable to other  
95 dietary carotenoids, with complete plasma clearance after 8 hours and for norbixin  
96 after 24 hours (Levy, 1997). While annatto intake is an important issue within the  
97 regulatory context, intake estimates for annatto have in the past provided ambiguous  
98 results largely due to the lack of reliable data on the colour principals (bixin/norbixin)

1  
2  
3 99 content of annatto extracts (Levy and Rivadeneira, 2000). In response to a request by  
4  
5  
6 100 the Joint FAO/WHO Expert Committee on Food Additives (JECFA) for information  
7  
8 101 relating to the toxicity, intake and specifications of annatto, the European annatto  
9  
10 102 producers consulted with the food industry to determine usage levels of specific  
11  
12 103 annatto extracts (JECFA, 2002). The data obtained were combined with the levels of  
13  
14 104 bixin/norbixin in particular extracts to provide an estimate of their concentration in  
15  
16 105 food. These data have been combined with food consumption data using various  
17  
18 106 methods to estimate consumer intakes, which ranged from 1 to 163% of the ADI  
19  
20 107 (Tennant and O'Callaghan, 2005). The actual levels of annatto in foodstuffs were well  
21  
22 108 below maximum limits prescribed under EU regulations and Codex standards, which  
23  
24 109 had been confirmed by an earlier analytical study (Scotter *et al.*, 2002).  
25  
26  
27  
28  
29  
30  
31

### 32 111 **Annatto chemistry**

33  
34 112 Elucidation of the chemical constitution of bixin (and thereafter norbixin) was first  
35  
36 113 put forward by Heiduschka and Panzer (1917) who suggested the correct molecular  
37  
38 114 formula for bixin ( $C_{25}H_{30}O_4$ ) as an unsymmetrical molecule. Herzig and Faltis (1923)  
39  
40 115 recognised that bixin was the monomethyl ester of an unsaturated dicarboxylic acid.  
41  
42 116 The results from their catalytic hydrogenation studies led them to conclude that bixin  
43  
44 117 contains 9 carbon double bonds, which, evidenced by the intense red colour of the  
45  
46 118 pigment, were conjugated. However, the unsymmetrical molecule hypothesis was  
47  
48 119 abandoned when proposed the now accepted structure was proposed (Kuhn and  
49  
50 120 Winterstein, 1932), later confirmed by Karrer *et al.* (1932). A new, higher melting  
51  
52 121 form termed  $\beta$ -bixin was obtained during the course of pigment isolation (Herzig and  
53  
54 122 Faltis, 1923), which was later proposed as the *trans*-isomer that the original form may  
55  
56 123 be *cis*-isomer (Karrer *et al.*, 1929). A stable form of bixin identical to the  $\beta$ -form by  
57  
58  
59  
60



1  
2  
3 124 treatment of the 'natural' (or 'labile') form with iodine was subsequently obtained  
4  
5 125 (Kuhn and Winterstein, 1932). From the results of these investigations that it was  
6  
7  
8 126 concluded that bixin was the first known naturally occurring *cis*-polyene. The  
9  
10 127 structural elucidation of bixin was confirmed from various oxidation and degradation  
11  
12 128 experiments (Karrer and Jucker, 1950). During investigations to determine the  
13  
14 129 stereochemical configuration of labile bixin, several stereoisomers were isolated  
15  
16 130 (Zechmeister, 1960). The consequences of *cis-trans* isomerism on the chemistry,  
17  
18 131 stability and analysis of annatto are significant and are discussed below.  
19  
20  
21 132  
22  
23 133 The major colouring component of annatto is confirmed as the apo-carotenoid 9'-*cis*-  
24  
25 134 bixin (methyl hydrogen 9'-*cis*-6,6'-diapocarotene-6,6'-dioate, C<sub>25</sub>H<sub>30</sub>O<sub>4</sub>), the  
26  
27 135 monomethyl ester of the dicarboxylic acid 9'-*cis*-norbixin, commonly referred to as  
28  
29 136 *cis*-bixin (Figure 1). 9'-*Cis*-bixin is soluble in most polar organic solvents to which it  
30  
31 137 imparts an orange colour but is largely insoluble in vegetable oil. It may be readily  
32  
33 138 converted to the all-*trans* isomer due to its instability in the isolated form in solution.  
34  
35 139 *Trans*-bixin is the more stable isomer and has similar properties to the *cis*-isomer but  
36  
37 140 exhibits a red colour in solution and is soluble in vegetable oil. Commercially,  
38  
39 141 isomerisation is achieved by heating a suspension of the *cis*-isomer in oil to 130°C *in*  
40  
41 142 *vacuo*. The water-soluble analogue 9'-*cis*-norbixin (C<sub>24</sub>H<sub>28</sub>O<sub>4</sub>) can be isolated from  
42  
43 143 annatto seeds by agitation in aqueous alkali at <70° C or formed by alkaline hydrolysis  
44  
45 144 of *cis*-bixin to give either the sodium or potassium salt. The dicarboxylic acid is  
46  
47 145 soluble in polar solvents to which it imparts an orange colour. 9'-*Cis*-norbixin is only  
48  
49 146 sparingly soluble in chloroform and 0.1M sodium hydroxide (Preston and Rickard,  
50  
51 147 1980). Under extraction conditions, 9'-*cis*-bixin undergoes isomerization to produce oil  
52  
53 148 solutions containing approximately 0.2 - 0.5% of pigment comprising a mixture of all-

1  
2  
3 149 *trans*- and 9'-*cis*-bixin in variable proportions and characteristic degradation products,  
4  
5 150 dependent upon extraction temperature and time (see below).  
6  
7

8 151

9  
10 152 [Insert Figure 1 about here]  
11

12 153

13  
14  
15 154 While it is reported that 80% of the carotenoids in the annatto seed coat comprise bixin  
16  
17 155 (Preston and Rickard, 1980; Lauro, 1991), traces of bixin diesters may be found  
18  
19 156 (Mercadante *et al.*, 1997b). The preparation and use of ethylbixin has been discussed  
20  
21 157 (Levy and Rivadeneira, 2000).  
22  
23

24 158

25  
26  
27 159 The presence of other minor carotenoids in annatto has also been postulated, which  
28  
29 160 included  $\beta$ -carotene, cryptoxanthin, lutein, zeaxanthin and methyl bixin (Tirimanna,  
30  
31 161 1980). The presence of a range of lycopenoate analogues and other minor carotenoids  
32  
33 162 in annatto has been reported in a series of papers by Mercadante and co-workers (1996,  
34  
35 163 1997a, 1997b and 1999) and has been reviewed by Mercadante (2001) and  
36  
37 164 Satyanarayana, Prabhakara Rao and Rao (2003). Bixin and three minor carotenoids  
38  
39 165 have been chemically synthesized using the Wittig reaction of the (Z)-terminus  
40  
41 166 followed by a Horner-Emmons reaction (Haberli and Pfander, 1999).  
42  
43

44 167

45  
46  
47 168 *Molecular properties*  
48

49  
50 169 It is the delocalisation of  $\pi$ -electrons along the polyene backbone that gives  
51  
52 170 carotenoids their characteristic electronic spectra and is largely responsible for the  
53  
54 171 photophysical and photochemical properties of these molecules, including *cis-trans*  
55  
56 172 photoisomerization. Detailed explanations of *cis-trans* isomerization may be found in  
57  
58 173 standard texts (Karrer and Jucker, 1950; Lunde and Zechmeister, 1954; Zechmeister,  
59  
60

1  
2  
3 174 1960; Kolher, 1995). Only the basic properties of linear conjugated molecules will be  
4  
5  
6 175 reviewed here along with a brief account of the simple concepts that apply to bixin  
7  
8 176 and norbixin, in order to provide a background for discussion on the UV-VIS  
9  
10  
11 177 spectroscopy of these compounds and to show how UV-VIS spectra are affected by  
12  
13 178 isomerization.

14  
15 179  
16  
17 180 The sufficiently large barriers to rotation about the formal double bonds in polyenes  
18  
19 181 or carotenoids allow double bond *cis*- and *trans*- isomers to be isolated as  
20  
21  
22 182 independent, distinct compounds. Since the differences in excitation energies for *cis*-  
23  
24 183 and *trans*- isomers of a given molecule are small compared to the change in excitation  
25  
26  
27 184 energy that is associated with adding or subtracting a conjugated double bond, the  
28  
29 185 basic electronic structure is *almost* independent of isomeric form. The four single  
30  
31 186 bonds that surround a carbon-carbon double bond all lie in the same plane. In  
32  
33  
34 187 consequence, each of the disubstituted and trisubstituted acyclic double bonds that  
35  
36 188 constitute the polyene chain of a carotenoid can exist in two forms i.e. geometric  
37  
38 189 isomers. Nomenclature of the *cis*- or *trans*- isomers is designated in accordance with  
39  
40  
41 190 IUPAC rules (Weedon and Moss, 1995a, 1995b). In recent years however, these  
42  
43 191 designations have been replaced largely by *Z* and *E* respectively.

44  
45  
46 192  
47  
48 193 Since each double bond in the polyene chain could, in principle, exist in one of two  
49  
50  
51 194 forms, a large number of geometric isomers are theoretically possible for any  
52  
53 195 carotenoid. However, in practice few of these isomers are encountered. An  
54  
55 196 explanation for this is provided by studying molecular models, which indicate that the  
56  
57 197 introduction of a *cis*- double bond normally results in steric hindrance thus rendering  
58  
59 198 the *cis*-isomer less stable than the *trans*- form. With both trisubstituted double bonds  
60

1  
2  
3 199 and disubstituted double bonds in the 15,15'-position, the effect is relatively small, as  
4  
5 200 it results from limited interference between two hydrogen atoms and hence these  
6  
7  
8 201 isomers may be formed quite readily. With other disubstituted double bonds the  
9  
10 202 adoption of the *cis*- configuration results in major interference between a hydrogen  
11  
12 203 atom and a methyl group. This renders such molecules less stable than the  
13  
14  
15 204 corresponding *trans*- form and hence less likely to be encountered (Karrer and Jucker,  
16  
17 205 1950; Zechmeister, 1960; Kolher, 1995; Weedon and Moss, 1995a).

18  
19 206  
20  
21  
22 207 Stereomutation studies, in which interconversion of geometrical isomers is  
23  
24 208 deliberately promoted, lend support to this theory. Interconversion generally produces  
25  
26 209 a 'set' of isomers that approximates to an equilibrium mixture of all possible geometric  
27  
28 210 forms proportional to their relative thermodynamic stabilities. The all *trans*- form  
29  
30 211 usually predominates, indicating that it is the most thermodynamically stable isomer.  
31  
32 212 A number of mono- and di-*cis*-isomers are usually also present, however those  
33  
34 213 isomers with more than two *cis*- double bonds and/or those that are sterically hindered  
35  
36 214 usually occur only in trace amounts, if at all. It is not surprising that most naturally  
37  
38 215 occurring carotenoids are predominantly in the all *trans*- form. However, bixin occurs  
39  
40 216 predominantly as the *cis*-isomer, which has a *cis*- configuration about the  
41  
42 217 9'-trisubstituted double bond. Since asymmetric bixin has nine alkene bonds ( $n=9$ ),  
43  
44 218 theoretically 512 (i.e.  $2^n = 2^9$ ) geometric isomers are possible, whereas symmetric  
45  
46 219 norbixin has only 272 (i.e.  $2^{(n-1/2)} \times (2^{(n-1/2)} + 1)$ ) possible isomers. However, the  
47  
48 220 presence of stable *cis*-isomers at positions 7, 11, 12' and 8' are sterically hindered,  
49  
50 221 hence the remaining 5 alkene bonds are capable of yielding 32 and 20 isomers for  
51  
52 222 bixin and norbixin respectively (Figure 1).  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 224 Provided that an adequate sample of the pure isomer is available or the selected  
4  
5 225 analytical technique is adequately sensitive, spectroscopic analysis will normally  
6  
7  
8 226 allow the unambiguous assignment of the geometrical configuration of any carotenoid  
9  
10 227 isomer. All linear polyenes, the carotenoids included, possess similar low-lying  
11  
12 228 excited singlet ( $S_1$ ) states (Hudson and Kohler, 1974; Kohler, 1977; Hudson, Kohler  
13  
14 229 and Schulten, 1982). This is critically important since virtually all photo processes in  
15  
16 230 linear polyenes originate in the lowest-energy singlet excited state, the correct  
17  
18 231 identification and characterization of which is therefore also important. As might be  
19  
20 232 anticipated from the similarities in electronic structure, the electronic absorption  
21  
22 233 spectrum of a given carotenoid closely resembles that of the unsubstituted polyene  
23  
24 234 with the same number of conjugated double bonds. There are however well  
25  
26 235 characterised principle differences due largely to the presence of methyl substituents  
27  
28 236 along the carotenoid skeleton, which affect the basic polyene electronic structure.

29  
30  
31  
32  
33  
34 237 These are:

- 35  
36 238 • a 10 to 30 nm shift of the lowest energy strong absorption band to  
37  
38 239 longer wavelength  
39  
40  
41 240 • decreased vibrational fine structure  
42

43  
44 241  
45  
46 242 Thus, *cis*- and *trans*- isomers may often be distinguished on the basis of their UV-VIS  
47  
48 243 absorption spectra, but the most important differences observed between isomers are  
49  
50 244 not related to excitation energies but to the relative intensities of high-energy  
51  
52 245 absorption bands i.e. '*cis*-' peaks (Dale, 1954; Zechmeister, 1960). It is well  
53  
54 246 established that the lowest excited state of linear polyenes (including carotenoids) is  
55  
56 247 the  $2^1A_g$  state and that the origin of the main absorption band is the strongly allowed  
57  
58 248  $2^1A_g \rightarrow 1^1B_u$  transition. The shapes of electronic absorption (and fluorescence) bands

1  
2  
3 249 are derived from the vibrational levels that are associated with the initial and final  
4  
5 250 electronic states. Thus, the typical three-peaked shape of the main absorption band of  
6  
7  
8 251 linear polyenes arises from transitions of the lowest vibrational level of the electronic  
9  
10 252 ground state to the lowest vibrational levels of the electronic excited states.  
11  
12 253 Broadening of these peaks is observed because of rotational levels and inhomogeneity  
13  
14 254 leading to peak overlap. This is particularly relevant for many carotenoids measured  
15  
16 255 as solutions at ambient temperatures (Kohler, 1995).  
17  
18  
19  
20 256

21  
22 257 The positions of the absorption maxima and the shape or fine structure of the UV-VIS  
23  
24 258 spectrum of carotenoids are therefore characteristic. But while the UV-VIS spectrum  
25  
26 259 gives information about the chromophores of the molecule, it yields nothing about  
27  
28 260 functional groups apart from conjugated carbonyl groups that form part of the  
29  
30 261 molecule (Scott, 1964; Britton, 1995a and 1995b). In the case of carotenoids, the  
31  
32 262 relevant transition is the  $\pi \rightarrow \pi^*$  transition. For such a conjugated system, in which the  
33  
34 263  $\pi$ -electrons are highly delocalized, the excited state is of comparatively low energy.  
35  
36 264 The energy required to bring about the transition is therefore relatively small and  
37  
38 265 corresponds to light in the visible region. While the transition responsible for the main  
39  
40 266 absorption band is strongly 'allowed', transitions from the ground state to higher  
41  
42 267 electronic states are also possible, providing they obey the symmetry selection rules.  
43  
44 268 These high energy transitions give rise to absorption bands in the UV region which  
45  
46 269 are usually weak, but are observed particularly in the spectra of compounds with  
47  
48 270 extended chromophores.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

271  
272 When the symmetry properties of a carotenoid change, absorption bands that are  
273 otherwise not detected may become a significant feature of the spectrum, as the

1  
2  
3 274 transitions that produce them become allowed (Britton, 1995a). For *trans*- isomers,  
4  
5 275 the electronic structure has a centre of symmetry and the ground state is a *g* state, so  
6  
7  
8 276 transitions to a higher *g* state are forbidden. Transition to a higher excited *g* state only  
9  
10 277 becomes allowed when at least one double bond becomes *cis*- and the original  
11  
12 278 symmetry is lost. This gives rise to an absorption band in the UV region, known as the  
13  
14 279 *cis*-band or *cis*-peak. The most important feature of the absorption spectrum of a  
15  
16 280 carotenoid is the main absorption band in the visible region. Several important pieces  
17  
18  
19 281 of information can be obtained from the spectrum:  
20  
21

22 282

- 23  
24 283 • The position of the main absorption band, specified by  $\lambda_{\max}$ , provides  
25  
26 284 structural information because it is determined by the chromophore of the  
27  
28 285 molecule  
29  
30  
31 286
- 32  
33 287 • The intensity of the absorption at  $\lambda_{\max}$ , (*A*) is related both to the structure and  
34  
35 288 to the concentration of the carotenoid in the sample, and provides the basis for  
36  
37 289 quantitative analysis  
38  
39  
40 290
- 41  
42 291 • The position or the intensity of the main absorption band of a carotenoid can  
43  
44 292 be influenced by a number of factors such as a change in the molecular  
45  
46 293 environment of the carotenoid e.g. solvent  
47  
48  
49 294

50 294

51  
52  
53 295 Since the structure of the carotenoid chromophore is related to the overall shape or  
54  
55 296 fine structure of the spectrum, the shape as well as the positions of the absorption  
56  
57 297 maxima may therefore be used as a diagnostic tool, especially when comparing  
58  
59 298 carotenoid spectra (Britton, 1995a). A numerical notation describing fine structure has  
60

1  
2  
3 299 proved convenient and removes the requirement for presenting all spectra as diagrams  
4  
5 300 (Kohler, 1995). In this notation, the baseline or zero value is taken as the minimum  
6  
7 301 between the two peaks, the peak height of the longest wavelength absorption band is  
8  
9 302 designated as III, that of the middle absorption band (usually  $\lambda_{\max}$ ) as II (Figure 2).  
10  
11 303 Spectral fine structure is then expressed as the ratio of the peak heights III/II, as a  
12  
13 304 percentage.  
14  
15  
16  
17  
18  
19

20 306 [Insert Figure 2 about here]  
21

22 307  
23

24 308 The annatto carotenoids bixin and norbixin are unusual in that they contain two  
25  
26 309 carbonyl (i.e. carboxyl) groups, one at either end of the conjugated system and in  
27  
28 310 conjugation with it, which formally extends the chromophore. This results in a  
29  
30 311 spectral shift to longer wavelength, usually accompanied by loss of spectral fine  
31  
32 312 structure but the degree of effect is solvent dependent. With dicarboxylic acids and  
33  
34 313 their esters such as bixin (and norbixin), the further extension of the chromophore  
35  
36 314 causes a large relative increase in  $\lambda_{\max}$ . The spectral shift depends on the polarizability  
37  
38 315 rather than the polarity of the solvent, and the frequency shifts to lower energy, i.e. a  
39  
40 316 spectral shift to longer wavelength, as the refractive index of the solvent increases  
41  
42 317 Britton, 1995a).  
43  
44  
45  
46  
47  
48

49 318  
50

51 319 While water can have a major effect on the spectra of carotenoids in water-miscible  
52  
53 320 solvents, both bixin and norbixin isomers are relatively polar carotenoid molecules,  
54  
55 321 which are significantly soluble in solvents with a high aqueous content and as such  
56  
57 322 similar solvent-related properties are not observed. *Cis*-norbixin is in fact highly  
58  
59 323 soluble in 0.1M sodium hydroxide solution, and both compounds are fairly soluble in  
60



1  
2  
3 324 mixtures of acetonitrile and dilute aqueous acetic acid, used as a mobile phase in  
4  
5 325 HPLC analysis.  
6  
7  
8 326  
9  
10 327 The differences that are observed consistently between the spectra of *trans*- and *cis*-  
11  
12 328 isomers of carotenoids are therefore diagnostic for structural assignment. A small  
13  
14  
15 329 hypsochromic shift in  $\lambda_{\max}$  of ca. 2 to 6 nm is usually observed for mono-*cis* isomers  
16  
17 330 along with a significant hypochromic effect and a reduction in vibrational fine  
18  
19  
20 331 structure. A new absorption band appears at a characteristic position about 142 nm  
21  
22 332 below the *longest* wavelength absorption maximum, often referred to as the *cis*-peak.  
23  
24  
25 333  
26  
27 334 For the all-*trans* form of bixin, the main absorption band ( $2^1A_g \rightarrow 1^1B_u$ ) is very  
28  
29 335 intense. In the 9'-*cis* isomer), the intensity of this main band decreases as a weak *cis*-  
30  
31 336 peak appears at 355 nm, corresponding to the transition to a higher energy level *g*  
32  
33 337 state. From their studies on  $\beta$ -carotene isomers, it has been shown that the 15-*cis*  
34  
35 338 isomer, in which the *cis* double bond is in the centre of the molecule, shows maximum  
36  
37 339 bending of the chromophore and a well developed *cis*-band, with corresponding  
38  
39 340 decrease in intensity of the main absorption band (Pettersson and Jonsson, 1990).  
40  
41  
42 341  
43  
44  
45 342 The intensity of the *cis*-band is essentially greater as the *cis* double bond is nearer the  
46  
47 343 centre of the molecule and is therefore empirically diagnostic. In a symmetrical di-*cis*  
48  
49 344 carotenoid, the centre of symmetry may be restored so that the *cis*-band again  
50  
51 345 becomes a weak feature. For di-*cis* and poly-*cis* carotenoids, a larger hypsochromic  
52  
53 346 shift in the main absorption band may be seen e.g. 13 nm for di-*cis* norbixin (Figure  
54  
55 347 3) (Scotter *et al.*, 1994).  
56  
57  
58 348  
59  
60

1  
2  
3 349 [Insert Figure 3 about here]  
4  
5  
6 350  
7

8 351 A numerical notation similar to the % III/II notation to indicate spectral fine structure,  
9  
10 352 has been adapted to designate the relative intensity of the *cis*-peak (Kohler, 1995).

11  
12 353 The intensity or absorbance of the *cis*-peak is expressed as a percentage of the  
13  
14  
15 354 absorbance of the middle main absorption band, which is usually the  $\lambda_{\max}$ . Scotter *et*  
16  
17 355 *al.* (1994) used this technique to study the spectra obtained by HPLC-photodiode  
18  
19  
20 356 array analysis of geometrical isomers of bixin and norbixin. The absorption intensity  
21  
22 357 at  $\lambda_{\max}$  (Figure 4, III) for each isomer was normalised and a comparison of relative  
23  
24  
25 358 intensity (REL (%)) made with the three other maxima at I, II and IV, to give REL(I),  
26  
27 359 REL(II) and REL(IV) respectively. In all cases,  $\lambda_{\max}$  (II) took the form of inflection  
28  
29  
30 360 rather than a peak, which made exact location of the wavelength maximum difficult.  
31  
32 361 However, post-run analysis of the spectral data allowed first derivative spectra to be  
33  
34  
35 362 taken which facilitated location of  $\lambda_{\max}$  (II), as shown in Figure 5.

36  
37 363

38  
39 364 [Insert Figure 4 about here]

40  
41 365 [Insert Figure 5 about here]  
42  
43  
44 366

### 45 46 367 ***Annatto stability***

47  
48  
49 368 It is well known that the polyene chain in carotenoids is responsible for their  
50  
51 369 instability i.e. their susceptibility to oxidation by various agents such as oxygen and  
52  
53 370 peroxides, addition of electrophiles including  $H^+$  and Lewis acids, and *cis*-/*trans*-  
54  
55  
56 371 isomerization due to various factors such as temperature and light as discussed above.  
57  
58 372 Other undesirable reactions may also be promoted by higher temperatures and the  
59  
60 373 light, and exposure to strong acids and alkalis should normally be avoided.

374

375 *Oxidative stability*

376 Annatto, especially norbixin, is susceptible to oxidation, particularly when applied in

377 powdered form due to the large surface area and when incorporated into foodstuffs;

378 although some foods can have a stabilizing effect (Berset and Marty, 1986; Collins,

379 1992; Levy and Rivadeneira, 2000). Spray-dried norbixin formulated with acaia gum

380 or maltodextrin as carriers have been reported to be particularly susceptible to

381 oxidation (Henry, 1992). The level of bixin prepared from annatto seeds and stored

382 for ca. 1 year at 30°C in packages comprising materials with different oxygen

383 transmission rates was reduced by ca. 10% during the first 2-3 weeks storage but

384 stabilized thereafter except for polyethylene film, which exhibited a degradation rate

385 of 0.04% per day, reflecting the permeability of the polyethylene (Carvalho *et al.*,

386 1993). Several mechanisms have been put forward for the effect of water activity on

387 the reduction of bixin oxidation in microcrystalline cellulose-based model systems to

388 simulate dehydrated foods (Gloria *et al.*, 1995). Bixin degradation followed first order

389 kinetics and the observed half-lives showed greater stability in systems of

390 intermediate and high water activity. It was postulated that this is because of the

391 ability of water to exclude oxygen from liposoluble materials by surface adsorption,

392 hydrogen bond with hydroperoxides, inactivate metal catalysts, reduce of free radicals

393 and lower the stability of singlet oxygen. Annatto has been shown to inhibit

394 hydroperoxide formation leading to triglyceride autoxidation by trapping peroxy

395 radicals (Haila *et al.*, 1996). Annatto was among a number of mediterranean spices

396 whose antioxidant capacities were compared with permitted food antioxidants in lipid

397 peroxidation (Martinez-Tomé *et al.*, 2001). Annatto was reported to have a greater

398 antioxidant capacity than either butylated hydroxyanisole (BHA) or butylated

1  
2  
3 399 hydroxytoluene (BHT) for preventing deoxyribose damage by hydroxyl radicals. In  
4  
5 400 aqueous media, annatto exhibited a lower antioxidant activity than propyl gallate but  
6  
7  
8 401 was more effective at peroxide scavenging than BHA or BHT. Annatto oleoresin  
9  
10 402 prepared by oil extraction of seeds was found to be more stable than a powdered  
11  
12 403 formulation during storage over ca. 1 year. Samples were stored in glass bottles with a  
13  
14  
15 404 3cm headspace of air. The greatest losses (60%) were observed for the powder at  
16  
17 405 ambient temperature in daylight, compared to ambient temperature in the dark (54%)  
18  
19 406 and at 5-8°C in the dark (23%). These results concur with the findings of Najar,  
20  
21 407 Bobbio and Bobbio, (1988) that light is the main degradation factor. Moreover,  
22  
23 408 photosensitized bixin is very reactive towards oxygen and thus may be considered as a  
24  
25 409 an oxygen quencher; the reaction of bixin with singlet oxygen is a related issue and is  
26  
27 410 discussed below.  
28  
29  
30  
31

32 411  
33  
34 412 Norbixin was the only carotenoid that inhibited the oxidative deterioration of lipids in  
35  
36 413 both olive oil and oil-in-water emulsions stored at 60°C and displayed a similar  
37  
38 414 activity to  $\delta$ -tocopherol in stored oil (Kiokias and Gordon, 2003). In olive oil-in-water  
39  
40 415 emulsions, norbixin reduced hydroperoxide formation and a synergistic effect  
41  
42 416 between norbixin and ascorbic acid or ascorbyl palmitate was observed.  
43  
44  
45  
46  
47

48 418 Bixin has been reported to be able to scavenge hydroxyl radicals generated by ferrous  
49  
50 419 ions ( $\text{Fe}^{2+}$ ) and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) but no mechanism was suggested (Zhao *et*  
51  
52 420 *al.*, 1998). Similarly, the behaviour of norbixin during in vivo plasmid DNA damage  
53  
54 421 induced by reactive oxygen species  $\text{Fe}^{2+}$ ,  $\text{Sn}^{2+}$  and  $\text{H}_2\text{O}_2$  have been studied and it has  
55  
56 422 proposed that since norbixin contains two free carboxyl moieties, its protective action  
57  
58  
59  
60

1  
2  
3 423 may rely on the formation of complexes (Kovary *et al.*, 2001). Norbixin showed a  
4  
5 424 stronger affinity for  $\text{Sn}^{2+}$  than for  $\text{Fe}^{2+}$  but was readily displaced by EDTA.  
6  
7

8 425

9  
10 426 During the isolation and analysis of carotenoids, the exclusion of atmospheric air by  
11  
12 427 inert gas or vacuum is strongly recommended in order to minimize the risk of  
13  
14 428 destruction or undesired reactions (Scheidt and Liaaen-Jensen, 1995) and annatto is  
15  
16 429 no different in this respect. The oxidation 'products' of bixin were identified  
17  
18 430 tentatively after transformation of bixin in corn oil at 125°C using spectrophotometry  
19  
20 431 and paper chromatography (McKeown and Mark, 1962). Evidence for the oxidative  
21  
22 432 decomposition of *cis*-bixin on TLC plates has been reported, where both powdered  
23  
24 433 colour formulations and chloroform solutions of *cis*-bixin exhibited decreased colour  
25  
26 434 content when stored in the dark in air at ambient temperature (Reith and Gielen,  
27  
28 435 1971). This was concluded to be due to the presence of oxygen, supported by  
29  
30 436 observing the relatively lower stability of cheese colour (norbixin in aqueous KOH)  
31  
32 437 compared to butter colour (bixin in vegetable oil), due to the presence of tocopherols  
33  
34 438 in the latter. The effects of light, air and pro-oxidants on the stability of annatto  
35  
36 439 extracts in chloroform over a 12 day period were monitored by spectrophotometry.  
37  
38 440 Air was much less effective at promoting loss of colour compared to light or to  
39  
40 441 benzoyl peroxide, a free-radical promoter (Najar, Bobbio and Bobbio, 1988). The  
41  
42 442 authors concluded that rapid loss of colour might occur whenever free radical  
43  
44 443 formation is promoted.  
45  
46  
47  
48  
49  
50  
51  
52

53 444

#### 54 445 *Reaction with singlet oxygen*

55  
56 446 Model studies on the photosensitized isomerization of *cis*-bixin show that while bixin  
57  
58 447 in the ground electronic state is stable to thermal isomerization, energy transfer via  
59  
60

1  
2  
3 448 photosensitization gives rise to the higher energy triplet state ( $^3\text{BIX}^*$ ) precursor,  
4  
5 449 which readily isomerizes to the *trans*- isomer (Montenegro *et al.*, 2004). The rate of  
6  
7 450 isomerization is dependent on several factors which compete for deactivation of  
8  
9  
10 451  $^3\text{BIX}^*$  e.g. ground state bixin and triplet oxygen ( $^3\text{O}_2$ ). Primary reaction products are  
11  
12 452 only degraded in the presence of air and under prolonged illumination, which is due to  
13  
14 453 the formation of oxidation products by reaction with singlet oxygen ( $^1\text{O}_2$ ). The  
15  
16 454 associated reaction mechanisms are discussed very elegantly by the authors. In a  
17  
18 455 similar study, the  $^3\text{BIX}^*$  energy level was calculated used laser-induced  
19  
20 456 photoacoustic calorimetry of bixin in methanol; acetonitrile solution (Rios *et al.*,  
21  
22 457 2007). The results of the study showed that bixin is a very efficient quencher of  $^1\text{O}_2$  in  
23  
24 458 fluid solutions due to an efficient energy-transfer process, and confirmed that that the  
25  
26 459  $^3\text{BIX}^*$  energy level is lower than that of  $^1\text{O}_2$  ( $18 \pm 2$  kcal/mol and 22.5 kcal/mol  
27  
28 460 respectively).  
29  
30  
31  
32  
33  
34  
35  
36  
37

#### 38 462 *Thermal stability*

39 463 While bixin and norbixin have good heat stability during food processing compared to  
40  
41 464 other carotenoids, 9'-*cis*-bixin undergoes a series of complex degradation reactions at  
42  
43 465 commercial extraction temperatures to produce a range of products coloured pale  
44  
45 466 yellow to orange (Iversen and Lam, 1953; Levy and Rivadeneira, 2000). Using paper  
46  
47 467 chromatography, the pigments in commercial annatto preparations were separated into  
48  
49 468 a series of bands that included a number of yellow bands comprising up to 40% of the  
50  
51 469 total pigments and including bright yellow fluorescent (*sic*) band (McKeown, 1961).  
52  
53 470 This band was thought to be the pale yellow breakdown product of bixin identified  
54  
55 471 previously (Iversen and Lam, 1953). The main thermal degradation product of 9'-*cis*-  
56  
57 472 bixin has since been isolated and identified using paper chromatography and UV/VIS  
58  
59  
60

1  
2  
3 473 spectrophotometry as the yellow coloured 17-carbon polyene 4,8,dimethyl-  
4  
5 474 tetradecaheptaenedioic acid monomethyl ester “C<sub>17</sub>” (McKeown and Mark, 1962;  
6  
7  
8 475 McKeown, 1963 and 1965; Preston and Rickard, 1980. The influence of heating time  
9  
10 476 on the thermal degradation of bixin in alkaline extracts of annatto showed that  
11  
12 477 pigment stability is related to the initial quantity of *cis*- and *trans*- bixin as well as to  
13  
14 478 the method used to obtain the extracts (Prentice-Hernández, Rusig and Carvalho,  
15  
16 479 1993).  
17  
18  
19 480  
20  
21  
22 481 The C<sub>17</sub> product has since been confirmed to be predominantly the *trans*- isomer and  
23  
24 482 that *cis*-isomerisation of bixin was prerequisite to its formation (Scotter, 1995).  
25  
26 483 However, this compound was shown to isomerise in solution to form small amounts  
27  
28 484 of *cis*- isomers and to be susceptible to hydrolysis thus forming a range of compounds  
29  
30 485 analogous to bixin and norbixin in terms of their chemical structures and  
31  
32 486 chromatographic properties. In the light of the results obtained, the mechanism of C<sub>17</sub>  
33  
34 487 formation originally suggested (McKeown, 1963) was postulated as a concerted  
35  
36 488 electrocyclic process followed by the elimination of *m*-xylene and, to a much lesser  
37  
38 489 extent toluene, toluic acid and toluic acid methyl ester, and the formation of C<sub>17</sub> which  
39  
40 490 can degrade further by a similar mechanism.  
41  
42  
43 491  
44  
45  
46 492 The analytical HPLC-photodiode array (PDA) method developed by Scotter *et al.*  
47  
48 493 (1994, 1995) provided superior qualitative and quantitative data compared with UV-  
49  
50 494 VIS spectroscopic methods (McKeown and mark, 1962; Smith, Blake and Porter,  
51  
52 495 1983) for determining the colour content (as bixin and norbixin) in 21 commercial  
53  
54 496 annatto formulations, particularly with respect to the coloured thermal degradation  
55  
56 497 products (Scotter *et al.*, 1998). Moreover, the levels of the all-*trans* and di-*cis*-  
57  
58  
59  
60

1  
2  
3 498 isomers of norbixin determined from chromatographic profiles of two different  
4  
5 499 norbixin formulations were found to be consistent with their known production  
6  
7  
8 500 history i.e. indicative of the degree of thermal treatment. The formulation obtained by  
9  
10 501 direct aqueous alkaline extraction contained higher levels of these isomers compared  
11  
12 502 to solvent pre-extracted bixin followed by alkaline hydrolysis obtained using lower  
13  
14 503 temperatures. However, the authors pointed out that while the isomer profiles  
15  
16 504 obtained by HPLC-PDA analysis support this, the different extraction procedures  
17  
18 505 might also give rise to different isomer profiles due differential solubilities and  
19  
20 506 stabilities in the extraction medium. The effects of light and oxygen may further  
21  
22 507 complicate this during extraction and handling, and by the nature of the source  
23  
24 508 material.  
25  
26  
27  
28  
29  
30  
31

32 510 In a follow-up study, a method was developed which used ambient alkaline hydrolysis  
33  
34 511 followed by solvent extraction and gas chromatography (GC), to analyse annatto  
35  
36 512 colour formulations for the main aromatic hydrocarbon thermal degradation products  
37  
38 513 m-xylene and toluene (Scotter *et al.*, 2000). Of the 20 samples analysed, 15 contained  
39  
40 514 <5 mg/kg toluene Four samples contained between 5 and 10 mg/kg and one sample  
41  
42 515 contained 12 m/kg toluene but these levels were not indicative of significant toluene  
43  
44 516 formation via thermal degradation of annatto. In contrast, 6 samples comprising both  
45  
46 517 bixin and norbixin formulations contained m-xylene in the range 30 – 200 mg/kg with  
47  
48 518 the highest level found in an oil-based bixin formulation. Moreover, the two norbixin  
49  
50 519 formulations of known production history analysed in the previous study (Scotter *et*  
51  
52 520 *al.*, 1998) differed markedly in m-xylene content, which appeared to be consistent  
53  
54 521 with the degree of thermal treatment.  
55  
56  
57  
58  
59  
60

522



1  
2  
3 523 For comparison with the alkaline hydrolysis-solvent extraction procedure, 7 of the  
4  
5 524 annatto formulations were submitted for headspace (HS) GC analysis for toluene and  
6  
7  
8 525 m-xylene in order to monitor the effects of heating in a closed controlled environment  
9  
10 526 (90°C for 20 minutes). An increase in m-xylene was observed, with the bixin in oil  
11  
12 527 formulations showing the highest rise in m-xylene concentration on heating. The  
13  
14  
15 528 authors anticipated that HS-GC could be used to monitor the thermal degradation of  
16  
17 529 annatto in food systems and thus conducted a number of experiments in combination  
18  
19 530 with HPLC-PDA and GC-MS to study this (Scotter *et al.*, 2001). Low levels (ca. 10 –  
20  
21 531 15 ug/kg) of m-xylene were detected in the headspace of annatto-coloured retail  
22  
23 532 samples of custard powder, extruded snacks, margarine and breadcrumbs but not in  
24  
25 533 control samples. Much higher levels of m-xylene were detected in annatto-coloured  
26  
27 534 smoked herring (kippers) at ca. 150-200 ug/kg and m-xylene was observed in the  
28  
29 535 headspace of heated Red Leicester cheese (not quantified). The C<sub>17</sub> coloured annatto  
30  
31 536 degradation product was also detected, indicating that thermal degradation of the  
32  
33 537 principal annatto colouring agent 9'-*cis*-bixin in model systems and foods is facile.  
34  
35 538 However, the degradation is complicated many competing isomerisation reactions  
36  
37 539 which proceed at different rates towards equilibrium. This is further complicated by  
38  
39 540 the simultaneous and irreversible formation of C<sub>17</sub> associated with the production of  
40  
41 541 m-xylene and to a lesser extent, toluene. While norbixin was reported to degrade  
42  
43 542 similarly but more slowly, the levels of m-xylene formation were nonetheless  
44  
45 543 consistent with bixin / norbixin concentration in the food and occurred more rapidly at  
46  
47 544 higher temperatures.  
48  
49 545  
50  
51 546 In order to better understand the kinetics and yields for the formation of both the  
52  
53 547 coloured and aromatic hydrocarbon thermal degradation products of annatto, the  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 548 authors carried out a number of experiments in model systems (*ibid.*). The thermal  
4  
5 549 stability of bixin at the boiling point of three homologous alcohol solvents was  
6  
7  
8 550 evaluated using HPLC-PDA to monitor the rate of loss of 9'-*cis*-bixin as well as the  
9  
10 551 appearance of a di-*cis*- and *trans*- isomer, and the C<sub>17</sub> degradation product. Loss of  
11  
12 552 linearity was observed at each temperature beyond 2 hours, suggesting that two or  
13  
14  
15 553 more competing reactions were taking place at different rates. From the rate constants  
16  
17 554 calculated for the initial phase of the reaction, the Arrhenius activation energy for the  
18  
19 555 loss of 9'-*cis*-bixin in refluxing alcohol solvent was 35.7 kJ.mol<sup>-1</sup>. Since the rate of  
20  
21 556 loss of 9'-*cis*-bixin was measured as a function of time regardless of reaction pathway  
22  
23 557 i.e. isomerisation vs. degradation), the authors concluded that the rate data represented  
24  
25 558 only the total (summed) values. Thus, several concurrent reaction pathways are  
26  
27 559 available hence deviation from first order kinetics at long observation times was not  
28  
29 560 unexpected as suggested in Figure 6.  
30  
31  
32  
33

34 561  
35  
36 562 [Insert Figure 6 about here]  
37  
38  
39 563

40  
41 564 Berset and Marty (1986) had reported previously an activation energy of 125 kJ.mol<sup>-1</sup>  
42  
43 565 for the thermal degradation of annatto pigments in petroleum jelly using a simple  
44  
45 566 first-order kinetic model for the complete decay. This disparity in values therefore  
46  
47 567 suggests a controversy in the kinetic analysis or a misinterpretation of the  
48  
49 568 experimental data. Interestingly, bixin was reported to be easily transformed to the all-  
50  
51 569 *trans*- isomer at ambient temperature in the presence of a photosensitizer and light,  
52  
53 570 where the activation energy for the excitation of bixin to an excited triplet state was  
54  
55 571 ca. 25 kJ.mol<sup>-1</sup> as discussed above (Montenegro *et al.*, 2004), which suggests strongly  
56  
57  
58  
59  
60

1  
2  
3 572 that a greater energy barrier may be anticipated for the thermal isomerisation of 9'-*cis*-  
4  
5 573 bixin to *trans*-bixin.  
6  
7

8 574

9  
10 575 A more detailed kinetic study on the thermal degradation of bixin in an aqueous  
11  
12 576 model system comprising water: ethanol (8:2) as a function of temperature has been  
13  
14 577 described, where HPLC was used to monitor the decay of 9'-*cis*-bixin and the  
15  
16 578 formation of the di-*cis*- and *trans*- isomers, as well as C<sub>17</sub> (Rios *et al.*, 2005). The  
17  
18 579 reactions were found not to follow first order rate characteristics but rather fitted well  
19  
20 580 to a biexponential model. The rate constants for the *formation* of the primary products  
21  
22 581 of bixin and the energy barriers for each step were calculated. Di-*cis*- isomers were  
23  
24 582 formed immediately (energy barrier ca. 63 kJ.mol<sup>-1</sup>) followed by a slow consumption  
25  
26 583 (with the associated decay of 9'-*cis*-bixin), indicating their role as reaction  
27  
28 584 intermediates. The di-*cis*- isomers can either revert readily to 9'-*cis*-bixin (ca. 13  
29  
30 585 kJ.mol<sup>-1</sup>) or yield the primary C<sub>17</sub> degradation product with a higher energy  
31  
32 586 requirement of ca. 27 kJ.mol<sup>-1</sup>). However, the isomerisation of 9'-*cis*-bixin to *trans*-  
33  
34 587 bixin requires ca. 100 kJ.mol<sup>-1</sup>, thereby explaining its relatively slow formation. The  
35  
36 588 Arrhenius plot obtained from the initial decay component for 9'-*cis*-bixin yielded an  
37  
38 589 activation energy of ca. 33 kJ.mol<sup>-1</sup>, which concurs with earlier data (Scotter *et al.*,  
39  
40 590 2001). In conclusion, while the activation energy obtained for the 9'-*cis*- → *trans*-  
41  
42 591 isomerisation of bixin is very similar to that reported for β-carotene, the value of ca.  
43  
44 592 155 kJ.mol<sup>-1</sup> for the summed isomerisation steps of bixin is much higher than those  
45  
46 593 reported for the thermal isomerisation of C<sub>40</sub> carotenoids (ca. 105 kJ.mol<sup>-1</sup>). Thus the  
47  
48 594 reaction scheme suggested by Scotter *et al.* (2001) and the greater relative stability of  
49  
50 595 bixin, especially during its isolation and manipulation were confirmed (Figure 7).  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60 596

1  
2  
3 597 [Insert Figure 7 about here]  
4  
5

6 598  
7

8 599 Thermogravimetric analysis has been used to investigate the thermal degradation of  
9  
10 600 bixin derived from annatto seeds at different heating rates over the 25 – 900°C  
11  
12 601 temperature range (Silva *et al.*, 2005). The results indicated that the decomposition of  
13  
14 602 solid 9'-*cis*-bixin occurs in the liquid phase and that four decomposition stages are  
15  
16 603 evident over the temperature range 205-545°C, with isomerisation to the *trans*- isomer  
17  
18 604 occurring between 200 and 240°C. The calculated activation energy was dependent  
19  
20 605 upon heating rate (i.e. 5, 10 or 15 K.min<sup>-1</sup>) at ca. 108, 147 and 128 kJ.mol<sup>-1</sup>  
21  
22 606 respectively compared to the value of ca. 100 kJ.mol<sup>-1</sup> reported by Rios *et al.*, (2005)  
23  
24 607 obtained in solution. In a similar follow up study, *cis*-norbixin was heated at rates of  
25  
26 608 5, 10 and 20°C.min<sup>-1</sup> over the temperature range 25 – 900°C, where the thermal  
27  
28 609 decomposition reactions occurred in the solid phase (Silva *et al.*, 2007). Using the  
29  
30 610 Coats-Redfern model, the calculated activation energy was dependent upon heating  
31  
32 611 rate at ca. 154, 131 and 99 kJ.mol<sup>-1</sup> at 5, 10 and 20°C.min<sup>-1</sup> respectively for the first-  
33  
34 612 order process.  
35  
36  
37  
38  
39  
40

41 613  
42

43 614 Heating solid non-purified extracts of annatto seeds as a thin film deposited on a  
44  
45 615 silicon wafer *in vacuo* and monitored using time of flight (ToF) secondary ion mass  
46  
47 616 spectrometry (SIMS), does not give the same results as heating in solution  
48  
49 617 (Bittencourt *et al.*, 2005. Principal component analysis revealed that the thermal  
50  
51 618 degradation of the annatto extracts under these conditions occurs in three distinct  
52  
53 619 temperature ranges; below 70°C, the extracts remain thermally stable but above this  
54  
55 620 temperature dimerization reactions occur and the signals attributed to bixin decrease.  
56  
57  
58 621 Near to 100°C, the bixin molecules begin to degrade, leading to fragmentation with  
59  
60

1  
2  
3 622 extensive degradation of bixin above 120°C. However, the nature of the degradation  
4  
5 623 mechanism described is not fully understood since there was no evidence for the  
6  
7  
8 624 formation of C<sub>17</sub> or related fragments from solid bixin.  
9

10 625

11  
12  
13 626 *Light stability*

14  
15 627 The effect of light at 900 lux intensity on the 30-day stability of a microencapsulated  
16  
17 628 water-miscible extract of bixin compared to that of a purified bixin extract have been  
18  
19 629 studied by measuring the loss of spectrophotometric absorbance at 470nm with time  
20  
21  
22 630 (Prentice-Hernández and Rusig, 1999). The degradation rate of bixin in the  
23  
24 631 microencapsulated extract was ca. 0.05% compared to 0.11% per day for the purified  
25  
26 632 extract.  
27

28  
29 633

30  
31 634 Ferreira *et al.*, (1999) submitted commercial water-soluble annatto (norbixin)  
32  
33 635 solutions to different time and temperature treatments to investigate colour stability.  
34  
35 636 The colour change was measured by spectrophotometry using the Hunter Lab System  
36  
37 637 and the results presented in terms of changes in the norbixin concentration and L, a, b  
38  
39 638 colour parameters. Data were analysed for reaction order and the temperature  
40  
41 639 dependence was explained by the Arrhenius model, with activation energy values  
42  
43 640 between 46 and 105 kJ.mol<sup>-1</sup>. The changes in colour showed an increase in lightness  
44  
45 641 and yellow colour and a decrease in red colour. Norbixin degradation reaction  
46  
47 642 followed second order kinetics whereas for other colour parameters, first order  
48  
49 643 kinetics was followed.  
50

51 644

52  
53 645 The light stability of spray-dried bixin encapsulated with gum Arabic or maltodextrin  
54  
55 646 plus Tween 80 surfactant has been reported, where the kinetic behaviour of bixin  
56  
57  
58  
59  
60

1  
2  
3 647 photo degradation in all systems was characterized by two first-order decays due to  
4  
5 648 the presence of bixin both inside and outside the microcapsules (Barbosa *et al.*, 2005).  
6  
7  
8 649 Unsurprisingly, approximately two orders of magnitude greater stability was observed  
9  
10 650 for bixin in the dark compared to illuminated conditions and in the absence of light,  
11  
12 651 and bixin stability in encapsulated solutions was ca. ten times greater than in non-  
13  
14 652 encapsulated systems. The effect of processing conditions used for the preparation of  
15  
16 653 traditional Indian foods on bixin stability including baking, frying, microwave  
17  
18 654 cooking and pressure cooking have been monitored by following losses using  
19  
20 655 spectrophotometry (Rao *et al.*, 2005). The losses of bixin under model processing  
21  
22 656 conditions was compared to the preparation of cakes, chegodis, biscuits and fried rice.  
23  
24 657 The greatest losses of bixin were observed in direct exposure to oven baking (54%  
25  
26 658 loss ) and deep fat frying (47%), whereas microwave cooking did not affect the colour  
27  
28 659 during direct exposure or in food products. The maximum loss of bixin (65%) was  
29  
30 660 observed for deep fried snack due largely to leaching of the dye into the oil. Pressure  
31  
32 661 cooking produced losses of between 25% and 33%. In cakes, the loss was 30% but  
33  
34 662 negligible losses were found for biscuits (1.5%).  
35  
36 663  
37  
38 664 Similarly, the combined effects of light and temperature on annatto extract under  
39  
40 665 different storage conditions were evaluated spectrophotometrically at 470nm in  
41  
42 666 chloroform over a period of 360 days (Balaswamy, Prabhakara Rao, Satyanarayana  
43  
44 667 and Rao, 2006). Annatto oleoresin was generally more stable during storage with  
45  
46 668 respect to bixin content than annatto powder obtained by solvent extraction of annatto  
47  
48 669 seeds. The bixin lost in oleoresin stored under cold (5-8°C), dark conditions was  
49  
50 670 minor (11%) throughout the study, whereas considerable losses were observed for the  
51  
52 671 powdered dye (23%). Likewise, the bixin lost after storage at ambient temperature in  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 672 the dark were 8% and 54% for oleoresin and powder respectively. Under diffused  
4  
5 673 daylight ambient and temperature the losses were 14% and 60% respectively, whereas  
6  
7  
8 674 bixin seed stored in jute sacks showed a loss of only 15%. As expected, the rate  
9  
10 675 constants for bixin degradation were much higher in powder compared to oleoresin  
11  
12 676 and were reported to follow second order kinetics. It was proposed that the colour is  
13  
14 677 protected from exposure to oxygen and light by the oleoresin compared to the dry  
15  
16 678 powder, which has a large surface area.  
17  
18  
19

20 679

21  
22 680 Bixin complexed with  $\alpha$ -cyclodextrin is also reported to be more resistant to the  
23  
24 681 damage caused by light and air (Lyng, Passos and Fontana, 2005).  
25  
26

27 682

28  
29 683 *Analytical methods for annatto*

30  
31 684 *Spectrophotometry*

32  
33 685 Historically, chloroform has been used as solvent for the spectrophotometric analysis  
34  
35 686 of bixin and dilute sodium hydroxide (ca. 0.1M) for norbixin. Absorbance  
36  
37 687 measurements at the two most intense spectral peaks (III and IV in Figure 4) are used  
38  
39 688 for quantitative analysis, where peak IV is preferred because it is less prone to  
40  
41 689 interference from yellow decomposition products. This interference was corrected by  
42  
43 690 using a factor related to the absorbances at  $\lambda_{\max}$  and at 404nm in determining the total  
44  
45 691 pigment content of annatto formulations (McKeown and Mark, 1962). In practice, the  
46  
47 692 spectrophotometric determination of annatto (as bixin or norbixin) is somewhat  
48  
49 693 confused by the use of conflicting extinction coefficients. This has been discussed in  
50  
51 694 detail and the published ( $E_{1\text{cm}}^{1\%}$ ) extinction coefficients for norbixin and bixin  
52  
53 695 summarized and compared to highlight disparities (Levy and Rivadeneira, 2000).  
54  
55 696 Depending upon the extinction coefficient used, large errors might be incurred and  
56  
57  
58  
59  
60

1  
2  
3 697 propose a practical conversion factor to correlate the relative absorbances at the two  
4  
5 698 peak maxima. This is based upon the increase in absorbance observed upon hydrolysis  
6  
7  
8 699 of bixin to norbixin at constant concentration – thus proving that the extinction value  
9  
10 700 for norbixin must be higher than that for bixin, which was also reported (Smith *et al.*,  
11  
12 701 1983). Furthermore, from data recorded by the authors from more than 1000  
13  
14 702 spectrophotometric measurements of different samples of bixin before and after  
15  
16 703 hydrolysis, the difference between the extinction values of bixin and norbixin was  
17  
18 704 reported to be of the order of 6%. When compared with a value of  $E_{1\text{cm}}^{1\%} = 3208$   
19  
20 705 reported for pure norbixin, this equates to an extinction coefficient for bixin of  $E_{1\text{cm}}^{1\%} =$   
21  
22 706 3016, which concurs with the values reported for purified bixin in chloroform (Scotter  
23  
24 707 *et al.*, 1994).

25  
26 708  
27  
28 709 However, these extinction values do not agree with those adopted for colour purity  
29  
30 710 specifications by the European Union (EC, 1995) or the FAO/WHO (1996), largely  
31  
32 711 due to misassumptions made regarding solvent effects. The discrepancy in published  
33  
34 712 extinction values might be traced back to the ‘erroneous’ coefficient reported by Reith  
35  
36 713 and Gielen (1971) that has been used subsequently as a reference value by various  
37  
38 714 other workers. Serious doubt is expressed over the validity of the extinction values for  
39  
40 715 norbixin in aqueous alkaline solution at 453nm (2850) and 482nm (2550). Moreover,  
41  
42 716 the same reservations were expressed over the value of 3473 at 453nm reported by the  
43  
44 717 FAO/WHO specification (FAO/WHO, 1981).

45  
46 718  
47  
48 719 An interesting and important aspect of the spectrophotometric analysis of bixin in  
49  
50 720 chloroform is its rapid rate of degradation when contained in a quartz cuvette, which  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 721 unlike glass cuvettes allows the transmission of ultraviolet light (i.e. < 300nm) (Levy  
4  
5 722 and Rivadeneira, 2000).

6  
7  
8 723

9  
10 724 *Planar chromatography*

11  
12 725 Prior to 1961, there were few references in the literature to paper and adsorption

13  
14 726 chromatography, which dealt mainly with the gross separation of different

15  
16 727 carotenoids, and from chlorophylls. The first paper chromatographic method for the

17  
18 728 direct separation of annatto colouring components used Whatman 3MM paper

19  
20 729 impregnated with 50% N,N-dimethylformamide (DMF) in acetone and which

21  
22 730 developed with cyclohexane: chloroform: DMF and acetic acid (85:10:3:2)

23  
24 731 (McKeown, 1961). This method was used in a number of subsequent studies on

25  
26 732 annatto and its main thermal degradation product, C<sub>17</sub> (McKeown and Mark, 1962;

27  
28 733 McKeown, 1963). The first thin layer method for separation of annatto colour and

29  
30 734 other fat-soluble dyes shortly thereafter, employed silica gel G, plaster of Paris and

31  
32 735 silicic acid media with amyl acetate mobile phase (Ramamurthy and Bhalerao, 1964).

33  
34 736 However, of the 30 solvent systems studied, bixin was reported to migrate from the

35  
36 737 base line only when acetic acid was present (Francis, 1965). The findings suggested

37  
38 738 that the amyl acetate solvent used by Ramamurthy and Bhalerao (1964) must have

39  
40 739 contained acetic acid as an impurity, which was proven by subsequent

41  
42 740 experimentation. Later methods used silica gel with various solvent systems

43  
44 741 containing acetic acid for the separation of bixin and norbixin in colour formulations

45  
46 742 (Dendy, 1966) and cheese colour i.e. norbixin (Reith and Gielen, 1971), who also

47  
48 743 employed cellulose media for the analysis of butter colour i.e. bixin. Other methods

49  
50 744 include those developed by Preston and Rickard (1980) and Corradi and Micheli

51  
52 745 (1981). Chao *et al.* (1991) used reverse phase (C<sub>18</sub>) plates with methanol:water mobile

1  
2  
3 746 phase to separate annatto pigments from supercritical CO<sub>2</sub> extractions of annatto  
4  
5  
6 747 seeds. More recently, TLC has been used for the detection of bixin and other food  
7  
8 748 colour carotenoids derived from red pepper (Mínguez-Mosquera, Hornero-Méndez  
9  
10 749 and Garrido-Fernández, 1995) and for the isolation and identification of new (trace)  
11  
12 750 apocarotenoids from annatto seeds (Mercadante, Steck and Pfander, 1997b) and in the  
13  
14 751 bioautographic detection of antimicrobial compounds in water-soluble annatto  
15  
16 752 extracts (Galindo-Cuspinera and Rankin, 2005). The various methods are summarized  
17  
18 753 in Table 2.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 754

30 755 [Insert Table 2 about here]

31 756

32 757 *HPLC*

33 758 As discussed above, developments in HPLC techniques have enabled more detailed  
34 759 studies of other bixin and norbixin isomers as well as their degradation products  
35 760 compared to TLC methods and have been utilized to gain a greater understanding of  
36 761 the stability of annatto and which in turn have been applied to the detection and  
37 762 measurement of annatto colour in foodstuffs (below).  
38  
39  
40  
41  
42  
43  
44  
45

46 763

47 764 Literature references on the application of HPLC to the separation of annatto colouring  
48 765 components are sparse. Early methods include the HPLC analysis of annatto extract  
49 766 (Nishizawa *et al.*, 1983) and Smith *et al.* (1983), who reported the use of an isocratic  
50 767 reverse-phase system employing an ODS column and methanol/aqueous acetic acid  
51 768 mobile phase. Using this system the *cis*- and *trans*-isomers of both bixin and norbixin  
52 769 were separated within 10 minutes. However, the *cis*- and *trans*-bixin peaks were not  
53 770 fully resolved and the peak shapes were generally very poor. A method for the reverse-

1  
2  
3 771 phase separation of bixin, norbixin and three curcuminoids using both isocratic and  
4  
5 772 gradient elution systems, comprising a Zorbax ODS column and water/THF mobile  
6  
7  
8 773 phase was later developed that gave improved chromatographic separation (Rouseff,  
9  
10 774 1988) developed. However, only separation of the 'main' annatto colouring  
11  
12 775 components were reported and no reference to stereoisomer separation was given.  
13  
14 776 Other approaches have been reported for the analysis of cheese extracts (Luf and  
15  
16 777 Brandl, 1988) and of foods after protease digestion (Chatani and Adachi, 1988). A  
17  
18 778 procedure similar to that reported by Smith *et al.*, (1983) has been developed and  
19  
20 779 applied to the determination of annatto in selected foodstuffs with reasonable success  
21  
22 780 (Lancaster and Lawrence, 1995).  
23  
24  
25  
26  
27  
28

29 782 The method developed by Scotter *et al.* (1994) has played a key role in the  
30  
31 783 advancement of HPLC capabilities for the separation and characterization of norbixin  
32  
33 784 and bixin isomers, and has been refined and adapted for the study of annatto stability  
34  
35 785 and for the determination of annatto colouring components in colour formulations,  
36  
37 786 foodstuffs and human plasma. These are summarized along with other published  
38  
39 787 methods in Table 3.  
40  
41  
42  
43  
44

45 789 [Insert Table 3 about here]  
46  
47  
48  
49

50  
51 791 While the development of column stationary phases been vital in allowing separation  
52  
53 792 of geometrical isomers of bixin and norbixin, C<sub>17</sub> analogues and other food  
54  
55 793 components, it is the power of the detection systems that have enabled the  
56  
57 794 development of highly useful qualitative and quantitative analyses. Many developed  
58  
59 795 methods utilise detection with fixed wavelength UV-visible (UV-VIS) detectors at  
60

1  
2  
3 796 wavelengths specific to bixin/norbixin isomer absorption maxima quite successfully.  
4  
5 797 However, photodiode-array (PDA) technology offers combined sensitivity and  
6  
7 798 specificity coupled to real-time qualitative (spectral) confirmatory analysis, thereby  
8  
9 799 enabling powerful isomer identification and measurement. PDA allows isomer peaks  
10  
11 800 with different  $\lambda_{\max}$  wavelengths to be monitored using a spectral bandwidth that  
12  
13 801 encompasses them. A reference wavelength can also be used to subtract background  
14  
15 802 absorbances and to allow for baseline drift, which is usually set outside of the  
16  
17 803 absorbance range of the main analyte and interfering peaks e.g. at 600nm x 4nm  
18  
19 804 bandwidth. The lack of availability of authenticated reference standards is the main  
20  
21 805 limiting factor in the HPLC analysis of annatto colouring components but methods are  
22  
23 806 available for the isolation, purification and characterization of the main bixin and  
24  
25 807 norbixin isomers (Scotter *et al.*, 1994) and for C<sub>17</sub> analogues (Scotter, 1995). Other  
26  
27 808 workers have exploited the use of PDA detection for the identification of trace levels  
28  
29 809 of other apocarotenoids in annatto seeds very successfully (Mercadante, Steck and  
30  
31 810 Pfander, 1997b). Figure 8 shows the HPLC separation of bixin and norbixin isomers  
32  
33 811 (Scotter *et al.*, 1994).  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 813 [Insert Figure 8 about here]  
44  
45  
46  
47  
48  
49

#### 815 *Mass spectrometry (MS)*

50  
51 816 A comprehensive review on the use of mass spectrometry in the study of carotenoids  
52  
53 817 in general may be found elsewhere (Enzell and Back, 1990). This work cites earlier  
54  
55 818 reviews and studies that consolidate the importance of the technique not only for  
56  
57 819 elucidation of structure but also for analytical research, not least those carried out by  
58  
59 820 Vetter *et al.*, (1971), Budzikiewicz, (1974) and Enzell and Wahlberg, (1980). The  
60

1  
2  
3 821 1990 review covers in detail ionization techniques, tandem MS, combined  
4  
5 822 chromatographic-MS techniques and elimination reactions of in-chain units and  
6  
7  
8 823 terminal groups. The first method for electrospray liquid chromatography-mass  
9  
10 824 spectrometry (LC-ES-MS) of carotenoids employed gradient reversed-phase HPLC  
11  
12 825 with PDA and MS detection in tandem (van Breemen, 1995). Molecular ions,  $M^{(+)}$ ,  
13  
14  
15 826 without evidence of any fragmentation, were observed in the ES mass spectra of both  
16  
17 827 xanthophylls and carotenes but neither bixin nor norbixin were studied.  
18  
19  
20 828  
21  
22 829 In common with other carotenoids, the MS spectra of bixin and norbixin are  
23  
24 830 characterized by fragmentation leading to losses of toluene and xylene from the  
25  
26  
27 831 polyene chain and the structural significance of the intensity ratio of the  $[M-92]^+$  and  
28  
29 832  $[M-106]^+$  ions (and to a lesser extent the  $[M-158]^+$  ion), which is related to the  
30  
31 833 number of conjugated double bonds. It is the apo-configuration that gives rise to  
32  
33 834 anomalous MS properties of bixin and norbixin that have diagnostic value i.e. the -  
34  
35 835  $CH_2-CH=CH-CH_2-COOH$  end group gives characteristic fragments at  $[M-44]^+$  and  
36  
37 836  $[M-99]^+$ , whereas the  $-CH_2-CH=CH-CH_2-COOCH_3$  end group gives characteristic  
38  
39 837 fragments at  $[M-31]^+$ ,  $[M-59]^+$  and  $[M-113]^+$ . Solid probe electron ionization (EI+)  
40  
41 838 was used to confirm the structures of isolated and purified bixin and norbixin isomers  
42  
43 839 (Scotter *et al.*, 1994). Both the 9'-*cis*- and *trans*- isomers gave a molecular ion at  $m/z$   
44  
45 840 394 (bixin) and  $m/z$  380 (norbixin), with major fragment ions at  $m/z$   $[M-106]$ , 106  
46  
47 841 (xylene), 105 (methyl tropylium) and 91. Using thermospray analysis,  $[M+H]^+$  was  
48  
49 842 identified as the base peak along with the presence of sodium and (possibly) water  
50  
51 843 adducts, and fragment ions corresponding to  $[M-H_2O]^+$  and  $[M-CH_3OH]^+$ . In a later  
52  
53 844 study, similar analytical conditions were used to characterize the 17-carbon major  
54  
55 845 thermal degradation product of annatto (Scotter, 1995). Solid probe EI revealed the  
56  
57  
58  
59  
60

1  
2  
3 846 molecular ion at  $m/z$  288 along with fragment ions at  $m/z$  [M-106], 106, 105 and 91,  
4  
5 847 and thermospray analysis identified the base peak as [M+H]<sup>+</sup> as well as sodium  
6  
7 848 adducts at  $m/z$  311 ([M+Na]<sup>+</sup>) and 333 [M-H+2Na]<sup>+</sup>.  
8  
9 849  
10  
11 850 Complementary to other analytical techniques, EI<sup>+</sup> and fast atom bombardment  
12  
13 851 (FAB) MS was used to determine the structure of the bixin family of apocarotenoids  
14  
15 852 (Kelly *et al.*, 1996). Both *cis*- and *trans*- bixin isomers gave EI<sup>+</sup> molecular ion  
16  
17 853 abundancies equivalent to ca. 30% of the base peaks at  $m/z$  59 or 91, and the [M+1]  
18  
19 854 and [M+2] ion intensities were consistent with predictions based upon calculated <sup>13</sup>C  
20  
21 855 isotope patterns. As expected, loss of xylene as a neutral group was most pronounced  
22  
23 856 for *cis*-bixin but no loss of neutral toluene was observed although the  $m/z$  91 peak was  
24  
25 857 prominent. Fast-atom bombardment (FAB<sup>+</sup>) spectra of *cis*- and *trans*- bixin gave the  
26  
27 858 molecular ion as the base peak but the abundance of the [M+1] peak exceeded the  
28  
29 859 calculated isotopic abundance by 55-75%, indicating a small contribution from  
30  
31 860 [M+H]. Small amounts of sodium adducts were observed but ions due to elimination  
32  
33 861 of toluene were not. However, significant amounts of  $m/z$  105, 115 and 165 were  
34  
35 862 observed. These observations were consistent with other FAB spectra of carotenoids  
36  
37 863 where odd electron molecular ions are frequently observed due presumably to their  
38  
39 864 lower ionization potentials (Vetter and Meister, 1985).  
40  
41 865  
42  
43 866 Bixin was among the polyenes studied using electrospray ionization (EI) and high  
44  
45 867 resolution (HR) matrix-assisted laser desorption ionization (MALDI) time-of-flight  
46  
47 868 (TOF) mass spectrometry (Guaratini *et al.*, 2004). In this study, the ability of neutral  
48  
49 869 organic molecules to give up an electron for oxidation was exploited, which is  
50  
51 870 governed by the energy of their highest occupied molecular orbital (HOMO) and can  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 871 be estimated by measurement of the half-wave potential for solution oxidation. Strong  
4  
5 872 evidence was reported for an ionization process that produces the molecular ion  $M^+$  in  
6  
7  
8 873 ESI and HR-MALDI MS of polyenes, and the correlation of the observed ions to the  
9  
10 874 oxidation potential. The formation of  $M^+$  and  $[M+H]^+$  species was shown to be  
11  
12  
13 875 dependent upon energetic variations and the presence of water or another protic  
14  
15 876 solvent. Neither the  $[M+H]^+$  nor the  $[M+H-H_2O]^+$  were detected as the major ions  
16  
17 877 from ESI analysis of bixin, whereas  $M^+$  was detected but only in the specific capillary  
18  
19 878 voltage range of 0.1 to 0.7 kV. The accurate mass measurement afforded by the HR-  
20  
21 879 MALDI-TOF analysis showed  $M^+$  for bixin at an observed mass of 394.2147 with  
22  
23 880 40% ion intensity, but  $[M+H]^+$  was not observed.  
24  
25  
26  
27 881

28  
29 882 The major carotenoid composition of *Bixa orellana* seeds has been ascertained using  
30  
31 883 TOF-MS with X-ray photoelectron spectroscopy (Felicissimo *et al.*, 2004). The  
32  
33 884 presence of bixin was revealed in the seed aril without any sample pretreatment from  
34  
35 885 the detection of ions attributable to  $[M+2H]^+$  at  $m/z$  396 with associated  $^{13}C$  isotope  
36  
37 886 analogues at  $m/z$  397 and 398. The presence of characteristic fragments at  $m/z$  337  
38  
39 887 was attributed to  $C_{23}H_{29}O_2^+$  obtained from the previous molecular ion with loss of a  
40  
41 888  $COOCH_3$  ester group, and at  $m/z$  281, a fragment compatible with loss of a  $C_6O_2H_8$   
42  
43 889 end group plus a hydrogen atom i.e.  $C_{19}H_{21}O_2^+$ . The characteristic presence of xylene  
44  
45 890 was confirmed via the detection of the  $C_8H_9^+$  ion at  $m/z$  105. Analysis of the coloured  
46  
47 891 interior of the seeds following cutting did not show any fragments consistent with  
48  
49 892 bixin. A methanol:chloroform extract of the seeds was analysed immediately after  
50  
51 893 preparation by blow-drying under nitrogen onto a silver substrate, and then after  
52  
53 894 exposure to ambient light for 3 months. TOF-MS analysis of the fresh extract was  
54  
55 895 dominated by the molecular peak at  $m/z$  396 along with all other characteristic  
56  
57  
58  
59  
60

1  
2  
3 896 fragments. As expected after 3 months exposure to light, the colour of the extract had  
4  
5 897 lightened to a more yellow shade with an associated 5-fold decrease in the intensity of  
6  
7  
8 898 the  $[M+2]^+$  ion and with a concomitant 2-fold increase in the intensity of the  $C_8H_9^+$   
9  
10 899 ion, indicating the formation of xylene via degradation. In a related study, Bittencourt  
11  
12 900 *et al.* (2005) analysed extracts of *Bixa orellana* using TOF-MS as a means of  
13  
14 901 characterising thermal effects. The spectrum was characterised by a large number of  
15  
16 902 peaks generated by the principal ions and their multiple fragmentation patterns but  
17  
18 903 also, more notably, by the presence of ions at  $m/z$  790 ( $[C_{50}H_{62}O_8]^+ = 2M+2H$ ), 804  
19  
20 904 ( $[C_{51}H_{64}O_8]^+ = 2M+2H+CH_2$ ) and 818 ( $[C_{52}H_{66}O_8]^+ = 2M+2H+2CH_2$ ) attributed to  
21  
22 905 the presence of dimers.  
23  
24  
25  
26  
27 906

28  
29 907 The confirmation of twelve different carotenoids used as food colorants was achieved  
30  
31 908 using positive atmospheric pressure chemical ionization (APCI)+ LC-MS (Breithaupt,  
32  
33 909 (2004). The  $[M+H]^+$  ions were monitored for norbixin and bixin at  $m/z$  381 and 395  
34  
35 910 respectively. Based on the presence of at least one carboxyl group, APCI  
36  
37 911 measurements in the negative mode were also carried out on bixin and norbixin but no  
38  
39 912 significant enhancement in sensitivity was observed. A similar approach has been  
40  
41 913 used for the analysis of water-soluble annatto extracts in both positive and negative  
42  
43 914 electrospray detection modes (Galindo-Cuspinera and Rankin, 2005). ES- detection  
44  
45 915 mode showed a major peak at  $m/z$  379 corresponding to  $[M-H]^-$  for norbixin, whereas  
46  
47 916 the major peak at  $m/z$  381 was found using ES+ mode. An ion at  $m/z$  117 was  
48  
49 917 identified in in the ES- spectrum of 9'-*cis*-norbixin but not in the spectrum of the  
50  
51 918 *trans*- isomer. Conversely, the *trans*- isomer showed an ion at  $m/z$  111.1 in ES+ mode  
52  
53 919 that was not present on the spectrum of 9'-*cis*-isomer. This was thought to be due to  
54  
55 920 differences in fragmentation patterns determined by stereochemical configuration.  
56  
57  
58  
59  
60



1  
2  
3 921 More recently, it has been shown that HPLC-PDA in combination with ion-trap  
4  
5 922 electrospray mass spectrometric confirmatory analysis can be used to identify and  
6  
7  
8 923 measure norbixin and bixin in meat products using precursor ions at  $m/z$  379 and 395  
9  
10 924 respectively and monitoring characteristic product ions at  $m/z$  253,291,310 and 335  
11  
12 925 (norbixin) and  $m/z$  317, 335, 345, 363 and 377 (bixin) (Noppe *et al.*, 2009).  
13  
14

926

927 *Nuclear magnetic resonance (NMR) spectroscopy*

17  
18 927  
19 928 A comprehensive review on the use of NMR spectroscopy in the study of carotenoids  
20  
21 929 in general is given by Englert (1995), in which a detailed treatise on the experimental  
22  
23 930 aspects, chemical shifts of end groups, chemical shifts and spin couplings,  
24  
25 931 stereoisomerization, and simple and multidimensional experiments are given for  $^1\text{H}$   
26  
27 932 and  $^{13}\text{C}$  nuclei.  
28  
29  
30  
31

933

32  
33 934 The earliest published use of NMR in the study of bixin stereochemistry used low  
34  
35 935 resolution (40 MHz) instrumentation to assign  $^1\text{H}$  frequencies and deduce that the *cis*-  
36  
37 936 bond of the methyl analogue of 'natural or  $\alpha$ -' bixin was in the 9'- (equivalent)  
38  
39 937 position (Barber *et al.*, 1961). The high frequency shift of the proton assigned to H-8'  
40  
41 938 was attributed to deshielding by the 11'-12' alkene bond when compared to the *trans*-  
42  
43 939 (or  $\beta$ -) isomer, which was confirmed via synthesis and more detailed structural  
44  
45 940 assignments (Pattenden, Way and Weedon, 1970). Fourier transform (FT) NMR was  
46  
47 941 used later to assign the  $^{13}\text{C}$  spectra of methyl *cis*- and *trans*-bixin using deuterated  
48  
49 942 compounds, however no experimental details were given and assignments were partly  
50  
51 943 derived from spectra of carotenoids with similar structural characteristics (Moss,  
52  
53 944 1976). The  $^1\text{H}$  FT-NMR spectrum of *cis*-bixin and *cis*-methyl bixin at 250Mhz has  
54  
55 945 been reported but is limited to assignment of the terminal acrylate moieties (Jondiko  
56  
57  
58  
59  
60

1  
2  
3 946 and Pattenden (1989). Proton NMR at 250MHz was used to confirm the structures of  
4  
5 947 purified *trans*- and 9'-*cis*- bixin, where the chemical shifts and coupling constants  
6  
7 948 associated with the change in stereochemistry were consistent with those reported  
8  
9 949 previously (Barber *et al.*, 1961) but afforded much higher resolution (Scotter *et al.*,  
10  
11 950 1994). A similar approach was used to confirm the structure of the principal thermal  
12  
13 951 degradation product of bixin as *trans*-4,8-dimethyltetradeca-hexaenedioic acid  
14  
15 952 monomethyl ester or C<sub>17</sub> (Scotter, 1995). The structure of a minor apocarotenoid  
16  
17 953 isolated from *Bixa orellana* was confirmed as methyl 9'-Z-apo-6'-lycopenate using  
18  
19 954 proton NMR at 500 MHz (Mercadante *et al.*, 1996) and a similar approach used to  
20  
21 955 identify apocarotenoids not previously found in annatto (Mercadante, Steck and  
22  
23 956 Pfander, 1997b; 1999). NMR (300 MHz <sup>1</sup>H) was used alongside TLC and HPLC in  
24  
25 957 the bioautographic detection of antimicrobial compounds in water-soluble annatto  
26  
27 958 extracts where peak assignments were reported to be consistent with previous reports  
28  
29 959 (Galindo-Cuspinera and Rankin, 2005).  
30  
31 960  
32  
33 961 The most comprehensive study to date on the determination of the structure of the  
34  
35 962 bixin family of apocarotenoids is by Kelly *et al.* (1996), who utilised a combination of  
36  
37 963 1D and 2D NMR techniques in conjunction with mass spectrometry and X-ray  
38  
39 964 diffraction analysis. Chemical shift, coupling constants and <sup>1</sup>H correlation data were  
40  
41 965 examined alongside the ion abundances and intensity ratios from standard electron  
42  
43 966 impact (EI+) and fast atom bombardment (FAB+) MS spectra, and bond  
44  
45 967 measurement, cell dimension and degree of hydrogen bonding from X-ray diffraction  
46  
47 968 data to elucidate and compare the crystal structures of the *cis*- and *trans*- isomers of  
48  
49 969 bixin and methyl bixin.  
50  
51 970  
52  
53  
54  
55  
56  
57  
58  
59  
60

971 *Other analytical techniques*

972 Notwithstanding where specific techniques have been discussed elsewhere in this  
973 review, there are several less widely known techniques that have been used in the  
974 study of annatto either alone or in conjunction with complementary techniques. These  
975 include infra-red spectroscopy, where the characteristic strong absorption due to the  
976 C=O stretching frequency between 1740 and 1700 cm<sup>-1</sup> and the complex bands in the  
977 1300-1050 cm<sup>-1</sup> region due to C-O single bond characteristic of esters and carboxylic  
978 acids has been used (Lunde and Zechmeister, 1954; Reith and Gielen, 1971; Chao *et*  
979 *al.*, 1991; Bernard and Grosjean, 1995). Photoacoustic spectrometry in the UV, VIS  
980 and IR regions has been used for the qualitative and quantitative analysis of annatto in  
981 commercial seasoning products (Haas and Vinha, 1995) and more recently in the  
982 determination of the triplet state energy of bixin (Rios *et al.*, 2007). X-ray  
983 photoelectron spectroscopy was used by Felicissimo *et al.* (2004) to ascertain the  
984 major carotenoid composition of *Bixa orellana* seeds and X-ray diffraction in  
985 conjunction with NMR and mass spectrometry has been used to determine of the  
986 structure of the bixin family of apocarotenoids (Kelly *et al.*, 1996).

987

988 *Analysis of foods*

989 Prior to 1970, there were very few published methods for the extraction of annatto  
990 from foods. The qualitative and quantitative analytical aspects of annatto extraction  
991 methods published prior to 1976 have been reviewed briefly (Aparnathi and Sharma,  
992 1991). These relatively simple methods generally involve extraction with solvent (e.g.  
993 chloroform, benzene, petroleum spirit or ether) with or without some form of sample  
994 pre-treatment such as protein precipitation, washing and adsorption onto an inert

1  
2  
3 995 substance. The foodstuffs analysed by these methods largely comprise dairy products,  
4  
5 996 which reflects the relatively narrow scope of annatto usage at that time.  
6  
7

8 997

9  
10 998 Annatto has been extracted from whey solids with dilute ammonium hydroxide where  
11  
12 999 proteins were precipitated by the addition of ethanol and phosphate buffer  
13

14  
15 1000 (Hammond, Chang and Reinhold, 1973), and from meats (McNeal, 1976). Annatto  
16

17 1001 may be analysed in milk and ice-cream by precipitation with boiling acetic acid and  
18

19  
20 1002 extraction of the whey with diethyl ether, and the colour extracted from macaroni and  
21

22 1003 noodles with 80% ethanol followed by back-extraction into diethyl ether under  
23

24 1004 alkaline conditions (AOAC, 1980).  
25

26  
27 1005

28  
29 1006 Rapid methods for the extraction of annatto from foods have been described where  
30

31 1007 drinks and syrups were dissolved in water, acidified with acetic acid and annatto was  
32

33  
34 1008 partitioned into diethyl ether (Corradi and Micheli, (1981). Products with a high fat  
35

36 1009 content *e.g.* butter and margarine, were dissolved in petroleum spirit and annatto was  
37

38 1010 partitioned into aqueous ammoniacal ethanol. Three extractions were required for  
39

40  
41 1011 quantitative extraction of the colour. The aqueous extracts were acidified with acetic  
42

43 1012 acid and back-extracted with diethyl ether. For foods containing fat and protein *e.g.*  
44

45 1013 yoghurt, cheese and pastries, samples were ground with sand and aqueous  
46

47  
48 1014 ammoniacal ethanol. The mixture was transferred to a centrifuge tube and the fat was  
49

50 1015 removed by agitation with petroleum spirit, centrifugation and siphoning off the  
51

52  
53 1016 petroleum spirit phase. The aqueous ammoniacal phase was retained, acidified with  
54

55 1017 acetic acid and the annatto partitioned into diethyl ether.  
56

57  
58 1018  
59  
60

1  
2  
3 1019 Methods for the extraction and determination of annatto in margarine, cheese and  
4  
5 1020 boiled sweets have been investigated using techniques similar to those described  
6  
7  
8 1021 previously, with modifications to enable measurement by spectrophotometry and  
9  
10 1022 HPLC (Smith *et al.*, 1983). Margarine samples were saponified to separate fat and to  
11  
12 1023 convert any bixin to norbixin, thereby facilitating its extraction into aqueous media  
13  
14 1024 and subsequent purification. However the reported HPLC conditions gave poor peak  
15  
16 1025 shapes and insufficient resolution. A method for the determination of annatto in  
17  
18 1026 cheese in which a simple acetone extraction was used, followed by concentration by  
19  
20 1027 rotary evaporation has been described (Luf and Brandl, 1988). Spectrophotometric  
21  
22 1028 (derivative) and HPLC techniques were used to quantify annatto in the presence of  
23  
24 1029 other carotenoids, based on the procedure described for the analysis of certain baked  
25  
26 1030 goods. However, the *cis*- and *trans*-isomers of bixin and norbixin were not identified  
27  
28 1031 separately under the stated conditions.  
29  
30  
31  
32  
33  
34 1032  
35  
36 1033 More recently, other workers have developed refined methods for the extraction of  
37  
38 1034 annatto from high-fat foods, dairy products and candy utilising solvent pre-extraction  
39  
40 1035 of fat and extraction of annatto into ethanolic aqueous ammonia (Lancaster and  
41  
42 1036 Lawrence, 1995) and to separate mixtures of bixin and norbixin from carminic acid in  
43  
44 1037 fruit beverages, yoghurt and candies (Lancaster and Lawrence, 1996). HPLC was  
45  
46 1038 used to measure both the *cis*- and *trans*-isomers of bixin and norbixin but no  
47  
48 1039 significant improvements in peak resolution were demonstrated compared to those  
49  
50 1040 reported previously (Smith *et al.*, 1983), and impure reference materials were used for  
51  
52 1041 calibration. Recovery of norbixin from spiked cheese samples was reported to average  
53  
54 1042 93% over the range 1 to 110 mg/kg, and the recovery of bixin from spiked wafers also  
55  
56  
57  
58  
59  
60

1  
2  
3 1043 averaged 93% over the range 0.1 to 445 mg/kg. The recovery of norbixin from  
4  
5 1044 laboratory-prepared hard candies averaged 88%.  
6  
7  
8 1045  
9  
10 1046 TLC and HPLC were used to determine bixin and other carotenoid colours in products  
11  
12 1047 derived from red pepper (Mínguez-Mosquera, Hornero-Méndez and Garrido-  
13  
14 1048 Fernández, 1995). A simple acetone extraction was used followed by partition with  
15  
16 1049 ether and sodium chloride solution and alkaline saponification. Back extraction with  
17  
18 1050 ether following acidification of the saponifying medium was necessary to recover the  
19  
20 1051 annatto colour (as norbixin). While good chromatographic separation of the  
21  
22 1052 carotenoids was obtained, no distinction between norbixin isomers was made.  
23  
24 1053 However, the method demonstrated the capability of detecting of colours added  
25  
26 1054 fraudulently to intensify the natural colour of paprika paste.  
27  
28  
29 1055  
30  
31 1056 Whilst remaining an uncommon analytical technique in food laboratories,  
32  
33 1057 photoacoustic spectrometry (PAS) has been used for the analysis of annatto products  
34  
35 1058 (Haas and Vinha, 1995). The method is limited to semi quantitative ( $\pm 1\%$  'annatto  
36  
37 1059 content') and qualitative analysis of commercial seasonings comprising mixtures of  
38  
39 1060 corn meal and powdered annatto seeds or annatto extract known as '*Colorifico du*  
40  
41 1061 *Urucum*'. The particle size of the samples has a strong influence on the amplitude of  
42  
43 1062 the PAS signal and therefore requires close control.  
44  
45  
46 1063  
47  
48 1064 Based on the methods described previously (Scotter *et al.*, 1994; Lancaster and  
49  
50 1065 Lawrence, 1995), HPLC and spectrophotometric methods have been developed for  
51  
52 1066 the simple and rapid determination of annatto in cheese and milk products (Bareth,  
53  
54 1067 Strohmar and Kitzelmann, 2002). Solid phase extraction (SPE) on amino phase was  
55  
56  
57  
58  
59  
60

1  
2  
3 1068 used to separate annatto components from fat and  $\beta$ -carotene. The choice of end  
4  
5  
6 1069 method was determined by the presence of other colouring materials i.e. curcumin or  
7  
8 1070  $\beta$ -apo-8'-carotenal but other food colours and emulsifiers did not affect the analysis.  
9  
10  
11 1071 The recovery of annatto colouring spiked into cheese, processed cheese, butter and  
12  
13 1072 ice-cream ranged between 80 and 100%. Nine samples of cheese were analysed in  
14  
15 1073 which norbixin was found in the range  $<0.15$  to  $11.89 \text{ mg.kg}^{-1}$ , whereas no bixin was  
16  
17 1074 detected ( $>0.15 \text{ mg.kg}^{-1}$ ).  
18  
19  
20 1075  
21  
22 1076 The methods described by Scotter *et al.* (1994 and 1998), Lancaster and Lawrence  
23  
24 1077 (1995) and Navaz Diaz and Peinado (1992) were further developed and consolidated  
25  
26 1078 to encompass a wide range of food commodities (Scotter *et al.*, 2002). Specific  
27  
28 1079 solvent extraction regimes were developed for specific sample matrices, with HPLC-  
29  
30 1080 PDA used for spectral confirmation and measurement of the main isomers of bixin  
31  
32 1081 and norbixin. The different extraction regimes are summarized in Table 4.  
33  
34  
35  
36 1082  
37  
38  
39 1083 [Insert Table 4 about here]  
40  
41 1084  
42  
43  
44 1085 With the exception of regime 5, samples were extracted essentially using  
45  
46 1086 ethanol:water:ammonia solution with or without a hexane partition to remove excess  
47  
48 1087 lipid. After centrifugation in the presence of Celite filter aid, the annatto colour was  
49  
50 1088 partitioned into chloroform:acetic acid solution, centrifuged and the solvent removed  
51  
52 1089 using vacuum-assisted rotary evaporation. To minimise analyte losses via oxidation, a  
53  
54 1090 0.1% solution of butylated hydroxyl toluene (BHT) was added. For regime 5 matrices,  
55  
56 1091 samples were mixed with Celite in the presence of dilute hydrochloric acid and  
57  
58 1092 extracted using a biphasic solvent system comprising hexane (to remove excess lipid)  
59  
60

1  
2  
3 1093 and acetonitrile, which was then concentrated using vacuum-assisted rotary  
4  
5 1094 evaporation.  
6  
7  
8 1095  
9  
10 1096 Using this method, comprehensive quantitative and qualitative data on 165 composite  
11  
12 1097 and 2 single food samples covering a wide range of foods at levels above the  
13  
14 1098 analytical reporting limit of 0.1 mg.kg<sup>-1</sup> were obtained. Quantitative results were  
15  
16 1099 given for those annatto colouring components for which reference standards were  
17  
18 1100 available (9'-*cis*-bixin, *trans*-bixin and 9'-*cis*-norbixin), whereas semi-quantitative  
19  
20 1101 results were given for other bixin and norbixin isomers. The method was single-  
21  
22 1102 laboratory validated by the repeat (n = 4 to 9) analysis of 12 different sample types of  
23  
24 1103 food commodity covering the permitted range of annatto content, spiked with annatto  
25  
26 1104 at levels of between 1.7 and 27.7 mg.kg<sup>-1</sup> and by the analysis of in-house reference  
27  
28 1105 matrices. Mean recoveries of between 61 and 96% were obtained from foods spiked  
29  
30 1106 with annatto.  
31  
32 1107  
33  
34 1108 Using response surface methodology to establish optimum conditions, a method for  
35  
36 1109 the determination of annatto colour in extruded corn snack products has been  
37  
38 1110 developed that exhibits improved accuracy and precision compared to the method  
39  
40 1111 described by Scotter *et al.* (2002) (Rios and Mercadante, 2004). However, pre-  
41  
42 1112 treatment of the samples with  $\alpha$ -amylase was necessary to remove starch and a total  
43  
44 1113 of 8 solvent extractions with ethyl acetate were required for complete extraction of the  
45  
46 1114 annatto colour. Lipids were removed using alkaline saponification therefore all of the  
47  
48 1115 bixin present was hydrolysed to norbixin and determined as such by HPLC.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58 1116  
59  
60



1  
2  
3 1117 Accelerated solvent extraction has been compared with manual solvent extraction to  
4  
5 1118 determine several food colouring carotenoids including bixin and norbixin in  
6  
7  
8 1119 processed foods (Breithaupt, 2004). Reverse-phase HPLC with a C<sub>30</sub> column  
9  
10 1120 successfully separated bixin and norbixin from 7 other carotenoids but the *cis*- and  
11  
12 1121 *trans*- isomers were not distinguishable. Due to its ostensibly higher polarity, lower  
13  
14 1122 recoveries of norbixin were reported for accelerated extraction ( $67 \pm 1.0 \text{ mg.kg}^{-1}$ )  
15  
16 1123 compared to manual extraction ( $88.7 \pm 6.2 \text{ mg.kg}^{-1}$ ). However, a similar difference in  
17  
18 1124 recoveries was reported for less polar bixin ( $91.0 \pm 2.7$  and  $98.0 \pm 1.7 \text{ mg.kg}^{-1}$   
19  
20  
21 1125 respectively) although bixin recovery was higher than norbixin with improved  
22  
23 1126 precision. The limit of quantitation for bixin and norbixin was in the range 0.53 – 0.79  
24  
25 1127  $\text{mg.kg}^{-1}$  for pudding mix and cereals. More recently, a method for the determination  
26  
27 1128 of norbixin and bixin in meat products using HPLC-PDA and LC-MS<sup>n</sup> that gives  
28  
29 1129 recoveries of between 99 and 102% and a limit of quantitation of 0.5 mg/kg. (Noppe  
30  
31 1130 *et al.*, 2009).

1131

1132 *Annatto as an illegal food dye*

1133 The illegal use of annatto to colour milk goes back as far as the early 20<sup>th</sup> century  
1134 where it was reported by UK Public Analyst laboratories (Richards, 1923;  
1135 Collingwood Williams, 1925). Amongst other specific food commodities, annatto is  
1136 currently permitted in the EU for the colouring of certain margarines and cheeses but  
1137 is not permitted for the colouring of milk cream or butter (EC 1994 as amended).  
1138 Moreover, while annatto is permitted for use in food commodities such as savoury  
1139 snack products, coated nuts, extruded products and flavoured breakfast cereals, it is  
1140 not permitted for use in spices. However, amongst other non-permitted dyes bixin was  
1141 detected in 18 of 893 samples of spices, sauces and oils by UK enforcement

1  
2  
3 1142 laboratories during 2005-2006 as part of the UK Imported Food Programme (Food  
4  
5 1143 Standards Agency, 2006). This has led directly to a need for analytical methods  
6  
7  
8 1144 capable of detecting very low levels of annatto in food ingredients and commodities in  
9  
10 1145 which it is not permitted, driven not only by the enforcement of regulations on a  
11  
12 1146 national scale (disseminated through the EU Rapid Alert System; EU 2008) but also  
13  
14 1147 by the need for the food manufacturing industry to ensure compliance, especially in a  
15  
16 1148 proactive manner and through the adoption of a 'zero tolerance' approach as applied  
17  
18 1149 to the monitoring of illegal dyes such as the Sudan Red group. Established HPLC  
19  
20 1150 methods capable of detecting bixin or norbixin at ca. 0.1 mg/kg in samples using UV-  
21  
22 1151 VIS or diode-array technology are not sufficiently sensitive. LC-MS/MS methodology  
23  
24 1152 is the obvious candidate but sufficiently detailed methods in peer-reviewed  
25  
26 1153 publications have not been forthcoming to date. Nevertheless, it is generally  
27  
28 1154 considered amongst analytical chemists working in this area that LC-MS/MS is  
29  
30 1155 capable of detecting bixin at ca. 0.01 mg/kg in certain commodities, but this is heavily  
31  
32 1156 dependent upon the degree of signal suppression caused by matrix effects. This can  
33  
34 1157 give rise to false negative results using a screening approach, which in turn identifies  
35  
36 1158 a need for suitable extract clean up regimes, and guarding against ion suppression by  
37  
38 1159 using the method of standard addition  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 1161 *Future aspects*

49  
50 1162 There is a clear requirement in the future for the development and validation of highly  
51  
52 1163 sensitive methods of analysis for annatto in food commodities and other food  
53  
54 1164 ingredients, driven by the need to ensure compliance with food quality regulations and  
55  
56 1165 especially in the light of the pursuit of suitable alternatives to synthetic food colours.  
57  
58  
59 1166 An in-depth understanding of the chemistry and stability of annatto is therefore  
60

1  
2  
3 1167 requisite and brings clear benefits to the production of annatto, and to the formulation  
4  
5 1168 and application of food colouring to a wide range of food commodities. Greater  
6  
7  
8 1169 understanding of the processes of degradation may also benefit studies in the areas of  
9  
10 1170 food safety, particularly in risk assessment, and biomarkers of exposure such as  
11  
12 1171 circulating (plasma) levels of norbixin. Here, complementary analytical techniques  
13  
14  
15 1172 such as HPLC-PDA, LC-MS/MS and NMR will play a vital role in the detection,  
16  
17 1173 confirmation and measurement of comparatively low levels of bixin and norbixin  
18  
19  
20 1174 isomers.  
21  
22 1175  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1176 **REFERENCES**

- 1177 Aparnathi KD, Sharma RS. 1991. Annatto colour for food: A review. Indian Food  
1178 Packer: March-April, 13-27.
- 1179
- 1180 AOAC 1980. Association of Official Analytical Chemists Official Methods of  
1181 Analysis, 11th Edition. Washington D.C: AOAC.
- 1182
- 1183 Balaswamy K, Prabhakara Rao, PGP, Satyanarayana A, Rao DG. 2006. Stability of  
1184 bixin in annatto oleoresin and dye powder during storage. LWT-Food Science and  
1185 Technology. 39: 952-956.
- 1186
- 1187 Barber MS., Hardisson A, Jackman LM, Weedon BCL. 1961. Studies in nuclear  
1188 magnetic resonance. Part IV. Stereochemistry of the bixins. Journal of the Chemical  
1189 Society: 1625-1630.
- 1190
- 1191 Barbosa MIMJ, Borsarelli CD, Mercadante, AZ. 2005. The light stability of spray-  
1192 fried bixin encapsulated with different edible polysaccharide preparations. Food  
1193 Research International. 38: 989-994.
- 1194
- 1195 Bareth A, Strohmar W, Kitzelmann E. 2002. HPLC and spectrophotometric  
1196 determination of annatto in cheese. European Food Research and Technology. 215:  
1197 359-364.
- 1198

- 1  
2  
3 1199 Bernard K, Grosjean M. 1995. Infrared spectroscopy. In: Britton G, Liaaen-Jensen, S,  
4  
5 1200 Pfander, editors. Carotenoids, Volume 1B: Spectroscopy. Basel: Birkhäuser-Verlag.  
6  
7  
8 1201 p117-134.  
9  
10 1202  
11  
12 1203 Berset C, Marty C. 1986. Potential use of annatto in extrusion cooking. Lebensmittel-  
13  
14 1204 Wissenschaft und Technologie. 19: 126-131.  
15  
16 1205  
17  
18 1206 Bittencourt C, Felicissimo MP, Pireaux J-J, Houssiau L. 2005. ToF-SIMS  
19  
20 1207 characterization of thermal modifications of bixin from *Bixa orellana* fruit. Journal of  
21  
22 1208 Agricultural and Food Chemistry. 53: 6195-6200.  
23  
24  
25 1209  
26  
27 1210 Breithaupt D. 2004. Simultaneous HPLC determination of carotenoids used as food  
28  
29 1211 coloring additives: applicability of accelerated solvent extraction. Food Chemistry.  
30  
31 1212 86: 449-456.  
32  
33 1213  
34  
35 1214 Britton G. 1995a. Chapter 2: UV/Visible spectroscopy. In: Britton G, Liaaen-Jensen,  
36  
37 1215 S, Pfander, editors. Carotenoids, Volume 1B: Spectroscopy. Basel: Birkhäuser-  
38  
39 1216 Verlag. p 13-62.  
40  
41 1217  
42  
43 1218 Britton G. 1995b. Structure and properties of carotenoids in relation to function.  
44  
45 1219 Federation of American Societies for Experimental Biology Journal. 9: 1551-1558.  
46  
47 1220  
48  
49 1221 Budzikiewicz, H. 1974. Mass spectra of carotenoids – labelling studies. Advances in  
50  
51 1222 Mass Spectrometry. 6: 155-165.  
52  
53  
54  
55  
56  
57  
58  
59  
60 1223

- 1  
2  
3 1224 Carvalho PRN, Sarantópoulos CIGJ, Shirose I, da Silva MG. .1993. Study of the  
4  
5 1225 shelf-life of the colour (bixin) from annatto seeds (*Bixa orellana*, L.). *Coltanea I*  
6  
7 1226 *Technologia*. 23: 98-104.  
8  
9  
10 1227  
11  
12 1228 Chao RR, Mulvaney SJ, Sansom DR, Hsieh F-H, Tempesta MS. .1991. Supercritical  
13  
14 1229 CO<sub>2</sub> extraction of annatto (*Bixa orellana*) pigments and some characteristics of the  
15  
16 1230 color extracts. *Journal of Food Science*. 56: 80-83.  
17  
18  
19 1231  
20  
21 1232 Chatani Y, Adachi T. 1988. Determination of norbixin in foods by HPLC after  
22  
23 1233 protease digestion. *Kyoto-fu Eisei Kenkyusho Nenpo*. 33: 36-40.  
24  
25  
26 1234  
27  
28 1235 Collins P. 1992. The role of annatto in food. *Food Ingredients and Processing*  
29  
30 1236 *International*. February: 23-27.  
31  
32  
33 1237  
34  
35 1238 Collingwood Williams, W. 1925. Annual report of the County Analyst for the year  
36  
37 1239 1925. *Notes from the Reports of Public Analysts*: 28-29.  
38  
39  
40 1240  
41  
42 1241 Corradi C, Micheli G. 1981. A rapid research and identification method for the natural  
43  
44 1242 colorant E160b bixin, norbixin (Oriana, annatto) in food products. *Industrie*  
45  
46 1243 *Alimentari*. May: 372-375.  
47  
48  
49 1244  
50  
51 1245 Dale J. 1954. Empirical relationships of the minor bands in the absorption spectra of  
52  
53 1246 polyenes. *Acta Chemica Scandinavica*. 8: 1235-1256.  
54  
55  
56 1247  
57  
58  
59  
60

- 1  
2  
3 1248 EC 1994. European Parliament and Council Directive 94/36/EC, of 30 June 1994 on  
4  
5 1249 colours for use in foodstuffs. Official Journal of the European Communities, No. L  
6  
7  
8 1250 237. 10 September 1994: 13-29.  
9  
10 1251  
11  
12 1252 EC 1995. Commission Directive 95/45/EC, of 26 July 1995 laying down specific  
13  
14  
15 1253 criteria of purity concerning colours for use in foodstuffs. Official Journal of the  
16  
17 1254 European Communities, No. L226. 22 September 1995: 1-44.  
18  
19  
20 1255  
21  
22 1256 EC 2006. Commission of the European Communities. Proposal for a Regulation of  
23  
24 1257 the European Parliament and of the Council establishing a common authorisation  
25  
26  
27 1258 procedure for food additives, food enzymes and food flavourings. Brussels 28.7.2006  
28  
29 1259 COM(2006) 423 Final. 2006/0143 (COD).  
30  
31  
32 1260  
33  
34 1261 Ellis Richards PA. 1923. Annual report of the County Analyst for the year 1923.  
35  
36 1262 Notes from the Reports of Public Analysts: 428.  
37  
38  
39 1263  
40  
41 1264 Englert G. 1995. NMR spectroscopy. In: Britton G, Liaaen-Jensen, S, Pfander,  
42  
43 1265 editors. Carotenoids, Volume 1B: Spectroscopy. Basel: Birkhäuser-Verlag. p 147-260.  
44  
45  
46 1266  
47  
48 1267 Enzell, CR., and Back, S., (1995), Mass Spectrometry. In: Britton G, Liaaen-Jensen,  
49  
50 1268 S, Pfander, editors. Carotenoids, Volume 1B: Spectroscopy. Basel: Birkhäuser-  
51  
52  
53 1269 Verlag. p 261-320.  
54  
55 1270  
56  
57  
58  
59  
60

- 1  
2  
3 1271 Enzell C., Wahlberg I. 1980. Carotenoids. In: Waller GR, Dermer OC, editors.  
4  
5 1272 Biomedical Applications of Mass Spectrometry – First supplementary volume.  
6  
7  
8 1273 Chichester: Wiley. p407-438.  
9  
10 1274  
11  
12 1275 EU 2008. European Union rapid Alert System [internet]. Available from:  
13  
14 1276 [http://ec.europa.eu/food/food/rapidalert/index\\_en.htm](http://ec.europa.eu/food/food/rapidalert/index_en.htm). Accessed 2008 January 09.  
15  
16  
17 1277  
18  
19 1278 FAO/WHO. 1981. Specifications for identity and purity of food additives (carrier  
20  
21 1279 solvents, emulsifiers and stabilizers, enzyme preparations, flavouring agents, food  
22  
23 1280 colours, sweetening agents, and other food additives). FAO Food and Nutrition Paper,  
24  
25 1281 No. 19.  
26  
27  
28 1282  
29  
30 1283 FAO/WHO. 1996. Compendium of food additive specifications, addendum 4. FAO  
31  
32 1284 Food and Nutrition Paper, No. 52, Add. 4.  
33  
34 1285  
35  
36 1286 Felicissimo MP, Bittencourt C, Houssiau L, Pireaux J.J. 2004. Time-of-flight  
37  
38 1287 secondary ion mass spectrometry and x-ray photoelectron spectroscopy analyses of  
39  
40 1288 Bixa orellana seeds. Journal of Agricultural and Food Chemistry. 52: 1810-1814.  
41  
42  
43 1289  
44  
45 1290 Ferreira VLP, Teixeira Neto RO, Moura SCSR, De Silva MS. 1999. Kinetics of  
46  
47 1291 colour degradation of water soluble commercial annatto solutions under thermal  
48  
49 1292 treatments. Ciencia-e-Tecnologia-de-Alimentos. 19: 37-42.  
50  
51  
52 1293  
53  
54 1294 FAO. 1990. Food and Agriculture Organization, Food and Nutrition Paper 49.  
55  
56  
57  
58  
59  
60



- 1  
2  
3 1295 Galindo-Cuspinera V, Rankin SA. 2005. Bioautography and chemical characterization  
4  
5  
6 1296 of antimicrobial compound(s) in commercial water-soluble annatto extracts. Journal  
7  
8 1297 of Agricultural and Food Chemistry. 53: 2524-2529.  
9  
10 1298  
11  
12 1299 Food Standards Agency [internet]. 2006. Study on illegal dyes in imported foods. 12th  
13  
14 1300 December. Available from: [www.food.gov.uk](http://www.food.gov.uk). Accessed 8 January 2009.  
15  
16  
17 1301  
18  
19 1302 Francis BJ. 1995. The separation of annatto pigments by thin-layer chromatography  
20  
21 1303 with special reference to the use of analytical-grade reagents. Analyst. 90: 374.  
22  
23  
24 1304  
25  
26 1305 Gloria MBA, Vale SR Bobbio PA. 1995. Effect of water activity on the stability of  
27  
28 1306 bixin in annatto extract-microcrystalline cellulose model system. Food Chemistry. 52:  
29  
30 1307 389-391.  
31  
32  
33 1308  
34  
35 1309 Guaratini T, Vessechi RL, Lavarda FC, Maia Campos, PMBG, Naal ., Gates PH,  
36  
37 1310 Lopes NP. 2004. New chemical evidence for the ability to generate radical molecular  
38  
39 1311 ions of polyenes from ESI and HR-MALDI mass spectrometry. Analyst. 129: 1223-  
40  
41 1312 1226.  
42  
43  
44 1313  
45  
46 1314 Haberli A, Pfander H. 1999. Synthesis of bixin and three minor carotenoids from  
47  
48 1315 annatto (*Bixa orellana*). Helvetica Chimica Acta. 82: 696-706.  
49  
50  
51 1316  
52  
53 1317 Haila KM, Lievonen SM, Heinonen MI. 1996. Effects of lutein, lycopene, annatto and  
54  
55 1318 g-tocopherol on autoxidation of triglycerides. Journal of Agricultural and Food  
56  
57 1319 Chemistry. 44: 2096-2100.  
58  
59  
60

- 1  
2  
3 1320  
4  
5  
6 1321 Hammond EG, Chang J, Reinbold G.W. 1975. Colorimetric method for residual  
7  
8 1322 annatto in dry whey. *Journal of Dairy Science*. 58: 1365.  
9  
10 1323  
11  
12 1324 Hass U, Vinha CA. 1995. Qualitative and semiquantitative analysis of annatto and its  
13  
14 1325 content in food additives by photoacoustic spectrometry. *Analyst*. 120: 351-354.  
15  
16 1326  
17  
18 1327 Heiduschka A, Panzer A. 1917. Zur Kenntnis des Bixins. *Berichte der Deutschen*  
19  
20 1328 *Chemischen Gesellschaft*. 50: 546-554 and 1525.  
21  
22 1329  
23  
24 1330 Henry BS. 1992. Natural food colours. In: Hendry GAF, Houghton JD, editors.  
25  
26 1331 *Natural Food Colorants*. London: Blackie, p39-78.  
27  
28 1332  
29  
30 1333 Herzig J, Faltis F. 1923. Zur Kenntnis des Bixins. *Justus Liebigs Annalen der Chemie*.  
31  
32 1334 431: 40-70.  
33  
34 1335  
35  
36 1336 Hudson BS, Kohler BE. 1974. Linear polyene electronic structure and spectroscopy.  
37  
38 1337 *Annual Review of Physical Chemistry*. 25: 437-460.  
39  
40 1338  
41  
42 1339 Hudson B., Kohler BE, Schulten K. 1982. Linear polyene electronic-structure and  
43  
44 1340 potential surfaces. *Excited States*. 6: 1-95.  
45  
46 1341  
47  
48 1342 Iversen S, Lam J. 1953. Uber den farbstoff in annatto-butterfarben. *Zeitschrift fur*  
49  
50 1343 *Lebensmittel Untersuchung und Forschung*. 97: 1-7.  
51  
52 1344  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1345 JECFA. 2002. Joint FAO/WHO Expert Committee on Food Additives, 2002, List of  
4  
5  
6 1346 substances scheduled for evaluation and request for data. Rome, 10-12.  
7  
8 1347  
9  
10 1348 Jondiko IJO, Pattenden G. 1989. Terpenoids and an apocarotenoid from seeds of *Bixa*  
11  
12 1349 *orellana*. *Phytochemistry*. 28: 3159-3162.  
13  
14 1350  
15  
16  
17 1351 Jukes DJ. 2008. FoodLaw-Reading, Department of Food Biosciences, University of  
18  
19 1352 Reading [internet]. Available from: <http://www.rdg.ac.uk/foodlaw/main.htm>. Accessed  
20  
21 1353 2008 December 8.  
22  
23 1354  
24  
25  
26 1355 Karrer P, Jucker E. 1950. Chapter V. Cis-trans isomerism of carotenoids. In:  
27  
28 1356 Carotenoids. Basel: Verlag Birkhäuser, pp 38-42.  
29  
30 1357  
31  
32 1358 Karrer P, Helfenstein A, Widmer R, van Itallie TB. 1929. Über Bixin. (XIII.  
33  
34 1359 Mitteilung über Pflanzenfarbstoffe). *Helvetica Chimica Acta*. 12: 741-756.  
35  
36 1360  
37  
38  
39 1361 Karrer P, Benz P, Morf R, Raudnitz H, Stoll M, Takahashi T. 1932. Konstitution des  
40  
41 1362 Safranfarbstoffs Crocetin, Synthese des Perhydro-bixin-äthylesters und Perhydro-  
42  
43 1363 norbixins. Vorläufige Mitteilung. *Helvetica Chimica Acta*. 15: 1218-1399.  
44  
45 1364  
46  
47  
48 1365 Kelly DR, Edwards AA, Parkinson JA, Olovsson G, Trotter J, Jones S, Abdul  
49  
50 1366 Malik KM, Hursthouse MB, Hibbs DE. 1996. NMR, MS and X-ray crystal structure  
51  
52 1367 determination of the bixin family of apocarotenoids. *Journal of Chemical Research*  
53  
54 1368 (M). 2640-2697.  
55  
56 1369  
57  
58  
59  
60

- 1  
2  
3 1370 Kiokias S, Gordon MH. 2003. Antioxidant properties of annatto carotenoids. Food  
4  
5 1371 Chemistry. 83: 523-529.  
6  
7 1372  
8  
9  
10 1373 Kohler BE. 1977. Visual chromophore electronic structure. Biophysics of Structure  
11  
12 1374 and Mechanism. 3: 101-106  
13  
14 1375  
15  
16  
17 1376 Kohler BE. 1995. Chapter 1: Electronic structure of carotenoids. In: Britton G,  
18  
19 1377 Liaaen-Jensen, S, Pfander, editors. Carotenoids, Volume 1B: Spectroscopy. Basel:  
20  
21 1378 Birkhäuser-Verlag. p 1-12.  
22  
23  
24 1379  
25  
26  
27 1380 Kovary K, Louvain TS, Costa e Silva MC, Albano F, Pires BBM, Laranj, GAT, Lage  
28  
29 1381 CLA, Felzenszwalb. 2001. I. Biochemical behaviour of norbixin during in vitro DNA  
30  
31 1382 damage induced by reactive oxygen species. British Journal of Nutrition. 85: 431-440.  
32  
33 1383  
34  
35  
36 1384 Kuhn R, Winterstein A. 1932. Die dihydroverbindung der isomeren bixine und die  
37  
38 1385 elektronen-konfiguration der polyene. Berichte der Deutschen Chemischen  
39  
40 1386 Gesellschaft. 65: 646-651.  
41  
42  
43 1387  
44  
45  
46 1388 Kuhn R, Winterstein A. 1933. Über einen lichtempfindlichen carotin-farbstoff aus  
47  
48 1389 safran. Berichte der Deutschen Chemischen gesellschaft. 66: 209-214.  
49  
50 1390  
51  
52  
53 1391 Lancaster FE, Lawrence JF. 1995. Determination of annatto in high-fat dairy  
54  
55 1392 products, margarine and hard candy by solvent extraction followed by high-  
56  
57 1393 performance liquid chromatography. Food Additives and Contaminants. 12: 9-19.  
58  
59 1394  
60

- 1  
2  
3 1395 Lancaster FE, Lawrence J.F. 1996. High performance liquid chromatographic  
4  
5 1396 separation of carminic acid,  $\alpha$ -bixin, and  $\alpha$ - and  $\beta$ -norbixin, and the determination of  
6  
7 1397 carminic acid in foods. *Journal of Chromatography A*. 732: 394-398.  
8  
9 1398  
10  
11 1399 Lauro GJ. 1991. A primer on natural colours. *Cereal Foods World*: 36: 949-953.  
12  
13 1400  
14  
15 1401 Levy LW, Regalado E, Navarette S, Watkins RH. 1997. Bixin and norbixin in human  
16  
17 1402 plasma: Determination and study of the absorption of a single dose of annatto food  
18  
19 1403 color. *Analyst*. 122: 977-980.  
20  
21 1404  
22  
23 1405 Levy LW, Rivadeneira DM. 2000. Annatto. In Lauro GJ, Francis, FJ, editors. *Natural*  
24  
25 1406 *Food Colorants: Science and Technology*. New York: Marcel Dekker Inc. p115-152.  
26  
27 1407  
28  
29 1408 Lewis GN, Calvin M. 1939. The color of organic substances. *Chemistry Reviews*. 25:  
30  
31 1409 273-328.  
32  
33 1410  
34  
35 1411 Luf W, Brandl E. 1988. The detection of the annatto dyestuff norbixin / bixin in  
36  
37 1412 cheese using derivative spectroscopy and high-performance liquid chromatography.  
38  
39 1413 *Zeitschrift für Lebensmittel-Untersuchung und –Forschung*. 186: 327-332.  
40  
41 1414  
42  
43 1415 Lunde K, Zechmeister L. 1954. Infrared spectra and cis-trans configurations of some  
44  
45 1416 carotenoid pigments. *Journal of the American Chemical Society*. 77: 1647-1653.  
46  
47 1417  
48  
49 1418 Lyng SM, Passos M, Fontana JD. 2005. Bixin and alpha-cyclodextrin inclusion  
50  
51 1419 complex and stability tests. *Process Biochemistry*. 40: 865-872.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1420  
4  
5  
6 1421 MAFF. 1989. Ministry of Agriculture, Fisheries and Food. Response to comments  
7  
8 1422 received on its Final Report on the Review of the Colouring Matter in Food  
9  
10 1423 Regulations 1973, Fd/FAC/6, Food Advisory Committee. London: HMSO.  
11  
12 1424  
13  
14  
15 1425 MAFF. 1993. Ministry of Agriculture, Fisheries and Food, Food Surveillance Paper  
16  
17 1426 No. 37, 1993. Dietary Intake of Food Additives in the UK: Initial Surveillance.  
18  
19 1427 London: HMSO.  
20  
21  
22 1428  
23  
24 1429 Martinez-Tomé M, Jiménez AM, Ruggieri S, Frega N, Strabbioli R, Murcia MA.  
25  
26 1430 2001. Antioxidant properties of Mediterranean spices compared with common food  
27  
28 1431 additives. Journal of Food Protection. 64: 1412-1419.  
29  
30  
31 1432  
32  
33 1433 McKeown GG. 1961. Paper chromatography of bixin and related compounds. Journal  
34  
35 1434 of the Association of Official Analytical Chemists. 45: 347-351.  
36  
37 1435  
38  
39 1436 McKeown GG. 1963. Composition of oil-soluble food colours. II. Thermal  
40  
41 1437 degradation of bixin. Journal of the Association of Official Analytical Chemists. 46:  
42  
43 1438 790-796.  
44  
45  
46 1439  
47  
48 1440 McKeown GG. 1965. Composition of oil-soluble annatto food colors. III. Structure of  
49  
50 1441 the yellow pigment formed by the thermal degradation of bixin. Journal of the  
51  
52 1442 Association of Official Analytical Chemists. 48: 835-837.  
53  
54  
55 1443  
56  
57  
58  
59  
60

- 1  
2  
3 1444 McKeown GG, Mark E. 1962. The composition of oil-soluble annatto food colors.  
4  
5 1445 Journal of the Association of Official Analytical Chemists. 45: 761-766.  
6  
7  
8 1446  
9  
10 1447 McNeal JE. 1976. Qualitative tests for added colouring matter in meat products.  
11  
12 1448 Journal of the Association of Official Analytical Chemists. 59: 570.  
13  
14 1449  
15  
16  
17 1450 Mercadante AZ. 2001. Composition of carotenoids from annatto. In: Ames, JM,  
18  
19 1451 Hofmann TS, editors. Chemistry and physiology of selected food colorants; ACS  
20  
21 1452 Symposium Series 775: Washington DC: p 92-101.  
22  
23  
24 1453  
25  
26  
27 1454 Mercadante AZ, Steck A, Pfander H. 1997a. Isolation and structure elucidation of  
28  
29 1455 minor carotenoids from annatto (*Bixa orellana*) seeds. *Phytochemistry*. 46: 1379-  
30  
31 1456 1383.  
32  
33  
34 1457  
35  
36 1458 Mercadante AZ, Steck A, Pfander H. 1997b. Isolation and identification of new  
37  
38 1459 apocarotenoids from annatto (*Bixa orellana*) seeds. *Journal of Agricultural and Food*  
39  
40 1460 *Chemistry*. 45: 1050-1054.  
41  
42  
43 1461  
44  
45  
46 1462 Mercadante AZ, Steck A, Rodriguez-Amaya D, Pfander H, Britton G. 1996. Isolation  
47  
48 1463 of methyl 9'Z-apo-6'-lycopenoate from *Bixa orellana*. *Phytochemistry*. 41: 1201-  
49  
50 1464 1203.  
51  
52  
53 1465  
54  
55 1466 Mercadante AZ, Steck A, Pfander, H. 1999. Three minor carotenoids from annatto  
56  
57 1467 (*Bixa orellana*). *Phytochemistry*. 52: 135-139.  
58  
59  
60 1468

- 1  
2  
3 1469 Mínguez-Mosquera MI, Jaren-Galan M. 1995. Kinetics of the decoloring of carotenoid  
4  
5 1470 pigments. *Journal of the Science of Food and Agriculture*. 67: 153-161.  
6  
7 1471  
8  
9 1472 Mínguez-Mosquera MI, Hornero-Méndez D, Garrido-Fernández J. 1995. Detection of  
10  
11 1473 bixin, lycopene, canthaxanthin and  $\beta$ -apo-8'-carotenal in products derived from red  
12  
13 1474 pepper. *Journal of the AOAC International*. 78: 491-496.  
14  
15 1475  
16  
17 1476 Montenegro MA, Rios A de O, Mercadante AZ, Nazareno MA, Borsarelli CD. 2004.  
18  
19 1477 Model studies on the photosensitized isomerisation of bixin. *Journal of Agricultural*  
20  
21 1478 *and Food Chemistry*. 52: 367-373.  
22  
23 1479  
24  
25 1480 Moss GP. 1976. Carbon-13 NMR spectra of carotenoids. *Pure and Applied*  
26  
27 1481 *Chemistry*. 47: 97-102.  
28  
29 1482  
30  
31 1483 Najar SV, Bobbio FO, Bobbio PA. 1988. Effects of light, air, anti-oxidants and pro-  
32  
33 1484 oxidants on annatto extracts (*Bixa orellana*). *Food Chemistry*. 29: 283-289.  
34  
35 1485  
36  
37 1486 Navas Diaz A, Ramos Peinado MC. 1992. Fluorimetric determination of curcumin in  
38  
39 1487 yogurt and mustard. *Journal of Agricultural and Food Chemistry*. 40: 56-59.  
40  
41 1488  
42  
43 1489 Nishizawa M, Chonan T, Sekijo I, Sugii Y. 1983. Studies on the analysis of natural  
44  
45 1490 dyes: Analysis of annatto extract and gardenia yellow dye in foods and natural dye  
46  
47 1491 preparations. *Hokkaidoritsu Eisei Kenkyusho*. 33: 32-34.  
48  
49 1492  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 1493 NNT. 2008. [Internet]. Nordic Working Group on Food Toxicology and Risk  
4  
5 1494 Assessment, Nordic Food Additives Database. Available from: <http://www.norfad.dk>.  
6  
7  
8 1495  
9  
10 1496 Noppe H, Abuin Martinez S, Verheyden K, Van Loco J, Companyó Beltran R, De  
11  
12 1497 Brabander HF. 2009. Determination of bixin and norbixin in meat using liquid  
13  
14 1498 chromatography and photodiode array detection. Food Additives and Contaminants.  
15  
16 1499 26: 17-24.  
17  
18  
19 1500  
20  
21 1501 Pattenden G, Way JE, Weedon BCL. 1970. Carotenoids and related compounds. Part  
22  
23 1502 XXVI. Synthesis of methyl natural bixin. Journal of the Chemical Society. 235-241.  
24  
25  
26 1503  
27  
28 1504 Pettersson A, Jonsson L. 1990. Separation of cis-trans isomers of alpha- and beta-  
29  
30 1505 carotene by adsorption HPLC and identification with diode-array detection. Journal of  
31  
32 1506 Micronutrient Analysis. 8: 23-41.  
33  
34  
35 1507  
36  
37 1508 Prentice-Hernández C, Rusig O. 1999. Effect of light on the stability of a  
38  
39 1509 microencapsulated extract obtained from annatto (*Bixa orellana* L.). Brazilian Journal  
40  
41 1510 of Food Technology. 2: 185-189.  
42  
43  
44 1511  
45  
46 1512 Prentice-Hernández C, Rusig O, Carvalho PRN. 1993. Influence of heating time on  
47  
48 1513 the thermal-degradation of bixin in alkaline extracts of annatto (*Bixa orellana*).  
49  
50 1514 Arquivos de Biologica e Tecnologica. 36: 819-828.  
51  
52  
53 1515  
54  
55 1516 Preston HD, Rickard MD. 1980. Extraction and chemistry of annatto. Food  
56  
57 1517 Chemistry. 5: 47-56.  
58  
59  
60




- 1  
2  
3 1518  
4  
5 1519 Ramamurthy MK, Bhalero VR. 1964. A thin-layer chromatographic method for  
6  
7 identifying annatto and other food colours. *Analyst*. 89: 740-744.  
8  
9 1520  
10 1521  
11  
12 1522 Rao PGP, Jyothirmayi T, Balaswamy K, Satyanarayana A, Rao DG. 2005. Effect of  
13  
14 processing conditions on the stability of annatto (*Bixa orellana* L.) dye incorporated  
15 1523  
16 into some foods. *LWT-Food Science and Technology*. 38: 779-784.  
17 1524  
18  
19 1525  
20  
21 1526 Reith JF, Gielen JW. 1971. Properties of bixin and norbixin and the composition of  
22  
23 annatto extracts. *Journal of Food Science*. 36: 861-864.  
24 1527  
25  
26 1528  
27  
28 1529 Rios ADO, Mercadante AZ. 2004. Novel method for the determination of added  
29  
30 annatto colour in extruded corn snack products. *Food Additives and Contaminants*.  
31 1530  
32 21: 125-133.  
33 1531  
34  
35 1532  
36  
37 1533 Rios ADO, Borsarelli CD, Mercadante AZ. 2005. Thermal degradation kinetics of  
38  
39 bixin in an aqueous model system. *Journal of Agricultural and Food Chemistry*. 53:  
40 1534  
41 2307-2311.  
42 1535  
43  
44 1536  
45  
46 1537 Rios ADO, Mercadante AZ, Borsarelli .D. 2007. Triplet state energy of the  
47  
48 carotenoid bixin determined by photoacoustic calorimetry. *Dyes and Pigments*. 74:  
49 1538  
50 561-565.  
51 1539  
52  
53 1540  
54  
55  
56  
57  
58  
59  
60


- 1  
2  
3 1541 Rouseff RL. 1988. High performance liquid chromatographic separation and spectral  
4  
5 1542 characterisation of the pigments in turmeric and annatto. Journal of Food Science. 53:  
6  
7 1543 1823-1826.  
8  
9 1544  
10  
11 1545 Satyanarayana A, Prabhakara rao PG, Rao DG. 2003. Chemistry, processing and  
12  
13 1546 toxicology of annatto (*Bixa orellana* L.). Journal of Food Science and Technology.  
14  
15 1547 40: 131-141.  
16  
17 1548  
18  
19 1549 Satyanarayana A, Prabhakara rao P, Balaswamy K, Velu V, Rao DG. 2006.  
20  
21 1550 Application of annatto dye formulations in different fruit and vegetable products.  
22  
23 1551 Journal of Food Service. 17: 1-5.  
24  
25 1552  
26  
27 1553 Schiedt K, Liaaen-Jensen S. 1995. Isolation and Analysis. In: Britton G, Liaaen-  
28  
29 1554 Jensen, S, Pfander, editors. Carotenoids Volume 1A, Isolation and Analysis. Basel:  
30  
31 1555 Birkhäuser Verlag. p81-108.  
32  
33 1556  
34  
35 1557 Scott AI. 1964. Chapter 8: Application of spectral data to the investigation of gross  
36  
37 1558 molecular structure. Interpretation of the ultraviolet spectra of natural products.  
38  
39 1559 International series of monographs on organic chemistry. London: Pergamon Press. p  
40  
41 1560 228-312.  
42  
43 1561  
44  
45 1562 Scotter MJ, Thorpe SA, Reynolds SL, Wilson LA, Strutt PR. 1994. Characterization  
46  
47 1563 of the principal colouring components of annatto using high performance liquid  
48  
49 1564 chromatography with photodiode array detection, Food Additives and Contaminants.  
50  
51 1565 11: 301-315.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1566  
4  
5  
6 1567 Scotter MJ. 1995. Characterisation of the coloured thermal degradation products of  
7  
8 1568 bixin from annatto and a revised mechanism for their formation, *Food Chemistry*. 53:  
9  
10 1569 177-185.  
11  
12 1570  
13  
14  
15 1571 Scotter MJ, Wilson LA, Appleton GP, Castle L. 1998. Analysis of annatto (*Bixa*  
16  
17 1572 *orellana*) food colouring formulations. 1. Determination of colouring components and  
18  
19 1573 coloured thermal degradation products by high-performance liquid chromatography  
20  
21 1574 with photodiode array detection. *Journal of Agricultural and Food Chemistry*. 46:  
22  
23 1575 1031-1038.  
24  
25  
26 1576  
27  
28  
29 1577 Scotter MJ, Wilson LA, Appleton GP, Castle L. 2000. Analysis of annatto (*Bixa*  
30  
31 1578 *orellana*) food colouring formulations. 2. Determination of aromatic hydrocarbon  
32  
33 1579 thermal degradation products by gas chromatography. *Journal of Agricultural and*  
34  
35 1580 *Food Chemistry*. 48: 484-488.  
36  
37  
38 1581  
39  
40  
41 1582 Scotter MJ, Appleton GP, Castle L. 2001. Kinetics and yields for the formation of  
42  
43 1583 coloured and aromatic thermal degradation products of annatto in foods. *Food*  
44  
45 1584 *Chemistry*. 74: 365-375.  
46  
47  
48 1585  
49  
50  
51 1586 Scotter MJ, Castle L, Honeybone C, Nelson C. 2002. Method development and  
52  
53 1587 analysis of retail foods for annatto food colouring material. *Food Additives and*  
54  
55 1588 *Contaminants*. 19: 205-222.  
56  
57  
58 1589  
59  
60

- 1  
2  
3 1590 Silva MCD, Bothelo JR, Conceição MM, Lira BF, Couthino MA, Dias AF, Souza  
4  
5 1591 AG, Filho PFA. 2005. Thermogravimetric investigations on the thermal degradation  
6  
7 1592 of bixin, derived from the seeds of annatto (*Bixa orellana* L.). Journal of Thermal  
8  
9 1593 Analysis and Calorimetry: 79: 277-281.  
10  
11 1594  
12  
13 1595 Silva GS, Souza AG, Botelho JR, Silva MCD, Silva TMS. 2007. Kinetics study of  
14  
15 1596 norbixin's first stage thermal decomposition, using dynamic method. Journal of  
16  
17 1597 Thermal Analysis and Calorimetry. 87: 871-874.  
18  
19 1598  
20  
21 1599 Smith PR, Blake CJ, Porter DC. 1983. Determination of added natural colours in  
22  
23 1600 foods. III. Annatto. Leatherhead Food R.A. Research Report No.431. Leatherhead:  
24  
25 1601 British Food Manufacturing Industries Research Association.  
26  
27 1602  
28  
29 1603 Tennant DR, O'Callaghan MO. 2005. Survey of usage and estimated intakes of  
30  
31 1604 annatto extracts. Food Research International. 38: 911-917.  
32  
33 1605  
34  
35 1606 Tirimanna ASL. 1980. Study of the carotenoid pigments of *Bixa orellana* L. seeds by  
36  
37 1607 thin layer chromatography. Mikrochimica Acta (Vienna). II: 11-16.  
38  
39 1608  
40  
41 1609 Van Breemen, RB. 1995. Electrospray liquid-chromatography mass-spectrometry of  
42  
43 1610 carotenoids. Analytical Chemistry. 67: 2004-2009.  
44  
45 1611  
46  
47 1612 Vetter W, Englert G, Rigassi , Schwieter U. 1971. Spectroscopic methods. In: Isler O,  
48  
49 1613 editor. Carotenoids. Basel: Birkhäuser. p 189-266.  
50  
51 1614  
52  
53  
54  
55  
56  
57  
58  
59  
60


- 1  
2  
3 1615 Vetter W, Meister W. 1985. Fast atom bombardment mass-spectrum of beta-carotene.  
4  
5 1616 Organic Mass Spectroscopy. 20: 266-267.  
6  
7 1617  
8  
9  
10 1618 Weedon BCL, Moss GP. 1995. Chapter 3: Structure and nomenclature. In: Britton G,  
11  
12 1619 Liaaen-Jensen, S, Pfander, editors. Carotenoids, Volume 1A: Isolation and analysis.  
13  
14 1620 Basel: Birkhäuser-Verlag. p 27-70.  
15  
16 1621  
17  
18 1622 Zechmeister L. 1960. Cis-trans isomeric carotenoid pigments. Progress in the  
19  
20 1623 chemistry of natural products. 18: 223-349.  
21  
22 1624  
23  
24 1625 Zhao W, Han YS, Zhao BL, Hirota S, Hou JW, Xin WJ. 1998. The effect of  
25  
26 1626 carotenoids on the respiratory burst of rat peritoneal macrophages. Biochimica et  
27  
28 1627 Biophysica Acta – General Subjects. 1381: 77-88.  
29  
30 1628  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60









1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Peer Review Only




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Peer Review Only



For Peer Review Only



For Peer Review Only



Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Legends for figures

Figure 1. Chemical structures of some bixin/norbixin isomers.  $R_1=H$ ,  $R_2=H$  = norbixin;

$R_1=H$ ,  $R_2=CH_3$  = bixin.

Figure 2. Spectral fine structure. Calculation of %III/II for a carotenoid (9'-*cis*-norbixin).

Figure 3. UV-VIS spectra of norbixin isomers (by HPLC-photodiode array): (a) 9'-*cis*-norbixin, (b) di-*cis*-norbixin and (c) *trans*-norbixin (Scotter et al., 1994).

Figure 4. HPLC-photodiode array spectra of bixin isomers showing the locations of  $\lambda_{\max}$  (I)-(IV) (Scotter et al., 1994).

Figure 5. HPLC-photodiode array spectrum of 9'-*cis*-bixin (broken line) and its first derivative spectrum (solid line) highlighting the inflection at  $\lambda_{\max}$  (II) (Scotter et al., 1994).

Figure 6. Suggested reaction pathways for the thermal degradation of 9'-*cis*-bixin (Scotter et al., 2001).

Figure 7. Coupled reaction scheme proposed for the degradation of bixin and the formation of its primary products (Rios et al., 2005).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 8. HPLC separation of bixin and norbixin isomers. Conditions: Column HRPB C<sub>8</sub>/C<sub>18</sub> 250 x 4.6mm, 5µm; Mobile phase Acetonitrile: 0.4% acetic acid (65:35) isocratic elution at 1 ml.min<sup>-1</sup> 35°C; Detection photodiode array at 455 x 10nm. Assignment of peaks: 1. *Trans*-norbixin. 2. *Di-cis*-norbixin. 3. 9'-*cis*-norbixin. 4. *Trans*-bixin. 5 and 9. *Di-cis*-bixin isomers. 6. 9'-*cis*-bixin. 7. 15-*cis*-bixin\*. 8. 13'-*cis*-bixin\*. (\*tentative).



Table 1. Permitted uses of annatto and maximum levels of addition (EC, 1994).

Food Commodity type	Maximum permitted level (mg/kg)*
Margarine, minarine, other fat emulsions, and fats essentially free from water	10
Decorations and coatings	20
Fine bakery wares	10
Edible ices	20
Liqueurs, including fortified beverages with less than 15% alcohol by volume	10
Flavoured processed cheese	15
Ripened orange, yellow and broken white cheese; unflavoured processed cheese	15
Desserts	10
'Snacks': dry, savoury potato, cereal or starch-based products: Extruded or expanded savoury snack products	20
Other savoury snack products and savoury coated nuts	10
Smoked fish	10
Edible cheese rinds and edible casings	20
Red Leicester cheese	50
Mimolette cheese	35
Extruded, puffed and/or fruit-flavoured breakfast cereals	25

[\* Refers to 100% bixin or norbixin]

Table 2. Summary of planar chromatographic methods for annatto colours.

Sample type	Adsorbent	Mobile phase	Reference(s)
Bixin, norbixin, C <sub>17</sub>	Paper	CHX:CHCl <sub>3</sub> :DMF:HOAc (85:10:3:2)	McKeown, 1961, 1963
Annatto and other fat soluble dyes	Silica gel G	Amyl acetate	Ramamurthy and Bhalerao, 1964
Annatto	Silica gel G	1% HOAc in amyl acetate	Francis, 1965
Bixin	Silica gel	CHCl <sub>3</sub> :ACE:HOAc (50:50:1)	Dendy, 1966
(1) Bixin (2) Norbixin	(1) Cellulose (2) Silica gel	(1) CHX:CHCl <sub>3</sub> :HOAc (65:5:1) (2) CHCl <sub>3</sub> :EtOH:HOAc (68:2:1)	Reith and Gielen, 1971
Annatto and other pigments	Silica gel G (2-dimensional)	(1) CHCl <sub>3</sub> :EtOAc (4:1) (2) Et <sub>2</sub> O	Tirimanna, 1980
Bixin and norbixin commercial formulations	Silica gel GF	PE:Et <sub>2</sub> O:HOAc (85:15:2.5)	Preston and Rickard, 1980
Ether extracts of foods	Silica gel	(1) CHCl <sub>3</sub> :HOAc (9:1) (2) Et <sub>2</sub> O:IPA (9:1)	Corradi and Micheli, 1981
Annatto seeds	KC18 reverse phase	MeOH:H <sub>2</sub> O (70:30)	Chao et al., 1991
Bixin and other carotenoids	Silica gel GF	(1) HEX:ACE (10:9) (2) DCM:Et <sub>2</sub> O (9:1) (3) PE: BZ (1:1) (4) PE	Mínguez-Mosquera, Hornero-Méndez and Garrido-Fernández (1995)
Annatto seeds	(1) Silica gel (2) MgO/Kieselguhr	(1) HEX:t-BME (90:10) (2) HEX:ACE (85:15)	Mercadante, Steck and Pfander, 1997b
Annatto formulations	Silica gel GF	CHCl <sub>3</sub> :HOAc:ACN:ACE (8:1:0.5:0.5)	Galindo-Cuspinera and Rankin, 2005

[Key: ACE = Acetone; ACN = Acetonitrile; BZ = Benzene; CHX = Cyclohexane, CHCl<sub>3</sub> = Chloroform; DMF = N,N-Dimethylformamide; EtOAc = Ethyl acetate; EtOH = Ethanol; Et<sub>2</sub>O = Diethyl ether; HEX = Hexane; HOAc = Acetic acid; IPA = isopropyl alcohol; PE = Petroleum ether; t-BME = tertiary butylmethyl ether]

Table 3. Summary of HPLC methods used for analysis of annatto.

Sample matrix	Analyte(s)	HPLC conditions			Reference(s)
		Column	Mobile phase	Detector	
Annatto colour	Bixin and norbixin isomers	HRPB C <sub>8</sub> /C <sub>18</sub> 250 x 4.6mm, 5µm	ACN: 2% HOAc (65:35) isocratic 1 ml.min <sup>-1</sup> 35°C	UV-VIS PDA 452, 460nm	Scotter et al., 1994
Annatto colour	Bixin and norbixin isomers and C <sub>17</sub> isomers	HRPB C <sub>8</sub> /C <sub>18</sub> 250 x 4.6mm, 5µm	ACN: 0.4% HOAc (65:35) isocratic 1 ml.min <sup>-1</sup> 35°C	UV-VIS PDA 435 x 60nm	Scotter, 1995; Scotter et al, 1998, 2001
Foods	<i>Cis/trans</i> bixin and norbixin	Supelco LC-18 250 x 4.6mm, 5µm	MeOH: 2%HOAc (9:1) isocratic 1 ml.min <sup>-1</sup>	UV-VIS 500nm	Lancaster and Lawrence, 1995
Foods	Bixin, norbixin and carminic acid	Supelco LC-18 250 x 4.6mm, 5µm	MeOH: 6%HOAc gradient 1 ml.min <sup>-1</sup>	UV-VIS 493nm	Lancaster and Lawrence, 1996
Plasma	Bixin and norbixin isomers	S5ODS1	ACN: 2% HOAc isocratic 1.5 ml.min <sup>-1</sup>	UV-VIS PDA 460nm	Levy et al., 1997
DNA	Bixin and norbixin	Supelco LC-8 250 x 4.6mm, 10µm	ACN: 0.08% CF <sub>3</sub> CO <sub>2</sub> H (85:15) isocratic 1 ml.min <sup>-1</sup>	UV-VIS 470nm	Kovary et al., 2001
Foods	Bixin and norbixin isomers	HRPB C <sub>8</sub> /C <sub>18</sub> 250 x 4.6mm, 5µm	ACN: 0.4% HOAc (65:35) isocratic 1 ml.min <sup>-1</sup> 35°C	UV-VIS PDA 455 x 10nm	Scotter et al., 2002
Cheese	<i>Cis/trans</i> bixin and norbixin	ODS2 C <sub>18</sub> 250 x 4mm, 5µm	ACN: 2%HOAc (75:25) isocratic 1 ml.min <sup>-1</sup>	UV-VIS 460nm	Bareth, Strohmar and Kitzelmann, 2002
Corn snacks	Norbixin	ODS2 C <sub>18</sub> 150 x 4mm, 3µm	ACN: 2%HOAc (65:35) isocratic 1 ml.min <sup>-1</sup> 29°C	UV-VIS PDA 450nm	O Rios and Mercadante, 2004
Food	Bixin, norbixin and other	YMC C30 250 x 4.6mm, 5µm	A: MeOH: H <sub>2</sub> O: TEA (90:10:0.1) B: MTBE: MeOH: H <sub>2</sub> O:TEA	UV-VIS PDA 450 x 4nm	Breithaupt, 2004

	carotenoids		(90:6:4:0.1) gradient 1 ml.min <sup>-1</sup> 35°C	+ LC-MS	
Bixin	Photodegradation products	(1) Vydac C18 250 x 4.6mm, 5µm (2) ODS2 C18 150 x 4.6 3µm	ACN: 2% HOAc: DCM (65:35:2) isocratic 1 ml.min <sup>-1</sup> 25°C	UV-VIS PDA 450nm	Montenegro et al., 2004
Aqueous model system	Bixin thermal degradation products	ODS2 C <sub>18</sub> 150 x 4mm, 3µm	ACN: 2%HOAc (65:35) or ACN: 2% HOAc: DCM (65:35:2) isocratic 1 ml.min <sup>-1</sup> 29°C	UV-VIS PDA 450nm	O Rios et al, 2005
Water soluble annatto	<i>Cis/trans</i> norbixin	Beckman C18 250 x 4.6mm, 5µm	ACN: 0.4% HOAc + 5% ACN isocratic and gradient 1 ml.min <sup>-1</sup>	UV-VIS PDA 250 - 600nm + LC-MS	Galindo-Cuspinera and Rankin, 2005

[Key: ACN = Acetonitrile; DCM = Dichloromethane; MeOH = Methanol; MTBE = Methyl tertiary butyl ether; TEA = Triethylamine]

Table 4. Summary of extraction regimes used for annatto in foods (Scotter et al., 2002).

<b>Regime</b>	<b>Matrices</b>
1	Cheese, cheese products and cheese-based compound foods
2	Custard powder and low-fat dessert dry mixes
3	Desserts, cake decorations, fine bakery wares, extruded snacks and breakfast cereals
4	Margarine, fat-based emulsions and spreads, butter and fat-based compound foods
5	Fish, ice cream and ice cream-based confectionery, yoghurt and other dairy desserts