



HAL
open science

Invariant differential operators on a class of multiplicity free spaces

Hubert Rubenthaler

► **To cite this version:**

Hubert Rubenthaler. Invariant differential operators on a class of multiplicity free spaces. Pacific Journal of Mathematics, 2014, 270 (2), pp.473-510. 10.2140/pjm.2014.270.473 . hal-00574585v2

HAL Id: hal-00574585

<https://hal.science/hal-00574585v2>

Submitted on 13 Feb 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INVARIANT DIFFERENTIAL OPERATORS ON A CLASS OF MULTIPLICITY FREE SPACES

HUBERT RUBENTHALER

ABSTRACT. If (G, V) is a multiplicity free space with a one dimensional quotient we give generators and relations for the non-commutative algebra $D(V)^{G'}$ of invariant differential operators under the semi-simple part G' of the reductive group G . More precisely we show that $D(V)^{G'}$ is the quotient of a Smith algebra by a completely described two-sided ideal.

AMS classification: 22E46, 16S32

Key words: multiplicity free space, invariant differential operator, Smith algebra.

1. INTRODUCTION

Let H be a reductive algebraic group over \mathbb{C} and let X be a smooth irreducible H -variety. Let $\mathbb{C}[X]$ be the algebra of regular functions on X and let $D(X)$ be the algebra of differential operators on X . Then the H -action on X extends naturally to $\mathbb{C}[X]$ and $D(X)$. Let $\mathbb{C}[X]^H$ (resp. $D(X)^H$) be the subalgebras of H -invariants in $\mathbb{C}[X]$ (resp. $D(X)$). The ring $\mathbb{C}[X]^H$ is the ring of regular functions on the categorical quotient $X//H$. The problem of determining the structure of $D(X)^H$ was investigated by several authors ([30], [34], [20]). On the other hand under the above mentioned hypothesis there exists a H -equivariant restriction map

$$\delta : D(X)^H \longrightarrow D(X//H).$$

obtained by applying elements in $D(X)^H$ to functions in $\mathbb{C}[X]^H$. It is expected that $D(X)^H$ as well as its image under δ (the so-called algebra of radial components) should share many properties of enveloping algebras([30], [19]). In this paper we obtain the precise structure of $D(V)^{G'}$ in the case where (G, V) is a so called *multiplicity free spaces with one dimensional quotient*, (here G is reductive and $G' = [G, G]$ is the derived group). These spaces are defined to be the multiplicity free spaces (G, V) for which the quotient $V//G'$ is one dimensional. To be more precise we show that the (non-commutative) algebra $D(V)^{G'}$ is a quotient of a generalized *Smith algebra*. Over \mathbb{C} this kind of algebras were introduced by S. P. Smith ([32]) as natural generalizations of the enveloping algebra of \mathfrak{sl}_2 . As a Corollary we describe by generators and relations the algebras of radial components attached to the G' -isotypic

Date: February 13, 2014.

components in the polynomial algebra $\mathbb{C}[V]$ (the image under δ above corresponds to the trivial representation of G').

According to the classification obtained by the author ([27]), the class of multiplicity free spaces with a one dimensional quotient is a rather large class inside the multiplicity free spaces. It contains both irreducible and non irreducible representations.

The representations $(Str(V), V)$ where V is a simple Jordan algebra over \mathbb{C} and where $StrV$ is the structure group of V are examples of irreducible multiplicity free spaces with a one dimensional quotient (see Remark 2.2.7 and Example 2.3.3 below). Among these there is the natural representation of $GL(n, \mathbb{C})$ on the space $Sym_n(\mathbb{C})$ of $n \times n$ symmetric matrices and also the irreducible 27-dimensional representation of $E_6 \times \mathbb{C}^*$.

The spin representation of $Spin(7) \times \mathbb{C}^*$ and the irreducible 7-dimensional representation of $G_2 \times \mathbb{C}^*$ are other irreducible examples.

The representation $(SL(n, \mathbb{C}) \times (\mathbb{C}^*)^2, \Lambda_1 \oplus \Lambda^2(\Lambda_1))$ (n odd and $n \geq 5$) where Λ_1 is the natural representation of $SL(n, \mathbb{C})$ and where $\Lambda^2(\Lambda_1)$ is its second exterior power, provides a non irreducible example.

Let us now give a more precise description of our paper.

In section 2 we first give the basic definitions, notations and properties of multiplicity free spaces, including multiplicity free spaces with one dimensional quotient. If (G, V) is a multiplicity free space then G has an open orbit on V (i.e. (G, V) is a prehomogeneous vector space). We also prove that in the so-called regular case the G -invariant differential operators on the open orbit of a multiplicity free space have always polynomial coefficients (in fact a slightly more general result is proved, see Theorem 2.2.6).

In section 3 we introduce the various algebras of differential operators we are interested in. We define their natural gradings and we define the so-called Bernstein-Sato polynomial of an homogeneous operator of any degree, not only for degree zero operators as usual. We obtain there the first results concerning these algebras. Using the Harish-Chandra isomorphism for multiplicity free spaces ([16]), we prove a key lemma on invariant polynomials under the so-called little Weyl group which enables us to prove that $D(V)^G$ is a polynomial algebra over the center $\mathcal{Z}(\mathcal{T})$ of $D(V)^{G'}$, with the Euler operator as generator (Theorem 3.2.6). We also give generators of the center $\mathcal{Z}(\mathcal{T})$ (Theorem 3.2.10) and obtain some specific results in the case of prehomogeneous vector spaces of commutative parabolic type (Theorem 3.3.1).

Section 4, which is the main section, is devoted to the structure of $D(V)^{G'}$. We first briefly define and study the Smith algebras over a commutative ring \mathbf{A} with unit and no zero divisors (the original definition by Smith was over \mathbb{C}). These algebras are defined by generators and relations (involving a polynomial in $\mathbf{A}[t]$), and their center is a polynomial algebra $\mathbf{A}[\Omega_1]$, where Ω_1 is a generalized Casimir element. Our main result asserts that $D(V)^{G'}$ is isomorphic to the quotient of a Smith algebra over its center $\mathcal{Z}(\mathcal{T})$ by the two-sided ideal generated by the element Ω_1 . Concretely, we give generators and relations for $D(V)^{G'}$ (see Theorem 4.2.2).

Section 5 is devoted to the study of the algebras of radial components. By radial component of a differential operator in $D(V)^{G'}$ we mean the restriction of D to a G' -isotypic component of $\mathbb{C}[V]$. As a corollary of the preceding results we prove that these algebras are quotients of "classical" Smith algebras, that is Smith algebras over \mathbb{C} (see Theorem 5.2.3). Of course the defining relations depend on the G' -isotypic component. We also give generators of the kernel

of the radial component map. In the case of the trivial representation of G' , the structure of the algebra of radial components was first obtained by Levasseur ([19]), by other methods.

Acknowledgment: I would like to thank Thierry Levasseur for providing me with the manuscript of [19]. I would also like to thank Sylvain Rubenthaler who provided me with a first proof of Proposition 4.1.9 which was important for my understanding.

2. MULTIPLICITY FREE SPACES WITH A ONE DIMENSIONAL QUOTIENT

2.1. Prehomogeneous Vector Spaces. Basic definitions and properties.

Let G be a connected algebraic group over \mathbb{C} , and let (G, ρ, V) be a rational representation of G on the (finite dimensional) vector space V . Then the triplet (G, ρ, V) is called a *prehomogeneous vector space* (abbreviated to *PV*) if the action of G on V has a Zariski open orbit $\Omega \subset V$. For the general theory of *PV*'s, we refer the reader to the book of Kimura [14] or to [31]. The elements in Ω are called *generic*. The *PV* is said to be *irreducible* if the corresponding representation is irreducible. The *singular set* S of (G, ρ, V) is defined by $S = V \setminus \Omega$. Elements in S are called *singular*. If no confusion can arise we often simply denote the *PV* by (G, V) . We will also write $g.x$ instead of $\rho(g)x$, for $g \in G$ and $x \in V$. It is easy to see that the condition for a rational representation (G, ρ, V) to be a *PV* is in fact an infinitesimal condition. More precisely let \mathfrak{g} be the Lie algebra of G and let $d\rho$ be the derived representation of ρ . Then (G, ρ, V) is a *PV* if and only if there exists $v \in V$ such that the map:

$$\begin{array}{ccc} \mathfrak{g} & \longrightarrow & V \\ X & \longmapsto & d\rho(X)v \end{array}$$

is surjective (we will often write $X.v$ instead of $d\rho(X)v$). Therefore we will call (\mathfrak{g}, V) a *PV* if the preceding condition is satisfied.

Let (G, V) be a *PV*. A rational function f on V is called a *relative invariant* of (G, V) if there exists a rational character χ of G such that $f(g.x) = \chi(g)P(x)$ for $g \in G$ and $x \in V$. From the existence of an open orbit it is easy to see that a character χ which is trivial on the isotropy subgroup of an element $x \in \Omega$ determines a unique relative invariant P . Let S_1, S_2, \dots, S_k denote the irreducible components of codimension one of the singular set S . Then there exist irreducible polynomials P_1, P_2, \dots, P_k such that $S_i = \{x \in V \mid P_i(x) = 0\}$. The polynomials P_i 's are unique up to nonzero constants, they are relative invariants of (G, V) and any nonzero relative invariant f can be written in a unique way $f = cP_1^{n_1}P_2^{n_2} \dots P_k^{n_k}$, where $n_i \in \mathbb{Z}$ and $c \in \mathbb{C}^*$. The polynomials P_1, P_2, \dots, P_k are called the *fundamental relative invariants* of (G, V) . Moreover if the representation (G, V) is irreducible then there exists at most one irreducible polynomial which is relatively invariant.

The prehomogeneous vector space (G, V) is called *regular* if there exists a relative invariant polynomial P whose Hessian $H_P(x)$ is nonzero on Ω . If G is reductive, then (G, V) is regular if and only if the singular set S is a hypersurface, or if and only if the isotropy subgroup of a generic point is reductive. If the *PV* (G, V) is regular, or if G is reductive, then the contragredient representation (G, V^*) is again a *PV*.

2.2. Multiplicity free spaces.

For the results concerning multiplicity free spaces we refer the reader to the survey by Benson and Ratcliff ([1]) or to [16]. Let (G, V) be a finite dimensional rational representation of a connected reductive algebraic group G . Let $\mathbb{C}[V]$ be the algebra of polynomials on V . Then

G acts on $\mathbb{C}[V]$ by

$$g \cdot \varphi(x) = \varphi(g^{-1}x) \quad (g \in G, \varphi \in \mathbb{C}[V]).$$

As the space $\mathbb{C}[V]^n$ of homogeneous polynomials of degree n is stable under this action, the representation $(G, \mathbb{C}[V])$ is completely reducible. Let $D(V)$ be the algebra of differential operators on V with polynomial coefficients. The group G acts also on $D(V)$ by

$$(g \cdot D)(\varphi) = g \cdot (D(g^{-1} \cdot \varphi)) \quad (g \in G, D \in D(V), \varphi \in \mathbb{C}[V]).$$

Recall the G -equivariant identifications between $\mathbb{C}[V]$ and the symmetric algebra $S(V^*)$ of the dual space V^* and between $\mathbb{C}[V^*]$ and the symmetric algebra $S(V)$ of V . The embedding

$V \rightarrow D(V)$
 $v \mapsto D_v$ where $D_v P(x) = \lim_{t \rightarrow 0} \frac{P(x+tv) - P(x)}{t}$ extends uniquely to an embedding $S(V) \rightarrow D(V)$ whose image is the ring of differential operators with constant coefficients. If $f \in S(V) \simeq \mathbb{C}[V^*]$ we denote by $f(\partial)$ the corresponding differential operator. Another way to construct $f(\partial)$ for $f \in \mathbb{C}[V^*]$ is to say that $f(\partial)$ is the unique differential operator on V satisfying

$$f(\partial_x) e^{\langle x, y \rangle} = f(y) e^{\langle x, y \rangle} \quad (x \in V, y \in V^*) \quad (2-2-1)$$

Recall also that the $\mathbb{C}[V]$ -module $D(V)$ can be identified with $\mathbb{C}[V] \otimes S(V)$ through the multiplication map

$$\begin{array}{ccc} m : \mathbb{C}[V] \otimes S(V) & \xrightarrow{\simeq} & D(V) \\ \varphi \otimes f & \mapsto & \varphi f(\partial) \end{array}$$

The preceding map is in fact G -equivariant and therefore the G -module $D(V)$ is isomorphic to the G -module $\mathbb{C}[V] \otimes S(V)$. The duality pairing $V \otimes V^* \rightarrow \mathbb{C}$ extends uniquely to the non-degenerate G -equivariant pairing

$$\begin{array}{ccc} S(V) \otimes S(V^*) \simeq \mathbb{C}[V^*] \otimes \mathbb{C}[V] & \rightarrow & \mathbb{C} \\ f \otimes \varphi & \mapsto & \langle f, \varphi \rangle = f(\partial)\varphi(0) \end{array} \quad (2-2-2)$$

which gives rise to an embedding $\mathbb{C}[V^*] \hookrightarrow \mathbb{C}[V]^*$. It is easy to see that $\langle \mathbb{C}^i[V^*], \mathbb{C}^j[V] \rangle = \{0\}$ if $i \neq j$.

Definition 2.2.1. *Let G be a connected reductive algebraic group, and let V be the space of a finite dimensional (complex) rational representation of G . The representation (G, V) is said to be multiplicity free (abbreviated to MF) if each irreducible representation of G occurs at most once in the representation $(G, \mathbb{C}[V])$.*

Remark 2.2.2. Historically the classification of MF spaces goes as follows. Kac ([13]) determined all the MF spaces where the representation (G, V) is irreducible. Brion ([7]) did the case where $G' = [G, G]$ is (almost) simple. Finally Benson-Ratcliff and Leahy classified independently, up to geometric equivalence, all the indecomposable saturated MF -spaces (see [1], [2], [18],[16])

The following theorem summarizes some basic results concerning MF spaces (see [1], [11], [16]):

Theorem 2.2.3.

1) *A finite dimensional representation (G, V) is MF if and only if (B, V) is a prehomogeneous vector space for any Borel subgroup B of G (and hence each MF space (G, V) is a PV).*

2) A finite dimensional representation (G, V) is MF if and only if the algebra $D(V)^G$ of invariant differential operators with polynomial coefficients is commutative.

3) If (G, V) is a MF space, then the dual space (G, V^*) is also MF.

Proof. The first assertion is due to Vinberg and Kimelfeld ([35]), another proof can be found in [16]. The second assertion is due to Howe and Umeda ([11], Theorem 7.1). For the third assertion note that as $\langle \mathbb{C}^i[V^*], \mathbb{C}^j[V] \rangle = \{0\}$ for $i \neq j$, we obtain that $f \mapsto \langle f, \cdot \rangle$ is a G -equivariant isomorphism between $\mathbb{C}^i[V^*]$ and $\mathbb{C}^i[V]^*$, and hence (G, V^*) is multiplicity free. \square

Let us be more precise about the decomposition of the polynomials under the action of the group G or a Borel subgroup. Therefore we need more notations. We can write $G = G'C$ where $G' = [G, G]$ is the subgroup of commutators, and where $C = Z(G)^\circ \simeq (\mathbb{C}^*)^p$ is the connected component of the center of G . Let T' be a maximal torus in G' , and let $B' = T'U$ be a Borel subgroup of G' , where U is the nilradical of B' . The group $T = T'C$ is a maximal torus in G and $B = TU$ is a Borel subgroup of G . We will denote by $\mathfrak{g}, \mathfrak{g}', \mathfrak{t}, \mathfrak{t}', \mathfrak{c}, \mathfrak{b}, \mathfrak{b}', \mathfrak{u}$ the corresponding Lie algebras. Let R be the set of roots of $(\mathfrak{g}', \mathfrak{t}')$, let $\Delta = \{\alpha_1, \dots, \alpha_\ell\}$ be the basis of simple roots corresponding to \mathfrak{b}' and let R^+ be the corresponding set of positive roots. Denote by Λ' the lattice of weights of $(\mathfrak{g}', \mathfrak{t}')$. We have $\Lambda' = \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2 \oplus \dots \oplus \mathbb{Z}\omega_\ell$ where the ω'_i 's are the fundamental weights. Let $\Lambda'^+ = \mathbb{N}\omega_1 \oplus \mathbb{N}\omega_2 \oplus \dots \oplus \mathbb{N}\omega_\ell$ be the the set of dominant weights. Denote by $X(C)$ the group of algebraic characters of C , which we will sometimes consider as linear forms on \mathfrak{c} . Set:

$$\Lambda = \Lambda' \oplus X(C), \quad \Lambda^+ = \Lambda'^+ \oplus X(C).$$

For $\lambda \in \Lambda^+$ (resp. $\lambda' \in \Lambda'^+$) let us denote by $V_{-\lambda}$ (resp. $V_{-\lambda'}$) an irreducible \mathfrak{g} -module (resp. \mathfrak{g}' -module) with highest weight λ (resp. λ'). We use this unusual notation because we want to index the modules occuring in $\mathbb{C}[V]$ by the character of their highest weight polynomial, rather than by the highest weight.

For a multiplicity free space (G, V) we have the decomposition:

$$\mathbb{C}[V] = \bigoplus_{\lambda \in \Lambda^+} V_{-\lambda}^{m(\lambda)}$$

where $m(\lambda) = 0$ or 1 . If $m(\lambda) = 1$, then there exists a uniquely defined positive integer $d(\lambda)$ such that $V_{-\lambda} \in \mathbb{C}[V]^{d(\lambda)}$. The integer $d(\lambda)$ is called the *degree* of λ . Let us denote by $\Delta_0, \Delta_1, \dots, \Delta_k, \dots, \Delta_r$ the fundamental relative invariants invariants of the PV (B, V) , indexed in such a way that $\Delta_0, \Delta_1, \dots, \Delta_k$ are the fundamental relative invariants of the PV (G, V) and such that the other invariants are ordered by decreasing degree. We denote by d_i the degree of Δ_i ($i = 0, \dots, r$). It is worthwhile noticing that at least Δ_r is of degree one as the highest weight vectors of the irreducible components of V^* must occur. Then any relative invariant of (B, V) is of the form $c\Delta^{\mathbf{a}}$ where $\mathbf{a} = (a_0, a_1, \dots, a_r) \in \mathbb{Z}^{r+1}$ and where $\Delta^{\mathbf{a}} = \Delta_0^{a_0} \dots \Delta_r^{a_r}$. The non negative integer $r + 1$ is called the *rank* of the MF space (G, V) . The algebra of U -invariants is the subalgebra generated by the Δ_i 's, i.e. $\mathbb{C}[V]^U = \mathbb{C}[\Delta_0, \dots, \Delta_r]$. As the polynomials Δ_i are algebraically independent, this latter algebra is a polynomial algebra. Let λ_i be the character of Δ_i (we use the same notation λ_i for the character of the group and for its derivative, which is an element of Λ^+). Hence the (infinitesimal) character of $\Delta^{\mathbf{a}}$ is $\lambda_{\mathbf{a}} = a_0\lambda_0 + \dots + a_r\lambda_r$. Of course by definition the elements $\Delta^{\mathbf{a}}$ ($a_i \geq 0, i = 0, \dots, r$) are the highest weights vectors in $\mathbb{C}[V]$. Due to the fact that the group action on $\Delta^{\mathbf{a}}$ is given by $g.\Delta^{\mathbf{a}}(x) = \Delta^{\mathbf{a}}(g^{-1}x)$, the infinitesimal highest weight of $\Delta^{\mathbf{a}}$ is $-\lambda_{\mathbf{a}} = -a_0\lambda_0 - \dots - a_r\lambda_r$.

If we set $V_{\mathbf{a}} = V_{-\lambda_{\mathbf{a}}}$, we therefore can write

$$\mathbb{C}[V] = \bigoplus_{a_0 \geq 0, \dots, a_r \geq 0} V_{\mathbf{a}} \quad (2-2-3)$$

Sometimes, if $\lambda = a_0\lambda_0 + \dots + a_r\lambda_r$, we simply write V_{λ} instead of $V_{\mathbf{a}}$. If we denote by d_i the degree of Δ_i , one can notice that all elements in $V_{\mathbf{a}}$ are of degree $d(\mathbf{a}) = a_0d_0 + a_1d_1 + \dots + a_rd_r$. It is also worthwhile noticing that we have:

$$V_{\mathbf{a}} = \Delta_0^{a_0} \Delta_1^{a_1} \dots \Delta_k^{a_k} V_{0, \dots, 0, a_{k+1}, \dots, a_r}. \quad (2-2-4)$$

The proof of the following lemma is straightforward.

Lemma 2.2.4.

Define $\mathcal{O} = \{x \in V \mid \Delta_i(x) \neq 0, i = 0, \dots, k\}$. Let $\mathbb{C}[\mathcal{O}]$ be the ring of regular functions on \mathcal{O} (elements of $\mathbb{C}[\mathcal{O}]$ are just rational functions whose denominators are of the form $\Delta_0^{a_0} \dots \Delta_k^{a_k}$, with $a_0, \dots, a_k \geq 0$). As the polynomials $\Delta_0, \dots, \Delta_k$ are relative invariants under G , the open set \mathcal{O} is G -stable, and therefore G acts on $\mathbb{C}[\mathcal{O}]$. Then $\mathbb{C}[\mathcal{O}]$ decomposes without multiplicities under the action of G . More precisely the decomposition into irreducibles is given by

$$\mathbb{C}[\mathcal{O}] = \bigoplus_{\substack{(a_0, \dots, a_k) \in \mathbb{Z}^{k+1} \\ (a_{k+1}, \dots, a_r) \in \mathbb{N}^{r-k}}} V_{\mathbf{a}}$$

where $V_{\mathbf{a}} = \Delta_0^{a_0} \Delta_1^{a_1} \dots \Delta_k^{a_k} V_{0, \dots, 0, a_{k+1}, \dots, a_r}$ is the irreducible subspace of $\mathbb{C}[\mathcal{O}]$ generated by the highest weight vector $\Delta^{\mathbf{a}} = \Delta_0^{a_0} \Delta_1^{a_1} \dots \Delta_r^{a_r}$.

Remark 2.2.5. We want to draw the attention of the reader to the fact that if (G, V) is not a regular PV , then the open set \mathcal{O} may be distinct from the open G -orbit Ω .

The preceding lemma has the following consequence.

Theorem 2.2.6.

Let (G, V) be a multiplicity free space. As before set $\mathcal{O} = \{x \in V \mid \Delta_i(x) \neq 0, i = 0, \dots, k\}$. Then $D(V)^G = D(\mathcal{O})^G$. In other words any G -invariant differential operator with coefficients in $\mathbb{C}[\mathcal{O}]$ has in fact polynomial coefficients.

Proof. Let $D \in D(\mathcal{O})^G$. As we know from the preceding Lemma that $\mathbb{C}[\mathcal{O}]$ decomposes without multiplicities under G , we obtain that D defines a G -equivariant endomorphism on each $V_{\mathbf{a}}$, $\mathbf{a} \in \mathbb{Z}^{k+1} \times \mathbb{N}^{r-k}$. Therefore D stabilizes $\mathbb{C}[V] = \bigoplus_{a_0 \geq 0, \dots, a_r \geq 0} V_{\mathbf{a}}$. It is easy to see that a differential operator with rational coefficients and which stabilizes the polynomials must have polynomial coefficients. □

Remark 2.2.7. Let V be a simple Jordan algebra over \mathbb{C} or \mathbb{R} . Let Ω be the set of invertible elements in V and let G be the structure group of V . It is known that (G, V) is a multiplicity free space with Ω as open G -orbit. Then the preceding theorem implies that $D(V)^G = D(\Omega)^G$. This result was already known in this context and is usually obtained by computing an explicit set of generators of $D(\Omega)^G$ (see [24], [37] or [8]). Through the so-called Kantor-Koecher-Tits construction there is a one-to-one correspondence between these spaces and the PV 's of commutative parabolic type (see example 2.3.3 below).

Proposition 2.2.8.

Let (G, V) be a MF space. For $\tilde{\mathbf{a}} = (a_{k+1}, \dots, a_r) \in \mathbb{N}^{r-k}$ we define $V_{\tilde{\mathbf{a}}} = V_{(0, \dots, 0, a_{k+1}, \dots, a_r)}$. Then for $\mathbf{a} = (a_0, \dots, a_k, a_{k+1}, \dots, a_r)$ the spaces $V_{\mathbf{a}} = \Delta_0^{a_0} \dots \Delta_k^{a_k} V_{\tilde{\mathbf{a}}}$ are G' -equivalent if $\tilde{\mathbf{a}}$ is fixed and if $(a_0, \dots, a_k) \in \mathbb{Z}^{k+1}$. If we define

$$U_{\tilde{\mathbf{a}}} = \bigoplus_{(a_0, \dots, a_k) \in \mathbb{N}^{k+1}} \Delta_0^{a_0} \dots \Delta_k^{a_k} V_{\tilde{\mathbf{a}}}, \quad W_{\tilde{\mathbf{a}}} = \bigoplus_{(a_0, \dots, a_k) \in \mathbb{Z}^{k+1}} \Delta_0^{a_0} \dots \Delta_k^{a_k} V_{\tilde{\mathbf{a}}},$$

then the decompositions of $\mathbb{C}[V]$ and $\mathbb{C}[\mathcal{O}]$ into G' -isotypic components are given by

$$\mathbb{C}[V] = \bigoplus_{\tilde{\mathbf{a}}} U_{\tilde{\mathbf{a}}}, \quad \mathbb{C}[\mathcal{O}] = \bigoplus_{\tilde{\mathbf{a}}} W_{\tilde{\mathbf{a}}}$$

Proof. The map $P \mapsto \Delta_0^{a_0} \dots \Delta_k^{a_k} P$ is a G' -equivariant isomorphism between $V_{\tilde{\mathbf{a}}}$ and $\Delta_0^{a_0} \dots \Delta_k^{a_k} V_{\tilde{\mathbf{a}}}$, hence all these spaces are G' -equivalent. To prove the second assertion it is enough to prove that if $\tilde{\mathbf{a}} \neq \tilde{\mathbf{b}}$, then the spaces $V_{\tilde{\mathbf{a}}}$ and $V_{\tilde{\mathbf{b}}}$ are not G' -equivalent. Suppose that this would be the case and let $\Delta^{\tilde{\mathbf{a}}}$ and $\Delta^{\tilde{\mathbf{b}}}$ be the corresponding highest weight vectors with characters $\lambda_{\tilde{\mathbf{a}}}$ and $\lambda_{\tilde{\mathbf{b}}}$ respectively. From the G' -equivalence we know that $\lambda_{\tilde{\mathbf{a}}|_{\mathfrak{t}'}} = \lambda_{\tilde{\mathbf{b}}|_{\mathfrak{t}'}}$ and hence $P = \frac{\Delta^{\tilde{\mathbf{a}}}}{\Delta^{\tilde{\mathbf{b}}}}$ is a relative invariant under B whose character is trivial on \mathfrak{t}' . Therefore it generates a one dimensional representation, hence P is a relative invariant under G . Finally we obtain that $\Delta^{\tilde{\mathbf{a}}} = \Delta_0^{a_0} \dots \Delta_k^{a_k} \Delta^{\tilde{\mathbf{b}}}$, and this is not possible if $\tilde{\mathbf{a}} \neq \tilde{\mathbf{b}}$. \square

As (G, V^*) is multiplicity free (Theorem 2.2.3) and as $\mathbb{C}^i[V^*] \simeq \mathbb{C}^i[V]^*$, we have

$$\mathbb{C}[V^*] = \bigoplus_{a_0 \geq 0, \dots, a_r \geq 0} V_{\mathbf{a}}^* \quad (2-2-5)$$

where $V_{\mathbf{a}}^*$ is the irreducible G -submodule of $\mathbb{C}[V^*]$ generated by a lowest weight vector $\Delta^{*\mathbf{a}} \in \mathbb{C}[V^*]$, defined up to a multiplicative constant, whose character with respect to the opposite Borel subgroup B^- is equal to $-\lambda_{\mathbf{a}} = -a_0\lambda_0 - \dots - a_r\lambda_r$. Let us fix a lowest weight vector Δ_i^* ($i = 0, \dots, r$) with character $-\lambda_i$ (with respect to B^-). Then we can choose $\Delta^{*\mathbf{a}} = \Delta_0^{*a_0} \Delta_1^{*a_1} \dots \Delta_r^{*a_r}$. Of course the module $V_{\mathbf{a}}^*$ is the dual module of $V_{\mathbf{a}}$ through $f \mapsto \langle f, \rangle$ (see (2-2-2)).

As $V_{\mathbf{a}}$ is a G -irreducible module, it is well known that the tensor G -module $V_{\mathbf{a}} \otimes V_{\mathbf{a}}^*$ contains up to constant, a unique G -invariant vector $R_{\mathbf{a}}$ and that $V_{\mathbf{a}} \otimes V_{\mathbf{b}}^*$ does not contain any non trivial G -invariant vector if $\mathbf{a} \neq \mathbf{b}$ (see for example [11]). To be more precise we define $R_{\mathbf{a}}$ to be the operator corresponding to the "unit matrix" in $V_{\mathbf{a}} \otimes V_{\mathbf{a}}^* \simeq \text{Hom}(V_{\mathbf{a}}, V_{\mathbf{a}})$. Moreover as $\mathbb{C}[V] \otimes \mathbb{C}[V^*]$ is G -isomorphic to $D(V)$, the element $R_{\mathbf{a}}$ can be viewed as a G -invariant differential operator with polynomial coefficients. The operators $R_{\mathbf{a}}$ are sometimes called *Capelli operators*. They are also called *unnormalized canonical invariants* in [1]. Moreover the family of elements $R_{\mathbf{a}}$ ($\mathbf{a} \in \mathbb{N}^{r+1}$) is a vector basis of the vector space $D(V)^G = D(\mathcal{O})^G$. The Capelli operators R_i corresponding to the space V_{λ_i} ($i = 0, \dots, r$) will be of particular importance because of the result below.

Theorem 2.2.9. (Howe-Umeda)

Let (G, V) be a MF space. The Capelli operators R_i ($i = 0, \dots, r$) are algebraically independent and $D(V)^G = \mathbb{C}[R_0, \dots, R_r]$.

Proof. See [11] (Theorem 9.1) or [1] (Corollary 7.4.4). \square

Remark 2.2.10.

- a) Recall that for $i = 0, 1, \dots, k$ the polynomials $\Delta_0, \Delta_1, \dots, \Delta_k$ are the fundamental relative invariants under the action of the full group G . Once these polynomials are fixed, let us define the polynomial $\Delta_i^* \in \mathbb{C}[V^*]$ as the unique fundamental relative invariant of (G, V^*) with character λ_i^{-1} , such that $\Delta_i^*(\partial)\Delta_i(0) = 1$, for $i = 0, \dots, k$. Then the Capelli operators R_i ($i = 0, \dots, k$) are given by $R_i = \Delta_i(x)\Delta_i^*(\partial)$, and the Capelli operator corresponding to the irreducible component $V_{a_0\lambda_0+\dots+a_k\lambda_k}$ is scalar multiple of $\Delta_0^{a_0}(x) \dots \Delta_k^{a_k}(x)\Delta_0^*(\partial)^{a_0} \dots \Delta_k^*(\partial)^{a_k}$. More generally the Capelli operator $R_{\mathbf{a}}$ corresponding to $V_{\mathbf{a}}$ where $\mathbf{a} = a_0\lambda_0 + \dots + a_k\lambda_k + a_{k+1}\lambda_{k+1} + \dots + a_r\lambda_r$ is a scalar multiple of $\Delta_0^{a_0}(x) \dots \Delta_k^{a_k}(x)R_{a_{k+1}\lambda_{k+1}+\dots+a_r\lambda_r}\Delta_0^*(\partial)^{a_0} \dots \Delta_k^*(\partial)^{a_k}$.
- b) Moreover, in the case where (G, V) is irreducible, as Δ_r is the highest weight vector in V^* , the operator R_r is nothing but the Euler operator E .
- c) More generally, if $V = V_1 \oplus \dots \oplus V_\ell$ where the representations (G, V_i) are irreducible, the various Euler operators E_i on V_i are the Capelli operators associated to the irreducible subspaces $V_i^* \in \mathbb{C}[V]$. Of course the global Euler operator E on V is given by $E = E_1 + \dots + E_\ell$. As the highest weight vectors of the spaces (G, V_i^*) occur as the ℓ last elements of $\Delta_0, \dots, \Delta_r$, we have $R_{r-\ell+1} = E_1, \dots, R_r = E_\ell$.
- d) According to b) and c) before, one can always take $\{R_0, R_1, \dots, R_{r-1}, E\}$ as a set of algebraically independent generators of $D(V)^G$.

2.3. Multiplicity free spaces with one dimensional quotient.

Let us now define the main objects this paper deals with, namely the MF -spaces with a one dimensional quotient, which were introduced by T. Levasseur.

Definition 2.3.1. (T. Levasseur, [19] sections 3.2 and 4.2)

- 1) A prehomogeneous vector space (G, V) is said to be of rank one* if there exists an homogeneous polynomial Δ_0 on V such that $\Delta_0 \notin \mathbb{C}[V]^G$ and such that $\mathbb{C}[V]^G = \mathbb{C}[\Delta_0]$.
- 2) A multiplicity free space (G, V) is said to have a one-dimensional quotient if it is a PV of rank one.

Remark 2.3.2.

- a) The classification of multiplicity free spaces with a one dimensional quotient has been obtained by the author ([27]).
- b) It can be shown that if (G, V) is a PV of rank one, then the polynomial Δ_0 is the unique fundamental relative invariant of (G, V) . More precisely a PV (G, V) is of rank one if and only if it has a unique fundamental relative invariant (see [27]). Hence in the notations of section 2.2 we have $k = 0$, in other words Δ_0 is the unique fundamental G relative invariant among the B relative invariants $\Delta_0, \Delta_1, \dots, \Delta_r$.

We give now some examples of MF -spaces with a one dimensional quotient.

Example 2.3.3. : PV 's of commutative parabolic type (for details we refer to [23], [28] is also relevant).

Let $\tilde{\mathfrak{g}}$ be a simple complex Lie algebra. Assume we are given a 3-grading of $\tilde{\mathfrak{g}}$:

$$\tilde{\mathfrak{g}} = V^- \oplus \mathfrak{g} \oplus V^+.$$

*It must be remarked that if (G, V) is also multiplicity free, then the rank as a PV is not at all the same as the rank as a MF space.

Then \mathfrak{g} is a reductive Lie subalgebra and it is well known that the representation (\mathfrak{g}, V^+) is prehomogeneous (here \mathfrak{g} acts on V^+ via the bracket). Let \tilde{G} be the adjoint group of $\tilde{\mathfrak{g}}$ and let G be the connected subgroup of \tilde{G} whose Lie algebra is \mathfrak{g} . Then the space (G, V^+) is multiplicity free. Moreover such a space has a one dimensional quotient if and only if it is a regular PV. Up to local isomorphism one obtains the following list:

- 1) $(SL(n, \mathbb{C}) \times SL(n, \mathbb{C}) \times \mathbb{C}^*, M_n(\mathbb{C}))$ acting by $(g_1, g_2, t).x = tg_1xg_2^{-1}$ $g_1, g_2 \in SL(n, \mathbb{C}), t \in \mathbb{C}^*, x \in M_n(\mathbb{C})$; here $\Delta_0(x) = \det(x)$.
- 2) $(O(n, \mathbb{C}) \times \mathbb{C}^*, \mathbb{C}^n)$ with the natural action. Here $\Delta_0(x) = Q(x) = \sum_{i=1}^{i=n} x_i^2$.
- 3) $(GL(n, \mathbb{C}), \text{Sym}_n(\mathbb{C}))$, where $\text{Sym}_n(\mathbb{C})$ denotes the $n \times n$ symmetric matrices, with the action $g.x = gx^tg$. Then $\Delta_0(x) = \det(x)$.
- 4) $(GL(n, \mathbb{C}), \text{Skew}_n(\mathbb{C}))$, n even, where $\text{Skew}_n(\mathbb{C})$ denotes the $n \times n$ skew-symmetric matrices, with the action $g.x = gx^tg$. Then $\Delta_0(x) = Pf(x)$, where $Pf(x)$ denotes the pfaffian of the even skew-symmetric matrix x .
- 5) $(E_6 \times \mathbb{C}^*, \mathbb{C}^{27})$ (the irreducible 27-dimensional representation of E_6). The fundamental relative invariant is of degree 3, it is known as the Freudenthal cubic.

Example 2.3.4. $(GL(2) \times Sp(n), \mathbb{C}^2 \otimes \mathbb{C}^{2n})$ (tensor product of the natural representations). Here the action is given by

$$(g_1, g_2).X = g_2X({}^t g_1), \quad g_1 \in SL(2), g_2 \in Sp(n), X \in M_{2n,2}$$

The relative invariant Δ_0 is given by $Pf({}^t X J X)$ where $J = \begin{pmatrix} 0 & Id_n \\ -Id_n & 0 \end{pmatrix}$, and where $Pf(\cdot)$ is the pfaffian of a 2×2 skew symmetric matrix. The rank is equal to 3 and it is a regular PV. For details see [11] (case 11.6) and [27] (case 4.1.7).

Example 2.3.5. $(GL(n) \times GL(n-1), M_{n,1} \oplus M_{n,n-1})$. The action is given by

$$(g_1, g_2)(v, x) = (g_1v, g_1xg_2^{-1}), \quad g_1 \in GL(n), g_2 \in GL(n-1), v \in M_{n,1}, x \in M_{n,n-1}.$$

The relative invariant Δ_0 is given by $\Delta_0(x) = \det(v; x)$ where $(v; x)$ is the $n \times n$ matrix obtained by putting the column vector v left to the $n \times n - 1$ matrix x . The rank is equal to $2n - 1$ and it is a regular PV. For details see [27] (case 4.2.5) and [3] (case 4.2.4).

3. ALGEBRAS OF DIFFERENTIAL OPERATORS

From now on we suppose that (G, V) is a MF space with a one dimensional quotient.

3.1. Gradings and Bernstein-Sato polynomials.

Recall that we denote by $\Delta_0, \Delta_1, \dots, \Delta_r$ the fundamental relative invariants under a fixed Borel subgroup B of G . As the space has a one dimensional quotient, Δ_0 is the unique polynomial among them which is relatively invariant under G (this means that $k = 0$ in the notations of section 2.2). We also set $\mathcal{O} = \{x \in V \mid \Delta_0(x) \neq 0\}$.

Of course the Euler operator E on V is defined for $P \in \mathbb{C}[V]$ by $EP(x) = \frac{\partial}{\partial t} P(tx)_{t=1} = P'(x)x$ is invariant by any element in $GL(E)$.

Once and for all we also define the following two elements in $D(V)$:

$$X = \Delta_0 \quad (\text{multiplication by } \Delta_0), \quad Y = \Delta_0^*(\partial).$$

The operator

$$X^{-1} \quad (\text{multiplication by } \Delta_0^{-1})$$

which belongs to $D(\mathcal{O})$ will also play an important role. From the definition of the G action on $\mathbb{C}[V]$ and on $D(V)$ we have

$$g.X = \lambda_0(g^{-1})X, \quad g.X^{-1} = \lambda_0(g)X^{-1}, \quad g.Y = \lambda_0(g)Y \quad (3-1-1)$$

and hence $X, Y \in D(V)^{G'}$ and $X^{-1} \in D(\mathcal{O})^{G'}$.

Let us now introduce the following notations that we will use in the rest of the paper:

$$\mathcal{T} = D(\mathcal{O})^{G'}, \quad \mathcal{T}_0 = D(V)^G = D(\mathcal{O})^G$$

(the last equality comes from Theorem 2.2.6). Remember that \mathcal{T}_0 is a polynomial algebra in $r+1$ variables (Theorem 2.2.9). We have the following inclusions:

$$\mathcal{T}_0 = D(V)^G = D(\mathcal{O})^G \subset D(V)^{G'} \subset \mathcal{T} = D(\mathcal{O})^{G'}.$$

An element D in \mathcal{T} is said to be of degree m if $[E, D] = mD$. As differential operators in \mathcal{T} have coefficients which are fractions whose denominators are homogeneous (powers of Δ_0), it is clear that \mathcal{T} is graded by its homogeneous components. But on the other hand any homogeneous element D in \mathcal{T} preserves the G' -isotypic components $W_{\mathbf{a}} = \bigoplus_{n \in \mathbb{N}} \Delta_0^n V_{\mathbf{a}}$ (see Proposition 2.2.8). Therefore an homogeneous element D maps $\Delta_0^n V_{\mathbf{a}}$ on $\Delta_0^{n+j} V_{\mathbf{a}}$ for some j and hence only multiples of d_0 (the degree of Δ_0) occur as homogeneous degrees in \mathcal{T} . If we define, for $p \in \mathbb{Z}$, $\mathcal{T}_p = \{D \in \mathcal{T} \mid [E, D] = pd_0D\}$, then

$$\mathcal{T} = \bigoplus_{p \in \mathbb{Z}} \mathcal{T}_p \quad (3-1-2)$$

(At this point it is not completely evident that the two definitions of \mathcal{T}_0 coincide, that is that $D(V)^G = \{D \in \mathcal{T} \mid [E, D] = 0\}$. This will be a consequence of the proof of Proposition 3.1.6 below).

Similarly if we define

$$D(V)_p^{G'} = \{D \in D(V)^{G'} \mid [E, D] = pd_0D\},$$

we have $D(V)^{G'} = \bigoplus_{p \in \mathbb{Z}} D(V)_p^{G'}$.

Definition 3.1.1. For $\mathbf{a} = (a_0, a_1, \dots, a_r)$ and $p \in \mathbb{N}$, we define $\mathbf{a} + p = (a_0 + p, a_1, \dots, a_r)$. Then if $D \in \mathcal{T}_p$, the Schur Lemma ensures that if $P \in V_{\mathbf{a}}$ we have $DP = b_D(\mathbf{a})X^pP$ where $b_D(\mathbf{a}) \in \mathbb{C}$. It is easy to see that b_D is a polynomial in the variables (a_0, a_1, \dots, a_r) (see for example [16], proof of Corollary 4.4). This polynomial is called the Bernstein-Sato polynomial of D .

Example 3.1.2. Relations (3-1-1) imply that $X \in \mathcal{T}_1$, $X^{-1} \in \mathcal{T}_{-1}$ and $Y \in \mathcal{T}_{-1}$. And of course $E \in \mathcal{T}_0$. Obviously, from the definition, we have $b_X(\mathbf{a}) = b_{X^{-1}}(\mathbf{a}) = 1$, $b_E(\mathbf{a}) = d_0a_0 + d_1a_1 + \dots + d_ra_r =$ the degree of $V_{\mathbf{a}}$ (recall that d_i is the degree of Δ_i). From (3-1-1) we obtain that $Y \in \mathcal{T}_{-1}$. The computation of b_Y is more difficult. However it is known in the case of PV 's of commutative parabolic type (see example 2.3.3). In this case, for $\mathbf{X} = (X_0, X_1, \dots, X_r)$ it is given by

$$b_Y(\mathbf{X}) = c \prod_{j=0}^r (X_0 + \dots + X_j + j \frac{d}{2}) \quad (3-1-3)$$

where the constant c can be made explicit (see [4], Théorème 3.19) and where $\frac{d}{2} = \frac{\dim(V) - d_0}{(d_0 - 1)d_0}$. This explicit computation of the polynomial b_Y in the particular case of PV 's of commutative parabolic type has been obtained by several authors, using distinct methods (see [4], [8], [17],

[36]). The constant d is the same as the constant d which is familiar to specialists of Jordan algebras.

The following Lemma is obvious, but useful.

Lemma 3.1.3.

Let $D_1, D_2 \in \mathcal{T}_p$. Then $D_1 = D_2$ if and only if $b_{D_1} = b_{D_2}$.

Definition 3.1.4. The automorphism τ of $\mathcal{T} = D(\mathcal{O})^{G'}$ is defined by

$$\forall D \in \mathcal{T}, \tau(D) = XDX^{-1}$$

Proposition 3.1.5.

The algebra \mathcal{T}_0 is stable under τ and for any $D \in \mathcal{T}_0$ we have

$$XD = \tau(D)X \quad (3-1-4)$$

$$DY = Y\tau(D) \quad (3-1-5)$$

Proof. By definition $\mathcal{T}_0 = D(V)^G$. From relations (3-1-1) we see that if D is G -invariant so is $\tau(D)$. Obviously $\tau(D) \in D(\mathcal{O})^G$. But $D(\mathcal{O})^G = D(V)^G$ by Theorem 2.2.6, hence \mathcal{T}_0 is τ -stable. Relation (3-1-4) is just the definition of τ . We will now prove that (3-1-5) holds on each subspace $V_{\mathbf{a}}$. Let b_D be the Bernstein-Sato polynomial of D . Then an easy calculation shows that the left and right side of (3-1-5) act on $V_{\mathbf{a}}$ by $b_D(\mathbf{a}-1)b_Y(\mathbf{a})X^{-1}$. Then Lemma 3.1.3 implies (3-1-5). \square

Let us denote by $\mathcal{T}_0[X, Y]$ the subalgebra of \mathcal{T} generated by \mathcal{T}_0 , X and Y . From the preceding Proposition and from the fact that XY and YX belong to \mathcal{T}_0 we know that any element $D \in \mathcal{T}_0[X, Y]$ can be written as a finite sum $D = \sum_{p, q \in \mathbb{N}} a_{p, q} X^p Y^q$ with $a_{p, q} \in \mathcal{T}_0$. Similarly, let $\mathcal{T}_0[X, X^{-1}]$ denote the subalgebra of \mathcal{T} generated by \mathcal{T}_0 , X and X^{-1} . Also any element D in $\mathcal{T}_0[X, X^{-1}]$ can be written as a finite sum $D = \sum_{p \in \mathbb{Z}} a_p X^p$. The following Proposition shows that $D(V)^{G'} = \mathcal{T}_0[X, Y]$ and that $\mathcal{T} = D(\mathcal{O})^{G'} = \mathcal{T}_0[X, X^{-1}]$ and makes the gradings more precise.

Proposition 3.1.6.

1) We have

$$D(V)^{G'} = \mathcal{T}_0[X, Y] = (\bigoplus_{p \in \mathbb{N}^*} \mathcal{T}_0 Y^p \bigoplus) \bigoplus \mathcal{T}_0 \bigoplus (\bigoplus_{p \in \mathbb{N}^*} \mathcal{T}_0 X^p)$$

(in particular $D(V)_p^{G'} = \mathcal{T}_0 X^p$ if $p \geq 0$, and $D(V)_p^{G'} = \mathcal{T}_0 Y^{-p}$ if $p < 0$). Equivalently we have

$$D(V)^{G'} = \mathcal{T}_0[X, Y] = (\bigoplus_{p \in \mathbb{N}^*} Y^p \mathcal{T}_0 \bigoplus) \bigoplus \mathcal{T}_0 \bigoplus (\bigoplus_{p \in \mathbb{N}} X^p \mathcal{T}_0).$$

2) We have $\mathcal{T} = D(\mathcal{O})^{G'} = \mathcal{T}_0[X, X^{-1}] = \bigoplus_{p \in \mathbb{Z}} \mathcal{T}_0 X^p = \bigoplus_{p \in \mathbb{Z}} X^p \mathcal{T}_0$.

3) Any element D in $\mathcal{T}_0[X, Y]$ can be written uniquely in the form

$$D = \sum_{i>0} u_i Y^i + \sum_{i \geq 0} v_i X^i \text{ or } D = \sum_{i>0} Y^i u_i + \sum_{i \geq 0} X^i v_i \text{ (finite sums)}$$

with $u_i, v_i \in \mathcal{T}_0$.

Any element $D \in \mathcal{T}$ can be written uniquely in the form

$$D = \sum_{i \in \mathbb{Z}} u_i X^i \text{ or } D = \sum_{i \in \mathbb{Z}} X^i u_i \text{ (finite sums)}$$

with $u_i \in \mathcal{T}_0$.

Proof. 1) For the moment we define \mathcal{T}_0 by $\mathcal{T}_0 = D(V)^G$. From Proposition 2.2.8 we know that the decomposition of $\mathbb{C}[V]$ into G' -isotypic components is given by

$$\mathbb{C}[V] = \bigoplus_{\tilde{\mathbf{a}} \in \mathbb{N}^r} U_{\tilde{\mathbf{a}}} \text{ where } U_{\tilde{\mathbf{a}}} = \bigoplus_{a_0 \in \mathbb{N}} \Delta_0^{a_0} V_{\tilde{\mathbf{a}}} \text{ and } \tilde{\mathbf{a}} = (0, a_1, \dots, a_r).$$

We will now use the technique of Howe and Umeda ([11]) which we have already mentioned before Theorem 2.2.9. As $\mathbb{C}[V] \otimes \mathbb{C}[V^*]$ is G' -isomorphic to $D(V)$, each subspace $\Delta^{a_0} V_{\tilde{\mathbf{a}}} \otimes (\Delta^{b_0} V_{\tilde{\mathbf{a}}})^*$ will give rise to a unique G' -invariant differential operator $R_{a_0, b_0, \tilde{\mathbf{a}}}$. Then by the same arguments as in Remark 2.2.10, it is easy to see that $R_{a_0, b_0, \tilde{\mathbf{a}}} = \Delta_0(x)^{a_0} R_{0,0,\tilde{\mathbf{a}}} \Delta_0^*(\partial)^{b_0} = X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0}$. The elements $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0}$ ($a_0, b_0 \in \mathbb{N}, \tilde{\mathbf{a}} \in \mathbb{N}^r$) form a vector basis of $D(V)^{G'}$. Remark now that $R_{0,0,\tilde{\mathbf{a}}}$ is in $D(V)^G = \mathcal{T}_0$. Then from Proposition 3.1.5, we get $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0} = \tau^{a_0}(R_{0,0,\tilde{\mathbf{a}}}) X^{a_0} Y^{b_0}$ and $\tau^{a_0}(R_{0,0,\tilde{\mathbf{a}}}) \in \mathcal{T}_0$. If now $a_0 \leq b_0$, then $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0} = R Y^{b_0 - a_0}$, where $R = \tau^{a_0}(R_{0,0,\tilde{\mathbf{a}}}) X^{a_0} Y^{a_0} \in \mathcal{T}_0$. If $a_0 > b_0$, then $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0} = R X^{a_0 - b_0}$ where $R = \tau^{a_0}(R_{0,0,\tilde{\mathbf{a}}}) \tau^{a_0 - b_0}(X^{b_0} Y^{b_0}) \in \mathcal{T}_0$. The first decomposition in assertion 1) is proved. The second decomposition is a consequence of relations (3-1-4) and (3-1-5).

2) A slight extension of (2-2-2) shows that $\mathbb{C}[\mathcal{O}] \otimes \mathbb{C}[V^*]$ is G -isomorphic to $D(\mathcal{O})$ through the map $\varphi \otimes f \mapsto \varphi f(\partial)$. Then the same proof as in 1) above shows that the elements $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{b_0}$ ($a_0 \in \mathbb{Z}, b_0 \in \mathbb{N}, \tilde{\mathbf{a}} \in \mathbb{N}^r$) form a vector basis of $D(\mathcal{O})^{G'} = \mathcal{T}$. Consider now an element $D \in \mathcal{T}$ such that $[E, D] = 0$. Then necessarily D is a linear combination of elements of the form $X^{a_0} R_{0,0,\tilde{\mathbf{a}}} Y^{a_0}$ with $a_0 \in \mathbb{N}$. Then, as announced previously, the two definitions of \mathcal{T}_0 coincide ($\mathcal{T}_0 = D(V)^G$ and $\mathcal{T}_0 = \{D \in \mathcal{T}, [E, D] = 0\}$). Now if $D \in \mathcal{T}_p$, then $D = D X^{-p} X^p$ and $D X^{-p} \in \mathcal{T}_0$. Hence $\mathcal{T}_p = \mathcal{T}_0 X^p = X^p \mathcal{T}_0$.

Assertion 3) is then obvious. □

Remark 3.1.7. The inclusion $D(V)^{G'} \subset D(\mathcal{O})^{G'}$ is obviously strict ($X^{-1} \in D(\mathcal{O})^{G'} \setminus D(V)^{G'}$), but the preceding results shows that these two graded algebras have the same "positive part" ($\bigoplus_{p \in \mathbb{N}} \mathcal{T}_0 X^p$).

The following proposition whose proof is straightforward shows that all the Bernstein-Sato polynomials are known if one knows the Bernstein-Sato polynomials of Y and of the elements of \mathcal{T}_0 .

Proposition 3.1.8.

Let $D = D_0 X^n$ ($n \in \mathbb{Z}$), resp. $D = D_0 Y^n$ ($n \in \mathbb{N}^*$), $D_0 \in \mathcal{T}_0$, be generic homogeneous elements in $\mathcal{T} = \mathcal{T}_0[X, X^{-1}]$ or $\mathcal{T}_0[X, Y]$. Then

$$b_D(\mathbf{a}) = b_{D_0}(\mathbf{a} + n), \text{ resp. } b_D(\mathbf{a}) = b_{D_0}(\mathbf{a} - n) b_Y(\mathbf{a}) b_Y(\mathbf{a} - 1) \dots b_Y(\mathbf{a} - n + 1).$$

3.2. The Harish-Chandra isomorphism and the center of \mathcal{T} .

The aim of this subsection is to describe $\mathcal{T}_0 = D(V)^G$ as a module over the center of \mathcal{T} . For this we will use the Harish-Chandra isomorphism for MF spaces due to F. Knop.

Let (G, V) be a MF space with a one dimensional quotient. Let B be a fixed Borel subgroup of G . Remember that (B, V) is a PV . Recall also that we denote by $\Delta_0, \Delta_1, \dots, \Delta_r$ the set of fundamental relative invariants of (B, V) and that Δ_0 is the unique fundamental relative invariant under G . We denote by d_i (resp. λ_i) the degree (resp. the infinitesimal character) of Δ_i . Let \mathfrak{b} be the Lie algebra of B , let $\mathfrak{t} \subset \mathfrak{b}$ be a Cartan subalgebra of \mathfrak{g} and let Σ be the

set of roots of the pair $(\mathfrak{g}, \mathfrak{t})$. Denote by W the Weyl group of Σ . Denote by Σ^+ the set of positive roots such that $\mathfrak{b} = \mathfrak{t} + \sum_{\alpha \in \Sigma^+} \mathfrak{g}^\alpha$. Let $\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} \alpha$. We define

$$\mathfrak{a}^* = \bigoplus_{i=0}^r \mathbb{C}\lambda_i \subset \mathfrak{t}^* \text{ and } A = \mathfrak{a}^* + \rho \subset \mathfrak{t}^*.$$

Let $\mathcal{Z}(\mathfrak{g})$ be the center of the enveloping algebra of \mathfrak{g} . Denote by $\mathbb{C}[\mathfrak{t}^*]^W$ the W -invariant polynomials on \mathfrak{t}^* . One knows that the classical Harish-Chandra isomorphism is an isomorphism $H : \mathcal{Z}(\mathfrak{g}) \rightarrow \mathbb{C}[\mathfrak{t}^*]^W$ which can be computed the following way. For any $\lambda \in \mathfrak{t}^*$, let V_λ be the irreducible highest weight module with highest weight λ . It is well known that $\mathcal{Z}(\mathfrak{g})$ acts by scalar multiplication on V_λ . The scalar by which an element $z \in \mathcal{Z}(\mathfrak{g})$ acts on V_λ is precisely $H(z)(\lambda + \rho)$.

The natural representation of G on $\mathbb{C}[V]$ extends to a representation of the enveloping algebra $\mathcal{U}(\mathfrak{g})$ on the same space $\mathbb{C}[V]$. Hence $z \in \mathcal{Z}(\mathfrak{g})$ acts on $V_{\mathbf{a}}$ by the scalar $H(z)(-\lambda_{\mathbf{a}} + \rho)^\dagger$ where $\lambda_{\mathbf{a}} = \sum_{i=0}^r a_i \lambda_i$ (remember that $\mathbf{a} = (a_0, \dots, a_r)$). Conversely if $\lambda = a_0 \lambda_0 + \dots + a_r \lambda_r$ we define $\mathbf{a}_\lambda = (a_0, \dots, a_r) \in \mathbb{C}^{r+1}$. By abuse of notation if b_D is the Bernstein-Sato polynomial of $D \in \mathcal{T}_0$, we set $b_D(\lambda) = b_D(\mathbf{a}_\lambda)$. Hence $H(z)(-\lambda + \rho) = b_D(\lambda)$.

On the other hand any $D \in D(V)^G = \mathcal{T}_0$ acts on each $V_{\mathbf{a}}$ by the scalar $b_D(\mathbf{a})$, where $b_D(\mathbf{a})$ is the Bernstein-Sato polynomial of D . This allows us to define the map:

$$\begin{aligned} h : D(V)^G &\longrightarrow \mathbb{C}[A] \\ D &\longmapsto h(D) : -\lambda + \rho \longmapsto h(D)(-\lambda + \rho) = b_D(\lambda) \end{aligned}$$

where $\mathbb{C}[A]$ denotes the algebra of polynomials on the affine space $A = \mathfrak{a}^* + \rho \subset \mathfrak{t}^*$.

Let $\pi(z)$ be the operator in $D(V)^G$ which represents the action of z on $\mathbb{C}[V]$ and let $r : \mathbb{C}[\mathfrak{t}^*]^W \rightarrow \mathbb{C}[A]$ be the restriction homomorphism. It is clear from the definitions that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{Z}(\mathfrak{g}) & \xrightarrow{H} & \mathbb{C}[\mathfrak{t}^*]^W \\ \pi \downarrow & & \downarrow r \\ D(V)^G & \xrightarrow{h} & \mathbb{C}[A] \end{array}$$

Theorem 3.2.1.

(Knop, see [16] Th. 4.8 and Corollary 4.9 or [1], Th. 9.2.1)

The homomorphism h is injective and there exists a finite group W_0 (sometimes called the little Weyl group) which is a subgroup of the stabilizer of A in W , such that the image of h is $\mathbb{C}[A]^{W_0}$. Hence h is an isomorphism between $D(V)^G$ and $\mathbb{C}[A]^{W_0}$. The isomorphism h is called the Harish-Chandra isomorphism for the MF space (G, V) . Moreover W_0 acts as a reflection group on \mathfrak{a}^ .*

Let us see what is the automorphism of $\mathbb{C}[A]^{W_0}$ which corresponds to the action of τ on $D(V)^G$ through the Harish-Chandra isomorphism h . Let $D \in D(V)^G$. Then $h(\tau(D))(-\lambda + \rho) = h(XDX^{-1})(-\lambda + \rho) = b_{XDX^{-1}}(\lambda) = b_D(\lambda - \lambda_0)$. This calculation proves of course that $\mathbb{C}[A]^{W_0}$ is stable under $P(\lambda + \rho) \mapsto P((\lambda - \lambda_0) + \rho)$. Therefore we make the following definition.

[†]The change of sign is due to the fact that we consider here characters of relative invariants instead of highest weights.

Definition 3.2.2. By abuse of notation τ will also denote the automorphism of $\mathbb{C}[A]^{W_0}$ which is defined by $\tau(P)(\lambda + \rho) = P((\lambda - \lambda_0) + \rho)$ ($P \in \mathbb{C}[A]^{W_0}$). Let $\mathbb{C}[A]^{W_0, \tau}$ denote the set of elements in $\mathbb{C}[A]^{W_0}$ which are invariant under τ .

Proposition 3.2.3.

Let $\mathcal{Z}(\mathcal{T})$ be the center of $\mathcal{T} = D(\mathcal{O})^{G'}$. Then $\mathcal{Z}(\mathcal{T})$ is also the center of $\mathcal{T}_0[X, Y] = D(V)^{G'}$. Moreover the following assertions are equivalent:

- i) $D \in \mathcal{Z}(\mathcal{T})$
- ii) $D \in \mathcal{T}_0$ and $\tau(D) = D$ (i.e. D commutes with X).
- iii) $D \in \mathcal{T}_0$ and the Bernstein-Sato polynomial $b_D(a_0, a_1, \dots, a_r)$ does not depend on a_0 .
- iv) $D \in \mathcal{T}_0$ and D commutes with Y .
- v) $D \in \mathcal{T}_0$ and $h(D) \in \mathbb{C}[A]^{W_0, \tau}$.

Proof. i) \Rightarrow ii): Let $D \in \mathcal{Z}(\mathcal{T})$. Then $[E, D] = 0$, hence $D \in \mathcal{T}_0$, and $[D, X] = 0$.

ii) \Rightarrow iii): Let $D \in \mathcal{T}_0$. If $XD = DX$ then, from the definitions we have $b_{XD}(a_0, a_1, \dots, a_r) = b_D(a_0, a_1, \dots, a_r) = b_{DX}(a_0, a_1, \dots, a_r) = b_D(a_0 + 1, a_1, \dots, a_r)$, hence $b_D(a_0, a_1, \dots, a_r)$ does not depend on a_0 .

iii) \Rightarrow i): Suppose that for $D \in \mathcal{T}_0$, the Bernstein-Sato polynomial does not depend on a_0 . Then the elements XD and DX in \mathcal{T}_1 have the same Bernstein-Sato polynomial. Hence $XD = DX$ (Lemma 3.1.3). Then from Proposition 3.1.6 2) we see that $D \in \mathcal{Z}(\mathcal{T})$.

iii) \Rightarrow iv): Let $D \in \mathcal{T}_0$ such that b_D does not depend on a_0 . Then $b_{DY}(a_0, a_1, \dots, a_r) = b_D(a_0 - 1, a_1, \dots, a_r)b_Y(a_0, a_1, \dots, a_r) = b_D(a_0, a_1, \dots, a_r)b_Y(a_0, a_1, \dots, a_r) = b_{YD}(a_0, a_1, \dots, a_r)$. Hence $DY = YD$.

iv) \Rightarrow iii): If $DY = YD$, then $b_{DY}(a_0, a_1, \dots, a_r) = b_{D(a_0 - 1, a_1, \dots, a_r)}b_Y(a_0, a_1, \dots, a_r) = b_{YD}(a_0, a_1, \dots, a_r) = b_Y(a_0, a_1, \dots, a_r)b_D(a_0, a_1, \dots, a_r)$. Hence b_D does not depend on a_0 .

The equivalence of iii) and v) is obvious as $h(D)(-\lambda + \rho) = b_D(\lambda)$.

From ii) we obtain that $\mathcal{Z}(\mathcal{T})$ is also the center of $\mathcal{T}_0[X, Y]$. □

Remark 3.2.4. As a consequence of the preceding Proposition it is worthwhile noticing that if $D \in D(V)^{G'}$ (or $D \in \mathcal{T}$) commutes with two operators among (X, E, Y) , then D commutes with the third one. This is a well known property if (X, E, Y) is an \mathfrak{sl}_2 -triple. But we know from [12] that except if Δ_0 is quadratic or linear the Lie algebra generated by (X, E, Y) is infinite dimensional. We will see that the associative algebra generated by (X, E, Y) over $\mathcal{Z}(\mathcal{T})$ is "similar" to $\mathcal{U}(\mathfrak{sl}_2(\mathcal{Z}(\mathcal{T})))$, see Theorem 4.2.2 below.

Define a linear form μ on \mathfrak{a}^* by

$$\mu(a_0\lambda_0 + \dots + a_r\lambda_r) = \sum_{i=0}^r a_i d_i = b_E(\mathbf{a}) \quad (\mathbf{a} = (a_0, \dots, a_r) \in \mathbb{C}^{r+1})$$

(μ is the *degree form*, as its value on $\mathbf{a} = (a_0, \dots, a_r) \in \mathbb{N}^{r+1}$ is equal to the degree of the polynomials in $V_{\mathbf{a}}$. Define also

$$\mathcal{M} = \{\lambda \in \mathfrak{a}^* \mid \mu(\lambda) = 0\} \text{ and } M = \mathcal{M} + \rho \subset A.$$

Note that $M = \{\lambda + \rho \in A \mid h(E)(-\lambda + \rho) = 0\}$. As $h(E)$ is W_0 -invariant, so is the set M . Set

$$I(M) = \{P \in \mathbb{C}[A]^{W_0} \mid P|_M = 0\}.$$

The key lemma is the following.

Lemma 3.2.5.

We have $I(M) = \mathbb{C}[A]^{W_0}h(E)$ and

$$\mathbb{C}[A]^{W_0} = \mathbb{C}[A]^{W_0, \tau} \oplus I(M).$$

Proof. Let $P \in I(M)$. It is a polynomial on the affine subspace $A \subset \mathfrak{t}^*$ which vanishes on M which is the set of zeros of the irreducible polynomial $h(E)$. Therefore $P = h(E)Q$. As P and $h(E)$ are W_0 -invariant, so is also the polynomial Q . Hence $I(M) \subset \mathbb{C}[A]^{W_0}h(E)$. As the reverse inclusion is obvious we get $I(M) = \mathbb{C}[A]^{W_0}h(E)$.

Let $F = \mathbb{C}\lambda_0 \subset \mathfrak{a}^*$. As obviously $\mathfrak{a}^* = \mathcal{M} \oplus F$, we have $A = M \oplus F$. Remember that $\mathfrak{t} = \mathfrak{c} \oplus \mathfrak{t}'$ where \mathfrak{c} is the center of \mathfrak{g} . The infinitesimal character λ_0 is a character of \mathfrak{g} , and is therefore trivial on $\mathfrak{t}' \subset \mathfrak{g}'$. As any $w_0 \in W_0$ fixes pointwise the center \mathfrak{c} of \mathfrak{g} , we see that F is pointwise fixed by W_0 .

Let $Q \in \mathbb{C}[M]^{W_0}$. Define

$$\tilde{Q}(m + f) = Q(m), \quad \text{for all } m \in M, f \in F.$$

From the preceding discussion we obtain that \tilde{Q} is W_0 -invariant, in other words $\tilde{Q} \in \mathbb{C}[A]^{W_0}$. But in fact \tilde{Q} is also τ -invariant: $\tau(\tilde{Q})(m + f) = \tilde{Q}(m + f - \lambda_0) = Q(m) = \tilde{Q}(m + f)$. Hence $\tilde{Q} \in \mathbb{C}[A]^{W_0, \tau}$, in other words any W_0 -invariant polynomial on M can be extended to a (W_0, τ) -invariant polynomial on A . This extension is in fact unique: for any τ -invariant extension $\tilde{\tilde{Q}}$ of Q we have $\tilde{\tilde{Q}}(m + x\lambda_0) = \tilde{\tilde{Q}}(m + (x+1)\lambda_0)$ and hence $\tilde{\tilde{Q}} = \tilde{Q}$. Hence we have proved that the restriction map:

$$\begin{array}{ccc} \mathbb{C}[A]^{W_0, \tau} & \longrightarrow & \mathbb{C}[M]^{W_0} \\ P & \longmapsto & P|_M \end{array}$$

is bijective (and therefore $\mathbb{C}[A]^{W_0, \tau} \cap I(M) = \{0\}$) and the inverse map is $Q \longmapsto \tilde{Q}$. Now for $P \in \mathbb{C}[A]^{W_0}$ we can write:

$$P = \widetilde{P|_M} + (P - \widetilde{P|_M}).$$

From the discussion above we have $\widetilde{P|_M} \in \mathbb{C}[A]^{W_0, \tau}$, and $(P - \widetilde{P|_M}) \in I(M)$. □

Theorem 3.2.6.

- 1) $\mathcal{T}_0 = D(V)^G = \mathcal{Z}(\mathcal{T}) \oplus E\mathcal{T}_0$
- 2) Any element $H \in D(V)^G$ can be uniquely written in the form

$$H = H_0 + EH_1 + E^2H_2 \cdots + E^kH_k$$

where $H_i \in \mathcal{Z}(\mathcal{T})$, $i = 1, 2, \dots, k \in \mathbb{N}$.

Proof. Through the Harish-Chandra isomorphism h , the algebra $D(V)^G = \mathcal{T}_0$ corresponds to $\mathbb{C}[A]^{W_0}$, the algebra $\mathcal{Z}(\mathcal{T})$ corresponds to $\mathbb{C}[A]^{W_0, \tau}$ and the ideal $E\mathcal{T}_0$ corresponds to $I(M)$. Therefore the first assertion is just the pull back by h of the decomposition obtained in Lemma 3.2.5.

An element $H \in D(V)^G$ can therefore be uniquely written $H = H_0 + EH^1$, with $H_0 \in \mathcal{Z}(\mathcal{T})$, and $H^1 \in \mathcal{T}_0$. By induction we obtain a decomposition $H = H_0 + EH_1 + E^2H_2 \cdots + E^{k-1}H_{k-1} + E^kH^k$ where $H_0, \dots, H_{k-1} \in \mathcal{Z}(\mathcal{T})$, and $H^k \in \mathcal{T}_0$. The process stops because if k is greater than the degree in a_0 of b_H , then necessarily $H^k = 0$ (see Proposition 3.2.3). □

From the preceding Theorem and Proposition 3.1.6 we obtain immediately the following Corollary.

Corollary 3.2.7.

1) Let $D \in \mathcal{T}$, then D can be written uniquely in the form:

$$D = \sum_{k \in \mathbb{Z}, \ell \in \mathbb{N}} H_{k,\ell} E^\ell X^k \text{ or } D = \sum_{k \in \mathbb{Z}, \ell \in \mathbb{N}} H_{k,\ell} X^k E^\ell \text{ (finite sums)}$$

where $H_{k,\ell} \in \mathcal{Z}(\mathcal{T})$

2) Let $D \in \mathcal{T}_0[X, Y]$, then D can be written uniquely in the form:

$$\begin{aligned} D &= \sum_{k \in \mathbb{N}^*, \ell \in \mathbb{N}} H_{k,\ell} E^\ell Y^k + \sum_{r \in \mathbb{N}, s \in \mathbb{N}} H'_{r,s} E^s X^r \text{ (finite sum)} \text{ or} \\ D &= \sum_{k \in \mathbb{N}^*, \ell \in \mathbb{N}} H_{k,\ell} Y^k E^\ell + \sum_{r \in \mathbb{N}, s \in \mathbb{N}} H'_{r,s} X^r E^s \text{ (finite sum)} \end{aligned}$$

where $H_{k,\ell}, H'_{r,s} \in \mathcal{Z}(\mathcal{T})$

Corollary 3.2.8. Let $P \in \mathbb{C}[A]^{W_0}$. Then P can be uniquely written in the form

$$P(-\lambda + \rho) = \sum_{i=0}^p \alpha_i(-\lambda + \rho) (a_0 d_0 + a_1 d_1 + \cdots + a_r d_r)^i$$

where $\alpha_i \in \mathbb{C}[A]^{W_0, \tau}$ and where $\lambda = a_0 \lambda_0 + a_1 \lambda_1 + \cdots + a_r \lambda_r \in \mathfrak{a}^*$.

Proof. As $h(E)(-\lambda + \rho) = a_0 d_0 + a_1 d_1 + \cdots + a_r d_r$, the preceding decomposition is just the image through the Harish-Chandra isomorphism of the decomposition in Theorem 3.2.6 2). \square

Remark 3.2.9. Let us make some remarks about the W_0 -action on A . In fact it is easy to see that as W_0 stabilizes the affine space $A = \mathfrak{a}^* + \rho$ it also stabilizes \mathfrak{a}^* (this is implicit in Theorem 3.2.1). Moreover if we denote by 0_ρ the barycenter of the W_0 -orbit of ρ , then 0_ρ is a fixed point of the W_0 -action on A which is in M . As $\mathbb{C}[A]^{W_0} = \mathbb{C}[\mathfrak{a}^* + \rho]^{W_0} = \mathbb{C}[\mathfrak{a}^* + 0_\rho]^{W_0} \simeq \mathbb{C}[\mathfrak{a}^*]^{W_0}$, and as $\mathcal{T}_0 = D(V)^G \simeq \mathbb{C}[A]^{W_0}$ is a polynomial algebra in $r + 1$ variables by Theorem 2.2.9, the group W_0 acts as a reflection group on \mathfrak{a}^* by the Shephard-Todd-Chevalley Theorem (this is a part of Knop's argument for Theorem 3.2.1). Hence by the Theorem of Chevalley, the $r + 1$ algebraically independent generators of the algebra $\mathbb{C}[A]^{W_0} \simeq \mathbb{C}[\mathfrak{a}^*]^{W_0}$ can be chosen to be homogeneous, either as functions on the vector space \mathfrak{a}^* , or as functions on A , for the vector space structure on A defined by taking 0_ρ as origin.

We will now describe more precisely the algebra $\mathcal{Z}(\mathcal{T})$.

Theorem 3.2.10.

1) $\mathcal{Z}(\mathcal{T})$ is a polynomial algebra in r variables. For $D \in \mathcal{T}_0$, let us denote by \overline{D} the projection of D on $\mathcal{Z}(\mathcal{T})$ according to the decomposition $\mathcal{T}_0 = \mathcal{Z}(\mathcal{T}) \oplus E\mathcal{T}_0$. Remember from Theorem 2.2.9 that the set R_0, \dots, R_{r-1}, R_r of Capelli operators associated to the invariants $\Delta_0, \Delta_1, \dots, \Delta_r$ ordered by decreasing degree is a set algebraically independent generators of \mathcal{T}_0 . Then $\{\overline{R_0}, \dots, \overline{R_{r-1}}\}$ is a set of algebraically independent generators of $\mathcal{Z}(\mathcal{T})$.

2) Let D be an element of \mathcal{T}_0 and let b_D be its Bernstein-Sato polynomial. Then the Bernstein-Sato polynomial of \overline{D} is given by

$$b_{\overline{D}}(a_0, a_1, \dots, a_r) = b_D\left(-\frac{(a_1 d_1 + \cdots + a_r d_r)}{d_0}, a_1, \dots, a_r\right).$$

Proof. 1) Let us remark first that $\mathcal{Z}(\mathcal{T})$ is already known to be a polynomial algebra from a result of Knop ([15]). He has proved that for a regular action of a reductive group on a smooth affine variety the center of the ring of invariant differential operators is always a polynomial algebra. We give here a direct proof and obtain some extra information. We know from Proposition 3.2.3 that $\mathcal{Z}(\mathcal{T})$ is isomorphic, through the Harish-Chandra isomorphism h , to $\mathbb{C}[A]^{W_0, \tau}$. From the proof of Lemma 3.2.5 we know that W_0 stabilizes M and that $\mathbb{C}[A]^{W_0, \tau} \simeq \mathbb{C}[M]^{W_0} = (\mathbb{C}[A]^{W_0})_{|_M}$. As W_0 is a reflection group on A (this means that it is generated by the reflections it contains), so is $W_0|_M$. Therefore $\mathbb{C}[M]^{W_0}$ (and hence $\mathcal{Z}(\mathcal{T})$) is a polynomial algebra in $r = \dim M$ variables by Chevalley's Theorem. We know from Remark 2.2.10 d) that $\{R_0, \dots, R_{r-1}, E\}$ is also a set algebraically independent generators of \mathcal{T}_0 , hence $\{h(R_0), \dots, h(R_{r-1}), h(E)\}$ is a set of algebraically independent generators of $\mathbb{C}[A]^{W_0}$. As $h(E)|_M = 0$ we obtain that $\mathbb{C}[M]^{W_0} = \mathbb{C}[h(R_0)|_M, \dots, h(R_{r-1})|_M]$. As the transcendence degree of $\text{Frac}(\mathbb{C}[M]^{W_0})$ over \mathbb{C} is r , the generators $h(R_0)|_M, \dots, h(R_{r-1})|_M$ are algebraically independent. Taking their inverse image under h gives the first assertion of the Theorem.

2) As we have seen the decomposition $\mathcal{T}_0 = \mathcal{Z}(\mathcal{T}) \oplus E\mathcal{T}_0$ is nothing else but the inverse image under h of the decomposition $\mathbb{C}[A]^{W_0} = \mathbb{C}[A]^{W_0, \tau} \oplus I(M)$. Let $D \in \mathcal{T}_0$. From the proof of Lemma 3.2.5 we have $h(\overline{D}) = \widetilde{h(D)|_M}$, where $\widetilde{h(D)|_M}$ is the unique (W_0, τ) -invariant extension to A of $h(D)|_M$. For $\lambda = a_0\lambda_0 + \dots + a_r\lambda_r \in \mathfrak{a}^*$, we have $h(E)(\lambda + \rho) = b_E(-\lambda) = -(a_0d_0 + \dots + a_r d_r) = -\mu(\lambda)$ (the degree form). Remember also that $\mathfrak{a}^* = \mathcal{M} \oplus F$, where $F = \mathbb{C}\lambda_0$, where $\mathcal{M} = \ker(\mu)$. Let us write $\lambda = m_\lambda + \alpha\lambda_0$, according to this decomposition. Then $b_E(\lambda) = \alpha b_E(\lambda_0) = \alpha d_0$. Hence $\alpha = \frac{\mu(\lambda)}{d_0}$ and $m_\lambda = \lambda - \frac{\mu(\lambda)}{d_0}\lambda_0$. Then we obtain:

$$\begin{aligned} b_{\overline{D}}(\lambda) &= h(\overline{D})(-\lambda + \rho) = \widetilde{h(D)|_M}(-\lambda + \rho) = \widetilde{h(D)|_M}(-\lambda + \frac{\mu(\lambda)}{d_0}\lambda_0 - \frac{\mu(\lambda)}{d_0}\lambda_0 + \rho) \\ &= \widetilde{h(D)|_M}(-\lambda + \frac{\mu(\lambda)}{d_0}\lambda_0 + \rho) \\ &= h(D)|_M(-\lambda + \frac{\mu(\lambda)}{d_0}\lambda_0 + \rho) = h(D)(-\lambda + \frac{\mu(\lambda)}{d_0}\lambda_0 + \rho) \\ &= b_D(\lambda - \frac{\mu(\lambda)}{d_0}\lambda_0). \end{aligned}$$

If we translate this into the (a_0, \dots, a_r) -variables we obtain the second assertion. □

Corollary 3.2.11.

Let b_Y be the Bernstein-Sato operator of Y . For any $\ell \in \mathbb{N}$ the element of $\text{End}(\mathbb{C}[V])$ which acts on each space $V_{\mathbf{a}}$ as the scalar multiplication by $b_Y(-\frac{(a_1d_1 + \dots + a_r d_r)}{d_0} + \ell, a_1, \dots, a_r)$ is the differential operator $\overline{X^{1-\ell}YX^\ell} \in \mathcal{Z}(\mathcal{T})$. Moreover, if (G, V^+) is a PV of commutative parabolic type, the differential operators $\overline{X^{1-\ell}YX^\ell}$ ($\ell = 0, 1, \dots, r$) are generators of $\mathcal{Z}(\mathcal{T})$.

Proof. As $b_{X^{1-\ell}YX^\ell}(a_0, \dots, a_r) = b_Y(a_0 + \ell, a_1, \dots, a_r)$, the first assertion is an immediate consequence of Theorem 3.2.10. If (G, V^+) is a PV of commutative parabolic type, we know from Theorem 3.3.1 that the operators $X^{1-\ell}YX^\ell$ ($\ell = 0, \dots, r$) are (algebraically independent) generators of \mathcal{T}_0 . □

3.3. The case of regular PV's of commutative parabolic type.

In the case where (G, V^+) is a regular PV of commutative parabolic type (see Example 2.3.3), we obtain some specific results.

Theorem 3.3.1.

Let (G, V^+) be a regular PV of commutative parabolic type.

- 1) The degree of Δ_0 is equal to $r + 1$ which is the rank of (G, V^+) as a MF space. More generally the degree of Δ_i is equal to $r + 1 - i$.
- 2) For $\ell \in \mathbb{Z}$ set $D_\ell = X^{1-\ell} Y X^\ell$. Then D_0, D_1, \dots, D_r are algebraically independent generators of $\mathcal{T}_0 = D(V^+)^G$ (i.e. $\mathcal{T}_0 = \mathbb{C}[D_0, D_1, \dots, D_r]$).
- 3) We have $\mathcal{T} = D(\Omega^+)^{G'} = \mathbb{C}[X, X^{-1}, Y]$ where $\mathbb{C}[X, X^{-1}, Y]$ is the associative subalgebra of $D(\Omega^+)$ generated by X, X^{-1}, Y .
- 4) We have $\mathcal{T}_0[X, Y] = D(V)^{G'} = \mathbb{C}[X, Y, R_1, \dots, R_r]$ where the R_i 's are the Capelli operators which were introduced before Theorem 2.2.9, and where $\mathbb{C}[X, Y, R_1, \dots, R_r]$ is the associative subalgebra of $D(V^+)$ generated by X, Y, R_1, \dots, R_r .

Proof. 1) This first assertion is proved in [23] (Prop. 2.16 and Lemme 3.7).

2) We need now to use some technical results from the structure theory of commutative PV's of parabolic type. For details see [23] and [28]. We need also results concerning the symmetric space structure of the open G orbit Ω^+ in V^+ , these can be found in [4]. Let \mathfrak{t} be a Cartan subalgebra of \mathfrak{g} , then \mathfrak{t} is also a Cartan subalgebra of $\tilde{\mathfrak{g}}$ (see the notation in Example 2.3.3) and let $\tilde{\Sigma}$ (resp. Σ) be the root system of $(\tilde{\mathfrak{g}}, \mathfrak{t})$ (resp. $(\mathfrak{g}, \mathfrak{t})$). We choose an order on $\tilde{\Sigma}$ such that the roots occurring in V^+ are positive. We know from Prop. 2.9. in [4] that the open G -orbit $\Omega^+ = \{x \in V^+ \mid \Delta_0(x) \neq 0\}$ is a symmetric space G/H where H is the isotropy subgroup of a point $I^+ \in \Omega^+$. The choice of I^+ can be made the following way. It is known that any maximal set of strongly orthogonal long roots occurring in V^+ has $r + 1 = rk(G, V^+)$ elements. There is a canonical way to construct such a maximal set, called the "descent", see [23], Th. 2.7. p.101. If $\{\alpha_0, \alpha_1, \dots, \alpha_r\}$ is such a maximal set of strongly orthogonal long roots, then the element $I^+ = X_{\alpha_0} + X_{\alpha_1} + \dots + X_{\alpha_r}$ is generic (here as usual the X_{α_i} 's are non zero root vectors). Let $\mathfrak{h} = Z_{\mathfrak{g}}(I^+)$ be the Lie algebra of H , and let \mathfrak{q} be the orthogonal complement of \mathfrak{h} in \mathfrak{g} with respect to the Killing form of $\tilde{\mathfrak{g}}$. Let $H_{\alpha_i} \in \mathfrak{t}$ be the co-root of α_i . Set $\mathfrak{a} = \sum_{i=0}^r \mathbb{C}H_{\alpha_i}$. Then \mathfrak{a} is a maximal abelian subspace of \mathfrak{q} ([4], Prop. 5.4) and the dual space \mathfrak{a}^* can be identified with the space of restrictions of the fundamental characters $\lambda_0, \lambda_1, \dots, \lambda_r$ ([4], Lemme 2.5). Hence this definition of \mathfrak{a}^* is coherent with the direct definition of \mathfrak{a}^* given in section 3.2 in the general case ($\mathfrak{a}^* = \sum_{i=0}^r \mathbb{C}\lambda_i$).

For $\lambda \in \mathfrak{t}^*$, we will denote by $\bar{\lambda}$ the restriction of λ to \mathfrak{a} . Through the "classical" Harish-Chandra isomorphism γ for symmetric spaces ([10], Part II, Theorem 4.3) the algebra \mathcal{T}_0 is isomorphic to $S(\mathfrak{a})^{W_R} = \mathbb{C}[\mathfrak{a}^*]^{W_R}$, where W_R is the Weyl group of the root system R of $(\mathfrak{g}, \mathfrak{a})$. This root system is known to be of type A_r (the proof is the same as for Theorem 3.11 in [5]). Hence W_R is the symmetric group of $r + 1$ variables and it acts by permutations on the $\bar{\alpha}_i$'s. We will choose an order on R such that $\bar{\Sigma}^+ \subset R^+$. As in [23] and [28] we consider here relative invariants $\Delta_0, \Delta_1, \dots, \Delta_r$ with respect to the Borel subgroup defined by Σ^- . Define $\rho = \frac{1}{2} \sum_{\beta \in R^-} \beta$. It is well known that for $D \in \mathcal{T}_0$ and $\lambda = \sum_{i=0}^r a_i \lambda_i \in \mathfrak{a}^*$, $\gamma(D)(-\bar{\lambda} + \rho)$ is equal to the eigenvalue of D acting on $\Delta_0^{a_0} \dots \Delta_r^{a_r}$. In other words we have:

$$\gamma(D)(-\bar{\lambda} + \rho) = b_D(\lambda).$$

From [29], Lemme 3.9 p. 155 we know that

$$\rho = \frac{d}{4} \sum_{i < j} (\bar{\alpha}_i - \bar{\alpha}_j) = \frac{d}{4} \sum_{i=0}^r (r-2i)\bar{\alpha}_i$$

and from [ibid.], Lemme 3.8 p. 155 we also have:

$$\bar{\lambda} = a_0\bar{\alpha}_0 + (a_0 + a_1)\bar{\alpha}_1 + \dots + (a_0 + \dots + a_r)\bar{\alpha}_r.^\ddagger$$

Let us now make the following change of variables:

$$s_i = a_0 + \dots + a_i, \text{ for } i = 0, \dots, r.$$

As $b_{D_\ell}(\lambda) = b_Y(s_0 + \ell, \dots, s_r + \ell) = c \prod_{i=0}^r (s_i + \ell + i\frac{d}{2})$ (see Example 3.1.2) we obtain

$$\begin{aligned} \gamma(D_\ell)(\bar{\lambda}) &= b_{D_\ell}(-\lambda + \rho) = b_{D_\ell}(\sum_{i=0}^r -s_i\bar{\alpha}_i + \frac{d}{4} \sum_{i=0}^r (r-2i)\bar{\alpha}_i) \\ &= c \prod_{i=0}^r (-s_i + \frac{d}{4}r + \ell). \end{aligned}$$

As expected the polynomials $\gamma(D_\ell)$ are symmetric in the s_i variables (i.e. invariant under W_R). Moreover it is easy to prove that these polynomials, for $\ell = 0, \dots, r$, are algebraically independent generators of the algebra of symmetric polynomials. This proves 2).

3) As $\mathcal{T} = \mathcal{T}_0[X, X^{-1}]$ (see Proposition 3.1.6), and as, from 2), the elements of \mathcal{T}_0 are polynomials in X, X^{-1}, Y we obtain that $\mathcal{T} \subset \mathbb{C}[X, X^{-1}, Y]$. The inverse inclusion is obvious.

4) The inclusion $\mathbb{C}[X, Y, R_1, \dots, R_r] \subset D(V^+)^{G'} = \mathcal{T}_0[X, Y]$ is obvious. Conversely, from Theorem 2.2.9 we have $\mathcal{T}_0[X, Y] = \mathbb{C}[R_0, R_1, \dots, R_r][X, Y]$. As $R_0 = XY$ (see Remark 2.2.10), we have $\mathcal{T}_0[X, Y] \subset \mathbb{C}[X, Y, R_1, \dots, R_r]$. □

Remark 3.3.2. According to Terras [33](II, p.208), the operators D_ℓ were first considered by Selberg on positive definite symmetric matrices. They appear also in Maass ([21]), in the same context of positive definite symmetric matrices. In the setting of symmetric cones, the analogue of assertion 2) of the preceding theorem can be found in [8](Corollary XIV.1.6).

Remark 3.3.3. Note that for PV 's of commutative parabolic type we have $R_r = E$. In the special case where $G \simeq SO(k) \times \mathbb{C}^*$ and $V^+ \simeq \mathbb{C}^k$, we have always $r = 1$, and assertion 4) of the preceding theorem yields

$$D(\mathbb{C}^k)^{SO(k)} = \mathbb{C}[Q(x), Q(\partial), E]$$

where $Q(x) = X = \sum_{i=1}^k x_i^2$, $Q(\partial) = Y = \sum_{i=1}^k \frac{\partial^2}{\partial x_i^2}$.

This was proved by S. Rallis and G. Schiffmann ([26], Lemma 5.2. p. 112).

4. THE STRUCTURE OF $D(V)^{G'}$

[‡]The change of sign with respect to Lemme 3.8 in [29] is again due to the fact that we consider here characters of relative invariants instead of highest weights.

4.1. Smith algebras over rings.

As usual if a, b are elements of an associative algebra we define $[a, b] = ab - ba$.

Definition 4.1.1. *Let \mathbf{A} be a commutative associative algebra over \mathbb{C} , with unit element 1 and without zero divisors. Let $f, u \in \mathbf{A}[t]$ be two polynomials in one variable with coefficients in \mathbf{A} . Let $n \in \mathbb{N}^*$.*

- 1) *The Smith algebra $S(\mathbf{A}, f, n)$ is the associative algebra over \mathbf{A} with generators (x, y, e) subject to the relations $[e, x] = nx$, $[e, y] = -ny$, $[y, x] = f(e)$.*
- 2) *The algebra $U(\mathbf{A}, u, n)$ is the associative algebra over \mathbf{A} with generators $(\tilde{x}, \tilde{y}, \tilde{e})$ subject to the relations $[\tilde{e}, \tilde{x}] = n\tilde{x}$, $[\tilde{e}, \tilde{y}] = -n\tilde{y}$, $\tilde{x}\tilde{y} = u(\tilde{e})$, $\tilde{y}\tilde{x} = u(\tilde{e} + n)$.*

Remark 4.1.2. 1) The algebras $S(\mathbb{C}, f, n)$ were introduced and intensively studied by S. P. Smith ([32]) who called them "algebras similar to $\mathcal{U}(\mathfrak{sl}_2)$ ", where $\mathcal{U}(\mathfrak{sl}_2)$ is the enveloping algebra of \mathfrak{sl}_2 . In fact they share many interesting properties with $\mathcal{U}(\mathfrak{sl}_2)$, in particular they have a very rich representation theory.

2) One can prove, as in [32], that if the degree of f is one and $n \neq 0$, and if the leading coefficient is invertible in \mathbf{A} , then $S(\mathbf{A}, f, n)$ is isomorphic to the enveloping algebra $\mathcal{U}(\mathfrak{sl}_2(\mathbf{A}))$.

Let \mathcal{R} be a ring and let $\sigma \in \text{Aut}(\mathcal{R})$. Let us recall that a σ -derivation of \mathcal{R} is an additive map $\delta : \mathcal{R} \rightarrow \mathcal{R}$ such that $\delta(su) = s\delta(u) + \delta(s)\sigma(u)$. Given a σ -derivation δ , the skew polynomial ring over \mathcal{R} determined by σ and δ is the ring $\mathcal{R}[t, \sigma, \delta] := \langle \mathcal{R}, t \rangle / \{st - t\sigma(s) - \delta(s) \mid s \in \mathcal{R}\}$, where $\langle \mathcal{R}, t \rangle$ stands for the ring freely generated by \mathcal{R} and an element t with the relations given by the ring structure on \mathcal{R} (for details see [22], section 1.2, p.15 or [9] p.34).

Proposition 4.1.3.

Let \mathfrak{b} the 2-dimensional Lie algebra over \mathbf{A} , with basis $\{\varepsilon, \alpha\}$ and relation $[\varepsilon, \alpha] = n\alpha$. Let $\mathcal{U}(\mathfrak{b})$ be the enveloping algebra of \mathfrak{b} . Define an automorphism σ of $\mathcal{U}(\mathfrak{b})$ by $\sigma(\alpha) = \alpha$ and $\sigma(\varepsilon) = \varepsilon - n$ and define also a σ -derivation δ of $\mathcal{U}(\mathfrak{b})$ by $\delta(\alpha) = f(\varepsilon)$ and $\delta(\varepsilon) = 0$. Then $S(\mathbf{A}, f, n) \simeq \mathcal{U}(\mathfrak{b})[t, \sigma, \delta]$.

Proof. The proof is almost the same as the one given by S. P. Smith ([32], Prop. 1.2.). The isomorphism $S(\mathbf{A}, f, n) \simeq \mathcal{U}(\mathfrak{b})[t, \sigma, \delta]$ is given by $e \mapsto \varepsilon$, $x \mapsto \alpha$ and $y \mapsto t$. □

Corollary 4.1.4.

$S(\mathbf{A}, f, n)$ is a noetherian domain with \mathbf{A} -basis $\{y^i x^j e^k, i, j, k \in \mathbb{N}\}$ (or any similar family of ordered monomials obtained by permutation of the elements (y, x, e)).

Proof. (compare with [32], proof of corollary 1.3 p. 288). We know from [22] Th.1.2.9, that as $\mathcal{U}(\mathfrak{b})$ is a noetherian domain, so is $S(\mathbf{A}, f, n) \simeq \mathcal{U}(\mathfrak{b})[t, \sigma, \delta]$. Since

$$\begin{aligned} \mathcal{U}(\mathfrak{b})[t, \sigma, \delta] &= \mathcal{U}(\mathfrak{b}) \oplus \mathcal{U}(\mathfrak{b})t \oplus \mathcal{U}(\mathfrak{b})t^2 \oplus \mathcal{U}(\mathfrak{b})t^3 \oplus \dots \oplus \mathcal{U}(\mathfrak{b})t^\ell \oplus \dots \\ &= \mathcal{U}(\mathfrak{b}) \oplus t\mathcal{U}(\mathfrak{b}) \oplus t^2\mathcal{U}(\mathfrak{b}) \oplus t^3\mathcal{U}(\mathfrak{b}) \oplus \dots \oplus t^\ell\mathcal{U}(\mathfrak{b}) \oplus \dots \end{aligned}$$

(direct sums of \mathbf{A} -modules) and since the Poincaré-Birkhoff-Witt Theorem is still true for enveloping algebras of Lie algebras which are free over rings (see [6]), the ordered monomials in (y, x, e) beginning or ending with y form a basis of the algebra $S(\mathbf{A}, f, n)$. To obtain the basis $\{e^i y^j x^k\}$ or $\{x^k y^j e^i\}$ it suffices to replace the algebra \mathfrak{b} by the algebra \mathfrak{b}_- which is generated by e and y . □

Remark 4.1.5. The adjoint action of e ($u \mapsto [e, u]$) on $S(\mathbf{A}, f, n)$ is semi-simple and gives a decomposition of $S(\mathbf{A}, f, n)$ into weight spaces:

$$S(\mathbf{A}, f, n) = \bigoplus_{\nu \in \mathbb{Z}} S(\mathbf{A}, f, n)^\nu$$

where $S(\mathbf{A}, f, n)^\nu = \{u \in S(\mathbf{A}, f, n), [e, u] = \nu nu\}$. As $[e, x^j y^i e^k] = n(j-i)y^i x^j e^k$, we obtain, using Corollary 4.1.4, that the ordered monomials of the form $x^i y^j e^k$ form an \mathbf{A} -basis for $S(\mathbf{A}, f, n)^0$. Moreover as $yx = xy + f(e)$, it is easy to see that $S(\mathbf{A}, f, n)^0 = \mathbf{A}[xy, e] = \mathbf{A}[yx, e]$, where $\mathbf{A}[xy, e]$ (resp. $\mathbf{A}[yx, e]$) denotes the \mathbf{A} -subalgebra generated by xy (resp. yx) and e .

The proof of the following Lemma is straightforward.

Lemma 4.1.6.

Let $n \in \mathbb{N}^*$ and let $f \in \mathbf{A}[t]$. There exists an element $u \in \mathbf{A}[t]$, which is unique up to addition of an element of \mathbf{A} , such that

$$f(t) = u(t+n) - u(t) \quad (5-1-1)$$

Proposition 4.1.7. (compare with [32], Prop. 1.5)

Let u be as in the preceding Lemma. Define

$$\Omega_1 = xy - u(e).$$

Then the center of $S(\mathbf{A}, f, n)$ is $\mathbf{A}[\Omega_1]$ which is isomorphic to the polynomial algebra $\mathbf{A}[t]$.

Proof. Let us now prove that Ω_1 is central. Obviously Ω_1 commutes with e .

From the defining relations of $S(\mathbf{A}, f, n)$ we have $ex = x(e+n)$ and therefore, for any $k \in \mathbb{N}$, $e^k x = x(e+n)^k$.

This implies of course that for any polynomial $P \in \mathbf{A}[t]$ we have

$$P(e)x = xP(e+n) \text{ or } P(e-n)x = xP(e). \quad (5-1-2)$$

Similarly one proves that

$$P(e)y = yP(e-n) \text{ or } P(e+n)y = yP(e). \quad (5-1-3)$$

Let us show that Ω_1 commutes with x . Using Lemma 4.1.6 and (5-1-2) we obtain:

$$\begin{aligned} x\Omega_1 &= x(xy - u(e)) = x^2y - xu(e) = x(yx - f(e)) - xu(e) \\ &= x(yx - u(e+n) + u(e)) - xu(e) = xyx - xu(e+n) = xyx - u(e)x \\ &= \Omega_1 x. \end{aligned}$$

A similar calculation using (5-1-3) shows that Ω_1 commutes also with y . Hence Ω_1 belongs to the center of $S(\mathbf{A}, f, n)$.

Let now z be a central element of $S(\mathbf{A}, f, n)$. Then $z \in S(\mathbf{A}, f, n)^0$. We have $S(\mathbf{A}, f, n)^0 = \mathbf{A}[xy, e] = \mathbf{A}[\Omega_1, e]$, and hence z can be written as follows:

$$z = \sum c_i(e)\Omega_1^i \quad (\text{finite sum})$$

where $c_i(e) \in \mathbf{A}[e]$.

We have:

$$\begin{aligned} 0 &= [z, x] = \left[\sum c_i(e) \Omega_1^i, x \right] = \sum [c_i(e), x] \Omega_1^i \\ &= \sum (c_i(e)x - xc_i(e)) \Omega_1^i = \sum x(c_i(e+n) - c_i(e)) \Omega_1^i \quad (\text{using } (5-1-2)) \\ &= x \left(\sum (c_i(e+n) - c_i(e)) \Omega_1^i \right) \end{aligned}$$

As the algebra $S(\mathbf{A}, f, n)$ has no zero divisors we get:

$$\sum (c_i(e+n) - c_i(e)) \Omega_1^i = 0$$

As $\Omega_1 = xy - u(e)$, we have $\Omega_1^i = x^i y^i$ modulo monomials of the form $e^k x^p y^p$ with $p < i$. Then from Corollary 4.1.4 above we obtain $c_i(e+n) - c_i(e) = 0$, for all i . As the elements e^k are free over \mathbf{A} (Corollary 4.1.4) we obtain from Lemma 4.1.6 that $c_i \in \mathbf{A}$, for all i . \square

Remark 4.1.8. Conversely let us start with $u \in \mathbf{A}[t]$. Define $f \in \mathbf{A}[t]$ by $f(t) = u(t+n) - u(t)$. Then, from the definitions we have:

$$U(\mathbf{A}, u, n) = S(\mathbf{A}, f, n) / (xy - u(e)) = S(\mathbf{A}, f, n) / (\Omega_1)$$

where $(xy - u(e)) = (\Omega_1)$ is the ideal generated by $xy - u(e) = \Omega_1$. Again, as for $S(\mathbf{A}, f, n)$, the adjoint action of \tilde{e} gives a decomposition of $U(\mathbf{A}, u, n)$ into weight spaces:

$$U(\mathbf{A}, u, n) = \bigoplus_{\nu \in \mathbb{Z}} U(\mathbf{A}, u, n)^\nu \quad (5-1-4)$$

where $U(\mathbf{A}, u, n)^\nu = \{\tilde{v} \in U(\mathbf{A}, u, n), [\tilde{e}, \tilde{v}] = \nu n \tilde{v}\}$.

Proposition 4.1.9.

Let $u \in \mathbf{A}[t]$ and $s \in \mathbb{N}$. The \mathbf{A} -linear mappings

$$\begin{array}{ll} \varphi : \mathbf{A}[t] & \longrightarrow U(\mathbf{A}, u, n) \\ P & \longmapsto \varphi(P) = \tilde{x}^s P(\tilde{e}) \end{array} \quad \begin{array}{ll} \psi : \mathbf{A}[t] & \longrightarrow U(\mathbf{A}, u, n) \\ P & \longmapsto \psi(P) = \tilde{y}^s P(\tilde{e}) \end{array}$$

are injective (in particular the subalgebra $\mathbf{A}[\tilde{e}] \subset U(\mathbf{A}, u, n)$ generated by \tilde{e} is a polynomial algebra).

Proof. Define $f(t) = u(t+n) - u(t)$. Every element of $S(\mathbf{A}, f, n)$ can be written uniquely in the form

$$\sum a_{k,\ell,m} e^k x^\ell y^m \quad (a_{k,\ell,m} \in \mathbf{A})$$

(Corollary 4.1.4). Therefore, from Remark 4.1.8, every element in $U(\mathbf{A}, u, n)$ can be written in the form:

$$\sum a_{k,\ell,m} \tilde{e}^k \tilde{x}^\ell \tilde{y}^m \quad (a_{k,\ell,m} \in \mathbf{A}).$$

Let $P(t) = \sum_{i=0}^p a_i t^i$, ($a_i \in \mathbf{A}$) be a polynomial such that $\tilde{x}^s P(\tilde{e}) = 0$ (i.e. $P \in \ker \varphi$). As $U(\mathbf{A}, u, n) = S(\mathbf{A}, f, n) / (\Omega_1)$, we see that there exists $\alpha \in S(\mathbf{A}, f, n)$ such that

$$x^s \sum_{i=0}^p a_i e^i = \alpha \Omega_1 = \alpha(xy - u(e)).$$

If $\alpha = \sum a_{k,\ell,m} e^k x^\ell y^m$, using the fact that $\Omega_1 = xy - u(e)$ is central and relation (5-1-2) we get:

$$\begin{aligned} x^s \sum_{i=0}^p a_i e^i &= \left(\sum_{k,\ell,m} a_{k,\ell,m} e^k x^\ell y^m \right) (xy - u(e)) = \sum_{k,\ell,m} a_{k,\ell,m} e^k x^\ell (xy - u(e)) y^m \\ &= \sum_{k,\ell,m} a_{k,\ell,m} e^k x^{\ell+1} y^{m+1} - \sum_{k,\ell,m} a_{k,\ell,m} e^k x^\ell u(e) y^m \\ &= \sum_{k,\ell,m} a_{k,\ell,m} e^k x^{\ell+1} y^{m+1} - \sum_{k,\ell,m} a_{k,\ell,m} e^k u(e - \ell n) x^\ell y^m \quad (*) \end{aligned}$$

Suppose now that $\alpha \neq 0$, then one can define

$$\ell_0 = \max\{\ell \in \mathbb{N}, \exists k, m, a_{k,\ell,m} \neq 0\}.$$

Let k_0, m_0 be such that $a_{k_0, \ell_0, m_0} \neq 0$. From (*) above we get

$$x^s \sum_{i=0}^p a_i e^i + \sum_{k,\ell,m} a_{k,\ell,m} e^k u(e - \ell n) x^\ell y^m = \sum_{k,\ell,m} a_{k,\ell,m} e^k x^{\ell+1} y^{m+1}.$$

Using again (5-1-2) we obtain:

$$\sum_{i=0}^p a_i (e - ns)^i x^s + \sum_{k,\ell,m} a_{k,\ell,m} e^k u(e - \ell n) x^\ell y^m = \sum_{k,\ell,m} a_{k,\ell,m} e^k x^{\ell+1} y^{m+1}.$$

The left hand side of the preceding equality does not contain the monomial $e^{k_0} x^{\ell_0+1} y^{m_0+1}$, whereas the right hand side does. As the elements $e^k x^\ell y^m$ are a basis over \mathbf{A} (Corollary 4.1.4), we obtain a contradiction. Therefore $\alpha = 0$, and hence $x^s \sum_{i=0}^p a_i e^i = 0$, and again from Corollary 4.1.4, we obtain that $a_i = 0$ for all i . This proves that $\ker \varphi = \{0\}$. The proof for ψ is similar. \square

Corollary 4.1.10. *Every element \tilde{u} in $U(\mathbf{A}, u, n)$ can be written uniquely in the form*

$$\tilde{u} = \sum_{\ell > 0, k \geq 0} \alpha_{k,\ell} \tilde{y}^\ell \tilde{e}^k + \sum_{m \geq 0, r \geq 0} \beta_{m,r} \tilde{x}^m \tilde{e}^r$$

with $\alpha_{k,\ell}, \beta_{m,r} \in \mathbf{A}$.

Proof. We have already noticed that any element in $U(\mathbf{A}, u, n)$ can be written (in a non unique way) as a linear combination, with coefficients in \mathbf{A} , of the elements $\tilde{x}^i \tilde{y}^j \tilde{e}^k$.

Suppose that $i \geq j$. Then we have $\tilde{x}^i \tilde{y}^j \tilde{e}^k = \tilde{x}^{i-j} \tilde{x}^j \tilde{y}^j \tilde{e}^k$. As $\tilde{x} \tilde{y} = u(\tilde{e})$, we see that $\tilde{x}^j \tilde{y}^j = Q_j(\tilde{e})$, where Q_j is a polynomial with coefficients in \mathbf{A} . Therefore $\tilde{x}^i \tilde{y}^j \tilde{e}^k = \sum_{\ell} \gamma_\ell \tilde{x}^{i-j} \tilde{e}^\ell$, with $\gamma_\ell \in \mathbf{A}$. Similarly one can prove that if $i < j$, we have $\tilde{x}^i \tilde{y}^j \tilde{e}^k = \sum_{\ell} \delta_\ell \tilde{y}^{j-i} \tilde{e}^\ell$, with $\delta_\ell \in \mathbf{A}$. This shows that any element \tilde{u} in $U(\mathbf{A}, u, n)$ can be written in the expected form.

Suppose now that:

$$\sum_{\ell > 0, k \geq 0} \alpha_{k,\ell} \tilde{y}^\ell \tilde{e}^k + \sum_{m \geq 0, r \geq 0} \beta_{m,r} \tilde{x}^m \tilde{e}^r = 0.$$

Then, as $\tilde{y}^\ell \tilde{e}^k \in U(\mathbf{A}, u, n)^{-\ell}$ and $\tilde{x}^m \tilde{e}^r \in U(\mathbf{A}, u, n)^m$, we deduce from (5-1-4) that

$$\forall \ell > 0, \sum_k \alpha_{k,\ell} \tilde{y}^\ell \tilde{e}^k = 0, \quad \forall m \geq 0, \sum_r \beta_{m,r} \tilde{x}^m \tilde{e}^r = 0.$$

Then from Proposition 4.1.9, we deduce that $\alpha_{k,\ell} = 0$ and $\beta_{m,r} = 0$. □

4.2. Generators and relations for $D(V)^{G'}$.

Let $\mathcal{Z}(\mathcal{T})[t]$ be the polynomials in one variable with coefficients in $\mathcal{Z}(\mathcal{T})$. From the commutation rules $[E, X] = EX - XE = d_0X$ and $[E, Y] = -d_0Y$, we easily deduce that for $P \in \mathcal{Z}(\mathcal{T})[t]$ we have

$$YP(E) = P(E + d_0)Y, \quad XP(E) = P(E - d_0)X. \quad (5-2-1)$$

From Proposition 3.2.6 above, we know that any element in $D(V)^G$ can be written uniquely as a polynomial in E with coefficients in $\mathcal{Z}(\mathcal{T})$. As XY and YX belong to $D(V)^G$, there exist therefore two uniquely determined polynomials u_{XY} and $u_{YX} \in \mathcal{Z}(\mathcal{T})[t]$ such that $XY = u_{XY}(E)$ and $YX = u_{YX}(E)$. From (5-2-1) we obtain that

$$YXY = u_{YX}(E)Y = Y u_{XY}(E) = u_{XY}(E + d_0)Y$$

and therefore

$$u_{YX}(E) = u_{XY}(E + d_0) \quad (5-2-2)$$

As the polynomial u_{XY} will play an important role in Theorem 4.2.2 below, let us emphasize the connection between u_{XY} and the Bernstein-Sato polynomial b_Y . Remark first that $b_Y = b_{XY}$. We know from Corollary 3.2.8 that

$$\begin{aligned} h(XY)(-\lambda + \rho) &= b_{XY}(\lambda) = b_Y(\lambda) \\ &= \sum_{i=0}^p \alpha_i(-\lambda + \rho)(a_0d_0 + a_1d_1 + \cdots + a_r d_r)^i = \sum_{i=0}^p \alpha_i(-\lambda + \rho)(h(E)(-\lambda + \rho))^i \end{aligned}$$

with uniquely defined polynomials $\alpha_i \in \mathbb{C}[A]^{W_{0,\tau}}$. Therefore we obtain:

Proposition 4.2.1.

Keeping the notations above, we have

$$u_{XY}(t) = \sum_{i=0}^p h^{-1}(\alpha_i)t^i$$

Theorem 4.2.2. *Let $f_{XY}(t) = u_{XY}(t + d_0) - u_{XY}(t)$. The mapping*

$$\tilde{x} \mapsto X, \quad \tilde{y} \mapsto Y, \quad \tilde{e} \mapsto E$$

extends uniquely to an isomorphism of $\mathcal{Z}(\mathcal{T})$ -algebras between $U(\mathcal{Z}(\mathcal{T}), u_{XY}, d_0)$ (which is isomorphic to $S(\mathcal{Z}(\mathcal{T}), f_{XY}, d_0)/(\Omega_1)$) and $D(V)^{G'} = \mathcal{T}_0[X, Y]$.

Proof. As $[E, X] = d_0X$, $[E, Y] = -d_0Y$, $XY = u_{XY}(E)$ and $YX = u_{XY}(E + d_0)$ (see (5-2-2)), and as from Proposition 3.2.6 the algebra $D(V)^{G'} = \mathcal{T}_0[X, Y]$ is generated over $\mathcal{Z}(\mathcal{T})$ by X, Y, E , we know (universal property) that the mapping

$$\tilde{x} \mapsto X, \quad \tilde{y} \mapsto Y, \quad \tilde{e} \mapsto E$$

extends uniquely to a surjective morphism of $\mathcal{Z}(\mathcal{T})$ -algebras:

$$\varphi : U(\mathcal{Z}(\mathcal{T}), u_{XY}, d_0) \longrightarrow D(V)^{G'}.$$

From Corollary 4.1.10 any element \tilde{u} in $U(\mathcal{Z}(\mathcal{T}), u_{XY}, d_0)$ can be written uniquely in the form

$$\tilde{u} = \sum_{\ell > 0, k \geq 0} \alpha_{k, \ell} \tilde{y}^\ell \tilde{e}^k + \sum_{m \geq 0, r \geq 0} \beta_{m, r} \tilde{x}^m \tilde{e}^r$$

with $\alpha_{k, \ell}, \beta_{m, r} \in \mathcal{Z}(\mathcal{T})$. Suppose now that $\tilde{u} \in \ker(\varphi)$, then

$$\varphi(\tilde{u}) = \sum_{\ell > 0, k \geq 0} \alpha_{k, \ell} Y^\ell E^k + \sum_{m \geq 0, r \geq 0} \beta_{m, r} X^m E^r = 0,$$

with $\alpha_{k, \ell}, \beta_{m, r} \in \mathcal{Z}(\mathcal{T})$. Then Corollary 3.2.7 implies that $\alpha_{k, \ell} = \beta_{m, r} = 0$. Hence φ is an isomorphism. \square

5. RADIAL COMPONENTS

5.1. Radial components and Bernstein-Sato polynomials.

Remember that for $\tilde{\mathbf{a}} = (a_1, a_2, \dots, a_r) \in \mathbb{N}^r$ we have defined $V_{\tilde{\mathbf{a}}} = V_{(0, a_1, \dots, a_r)}$. Remember also that for $\mathbf{a} = (a_0, a_1, \dots, a_r)$ we have $V_{\mathbf{a}} = \Delta^{a_0} V_{\tilde{\mathbf{a}}}$. We know from Proposition 2.2.8 that the spaces $U_{\tilde{\mathbf{a}}} = \bigoplus_{a_0 \in \mathbb{N}} \Delta_0^{a_0} V_{\tilde{\mathbf{a}}}$ are the G' -isotypic components of $\mathbb{C}[V]$ and that the spaces $W_{\tilde{\mathbf{a}}} = \bigoplus_{a_0 \in \mathbb{Z}} \Delta_0^{a_0} V_{\tilde{\mathbf{a}}}$ are the G' -isotypic components of $\mathbb{C}[\mathcal{O}]$. Therefore the algebra $D(V)^{G'} = \mathcal{T}_0[X, Y]$ stabilizes each space $U_{\tilde{\mathbf{a}}}$ and the algebra $D(\mathcal{O})^{G'} = \mathcal{T}_0[X, X^{-1}] = \mathcal{T}$ stabilizes each space $W_{\tilde{\mathbf{a}}}$.

Let us consider the restriction map:

$$\begin{aligned} D(\mathcal{O})^{G'} &\longrightarrow \text{End}(W_{\tilde{\mathbf{a}}}) \\ D &\longmapsto r_{\tilde{\mathbf{a}}}(D) = D|_{W_{\tilde{\mathbf{a}}}} \end{aligned}$$

Definition 5.1.1. Let $D \in D(\mathcal{O})^{G'} = \mathcal{T}_0[X, X^{-1}] = \mathcal{T}$. The operator $r_{\tilde{\mathbf{a}}}(D) = D|_{W_{\tilde{\mathbf{a}}}}$ is called the radial component of D with respect to $\tilde{\mathbf{a}}$.

Example 5.1.2. Consider the case where $\tilde{\mathbf{a}} = 0$. Then $W_{\tilde{\mathbf{a}}} = \mathbb{C}[\Delta_0, \Delta_0^{-1}]$, and $r_0(D)$ is the endomorphism of $\mathbb{C}[t, t^{-1}]$ defined by $D(\varphi \circ \Delta_0) = r_0(D)(\varphi) \circ \Delta_0$. The operator $r_0(D)$ is the usual radial component of D (we will see below that it is a differential operator).

Notice now that the space $W_{\tilde{\mathbf{a}}} = \bigoplus_{a_0 \in \mathbb{Z}} \Delta_0^{a_0} V_{\tilde{\mathbf{a}}}$ can be viewed as the space of Laurent polynomials in Δ_0 , with coefficients in $V_{\tilde{\mathbf{a}}}$, in other words any $P \in W_{\tilde{\mathbf{a}}}$ can be written uniquely under the form

$$P = \sum \Delta_0^p \gamma_p$$

with $\gamma_p \in V_{\tilde{\mathbf{a}}}$. This can also be written as $P = \varphi \circ (\Delta_0)$, with $\varphi(t) = \sum t^p \gamma_p \in V_{\tilde{\mathbf{a}}}[t, t^{-1}]$ (where $V_{\tilde{\mathbf{a}}}[t, t^{-1}]$ is precisely the set of linear combinations $\sum t^p \gamma_p$, with $\gamma_p \in V_{\tilde{\mathbf{a}}}$).

There is a natural action of $D(\mathbb{C}^*) = \mathbb{C}[t, t^{-1}, t \frac{d}{dt}]$ on $V_{\tilde{\mathbf{a}}}[t, t^{-1}]$ given by $\frac{d}{dt} t^p \gamma_p = p t^{p-1} \gamma_p$.

Proposition 5.1.3.

Let $D \in \mathcal{T}_n$ a homogeneous element of degree n . Let b_D be its Bernstein-Sato polynomial. Let $\varphi \in V_{\tilde{\mathbf{a}}}[t, t^{-1}]$. Then $D(\varphi \circ \Delta_0) = (t^n b_D(t \frac{d}{dt}, a_1, \dots, a_r) \varphi) \circ \Delta_0$, in other words $r_{\tilde{\mathbf{a}}}(D) = t^n b_D(t \frac{d}{dt}, a_1, \dots, a_r)$.

Proof. It is enough to show that the two operators coincide on elements of the form $\Delta_0^p \gamma_p$, with $\gamma_p \in V_{\tilde{\mathbf{a}}}$. Then $\varphi = t^p \gamma_p$. Let us write

$$b_D(\mathbf{a}) = \sum_k c_k(a_1, \dots, a_r) a_0^k$$

We have:

$$\begin{aligned} (t^n b_D(t \frac{d}{dt}, a_1, \dots, a_r) \varphi) \circ \Delta_0 &= t^n (\sum_k c_k(a_1, \dots, a_r) (t \frac{d}{dt})^k \varphi) \circ \Delta_0 \\ &= t^n (\sum_k c_k(a_1, \dots, a_r) p^k t^p \gamma_p) \circ \Delta_0 \\ &= (t^n b_D(p, a_1, \dots, a_r) t^p \gamma_p) \circ \Delta_0 = b_D(p, a_1, \dots, a_r) \Delta_0^{p+n} \gamma_p \\ &= D(\Delta_0^p \gamma_p) \end{aligned}$$

□

Corollary 5.1.4.

If (G, V) is a PV of commutative parabolic type of rank $r + 1$, then the radial component of Y is given by

$$r_{\tilde{\mathbf{a}}}(Y) = ct^{-1} \prod_{j=0}^r (t \frac{d}{dt} + a_1 + \dots + a_j + j \frac{d}{2})$$

Proof. This is just a consequence of the formula for b_Y given in Example 3.1.2

□

Example 5.1.5. Consider the case 1) in example 2.3.3. In this case $G = (SL(n) \times SL(n)) \times \mathbb{C}^*$ acting on $x \in V = M_n(\mathbb{C})$ by $(g_1, g_2, t).x = tg_1 x g_2^{-1}$. Then $\Delta_0 = X = \det x$ and

$$Y = \Delta_0^*(\partial) = \det\left(\frac{\partial}{\partial x_{ij}}\right)$$

where x_{ij} are the coefficients of the matrix X . As in this case $\frac{d}{2} = 1$ (see [23], table 2 p. 122), we have $b_Y(a_0, a_1, \dots, a_{n-1}) = \prod_{j=0}^{n-1} (a_0 + a_1 + \dots + a_j + j)$. Therefore the radial component $r_0(Y)$ defined by $\det\left(\frac{\partial}{\partial x_{ij}}\right)(\varphi \circ \det) = (r_0(Y)\varphi) \circ \det$ is given by

$$r_0(Y) = t^{-1} \prod_{j=0}^{n-1} (t \frac{d}{dt} + j).$$

This radial component has already been calculated by Raïs ([25], p.22), by other methods. He obtained that $r_0(Y) = [\prod_{j=2}^{n-1} (t \frac{d}{dt} + j)] \frac{d}{dt}$. A simple calculation shows that the two operators are the same.

5.2. Algebras of radial components.

Definition 5.2.1. The radial component algebra $R_{\tilde{\mathbf{a}}}$ is the image of $D(V)^{G'} = \mathcal{T}_0[X, Y]$ under the map $D \mapsto r_{\tilde{\mathbf{a}}}(D)$.

Remember from Proposition 3.2.3 that the elements D in $\mathcal{Z}(\mathcal{T})$ are characterized by the fact that the corresponding Bernstein-Sato polynomial b_D does not depend on the a_0 variable. Therefore such a D acts by the scalar $b_D(0, \tilde{\mathbf{a}})$ on $W_{\tilde{\mathbf{a}}}$, that is $r_{\tilde{\mathbf{a}}}(D) = b_D(0, \tilde{\mathbf{a}}) Id_{W_{\tilde{\mathbf{a}}}}$.

Let us consider the polynomial $u_{XY} \in \mathcal{Z}(\mathcal{T})[t]$ which was introduced in section 4.2. If $u_{XY} = \sum_j c_j t^j$, with $c_j \in \mathcal{Z}(\mathcal{T})$, we define

$$r_{\bar{\mathbf{a}}}(u_{XY}) = \sum_j r_{\bar{\mathbf{a}}}(c_j) t^j \in \mathbb{C}[t].$$

Lemma 5.2.2.

Let $\mathbf{a} = (a_0, a_1, \dots, a_r) \in \mathbb{N}^{r+1}$. Suppose that $a_0 > 0$. Then the map $P \mapsto YP$ from $V_{\mathbf{a}}$ to $V_{\mathbf{a}-1}$ is a G' -equivariant isomorphism.

Proof. (Sketch) It is enough to prove that this map is not 0. As $\Delta_0^{*a_0} \dots \Delta_r^{*a_r}$ is the lowest weight vector of $V_{\mathbf{a}}^* \subset \mathbb{C}[V^*]$, we have $\Delta_0^*(\partial)^{a_0} \dots \Delta_r^*(\partial)^{a_r} \Delta_0^{a_0} \dots \Delta_r^{a_r}(0) \neq 0$. Hence $\Delta_0^*(\partial) \Delta_0^{a_0} \dots \Delta_r^{a_r} \neq 0$. \square

Theorem 5.2.3.

The radial component algebra $R_{\bar{\mathbf{a}}}$ is isomorphic, as an associative algebra over \mathbb{C} , to the algebra $U(\mathbb{C}, r_{\bar{\mathbf{a}}}(u_{XY}), d_0)$ introduced in Definition 4.1.1.

Proof. The algebra $R_{\bar{\mathbf{a}}}$ is generated over \mathbb{C} by the elements $r_{\bar{\mathbf{a}}}(E), r_{\bar{\mathbf{a}}}(X), r_{\bar{\mathbf{a}}}(Y)$. The defining relations of $U(\mathbb{C}, r_{\bar{\mathbf{a}}}(u_{XY}), d_0)$ are verified:

- $[r_{\bar{\mathbf{a}}}(E), r_{\bar{\mathbf{a}}}(X)] = r_{\bar{\mathbf{a}}}([E, X]) = d_0 r_{\bar{\mathbf{a}}}(X)$
- $[r_{\bar{\mathbf{a}}}(E), r_{\bar{\mathbf{a}}}(Y)] = r_{\bar{\mathbf{a}}}([E, Y]) = -d_0 r_{\bar{\mathbf{a}}}(Y)$
- $r_{\bar{\mathbf{a}}}(X) r_{\bar{\mathbf{a}}}(Y) = r_{\bar{\mathbf{a}}}(XY) = r_{\bar{\mathbf{a}}}(u_{XY})(r_{\bar{\mathbf{a}}}(E))$
- $r_{\bar{\mathbf{a}}}(Y) r_{\bar{\mathbf{a}}}(X) = r_{\bar{\mathbf{a}}}(YX) = r_{\bar{\mathbf{a}}}(u_{XY})(r_{\bar{\mathbf{a}}}(E) + d_0)$.

Therefore the mapping

$$\tilde{x} \mapsto r_{\bar{\mathbf{a}}}(X), \quad \tilde{y} \mapsto r_{\bar{\mathbf{a}}}(Y), \quad \tilde{e} \mapsto r_{\bar{\mathbf{a}}}(E)$$

extends uniquely to a surjective morphism of \mathbb{C} -algebras

$$\varphi_{\bar{\mathbf{a}}} : U(\mathbb{C}, r_{\bar{\mathbf{a}}}(u_{XY}), d_0) \longrightarrow R_{\bar{\mathbf{a}}}.$$

From Corollary 4.1.10 any element \tilde{u} in $U(\mathbb{C}, r_{\bar{\mathbf{a}}}(u_{XY}), d_0)$ can be written uniquely in the form

$$\tilde{u} = \sum_{\ell > 0, k \geq 0} \alpha_{k, \ell} \tilde{y}^\ell \tilde{e}^k + \sum_{m \geq 0, s \geq 0} \beta_{m, s} \tilde{x}^m \tilde{e}^s$$

with $\alpha_{k, \ell}, \beta_{m, s} \in \mathbb{C}$. Suppose now that $\tilde{u} \in \ker(\varphi_{\bar{\mathbf{a}}})$, then

$$\varphi_{\bar{\mathbf{a}}}(\tilde{u}) = \sum_{\ell > 0, k \geq 0} \alpha_{k, \ell} r_{\bar{\mathbf{a}}}(Y)^\ell r_{\bar{\mathbf{a}}}(E)^k + \sum_{m \geq 0, s \geq 0} \beta_{m, s} r_{\bar{\mathbf{a}}}(X)^m r_{\bar{\mathbf{a}}}(E)^s = 0.$$

Applying this operator to a function of the form $\Delta^{a_0} P$, with $P \in V_{\bar{\mathbf{a}}}$, we obtain:

$$\sum_{\ell > 0} Y^\ell \left(\sum_{k \geq 0} \alpha_{k, \ell} E^k \Delta^{a_0} P \right) + \sum_{m \geq 0} X^m \left(\sum_{s \geq 0} \beta_{m, s} E^s \Delta^{a_0} P \right) = 0.$$

As the operators X and Y have degree d_0 and $-d_0$ respectively, this implies that

$$\forall \ell, \quad Y^\ell \left(\sum_{k \geq 0} \alpha_{k, \ell} E^k \Delta^{a_0} P \right) = 0, \quad \text{and} \quad \forall m, \quad X^m \left(\sum_{s \geq 0} \beta_{m, s} E^s \Delta^{a_0} P \right) = 0.$$

Therefore, by Lemma 5.2.2 we obtain that

$$\forall \ell, a_0 > \ell \quad \sum_{k \geq 0} \alpha_{k, \ell} E^k \Delta^{a_0} P = 0, \quad \text{and} \quad \forall m, \forall a_0 \quad \sum_{s \geq 0} \beta_{m, s} E^s \Delta^{a_0} P = 0.$$

As $E\Delta^{a_0}P = (a_0d_0 + d(\tilde{\mathbf{a}}))\Delta^{a_0}P$, where $d(\tilde{\mathbf{a}}) = a_1d_1 + \cdots + a_r d_r$, we have:

$$\forall \ell, \text{ and } a_0 > \ell, \sum_{k \geq 0} \alpha_{k,\ell} (a_0d_0 + d(\tilde{\mathbf{a}}))^k \Delta^{a_0}P = 0, \quad \text{and } \forall m, \forall a_0 \sum_{s \geq 0} \beta_{m,s} (a_0d_0 + d(\tilde{\mathbf{a}}))^s \Delta^{a_0}P = 0.$$

Hence

$$\forall \ell, \text{ and } a_0 > \ell \sum_{k \geq 0} \alpha_{k,\ell} (a_0d_0 + d(\tilde{\mathbf{a}}))^k = 0, \quad \text{and } \forall m, \forall a_0 \sum_{s \geq 0} \beta_{m,s} (a_0d_0 + d(\tilde{\mathbf{a}}))^s = 0.$$

This implies that $\forall(\ell, k)$ and $\forall(m, s)$, we have $\alpha_{k,\ell} = 0$ and $\beta_{m,s} = 0$. Hence $\tilde{u} = 0$ and $\varphi_{\tilde{\mathbf{a}}}$ is injective. \square

Remark 5.2.4. For $\tilde{\mathbf{a}} = 0$, the preceding result was first obtained by T. Levasseur ([19]), by other methods.

Define now $J_{\tilde{\mathbf{a}}} = \ker(r_{\tilde{\mathbf{a}}}|_{D(V)^{G'}})$. $J_{\tilde{\mathbf{a}}}$ is a two-sided ideal of $D(V)^{G'} = \mathcal{T}_0[X, Y]$. Remember from Proposition 3.1.6 that any $D \in D(V)^{G'}$ can be written uniquely in the form:

$$D = \sum_{k \in \mathbb{N}^*} u_k Y^k + \sum_{n \in \mathbb{N}} v_n X^n \text{ (finite sum)}$$

where $u_k, v_n \in \mathcal{T}_0 = D(V)^G$.

Lemma 5.2.5.

$$J_{\tilde{\mathbf{a}}} = \{D = \sum_{k \in \mathbb{N}^*} u_k Y^k + \sum_{n \in \mathbb{N}} v_n X^n \mid u_k, v_n \in J_{\tilde{\mathbf{a}}} \cap \mathcal{T}_0\}.$$

Proof. From Theorem 5.2.3 the algebra $R_{\tilde{\mathbf{a}}}$ is isomorphic to $U(\mathbb{C}, r_{\tilde{\mathbf{a}}}(u_{XY}), d_0)$. If $r_{\tilde{\mathbf{a}}}(D) = \sum_{k \in \mathbb{N}^*} r_{\tilde{\mathbf{a}}}(u_k) r_{\tilde{\mathbf{a}}}(Y)^k + \sum_{n \in \mathbb{N}} r_{\tilde{\mathbf{a}}}(v_n) r_{\tilde{\mathbf{a}}}(X)^n = 0$, then, from Corollary 4.1.10, we obtain that $r_{\tilde{\mathbf{a}}}(u_k) = 0$ and $r_{\tilde{\mathbf{a}}}(v_n) = 0$ for all k and all n . \square

Let us now give a set of generators for the ideal $\ker(r_{\tilde{\mathbf{a}}})$ in $D(V)^{G'} = \mathcal{T}_0[X, Y]$. From Proposition 5.1.3 we obtain that $r_{\tilde{\mathbf{a}}}(E) = d_0(t \frac{d}{dt}) + d(\tilde{\mathbf{a}})$. Therefore $r_{\tilde{\mathbf{a}}}(t \frac{d}{dt}) = t \frac{d}{dt} - d(\tilde{\mathbf{a}})$. Define

$G_i^{\tilde{\mathbf{a}}} = R_i - b_{R_i}(t \frac{d}{dt}, \tilde{\mathbf{a}})$ where the R_i 's are the Capelli operators introduced in section 2.2. Obviously $G_i^{\tilde{\mathbf{a}}} \in D(V)^G = \mathcal{T}_0$. Using Proposition 5.1.3 again we obtain

$$\begin{aligned} r_{\tilde{\mathbf{a}}}(G_i^{\tilde{\mathbf{a}}}) &= r_{\tilde{\mathbf{a}}}(R_i - b_{R_i}(t \frac{d}{dt}, \tilde{\mathbf{a}})) = r_{\tilde{\mathbf{a}}}(R_i) - b_{R_i}(r_{\tilde{\mathbf{a}}}(t \frac{d}{dt}), \tilde{\mathbf{a}}) \\ &= b_{R_i}(t \frac{d}{dt}, \tilde{\mathbf{a}}) - b_{R_i}(t \frac{d}{dt}, \tilde{\mathbf{a}}) = 0. \end{aligned}$$

Hence the elements $G_i^{\tilde{\mathbf{a}}}$ belong to $J_{\tilde{\mathbf{a}}} \cap \mathcal{T}_0$.

Theorem 5.2.6.

The elements $G_i^{\tilde{\mathbf{a}}}$ are generators of $J_{\tilde{\mathbf{a}}}$:

$$J_{\tilde{\mathbf{a}}} = \ker(r_{\tilde{\mathbf{a}}}|_{D(V)^{G'}}) = \sum_{i=0}^r D(V)^{G'} G_i^{\tilde{\mathbf{a}}} = \sum_{i=0}^r G_i^{\tilde{\mathbf{a}}} D(V)^{G'}.$$

Proof. From Lemma 5.2.5, it is now enough to prove that

$$J_{\tilde{\mathbf{a}}} \cap \mathcal{T}_0 \subset \sum_{i=0}^r D(V)^G G_i^{\tilde{\mathbf{a}}} = \sum_{i=0}^r \mathcal{T}_0 G_i^{\tilde{\mathbf{a}}}.$$

Let $D \in J_{\tilde{\mathbf{a}}} \cap \mathcal{T}_0$. As $\mathcal{T}_0 = \mathbb{C}[R_0, \dots, R_r]$ (Proposition 2.2.9), we have also $\mathcal{T}_0 = \mathbb{C}[G_0^{\tilde{\mathbf{a}}}, \dots, G_r^{\tilde{\mathbf{a}}}, E]$. Therefore $D = \sum Q_i E^i$, where $Q_i \in \mathbb{C}[G_0^{\tilde{\mathbf{a}}}, \dots, G_r^{\tilde{\mathbf{a}}}]$. Hence $Q_i \in Q_i(0) + \sum_{i=0}^r D(V)^G G_i^{\tilde{\mathbf{a}}}$. Then

$$0 = r_{\tilde{\mathbf{a}}}(D) = \sum_i Q_i(0) r_{\tilde{\mathbf{a}}}(E^i) = \sum_i Q_i(0) (d_0(t \frac{d}{dt}) + d(\tilde{\mathbf{a}}))^i.$$

Therefore $Q_i(0) = 0$ ($i = 0, \dots, r$). Hence $Q_i \in \sum_{i=0}^r D(V)^G G_i^{\tilde{\mathbf{a}}}$, which yields $D \in \sum_{i=0}^r D(V)^G G_i^{\tilde{\mathbf{a}}}$. \square

Remark 5.2.7. For $\tilde{\mathbf{a}} = 0$, the result of the preceding Theorem is due to T. Levasseur ([19], Theorem 4.11. (v)).

5.3. Rational radial component algebras.

Definition 5.3.1. *The rational radial component algebra $R_{\tilde{\mathbf{a}}}^r$ is the image of $D(\mathcal{O})^{G'} = \mathcal{T}_0[X, X^{-1}] = \mathcal{T}$ under the map $D \mapsto r_{\tilde{\mathbf{a}}}(D)$.*

In fact as shown in the following proposition the structure of the algebras $R_{\tilde{\mathbf{a}}}^r$ is more simpler than the structure of $R_{\tilde{\mathbf{a}}}$, and the ideal $I_{\tilde{\mathbf{a}}} = \ker(r_{\tilde{\mathbf{a}}}) \subset \mathcal{T}$ has the same generators as $J_{\tilde{\mathbf{a}}}$.

Proposition 5.3.2.

- 1) For all $\tilde{\mathbf{a}}$, the rational radial component algebra $R_{\tilde{\mathbf{a}}}^r$ is isomorphic to $\mathbb{C}[t, t^{-1}, t \frac{d}{dt}]$.
- 2) $I_{\tilde{\mathbf{a}}} = \ker(r_{\tilde{\mathbf{a}}}) = \sum_{i=0}^r \mathcal{T} G_i^{\tilde{\mathbf{a}}} = \sum_{i=0}^r G_i^{\tilde{\mathbf{a}}} \mathcal{T}$.

Proof. 1) We have $\mathcal{T} = \mathcal{T}_0[X, X^{-1}]$. And $\mathcal{T}_0 = \mathcal{Z}(\mathcal{T})[E]$, from Proposition 3.2.6. Therefore $\mathcal{T} = \mathcal{Z}(\mathcal{T})[X, X^{-1}, E]$. On the other hand we have $r_{\tilde{\mathbf{a}}}(\mathcal{Z}(\mathcal{T})) = \mathbb{C}$, $r_{\tilde{\mathbf{a}}}(X) = t$, $r_{\tilde{\mathbf{a}}}(X^{-1}) = t^{-1}$ and $r_{\tilde{\mathbf{a}}}(E) = d_0(t \frac{d}{dt}) + d(\tilde{\mathbf{a}})$. Hence $R_{\tilde{\mathbf{a}}}^r = r_{\tilde{\mathbf{a}}}(\mathcal{T}) = \mathbb{C}[t, t^{-1}, d_0(t \frac{d}{dt}) + d(\tilde{\mathbf{a}})] = \mathbb{C}[t, t^{-1}, t \frac{d}{dt}]$.

2) Obviously $\sum_{i=0}^r \mathcal{T} G_i^{\tilde{\mathbf{a}}} \subset I_{\tilde{\mathbf{a}}}$. As $I_{\tilde{\mathbf{a}}}$ is a two-sided ideal of \mathcal{T} , it is easily seen to be graded. If $D \in I_{\tilde{\mathbf{a}}} \cap \mathcal{T}_p$, then $X^{-p} D \in \mathcal{T}_0 \cap I_{\tilde{\mathbf{a}}} = \mathcal{T}_0 \cap J_{\tilde{\mathbf{a}}} = \sum_{i=0}^r \mathcal{T}_0 G_i^{\tilde{\mathbf{a}}}$. Therefore $D \in \sum_{i=0}^r \mathcal{T} G_i^{\tilde{\mathbf{a}}}$. \square

REFERENCES

- [1] C. BENSON, G. RATCLIFF — *On multiplicity free actions*, in "Representations of Real and p-adic groups", 221-304. Lecture Notes Series 2. Institute of Mathematical Sciences, National University of Singapore, Singapore University Press, World Scientific, 2004
- [2] C. BENSON, G. RATCLIFF — *A Classification of Multiplicity Free Actions*, J. Alg. 181, 152-186 (1996).
- [3] C. BENSON, G. RATCLIFF — *Rationality of the generalized binomial coefficients for a multiplicity free action*, J. Austral. Math. Soc. Ser. A 68 (2000), 387-410.
- [4] N. BOPP, H. RUBENTHALER — *Fonction zêta associée à la série principale sphérique de certains espaces symétriques*, Ann. Sci. Ecole Norm. Sup. 26 (1993) n°6, 701-745.
- [5] N. BOPP, H. RUBENTHALER — *Local Zeta functions attached to the minimal spherical series for a class of symmetric spaces*, Mem. Amer. Math. Soc. 174 (2005), n° 821, viii+233 pp

- [6] N. BOURBAKI — *Groupes et Algèbres de Lie, Chapitre 1*, C.C.L.S, Paris, 1971.
- [7] M. BRION — *Représentations exceptionnelles des groupes semi-simples*, Ann. Sci. Ecole Norm. Sup. 18 (1985), 345-387.
- [8] J. FARAUT, A. KORANYI — *Analysis on Symmetric Cones*, Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1994. xii+382 pp.
- [9] K. R. GOODEARL, R. B. WARFIELD, — *An Introduction to Noncommutative Noetherian Rings*, Second edition. London Mathematical Society Student Texts, 61. Cambridge University Press, Cambridge, 2004. xxiv+344 pp.
- [10] G. HECKMANN, H. SCHLICHTKRULL — *Harmonic Analysis and Special Functions on Symmetric Spaces*, Perspectives in Mathematics Vol. 16, Academic Press (1994).
- [11] R. HOWE, T. UMEDA — *The Capelli identity, the double commutant theorem, and multiplicity-free actions*, Math. Ann. 290 (1991), n°3, 565-619.
- [12] J. IGUSA — *On Lie Algebras Generated by Two Differential Operators*, Manifolds and Lie Groups , Progr. Math. 14 (1981) Birkhäuser, Boston, Mass., 187-195.
- [13] V. KAC — *Some remarks on nilpotent orbits*, J. Alg. 64 (1980), 190-213.
- [14] T. KIMURA — *Introduction to Prehomogeneous Vector Spaces*, Transl. Math. Monogr., vol 215, American Mathematical Society, Providence, RI, 2003.
- [15] F. KNOP — *A Harish-Chandra homomorphism for reductive group actions*, Ann. of Math. (2) 140 (1994), no. 2, 253-288.
- [16] F. KNOP — *Some remarks on multiplicity free spaces*, Representation theories and algebraic geometry (Montreal, PQ, 1997), 301-317, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Kluwer Acad. Publ., Dordrecht, 1998.
- [17] B. KOSTANT, S. SAHI — *The Capelli identity, Tube Domains and the Generalized Laplace Transform*, Adv. Math. 87 (1991), 71-92.
- [18] A. S. LEAHY — *A Classification of Multiplicity Free Representations*, Journal of Lie Theory, volume 8 (1998) 367-391.
- [19] T. LEVASSEUR — *Radial components, prehomogeneous vector spaces, and rational Cherednik algebras*, Int. Math. Res. Not. IMRN 2009, no. 3, 462-511.
- [20] T. LEVASSEUR and J.T. STAFFORD — *Rings of differential operators on classical rings of invariants*, Memoirs of the AMS, volume 81, n° 412 (1989).
- [21] H. MAASS — *Siegel's Modular Forms and Dirichlets series*, Lecture Notes in Mathematics, Vol. 216. Springer-Verlag, Berlin-New York, 1971. v+328 pp.
- [22] J.C. McCONNEL and J.C. ROBSON — *Non Commutative Noetherian Rings*, Wiley-interscience, Chichester, 1987.
- [23] I. MULLER, H. RUBENTHALER and G. SCHIFFMANN — *Structure des espaces préhomogènes associés à certaines algèbres de Lie graduées*, Math. Ann. 274 (1986), 95-123.
- [24] T. NOMURA, — *Algebraically independent generators of invariant differential operators on a symmetric cone*, J. reine angew. Math. 400 (1989), 122-133.
- [25] M. RAÏS — *Distributions homogènes sur des espaces de matrices*, Bull. Soc. Math. France, Mémoire 30 (1972)
- [26] S. RALLIS and G. SCHIFFMANN — *Weil Representation. I Intertwining distributions and discrete spectrum*, Mem. Amer. Math. Soc. 25 (1980) n°231.
- [27] H. RUBENTHALER — *Multiplicity Free Spaces with a One-Dimensional Quotient*, Journal of Lie Theory 23 (2013), n°2, 433-458.

- [28] H. RUBENTHALER and G. SCHIFFMANN — *Opérateurs différentiels de Shimura et espaces préhomogènes*, Invent. Math. 90 (1989), 409-442.
- [29] H. RUBENTHALER and G. SCHIFFMANN — *SL_2 -triplet associé à un polynôme homogène*, J. reine angew. math. 408 (1990), 136-158.
- [30] G. SCHWARZ — *Finite-dimensional representations of invariant differential operators*, J. Alg. 258 (2002), 160-204.
- [31] M. SATO – T. KIMURA — *A classification of irreducible prehomogeneous vector spaces and their relative invariants*, Nagoya Math. J. 65 (1977), 1–155.
- [32] S. P. SMITH — *A class of algebras similar to the enveloping algebra of $\mathfrak{sl}(2)$* , Trans. Amer. Math. Soc, 322, 1 (1990), 285-314.
- [33] A. TERRAS — *Harmonic analysis on symmetric spaces and applications I, II*, Springer (1988).
- [34] M. VAN DEN BERGH — *Some rings of differential operators for SL_2 -invariants are simple*, J. Pure App. Algebra 107 (1996) 309-335.
- [35] E. VINBERG and B. KIMELFELD — *Homogeneous domains on flag manifolds and spherical subsets of semi-simple Lie groups*, Funktsional. Anal. i Prilozhen. 12 (1978), no. 3, 12–19, 96.
- [36] N. WALLACH — *Polynomial Differential Operators Associated with Hermitian Symmetric Spaces*, Representation Theory of Lie Groups and Lie Algebras, World Sci. Publishing, River Edge, NJ (1992), 76-94.
- [37] Z. YAN — *Invariant Differential Operators and Holomorphic Function Spaces*, J. Lie Theory 10 (2000) n°1, 1-31.

HUBERT RUBENTHALER, INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE, E-MAIL: rubenth@math.unistra.fr