



HAL
open science

On domination of Cartesian product of directed cycles

Michel Mollard

► **To cite this version:**

| Michel Mollard. On domination of Cartesian product of directed cycles. 2011. hal-00576481v1

HAL Id: hal-00576481

<https://hal.science/hal-00576481v1>

Preprint submitted on 14 Mar 2011 (v1), last revised 24 May 2012 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On domination of Cartesian product of directed cycles

Michel Mollard*

Institut Fourier
100, rue des Maths
38402 St martin d'hères Cedex FRANCE
michel.mollard@ujf-grenoble.fr

Abstract

Let $\gamma(C_m \square C_n)$ be the domination number of the Cartesian product of directed cycles C_m and C_n for $m, n \geq 2$. In [11] and [12] Liu and al. determined the value of $\gamma(C_m \square C_n)$ when $m \leq 6$ and when both m and $n \equiv 0 \pmod{3}$. In this article we give, in general, the value of $\gamma(C_m \square C_n)$ when $m \equiv 2 \pmod{3}$ and improve the known lower bound for most of the remaining cases. We also disprove the conjectured formula for the case $m \equiv 0 \pmod{3}$ appearing in [12].

Keywords: Graph, Cartesian product, Domination number, Combinatorial problems.

1 Introduction and definitions

Let $G = (V, E)$ be a finite directed graph (digraph for short) without loops or multiple arcs.

A vertex u *dominates* a vertex v if $u = v$ or $uv \in E$. A set $S \subset V$ is a *dominating* set of G if any vertex of G is dominated by at least a vertex of S . The *domination number* of G , denoted by $\gamma(G)$ is the minimal cardinality of a dominating set. The set V is a dominating set thus $\gamma(G)$ is finite. These definitions extend to digraphs the classical domination notion for undirected graphs.

The determination of domination number of a directed or undirected graph is, in general, a difficult question in graph theory. Furthermore this problem has connections with information theory. For example the domination number of Hypercubes is linked to error-correcting codes. Among the lot of related works ([7], [8]) mention the special case of domination of Cartesian product of undirected paths or cycles ([1] to [6], [9], [10]).

For $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ two digraphs, the *Cartesian product* $G_1 \square G_2$ is the digraph with vertex set $V_1 \times V_2$ and $(x_1, x_2)(y_1, y_2) \in E(G_1 \square G_2)$ if and only

*CNRS Université Joseph Fourier

if $x_1y_1 \in E_1$ and $x_2 = y_2$ or $x_2y_2 \in E_2$ and $x_1 = y_1$. Note that $G \square H$ is isomorph to $H \square G$.

In two articles published in this journal ([11], [12]) Liu and al. consider the domination number of Cartesian product of two directed cycles. They give the value of $\gamma(C_m \square C_n)$ when $m \leq 6$ and when both m and $n \equiv 0 \pmod{3}$. Furthermore they propose lower and upper bounds for the general case.

In this paper we are able to give, in general, the value of $\gamma(C_m \square C_n)$ when $m \equiv 2 \pmod{3}$ and we improve the lower bounds for most of the still unknown cases. We also disprove the conjectured formula appearing in [12] for the case $m \equiv 0 \pmod{3}$.

We denote the vertices of a directed cycle C_n by the integers $\{0, 1, \dots, n-1\}$ considered modulo n . Thus, when used for vertices labelling, $a+b$ and $a-b$ will denote the vertices $a+b$ and $a-b \pmod{n}$. Furthermore there is an arc xy from x to y in C_n if and only if $y = x+1 \pmod{n}$. For any i in $\{0, 1, \dots, n-1\}$ we will denote by C_m^i the subgraph of $C_m \square C_n$ induced by the vertices $\{(k, i) / k \in \{0, 1, \dots, m-1\}\}$. Note that C_m^i is isomorph to C_m .

2 Main results

Proposition 1 *Let S be a dominating set of $\gamma(C_m \square C_n)$ then for all i in $\{0, 1, \dots, n-1\}$ considered modulo n we have $|S \cap C_m^{i-1}| + 2|S \cap C_m^i| \geq m$.*

Proof : The m vertices of C_m^i can only be dominated by vertices of $S \cap C_m^i$ and $S \cap C_m^{i-1}$. Each of the vertices of $S \cap C_m^i$ and $S \cap C_m^{i-1}$ dominates respectively two and one vertex in C_m^i . The result follows. □

Theorem 2 *Let $m, n \geq 2$ and $k_1 = \lfloor \frac{m}{3} \rfloor$ then*

- *if $m \equiv 0 \pmod{3}$ then $\gamma(C_m \square C_n) \geq nk_1$*
- *if $m \equiv 1 \pmod{3}$ then $\gamma(C_m \square C_n) \geq nk_1 + \frac{n}{2}$*
- *if $m \equiv 2 \pmod{3}$ then $\gamma(C_m \square C_n) \geq nk_1 + n$.*

Proof :

Let S be a dominating set of $C_m \square C_n$ and for any i in $\{0, 1, \dots, n-1\}$ let $a_i = |S \cap C_m^i|$.

Assume first $m = 3k_1$. Let J be the set of $i \in \{0, 1, \dots, n-1\}$ such that $a_i \leq k_1 - 1$. Let $J' = \{i \mid i+1 \pmod{n} \in J\}$. By proposition 1, we have $a_{i-1} + 2a_i \geq 3k_1$ thus, if i belongs to J , $a_{i-1} + a_i \geq 2k_1 + 1$. A first consequence is that $J \cap J' = \emptyset$. Furthermore $\sum_{i \in J \cup J'} a_i \geq |J|(2k_1 + 1)$. If $i \notin J \cup J'$ we have $a_i \geq k_1$ therefore $\sum_{i \in \{0, 1, \dots, n-1\}} a_i \geq |J|(2k_1 + 1) + (n - 2|J|)k_1 = |J| + nk_1 \geq nk_1$.

Assume now $m = 3k_1 + 1$. Let J be the set of $i \in \{0, 1, \dots, n-1\}$ such that $a_i \leq k_1$. Let $J' = \{i \mid i+1 \pmod{n} \in J\}$. If i belongs to J we have $a_{i-1} + 2a_i \geq 3k_1 + 1$ thus $a_{i-1} + a_i \geq 2k_1 + 1$. Then $J \cap J' = \emptyset$ and $\sum_{i \in J \cup J'} a_i \geq |J|(2k_1 + 1)$. Therefore $\sum_{i \in \{0, 1, \dots, n-1\}} a_i \geq |J|(2k_1 + 1) + (n - 2|J|)(k_1 + 1) = nk_1 + n - |J| \geq nk_1 + \frac{n}{2}$.

Assume then $m = 3k_1 + 2$. Let J be the set of $i \in \{0, 1, \dots, n-1\}$ such that $a_i \leq k_1$. Let $J' = \{i \mid i+1 \pmod{n} \in J\}$. If $i \in J$ we have $a_{i-1} + 2a_i \geq 3k_1 + 2$

70 thus $a_{i-1} + a_i \geq 2k_1 + 2$. Then $J \cap J' = \emptyset$ and $\sum_{i \in J \cup J'} a_i \geq |J| (2k_1 + 2)$. Therefore
 72 $\sum_{i \in \{0, 1, \dots, n-1\}} a_i \geq |J| (2k_1 + 2) + (n - 2|J|)(k_1 + 1) \geq n(k_1 + 1)$. □

74 Let us now study in details the case $m \equiv 2 \pmod{3}$. Assume $m = 3k_1 + 2$. Let A
 be the set of $k_1 + 1$ vertices of C_m defined by $S = \{0\} \cup \{2 + 3p \mid p = 0, 1, \dots, k_1 - 1\} =$
 76 $\{0\} \cup \{2, 5, \dots, m - 6, m - 3\}$. For any i in $\{0, 1, \dots, m - 1\}$ let A_i be the translate,
 considered modulo m , of A by i ; thus $A_i = \{i\} \cup \{i + 2, i + 5, \dots, i - 6, i - 3\}$ (see
 Figure 1).

78 We will call a set S of vertices of $C_m \square C_n$ a A -set if for any j in $\{0, 1, \dots, n - 1\}$
 we have $S \cap C_m^j = A_{i_j}$ for some i_j . We have $|S| = n(k_1 + 1)$, thus if a set is
 80 both a A -set and a dominating set, by theorem 2, it is minimum and we have
 $\gamma(C_m \square C_n) = n(k_1 + 1)$.




Figure 1: A_i, A_{i-1} and A_{i+2}

Lemma 3 Let $m = 3k_1 + 2$. Let S be a A -set and for any j in $\{0, 1, \dots, n - 1\}$ let
 i_j such that $S \cap C_m^j = A_{i_j}$. Assume that:

$$\begin{cases} \text{for any } j \in \{1, \dots, n - 1\} \ i_j = i_{j-1} + 1 \pmod{m} \text{ or } i_j = i_{j-1} - 2 \pmod{m} \\ \text{and} \\ i_0 = i_{n-1} + 1 \pmod{m} \text{ or } i_0 = i_{n-1} - 2 \pmod{m}. \end{cases}$$

82 then S is a dominating set of $C_m \square C_n$.

Proof :

84 Note first that for any i in $\{0, 1, \dots, m-1\}$ the set of non dominated vertices
of C_m by A_i is $T = \{i+4, i+7, \dots, i-4, i-1\}$. Note also that $A_{i+2} = \{i+2\} \cup$
86 $\{i+4, i+7, \dots, i-4, i-1\}$ and $A_{i-1} = \{i-1\} \cup \{i+1, i+4, \dots, i-7, i-4\}$. Thus
 $T \subset A_{i+2}$ and $T \subset A_{i-1}$.

88 Let j in $\{1, \dots, n-1\}$. Let us prove that the vertices of C_m^j are dominated.
Indeed, by the previous remark and the lemma hypothesis, the vertices non domi-
90 nated by $S \cap C_m^j$ are dominated by $S \cap C_m^{j-1}$ (see Figure 1). For the same reasons
the vertices of C_m^0 are dominated by those of $S \cap C_m^0$ and $S \cap C_m^{-1}$.
92 □

Lemma 4 *Let $m = 3k_1 + 2$. If there exist integers $a, b \geq 0$ such that*

$$\begin{cases} a + b = n - 1 \\ \text{and} \\ a - 2b \equiv 2 \pmod{m} \text{ or } a - 2b \equiv m - 1 \pmod{m} \end{cases}$$

then $\gamma(C_m \square C_n) = n(k_1 + 1)$.

94 **Proof :** Consider a word $w = w_1 \dots w_{n-1}$ on the alphabet $\{1, -2\}$ with a
occurrences of 1 and b of -2 . Such a word exists, for example $w = 1^a(-2)^b$. We
96 can associate to w a set S of vertices of $C_m \square C_n$ by the following algorithm:

- $S \cap C_m^0 = A_0$
- 98 • For $i = 1$ to $n - 1$ do
begin
100 Let k such that $S \cap C_m^{i-1} = A_k$
If $w_i = 1$ let $k' = k + 1 \pmod{m}$ else $k' = k - 2 \pmod{m}$
102 $S \cap C_m^i := A_{k'}$
end

104 By construction S is a A -set. Notice that we have $S \cap C_m^{n-1} := A_{i_{n-1}}$ where $i_{n-1} \equiv$
 $\sum_{k=1}^{n-1} w_k \equiv a - 2b \pmod{m}$. Thus $i_{n-1} = 2$ or $i_{n-1} = m - 1 \pmod{m}$. By lemma 3
106 S is a dominating set. Furthermore, because S is a A -set, $|S| = n(k_1 + 1)$; thus by
theorem 2 it is minimum and we have $\gamma(C_m \square C_n) = n(k_1 + 1)$. □

108 **Theorem 5** *Let $m, n \geq 2$ and $m \equiv 2 \pmod{3}$. Let $k_1 = \lfloor \frac{m}{3} \rfloor$ and $k_2 = \lfloor \frac{n}{3} \rfloor$.*

- if $n = 3k_2$ then $\gamma(C_m \square C_n) = n(k_1 + 1)$
- 110 • if $n = 3k_2 + 1$ and $2k_2 \geq k_1$ then $\gamma(C_m \square C_n) = n(k_1 + 1)$
- if $n = 3k_2 + 1$ and $2k_2 < k_1$ then $\gamma(C_m \square C_n) > n(k_1 + 1)$
- 112 • if $n = 3k_2 + 2$ and $n \geq m$ then $\gamma(C_m \square C_n) = n(k_1 + 1)$.
- if $n = 3k_2 + 2$ and $n \leq m$ then $\gamma(C_m \square C_n) = m(k_2 + 1)$.

Proof : We will use lemma 4 considering the following integer solutions of

$$\begin{cases} a, b \geq 0 \\ a + b = n - 1 \\ a - 2b \equiv 2 \pmod{m} \text{ or } a - 2b \equiv m - 1 \pmod{m}. \end{cases}$$

- 114 • if $n = 3k_2$ then $k_2 \geq 1$. Take $a = 2k_2$ and $b = k_2 - 1$
 - if $n = 3k_2 + 1$ and $2k_2 \geq k_1$ then take $a = 2k_2 - k_1$ and $b = k_2 + k_1$
 - 116 • if $n = 3k_2 + 1$ and $2k_2 < k_1$ then $\gamma(C_m \square C_n) = \gamma(C_n \square C_m) \geq \frac{(2k_2+1)m}{2}$ by
theorem 2. Furthermore $\frac{(2k_2+1)m}{2} - n(k_1 + 1) = \frac{k_1}{2} - k_2 > 0$
 - 118 • if $n = 3k_2 + 2$ and $k_2 \geq k_1$ then take $a = 2k_2 - 2k_1$ and $b = k_2 + 2k_1 + 1$
 - if $n = 3k_2 + 2$ and $k_2 \leq k_1$ then use $\gamma(C_m \square C_n) = \gamma(C_n \square C_m)$.
- 120 □

3 The case $m \equiv 0 \pmod{3}$

122 In [12] Liu and al. conjectured the following formula:

Conjecture 6 *Let $k \geq 2$. Then $\gamma(C_{3k} \square C_n) = k(n + 1)$ for $n \not\equiv 0 \pmod{3}$.*

124 Our theorem 5 confirms the conjecture when $n \equiv 2 \pmod{3}$. Unfortunately the
formula is not always valid when $n \equiv 1 \pmod{3}$.

126 Indeed consider $C_{3k} \square C_4$. In [11] the following result is proved:

Theorem 7 *Let $n \geq 2$. Then $\gamma(C_4 \square C_n) = \frac{3n}{2}$ if $n \equiv 0 \pmod{8}$ and $\gamma(C_4 \square C_n) =$
128 $n + \lceil \frac{n+1}{2} \rceil$ otherwise*

130 We have thus $\gamma(C_{3k} \square C_4) = \gamma(C_4 \square C_{3k}) = 3k + \lceil \frac{3k+1}{2} \rceil$ when $k \not\equiv 0 \pmod{8}$.
On the other side the conjecture 6 proposes the value $\gamma(C_{3k} \square C_4) = 5k$. These two
numbers are different when $k \geq 3$.

132 References

- 134 [1] T.Y. Chang, W.E. Clark: “The Domination numbers of the $5 \times n$ and $6 \times n$
grid graphs”, *J. Graph Theory*, **17** (1993) 81-107.
- 136 [2] M. El-Zahar, C.M. Pareek: “Domination number of products of graphs”, *Ars
Combin.*, **31** (1991) 223-227.
- 138 [3] M. El-Zahar, S. Khamis, Kh. Nazzal: “On the Domination number of the
Cartesian product of the cycle of length n and any graph”, *Discrete App.
Math.*, **155** (2007) 515-522.
- 140 [4] R.J. Faudree, R.H. Schelp: “The Domination number for the product of
graphs”, *Congr. Numer.*, **79** (1990) 29-33.
- 142 [5] S. Gravier, M. Mollard: “On Domination numbers of Cartesian product of
paths”, *Discrete App. Math.*, **80** (1997) 247-250.
- 144 [6] B. Hartnell, D. Rall: “On dominating the Cartesian product of a graph and
 K_2 ”, *Discuss. Math. Graph Theory*, **24(3)** (2004) 389-402.
- 146 [7] T.W. Haynes, S.T. Hedetniemi, and P.J. Slater: *Fundamentals of Domination
in Graphs*, Marcel Dekker, Inc. New York, 1998.

- 148 [8] T.W. Haynes, S.T. Hedetniemi, and P.J. Slater eds.: *Domination in Graphs: Advanced Topics*, Marcel Dekker, Inc. New York, 1998.
- 150 [9] M.S. Jacobson, L.F. Kinch: “On the Domination number of products of graphs I”, *Ars Combin.*, **18** (1983) 33-44.
- 152 [10] S. Klavžar, N. Seifter: “Dominating Cartesian products of cycles”, *Discrete App. Math.*, **59** (1995) 129-136.
- 154 [11] J. Liu, X.D. Zhang, X. Chen, J.Meng: “The Domination number of Cartesian products of directed cycles”, *Inf. Process. Lett.*, **110(5)** (2010) 171-173.
- 156 [12] J. Liu, X.D. Zhang, X. Chen, J.Meng: “On Domination number of Cartesian product of directed cycles”, *Inf. Process. Lett.*, **111(1)** (2010) 36-39.