



Natural variation of root hydraulics in *Arabidopsis* grown in normal and salt-stressed conditions.

Moira Sutka, Guowei Li, Julie Boudet, Yann Boursiac, Patrick Doumas,
Christophe Maurel

► To cite this version:

Moira Sutka, Guowei Li, Julie Boudet, Yann Boursiac, Patrick Doumas, et al.. Natural variation of root hydraulics in *Arabidopsis* grown in normal and salt-stressed conditions.. *Plant Physiology*, 2011, 155 (3), pp.1264-76. 10.1104/pp.110.163113 . hal-00576512


HAL Id: hal-00576512

<https://hal.science/hal-00576512>

Submitted on 29 May 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Supplemental Figure 1. Relationship among the 13 accessions between Lp_r and its relative inhibition by mercury (\blacktriangle), propionic acid (\square), and azide (\bullet).

The graph shows a plot of relative inhibition data (Figure 2) against mean Lp_r (Figure 1).



Supplemental Figure 2. Absolute inhibition (black bars) and residual values (grey bars) of LPr after exposure of roots to mercury, propionic acid or azide inhibiting treatments (see text and Figure 2 for experimental details).

Absolute values were recalculated from mean LPr (Figure 1) and relative inhibition (Figure 2) data.



Supplemental Figure 3. Relative transcript abundance of *PIPs* in 13 accessions.

Data complimentary to Figure 4, showing the complement of expression data of the 13 *PIP* genes. Same legend as in Figure 4. *PIP1;1* : $\sigma_{RE} = 0.28$, *PIP1;2* : $\sigma_{RE} = 0.20$, *PIP1;4* : $\sigma_{RE} = 0.27$, *PIP1;5* : $\sigma_{RE} = 0.41$, *PIP2;3* : $\sigma_{RE} = 0.32$, *PIP2;4* : $\sigma_{RE} = 0.36$, *PIP2;5* : $\sigma_{RE} = 0.35$, *PIP2;7* : $\sigma_{RE} = 0.50$, *PIP2;8* : $\sigma_{RE} = 0.50$.

Supplemental Table 1. Correlation analysis of PIP transcript abundance vs. L_p

The table shows the Pearson correlation coefficient (R_{Pearson}) derived from a linear regression between L_p values and relative transcript abundance of the indicated genes. The corresponding Spearman's rank correlation coefficients (R_{Spearman}) are also indicated. Significant correlations ($P < 0.05$) are indicated in bold red.

Gene	R_{Pearson}	R_{Spearman}
<i>PIP1;1</i>	0.105	-0.066
<i>PIP1;2</i>	0.267	0.148
<i>PIP1;3</i>	-0.095	0.060
<i>PIP1;4</i>	-0.191	-0.165
<i>PIP1;5</i>	-0.245	-0.236
<i>PIP2;1</i>	0.016	0.137
<i>PIP2;2</i>	0.277	0.286
<i>PIP2;3</i>	0.037	0.088
<i>PIP2;4</i>	-0.154	-0.165
<i>PIP2;5</i>	-0.001	-0.0165
<i>PIP2;6</i>	-0.435	-0.445
<i>PIP2;7</i>	-0.035	0.027
<i>PIP2;8</i>	-0.589	-0.473
<i>PIP2;6 + PIP2;8</i>	-0.601	-0.637

Supplemental Table 2. Water relation and morphological parameters of root cortical cells under control and salt stress conditions.

Accession	Condition ^a	P_e ^b (MPa)	$T_{1/2}$ ^c (s)	ε ^d (MPa)	Lp_{cell} ^e ($10^{-6} \text{ m s}^{-1} \text{ MPa}^{-1}$)	n ^f	cell length ^g (μm)	cell diameter ^g (μm)
Blh-1	control	0.40 ± 0.01	0.79 ± 0.06	10.18 ± 0.96	1.02 ± 0.12	20	240.7 ± 8.1	42.1 ± 0.9
	salt	0.20 ± 0.02	2.00 ± 0.25	6.24 ± 0.52	0.71 ± 0.09	21	(45)	(45)
Col-0	control	0.32 ± 0.01	1.07 ± 0.07	5.48 ± 0.05	1.14 ± 0.08	22	155.0 ± 4.5	38.7 ± 0.8
	salt	0.16 ± 0.01	2.42 ± 0.18	2.70 ± 0.17	0.90 ± 0.09	20	(52)	(52)
Ct-1	control	0.53 ± 0.15	1.09 ± 0.09	6.55 ± 0.54	1.05 ± 0.18	17	170.8 ± 5.7	44.1 ± 1.0
	salt	0.19 ± 0.02	1.10 ± 0.15	4.14 ± 0.39	1.43 ± 0.19	14	(38)	(38)
Di-M	control	0.36 ± 0.01	1.02 ± 0.07	7.72 ± 0.058	0.97 ± 0.07	27	194.4 ± 7.2	42.6 ± 0.5
	salt	0.22 ± 0.01	1.98 ± 0.18	5.38 ± 0.26	0.71 ± 0.06	21	(36)	(36)
Mr-0	control	0.38 ± 0.02	0.98 ± 0.09	6.09 ± 0.70	1.43 ± 0.24	16	182.8 ± 5.4	43.6 ± 0.9
	salt	0.16 ± 0.01	1.09 ± 0.13	4.15 ± 0.43	1.66 ± 0.024	17	(54)	(54)

-

^aControl: plants were maintained in a standard hydroponic culture medium and the excised root segment to be punctured was transferred into a standard cell pressure probe solution for 10-40 min prior to measurement. Salt: roots of intact plants were bathed for 4 h in a hydroponic culture medium complemented with 100 mM NaCl and the excised root segment was bathed in a cell pressure probe solution complemented with 100 mM NaCl for 10-40 min prior to measurement.

^bMean stationary turgor pressure ± SE.

^cHalf-time of water exchange ± SE.

^dMean volumetric elastic modulus ± SE.

^eCell hydraulic conductivity ± SE.

^fNumber of cells used for measurements of water relation parameters in the indicated conditions.

^gMean morphological parameters ± SE measured on the indicated number of cells (in bracket) in roots bathing in a standard hydroponic culture medium. These parameter values were used for calculation of Lp_{cell} in both standard and salt stress conditions.

Supplemental Table 3 Sequences of TIP-specific primer pairs used for real time RT-PCR amplification.

Gene	Primer	Amplicon size (bp)
<i>TIP1;1</i>	Forward*: 5'- ⁸²⁷ CTCCCAACCACAGACTACTGAA ⁸⁴⁸ -3' Reverse: 5'- ⁹⁶² GCACGATCATAAACCCCTTG ⁹⁴³ -3'	136
<i>TIP1;2</i>	Forward: 5'- ⁹⁹⁴ GCATCGTAATGGGTTCTGG ¹⁰¹³ -3' Reverse: 5'- ¹¹²¹ TACAATTGCACAAAAGCCTTCC ¹¹⁰⁰ -3'	128
<i>TIP2;2</i>	Forward: 5'- ⁹⁰⁶ AGCTCCCACCAACAGAAAGCTA ⁹²⁶ -3' Reverse: 5'- ¹⁰⁰⁸ TTTGGAAAGAAACGAGGACCA ⁹⁸⁹ -3'	103
<i>TIP2;3</i>	Forward: 5'- ⁹⁴² GTGAGATCCGAGTGTATTGACTG ⁹⁶⁵ -3' Reverse: 5'- ¹⁰⁶⁰ TTTTCTTTCTCTACATACAATCTTGC ¹⁰³⁴ -3'	119

All primers were designed within or close to the 3'untranslated transcribed region. The numberings refer to positions from the initiating ATG codon in the genomic sequences.