



HAL
open science

Nonagenarian Surgical Admissions for the acute abdomen: Who benefits?

Zaher Toumi, Annabel Kesterton, Arnab Bhowmick, Alan Beveridge, Nigel A. Scott

► **To cite this version:**

Zaher Toumi, Annabel Kesterton, Arnab Bhowmick, Alan Beveridge, Nigel A. Scott. Nonagenarian Surgical Admissions for the acute abdomen: Who benefits?. *International Journal of Clinical Practice*, 2010, 64 (11), pp.1570. 10.1111/j.1742-1241.2010.02468.x . hal-00576625

HAL Id: hal-00576625

<https://hal.science/hal-00576625>

Submitted on 15 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Nonagenarian Surgical Admissions for the acute abdomen:
Who benefits?**

Journal:	<i>International Journal of Clinical Practice</i>
Manuscript ID:	IJCP-02-10-0099.R1
Wiley - Manuscript type:	Original Paper
Date Submitted by the Author:	08-May-2010
Complete List of Authors:	Toumi, Zaher; Pennine Acute Hospitals, General Surgery Kesterton, Annabel; Royal Preston Hospital, General Surgery Bhowmick, Arnab; Royal Preston Hospital, Colorectal Surgery Beveridge, Alan; Royal Preston Hospital, Colorectal Surgery Scott, Nigel; Royal Preston Hospital, Colorectal Surgery
Specialty area:	



Title page**Title of article:**

Nonagenarian Surgical Admissions for the acute abdomen: Who benefits?

Authors:

Z Toumi¹, A Kesterton², A Bhowmick², AJ Beveridge², NA Scott²

Institutions:

Department of Surgery at the Royal Preston Hospital

Correspondence to:

Mr Nigel Scott

Royal Preston Hospital

Preston

PR2 9HT

nigel.scott@lthtr.nhs.uk – 01772 522261

Sources of financial support:

None

Category:

Original article

Keywords:

Emergency Surgery, Nonagenarian, acute abdomen.

Conflicts of interest:

The authors confirm that they do not have any potential or actual personal, financial or political interest in the material.

Funding:

No funding was sought or received.

Word Count

868 (excluding abstract)

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction

Patients 90 years and older, form an increasing proportion of the general population. Outcomes of their acute surgical admissions are not well documented.

Methods and Materials.

Surgical management of 49 consecutive nonagenarian admissions (median age 92yrs) with an acute abdomen compared to the management and outcome of 50 younger patients (median age 53.5) admitted with a suspected acute abdomen over the same period.

Results

Nonagenarian group were mainly female (71% v 50% $p=0.003$). The use of laboratory investigations and imaging was similar for the patients aged over 90 and the younger patients – although proportionately fewer nonagenarians were investigated by abdominal CT scan (8% v 24%).

Of the 49 nonagenarian patients admitted, only 4% ($n=2$) were operated on. By contrast, 38% ($n=19$) of patients aged 50-59 ($P=0.0001$) underwent a surgical intervention.

A much greater proportion of nonagenarians died in hospital than patients in the 50-59 comparator group (16% nonagenarians versus 4% comparator patients; $p=0.04$). The very large majority of survivors in both age groups were

1
2
3 discharged back to their pre-admission domicile (39 (95%) nonagenarians
4
5 versus 46 (96%) comparator 50-59 year group).
6
7
8
9

10 **Conclusions**

11
12 In this study, when compared with younger patients very few nonagenarian
13 patients (2%) with a suspected acute abdomen benefited from surgical
14
15 admission. Instead the large majority of nonagenarians either died or were
16
17 discharged back to their home address without surgery.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

What is known about the topics?

- Nonagenarians represent a growing part of the population.
- Nonagenarian acute surgical admissions have increased and are likely to increase further.
- Selected patients in their nineties can be successfully operated on as emergencies and discharged from hospital alive.

What does this study add?

- Most Nonagenarians are unlikely to benefit from hospital surgical admission.
- Palliative treatment must remain a major consideration for nonagenarians with significant comorbidity and/or severe surgical illness.
- Future admission protocols for nonagenarians with a suspected acute abdomen should usefully incorporate assessment of "illness severity" before the decision to admit.

1
2
3 **Text:**
4
5
6
7

8 **INTRODUCTION**

10 The number of nonagenarians in England and Wales continues to increase
11
12 (1). An inevitable consequence of this increase in life expectancy is an
13 increasing workload of surgical illness amongst the very elderly. (2) While the
14 outcomes of surgery in selected elderly patients are well documented (3-6),
15 less clear is whether the overall management of the nonagenarian acute
16 abdomen, including the frequency of the decision to operate, is the same as in
17 younger patients.
18
19
20
21
22
23
24
25
26
27
28

29 In this study we documented the surgical management of nonagenarians and
30 a younger (50-59 years old) comparator population admitted with a diagnosis
31 of an acute abdomen to the same hospital over the same time period. Our
32 aim was to determine if the management and outcomes of the nonagenarian
33 acute abdomen differ significantly from those of younger patients.
34
35
36
37
38
39
40
41
42

43 **METHODS AND MATERIALS**

44
45 49 consecutive patients aged 90 years or over with a clinical diagnosis of
46 acute abdomen, were admitted during a nine month period (2007) to the
47 Surgical Assessment Unit at Royal Preston Hospital. Over the same period
48 306 patients aged between 50 and 59 years of age were also admitted with an
49 acute abdomen. For the purposes of this study, a "young" comparator
50 population (n=50; median age 53.5 yrs) was selected from every sixth
51 admission in this latter age group.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 Each patient had details of age, gender and pre-admission domicile recorded.
7
8 In addition details of the use of laboratory and radiological investigations,
9
10 surgical operations, length of stay, outcome and discharge destination were
11
12 collected. The nonagenarian population and 50-59 year comparator
13
14 population were compared, to determine any differences in the use of
15
16 investigations, surgical intervention and outcome, including final discharge
17
18 address from the hospital. Chi square test was used for comparison between
19
20 the two groups.
21
22
23
24

25 RESULTS

26 Patients and Management.

27
28
29
30
31 The nonagenarian group had significantly more females (71%) than the
32
33 comparator group (50%). The frequency of haematological and biochemical
34
35 investigations was the same for patient groups as was the use of chest
36
37 radiographs, abdominal films and abdominal ultrasound. However,
38
39 significantly fewer nonagenarians underwent CT scan. (Table 1)
40
41
42 Of the 49 nonagenarian patients admitted, only 4% (n=2) were operated on,
43
44 one had a Hartmann's procedure for perforated diverticular disease and one
45
46 died after a repair of an incarcerated femoral hernia. By contrast, 38% (n=19)
47
48 of patients aged 50-59 (P=0.0001) underwent a surgical operation.
49
50
51

52 Outcome

53
54
55 The median hospital stay for the nonagenarians was very similar to that of the
56
57 50-59 age group (Table 2). However, a much greater proportion of
58
59 nonagenarians died in hospital than patients in the 50-59 comparator group
60

1
2
3 (16% nonagenarians versus 4% comparator patients; $p=0.04$). The same
4
5 proportion of surviving nonagenarians and younger comparator patients were
6
7 discharged back to their original home address. Table 3 summarises the pre-
8
9 admission and post-discharge domicile of the nonagenarian group. The
10
11 discharge diagnoses of the nonagenarian group are summarised in Table 4.
12
13
14

15 16 17 **DISCUSSION**

18
19 Not surprisingly, female patients predominate amongst very elderly patients
20
21 admitted with a suspected acute abdomen (6). In addition, we found the use
22
23 of laboratory and most radiological investigations in nonagenarians, to
24
25 determine the cause of the acute abdomen, was both similar to their use in
26
27 the younger comparator population and to their use in previously reported
28
29 elderly surgical groups (7). An exception was that significantly fewer
30
31 nonagenarians had an abdominal CT scan.
32
33
34
35
36
37

38
39 Previous studies tend to emphasise that selected patients in their nineties can
40
41 be successfully operated on as emergencies and discharged from hospital
42
43 alive (2-6). In favourable reports, the operative mortality of emergency
44
45 surgery in nonagenarians is between 20 and 30% (3-6). By contrast, the
46
47 striking finding here was the rarity of surgical operation amongst the
48
49 nonagenarians (4% v 38% of comparator group; $p=0.001$) – despite the higher
50
51 nonagenarian death rate after admission (16% v 4% of comparator group;
52
53 $p=0.04$). Others have also found that a smaller proportion of very elderly with
54
55 an acute abdomen are suitable for surgical intervention when compared to
56
57 younger patients (8).
58
59
60

1
2
3
4
5
6 Catastrophic outcomes after abdominal surgery in the elderly are closely
7
8 related to advanced comorbidity (6, 9) – NCEPOD (10) described 82% of
9
10 nonagenarian surgical deaths as being associated with ASA III or worse.
11
12 Thus significant comorbidity and poor performance status (11) means that
13
14 surgical intervention for the acute abdomen in many very elderly patients is
15
16 unlikely to be of benefit. This perception of poor patient outcomes in our
17
18 experience, led to 98% of nonagenarians admitted with an acute abdomen
19
20 either dying in hospital without surgical intervention or after a median stay of 4
21
22 days, being discharged back to their admission address.
23
24
25
26
27
28

29
30 It is strongly our view that neither age alone nor subjective rating systems (12,
31
32 13) can determine the appropriateness of surgical intervention in the individual
33
34 elderly patient. However, the place of palliative treatment must remain a
35
36 major consideration for nonagenarians with significant comorbidity and/or
37
38 severe surgical illness (9). Future admission protocols for nonagenarians with
39
40 a suspected acute abdomen would usefully incorporate assessment of “illness
41
42 severity” before the decision to admit –the aim being to avoid the upset and
43
44 distress of admission for the very many elderly patients who are unlikely to
45
46 benefit from hospital surgical admission. We plan to study prospectively and
47
48 comprehensively the management and outcome of this growing group of
49
50 patients in order to help in creating appropriate admission protocols.
51
52
53
54
55
56
57
58
59
60

REFERENCES

- 1) National Statistics Online <http://www.statistics.gov.uk/default.asp>
- 2) Warner MA, Hosking MP, Lobdell CM, Offord KP, Melton LJ 3rd. Surgical procedures among those > 90 years of age. A population-based study in Olmsted County, Minnesota, 1975–1985. *Ann Surg* 1988;270:380–6.
- 3) Cohen JR, Johnson H, Eaton S, Sterman H, Wise L. Surgical procedures in patients during the tenth decade of life. *Surgery* 1988;104(4):646–51.
- 4) Damhuis RAM, Meurs CJC, Meijer WS. Postoperative mortality after cancer surgery in octogenarians and nonagenarians: results from a series of 5 390 patients. *World J Surg Oncol* 2005;3:71.
- 5) Arenal JJ, de Teresa G, Tinoco C, Toledano M, Said A. Abdominal Surgery in Nonagenarians: Short-Term Results *Surg Today*. 2007;37(12):1064-7.
- 6) Arenal JJ, Bengoechea-Beeby M. Mortality associated with emergency abdominal surgery in the elderly. *Can J Surg*. 2003 Apr;46(2):111-6
- 7) Geloven AAW, Biesheuvel TH, Luitse JSK et al. Hospital Admissions of Patients aged over 80 with acute abdominal complaints. *Eur J Surg* 2000; 166:866-871

1
2
3 8 Walsh TH. Audit outcome of major surgery in the elderly. Br J Surg 1996;
4
5 83(1):92-7
6
7

8
9 9) Zerbib P, Kulick JF, Lebuffe G, Khoury-Helou A, Plenier I, Chambon JP.
10
11 Emergency major abdominal surgery in patients over 85 years of age. World
12
13 J Surg. 2005 Jul;29(7):820-5
14
15

16
17
18 10) Extremes of Age – The 1999 Report of the National Confidential Enquiry
19
20 inot Perioperative Death. NCEPOD Published 17 November 1999 by the
21
22 National Confidential Enquiry into Perioperative Deaths 35-43 Lincoln's Inn
23
24 Fields, London WC2A 3PN
25
26
27

28
29
30 11) Masuo K, Kumagai K, Tanaka T et al “Physiological” age as an outcome
31
32 predictor fo abdominal surgery in elderly patients. Surgery Today
33
34 1998;28(10): 997-1000.
35
36
37

38
39
40 12) Farquharson SM, Gupta R, Heald RJ, Moran BJ. Surgical decisions in the
41
42 elderly: the importance of biological age. J R Soc Med 2001: 94;232-235
43
44
45

46
47 13) Seymour DG Letter “Surgical decisions in the elderly” J R Soc Med 2001:
48
49 94;370-371
50
51
52
53
54
55
56
57
58
59
60

Tables:

Table 1 – Investigation and Management

	Nonagenarians n=49	Comparator Group n=50	p
Gender	14M:35F	25M:25F	0.003
Age Median (range)	92 (90-99)	53.5 (50-59)	<0.0001 (Mann Whitney)
Biochemical Investigations	44 (90%)	47 (94%)	ns
Haematological	45 (92%)	48 (96%)	ns
Chest Radiograph	26 (53%)	25 (50%)	ns
Plain abdominal film	34 (69%)	26 (52%)	ns
Ultrasound scan	9 (18%)	10 (20%)	ns
CT abdomen and pelvis	4 (8%)	12 (24%)	0.03
Surgical Operation	2 (4%)**	19(38%)	0.0001

(** 1 patient died after surgery)

Table 2 – Outcomes

	Nonagenarians n=49	Comparator Group n=50	p
In-patient Mortality	8(16%)*	2(4%)**	0.04 [Fisher's Exact Probability Test}
Hospital Stay (days) Median (range)	4 (2-72)	3 (0.1-90)	ns
Discharge to preadmission address	39/41 (95%)	46/48 (96%)	ns
Discharge to a new nursing facility	2/41 (5%)	2/48 (4%)	ns

(*Nonagenarians - 1 small bowel infarction, 1 CVA, 1 obstruction, 1

pneumonia, 2 cases of peritonitis, 1 postoperative death: 1 septicaemia

**Comparator group 1 pneumonia & cirrhosis, 1 carcinomatosis)

Table 3- Pre-admission and post- discharge domicile of the nonagenarian group

Pre-admission domicile	Number of patients	Outcome		
		Death	Discharge destination	
			Pre-admission address	Higher level of care
Nursing home	3	1	2	0
Residential home	15	2	13	0
Sheltered housing	3	1	2	0
Patient's own home	28	4	22	2
Total	49	8	39	2

Table 4- Final diagnoses of discharged nonagenarians*

Method of diagnosis	Number of patients	Diagnosis
CT**	4	2 diverticular disease, 1 cholangiocarcinoma, 1 small bowel obstruction
USS**	5	3 cholelithiasis, 1 pancreatic Ca, 1 normal
Endoscopy**	2	1 Sigmoid volvulus, 1 colon cancer
Operation**	1	1 Perforated diverticular disease
Laboratory investigations**	3	3 Pancreatitis
Plain imaging**	2	1 Pseudo-obstruction, 1 constipation
Clinical assessment only	10	4 uncomplicated hernia, 4 rectal prolapse, 2 haemorrhoids
Total	27 *	

*14 patients did not have investigations to positively confirm the discharge diagnoses (6 non-specific abdominal pain, 4 diverticular disease, 2 constipation, 1 biliary colic, 1 adhesions)

** In combination with clinical assessment