

Fumonisin B2 production by Aspergillus niger from Thai coffee beans

Paramee Noonim, Warapa Mahakarnchanakulb, Kristian F Nielsen, Jens

Frisvad, Robert A Samson

▶ To cite this version:

Paramee Noonim, Warapa Mahakarn
chanakulb, Kristian F Nielsen, Jens Frisvad, Robert A Samson. Fumonisin B2 production by Aspergillus niger from Thai coffee beans. Food Additives and Contaminants, 2009, 26 (01), pp.94-100. 10.1080/02652030802366090 . hal-00577319

HAL Id: hal-00577319 https://hal.science/hal-00577319

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Food Additives and Contaminants



Fumonisin B2 production by Aspergillus niger from Thai coffee beans

Journal:	Food Additives and Contaminants
Manuscript ID:	TFAC-2008-013.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	25-Jun-2008
Complete List of Authors:	Noonim, Paramee; Kasetsart University, Department of Food Science and Technology; Prince of Songkla University, Surat Thani Campus, Faculty of Technology and Management Mahakarnchanakulb, Warapa; Kasetsart University, Department of Food Science and Technology Nielsen, Kristian; Center for Microbial Biotechnology, Department of Systems Biology Frisvad, Jens; Center for Microbial Biotechnology, Department of Systems Biology Samson, Robert; CBS Fungal biodiversity centre
Methods/Techniques:	HPLC
Additives/Contaminants:	Fumonisins
Food Types:	Coffee

SCHOLARONE[™] Manuscripts

2		
3 4	1	Fumonisin B ₂ production by Aspergillus niger from Thai coffee beans
5 6 7	2	
8 9	3	
10 11 12	4	Paramee Noonim ^{a,b,c} , Warapa Mahakarnchanakul ^b , Kristian F. Nielsen ^d , Jens C. Frisvad ^d
12 13 14	5	and Robert A. Samson ^a
15 16 17	6	
18 19	7	
20 21	8	^a CBS Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands
22 23 24	9	^b Department of Food Science and Technology, Faculty of Agro-Industry, Kasetsart University,
25 26	10	10900 Bangkok, Thailand
27 28	11	^c Faculty of Technology and Management, Prince of Songkla University, Surat Thani Campus,
29 30 31	12	84100 Surat Thani, Thailand
32 33	13	^c Center for Microbial Biotechnology, Department of Systems Biology, Building 221, Technical
34 35	14	University of Denmark, DK-2800 Kgs Lyngby, Denmark
36 37 38	15	
39 40	16	
41 42 43	17	
44 45	18	
46 47	19	
48 49	20	Corresponding author: RA Samson
50 51	21	CBS Fungal Biodiversity Centre,
51 52	22	P.O. Box 85167, 3508 AD Utrecht, The Netherlands
53 54	23	Telephone + 31 30 2122656
55 56	24	Telefax + 31 30 2512097
57 58 59 60	25	E-mail: <u>samson@cbs.knaw.nl</u>

Abstract

During 2006 and 2007, a total of 64 Thai dried coffee bean samples (Coffea arabica) from two growing sites of Chiangmai Province, and 32 Thai dried coffee bean samples (*Coffea canephora*) from two growing sites of Chumporn Province, Thailand, were collected and assessed for fumonisin contamination by black Aspergilli. No Fusarium species known to produce fumonisin were detected, but black Aspergilli had high incidences on both Arabica and Robusta Thai coffee beans. Liquid chromatography (LC) with high resolution mass spectrometric (HRMS) detection showed that 67% of A. niger isolates from coffee beans were capable of producing fumonisins B₂ (FB₂) and B₄ when grown on Czapek Yeast Agar with 5% NaCl. Small amounts (1-9.7 ng/g) of FB₂ were detected in 7 of 12 selected coffee samples after ion-exchange purification and LC-MS/MS detection. Two samples also contained FB₄. This is the first record of freshly isolated Aspergillus niger strains producing fumonisins and the first report on the natural occurrence of FB₂ and FB₄ in coffee.

Key words: Fumonisin B₂, Coffee beans, *Aspergillus niger*

1 Introduction

Fumonisins are carcinogenic mycotoxins produced by several *Fusarium* species (Gelderblom *et al.*, 1988; Marin *et al.*, 2004) and have been reported in many food commodities especially corn (Marin *et al.*, 2004). Fumonisins has been reported to cause a fatal disease in horses (leukoencephalomalacia) (Marasas *et al.*, 1988), pulmonary edema in pigs (Haschek *et al.*, 2001) and possibly esophageal cancer in humans (Yoshizawa *et al.*, 1994). Even though fumonisins are less acutely toxic compared to aflatoxins, they could be found in high concentration of mg/kg in corn compared to µg/kg for aflatoxins.

Coffee is one of the most consumed beverages in the world, and have been reported to be contaminated with ochratoxin A. Aflatoxin contamination in coffee beans has also been reported (Soliman, 2002). However, there have been many publications discussing the ecology of ochratoxin-producing fungi, manipulation of environmental factors (Batista et al., 2003; Palacios-Cabrera et al., 2004; Esteban et al., 2006) and control strategies to prevent or reduce ochratoxin contamination (Suarez-Quiroz, 2005). Ochratoxigenic species in coffee beans are generally known from Aspergillus species of the section Circumdati and Nigri (Joosten et al., 2001; De Moraes et al., 2003; Martins, 2005; Leong et al., 2007; Ilic et al., 2006; Taniwaki et al., 2006) and various methods have been developed for the detection of ochratoxin-producing fungi and ochratoxin contamination in coffee beans (Patiño et al., 2005; Lobeau et al., 2005; Satori et al., 2006).

Recently, fumonisin B₂ was detected in agar cultures of four important isolates of *Aspergillus niger* (Frisvad *et al.*, 2007) including the culture ex type and three full genome sequenced
cultures (Baker, 2006). It was found that while *Fusarium verticillioides* produces fumonisin B₁,
B₂ and B₃ on plant extract agars, but *A. niger* produces fumonisin B₂ only on agar media with

high amounts of carbohydrate or NaCl.

As the results of a survey on ochratoxigenic species in Thai coffee beans, we found that black Aspergilli including *A. niger* were the predominant contaminating fungi. In this study, we investigate the presence of fumonisin producing black Aspergilli on coffee beans, as well as fumonisin production in the beans them selves.

- 8 Material and Methods
- 9 Sampling

There are two coffee growing regions in Thailand, the Northern and Southern region, which are different in varieties of coffee grown, geographical condition and climate during harvesting. In this study, two types of Arabica coffee beans, parchment and green coffee beans, from the North were collected from two selected farms in two different growing sites. Two types of Robusta coffee bean, dried coffee cherries and green coffee beans, from the South were collected from two selected farms in two different growing sites. Four samples of 0.5-1 kg of each type per farm were collected. A total of 64 samples were collected during the two harvesting year 2006 and 2007.

19 Mycological analysis

A total of 50 beans per sample were plated directly (5 per plate) onto Dichloran 18% Glycerol Agar (DG18) plates and Malt Extract Agar (MEA) plates (Samson *et al.*, 2004a) with and without surface sterilization. The plates were incubated for 5-7 days at 25°C, and then inspected for fungal growth. Of the many species encounter on these plates (Noonim et al., in prep.) potentially fumonisin producing species of *Aspergillus* section *Nigri* were isolated and identified to species level using morphology, physiology and molecular characteristics (Samson *et al.*,

1 2007) and kept in collection for further studies.

Determination of extrolite production by liquid chromatography-UV-mass spectrometry of fungal cultures

5 Representative isolates of each *Aspergillus* species in section *Nigri* were inoculated in Czapek
6 Yeast Agar with 5% salt (NaCl) (CYAS) medium (Frisvad and Samson, 2004) and incubated for
7 days at 25°C. Subsequently, 5 plugs of culture (1 cm²) were sampled, and moved to a 2-ml vial,
8 where it was extracted using ultra-sonication for 60 min with 0.75 ml 75% methanol, and
9 subsequently filtered trough a 0.45 µm syringe filter (Frisvad *et al.* 2007).

Solvents were HPLC grade and all other chemicals were analytical grade unless otherwise stated. Water was purified from a Milli-Q system (Millipore, Bedford, MA). LC-DAD-HRMS was performed on an Agilent 1100 system equipped with a photo diode array detector (DAD) and a 50×2 mm i.d., 3 µm, Luna C₁₈ II column (Phenomenex, Torrance, CA). The LC system was coupled to a LCT orthogonal time-of-flight mass spectrometer (Waters-Micromass, Manchester, UK), with a Z-spray ESI source (2).

Samples were analyzed in ESI⁺ using a water-CH₃CN gradient system starting with 0.3 ml/min flow of 30% CH₃CN which was increased linear to 60% in 5 min, then increased to 100% in 1 min while also increasing the flow to 0.5 ml/min, holding this for 2 min. The water was buffered with 10 mM ammonium formate and 20 mM formic acid and the CH₃CN with 20 mM formic acid (Nielsen and Smedsgaard, 2003; Nielsen et al., 2005). One scan function (1 s) was used with a potential difference of 50 V between the skimmers and using a scan range of m/z 100 to 900. Reference standards of fumonisin B₁, B₂ and B₃, AAL toxin TB₁ and TA₁, Malformins A, B and C, ochratoxin A, and Asperazine were also co-analyzed in the sequences. Source of 1 reference standards: Certified standards of 50 μ g/mL of FB₁ and fumonisin B₂ were obtained 2 from Biopure, Tulln, Austria. FB₃ was a gift from Dr. Michael Sulyok, Center for Analytical 3 Chemistry (IFA-Tulln, Austria) and other reference standards were available from previous 4 studies in our laboratory (Nielsen and Smedsgaard, 2003).

6 The presence of fumonisin B₂ was detected in ESI⁺ from the reconstructed ion chromatograms of 7 the $[M+H]^+$ ion at m/z 706.39-706.41 (calc. mass 706.4014). Other metabolites were detected as 8 the predominant ion in extracted ion chromatograms (± m/z 0.01).

9 A few samples were also analyzed by LC-tandem MS as described for the coffee samples below,
10 except that the MS was operated in ESI⁺ daughter ion scan mode using fragmentation potentials
11 from 20 to 50 V.

13 ELISA Screening of fumonisins in coffee beans

14 Coffee bean samples were assessed for fumonisin contamination by using RIDASCREEN[®] 15 Fumonisin ELISA test kits (r-biopharm) using the protocol for corn. All sample preparation and 16 test procedures were according to manufacturer's instructions. Specificity for fumonisin B_1 , B_2 17 and B_3 are 100, 40 and 100, respectively. The lower detection limit of the test kit was specified 18 as 25 µg/kg. Assay was not validated nor tested on spiked samples.

20 LC-MS/MS of fumonisins in coffee beans

The coffee bean samples were frozen by liquid nitrogen and grinded for 2 minus in a domestic electrical coffee grinder. Subsamples of 1.0 g were then shaken with 10.0 ml methanol-water (7:3 v/v) in a falcon tube for 15 min and centrifuged at 9000 g for 3 min. Then a 5 ml subsample transferred to a 100 mg Strata SAX column (Phenomenex) which had previously been sequentially conditioned with 1 ml methanol and 1 ml methanol-water (7:3 v/v). Columns were

Food Additives and Contaminants

washed with 3 ml methanol-water (7:3 v/v) and 2 ml methanol, and the fumonisins eluted with 2.4 ml methanol containing 1% acetic acid. Samples were then evaporated to dryness with nitrogen flow and redissolved in 250 μ l acetonitrile-water (3:7 v/v) (modified from the EN 13585:2001).

Sub-samples of 5 µl were analyzed by LC-MS/MS on an Agilent HP 1100 liquid chromatograph system (Waldbronn, Germany) coupled to a Quattro Ultima triple mass spectrometer (Micromass, Manchester, UK) with ESI source. The separations was performed on a Gemini C6-phenyl column (Phenomenex, 50×2 mm, 3 µm) fitted with a security guard system and using a linear gradient starting from 20 % acetonitrile in water (both 20 mM formic acid) to 55% acetonitrile for 6 minutes at a flow rate of 300 µL/min, which was then increase to 100% acetonitrile in 30 sec and a flow of 0.5 ml/min keeping this fro 3.5 min before returning to the start conditions in 6 min. Tandem mass spectrometry was performed in ESI⁺ at a source flow at 700 L/hr nitrogen at 350°C. Nitrogen was also used as collision gas, and the MS operated in MRM mode at the following transitions: FB₂ quantifier m/z 706 \rightarrow 336 cone 50V, collision 40 V, dwell time 50 ms, qualifier m/z 706 \rightarrow 512 a, cone 50V, collision 25 V, dwell time 100 ms; FB₄ quantifier m/z 690 \rightarrow 320 cone 50V, collision 35 V, dwell time 50 ms, qualifier m/z 690 \rightarrow 514 a, cone 50V, collision 30 V, dwell time 100 ms; and FB₁ and position analogues quantifier m/z 722 \rightarrow 334 cone 50V, collision 40 V, dwell time 50 ms, qualifier m/z 722 \rightarrow 528 a, cone 50V, collision 25 V, dwell time 100 ms.

Quantification was done from spiked samples, which were spiked with 30 to 100 µl acetonitrile solutions to final concentrations of 50, 25, 12.5, 5.0, 3.75, 2.50, 1.25, 0.500, 0.375, 0.00 ng/g grinded coffee and stored for 1-4 days prior to extraction. Samples and spiked samples were extracted and analyzed 4 times on different days.

Results and Discussion

3 Mycological analysis and identification of fungal isolates

Of all coffee bean samples analyzed, none of typical fumonisin-producing species (*Fusarium* spp.) were detected. Besides Penicillia and other saprophytes, *Aspergillus* spp. of section *Circumdati* and *Nigri* were the predominant species in the Arabica coffee samples with 77 and 75% infestation, respectively. In the Robusta coffee samples, *Aspergillus* spp. section *Nigri* was the predominant one with approximately 100% infection.

A diversity of black Aspergilli were observed (Fig. 1) in Thai coffee beans, including, *A. niger*, *A. carbonarius*, *A. tubingensis*, *A. foetidus*, *A. aculeatinus* and *A. sclerotiicarbonarius*. The latter
two species were found to unrelated to the known taxa and proposed as new taxa (Noonim et al.,
2008).

15 Considering each type of coffee beans, there are differences in the mycobiota observed. Arabica 16 coffee had a higher incidence of *A. niger* and related taxa while in Robusta coffee, both *A.* 17 *carbonarius* and *A. niger* were the dominant species. *A. carbonarius* and *A. sclerotiicarbonarius* 18 were found only in Robusta coffee from Southern Thailand while *A. foetidus* was found only in 19 Arabica coffee from the Northern region (Table 1.). These differences could be due to 20 differences in the geography, climate and methods used for coffee processing in the two regions.

22 Mycotoxigenic potential of the Aspergillus species

Using LC-HRMS (Frisvad *et al.*, 2007) a total of 82 isolates from 6 species were analyzed. Only A. *niger* isolates were found to produce fumonisin B_2 as well as fumonisin B_4 (same retention time and tandem spectrum as from a *Fusarium* extract). FB₄ were relative to FB₂ levels in the

Food Additives and Contaminants

range 0-40% with most being in the 10-20% range. Most of the tested A. niger isolated from Thai coffee beans (13 out of 17 isolates) produced fumonisin B₂ in the CYAS culture medium in amounts of 0.4 to 2 μ g/cm² (Table 2). This indicates that these A. niger isolates may also produce fumonisin B₂ in coffee cherries or beans. All A. niger isolated from Northern Arabica coffee bean samples could produce fumonisin B₂, while in some isolates from Southern Robusta coffee samples no fumonisin B₂ was found. More molecular studies are needed in order to compare the differences in these isolates at the genotypic level. A high percentage of infection by A. niger as determined after surface disinfection of the green coffee beans indicated that A. niger actually grows actively in the coffee beans.

LC-HRMS detection of fumonisin B_2 has been shown to have detection limit of ca. 25 ng/cm² culture, and a relative standard deviation better than 30%, and an apparent recovery better than 80% (Frisvad, Nielsen et al. unpublished). A chromatogram example of fumonisin B₂ detection in A. niger 1F6 is shown in figure 2. Figure 3 shows the comparison of the tandem spectrum of FB₂ from A. niger and reference standard.

In agreement with Samson et al. (2007) most of the isolates of Aspergillus niger from coffee beans produced funalenone, kotanin, orlandin, aurasperone B and other naphtho-y-pyrones, tensidol B and pyranonigrin A (Table 3). Two isolates produced ochratoxin A (1F1 and 1F6) in addition to FB₂ and thus could produce two important mycotoxins.

Analysis of coffee bens

Screening for B type fumonisins with the RIDASCREEN® ELISA test kits also indicated that FB₂ was present in some of the coffee bean samples (results not shown) in levels up to 77 ng/g and it was thus decided to confirm this by LC-MS/MS which is much more specific. Since both

a quantifier and a qualifier ion were used the method earns 4 identification-points accordingly to
 Council Directive 96/23/EC, which is required for forbidden compounds.

4 The detection limit of the LC-MS/MS method was approx 0.5 ng/g in spiked sample and the 5 limit of quantification (LOQ) 1.25 ng/g (RSD < 30% for 4 replicates at this level). R² from the 6 calibration curves were in the 4 experiments >0.985 (7 detected levels).

LC-MS/MS showed that the ELISA results were false positives, and of the 12 samples analyzed (4 times each), the most contaminated one (Robusta, R1) contained 9.7 ng/g, and the 3 other (SO.1C, Ch.3P, R8) between 1.9 and 1.3 ng/g (RSD ca. 40%, 95% level), while 2 were positive but below LOQ (R7 and R9). Chromatographic separation between FB₁ and FB₂ was 0.98 min. R1 additionally contained FB4 as shown in figure 4 where the un-smoothed chromatographic profiles can be seen from the FB₂ and FB₄ identifications.

More experiments are needed for the detection of the *Aspergillus* fumonisin in food commodities.
An extensive survey of fumonisin producing black Aspergilli from other sources is in progress
(Frisvad, Nielsen and Samson, personal communication).

One of the great concerns with fumonisin contamination of maize is that very large amounts of fumonisin may be produced by *Fusarium* species, but in accordance with the results of Frisvad *et al.* (2007), *Aspergillus niger* needs a relatively large amount of carbohydrate in the substrate to produce high amounts of fumonisin B_2 . In contrast *Fusarium verticillioides* needs less carbohydrate to produce substantial amounts of fumonisins. Green coffee beans contain small amounts of carbohydrates, while coffee cherries contain some carbohydrate. Thus, widespread infection of *A. niger* in coffee beans does not necessarily represent a high risk for fumonisin

1							
2 3	1	contamination, which is also indicated from the concentrations detected in very limited number					
4 5	1	containination, which is also indicated from the concentrations detected in very initied number					
6 7	2	of samples analyzed.					
8 9	3						
10 11	4	References					
12 13	5	Baker, S. E., 2006. Aspergillus niger genomics: Past, present and into the future. Medical					
14 15 16	6	Mycology, 44, S17-S21.					
17 18	7	Batista, L. R., Chalfoun, S. M., Prado, G., Schwan, R. F., and Wheals, A. E, 2003. Toxigenic					
19 20	8	fungi associated with processed (green) coffee beans (Coffea arabica L.). International					
21 22 23	9	Journal of Food Microbiology, 85, 293-300.					
24 25	10	Commission of the European Communities. COMMISSION DECISION of 12 August 2002					
26 27 28	11	implementing Council Directive 96/23/EC concerning the performance of analytical					
20 29 30	12	methods and the interpretation of results. Brussels, Belgium: 2002.					
31 32	13 De Moraes, M. H. P., and Luchese, R. H., 2003. Ochratoxin A in coffee: Influen						
33 34 35	14	and drying processing procedures. Journal of Agricultural and Food Chemistry, 51,					
36 37	15	5824-5828.					
38 39 40	16	EN 13585:2001, Foodstuffs - Determination of fumonisins B_1 and B_2 in maize - HPLC method					
40 41 42	17	with solid phase extraction clean-up, CEN/TC 275 - Food analysis - Horizontal methods,					
43 44	18	European Committee for Standardization. Brussels, Belgium: 2001.					
45 46 47	19	Esteban, A., Abarca, M. L., Bragulat, M. R., and Cabanes, F. J., 2006. Study of the effect of					
48 49	20	water activity and temperature on ochratoxin A production by Aspergillus carbonariu					
50 51	21	Food Microbiology, 23, 634-640.					
52 53 54	22	Frisvad, J. C., and Samson, R. A., 2004. Polyphasic taxonomy of Penicillium subgenus					
55 56	23	Penicillium: A guide to identification of food and air-borne terverticillate Penicillia and					
57 58 59	24	their mycotoxins. Studies in Mycology, 49, 1-173.					
60	25	Frisvad, J. C., Smedsgaard, J., Samson, R. A., Larsen, T. O., and Thrane, U., 2007. Fumonisin					

1		
2 3	1	B ₂ production by Aspergillus niger Journal of Agricultural and Food Chemistry 55
4 5	1	D ₂ production by hisperginus higer, souther of rightendular and rood cholinsity, 55,
6 7	2	9727-9732.
8 9	3	Gelderblom, W. C. A., Jaskiewicz, K., Marasas, W. F. O., Thiel, P. G., Horak, R. M., Vleggaar,
10 11	4	R. and Kriek, N. P. J., 1988. Fumonisins - novel mycotoxins with cancer-promoting
12 13	5	activity produced by Fusarium moniliforme. Applied and Environmental Microbiology,
15 16	6	54, 1806-1811.
17 18	7	Haschek, W. M., Gumprecht, L. A., Smith, G., Tumbleson, M. E., and Constable, P. D., 2001.
19 20 21	8	Fumonisin toxicosis in swine: an overview of porcine pulmonary edema and current
22 23	9	perspectives. Enrironmental Health Perspectives, 109, 215-257.
24 25	10	Ilic, Z., Bui, T., Tran-Dinh, N., Dang, V., Kennedy, I., and Carter, D., 2007. Survey of
26 27 28	11	Vietnamese coffee beans for the presence of ochratoxigenic Aspergilli. Mycopathologia,
29 30	12	163, 177-182.
31 32	13	Joosten, H. M. L. J., Goetz, J., Pittet, A., Schellenberg, M., and Bucheli, P., 2001. Production of
34 35	14	ochratoxin A by Aspergillus carbonarius on coffee cherries. International Journal of
36 37	15	Food Microbiology, 65, 39-44.
38 39 40	16	Leong, S. L., Hien, L. T., An, T. V., Trang, N. T., Hocking, A. D., and Scott, E. S., 2007.
41 42	17	Ochratoxin A-producing Aspergilli in Vietnamese green coffee beans. Letters in Applied
43 44 45	18	Microbiology, 45, 301-306.
46 47	19	Lobeau, M., De Saeger, S., Sibanda, L., Barna-Vetró, I., and Van Peteghem, C., 2005.
48 49	20	Development of a new clean-up tandem assay column for the detection of ochratoxin A
50 51 52	21	in roasted coffee. Analytica Chimica Acta, 538, 57-61.
53 54	22	Marasas, W. F. O., Kellerman, T. S., Gelderblom, W. C. A., Coetzer, J. A. W., Thiel, P. G. and
55 56	23	van der Lugt, J. J., 1988. Leukoencephalomalacia in a horse induced by fumonisin B_1
ວ7 58 59	24	isolated from Fusarium moniliforme. Onderstepoort Journal of Veterinary Research, 55,
60	25	197-203.

Food Additives and Contaminants

1 2	
3 4 5	
6 7	
8 9	
10 11 12	
13 14	
15 16	
17 18 10	
20 21	
22 23	
24 25 26	1
26 27 28	1
29 30	1
31 32	1
33 34 35	1
36 37]
38 39	1
40 41 42]
43 44	1
45 46	1
47 48 49	2
50 51	2
52 53	2
54 55 56	2
57 58	2
59 60	

Marin, S., Magan, N., Ramos, A. J. and Sanchis, V. 2004. Fumonisin-producing strains of
 Fusarium: a review of their ecophysiology. Journal of Food Protection, 67, 1792-1805.

Martins, M. L., Martins, H. M., and Gimeno, A., 2003. Incidence of microflora and of
ochratoxin A in green coffee beans (*Coffea arabica*). Food Additives and Contaminants,
20, 1127-1131.

Nielsen, K. F.; and Smedsgaard, J., 2003. Fungal metabolite screening: database of 474 mycotoxins and fungal metabolites for de-replication by standardised liquid chromatography-UV-mass spectrometry methodology. Journal of Chromatography A, 1002, 111-136.

- Nielsen, K. F., Graefenhan, T., Zafari, D., and Thrane, U., 2005. Trichothecene Production by
 Trichoderma brevicompactum. Journal of Agricultural and Food Chemistry, 53, 8190 8196.
- Noonim, P., Mahakarnchanakul, W., Varga, J., Frisvad, J. C., and Samson, R. A., 2008. Two
 novel species of *Aspergillus* section *Nigri* from Thai coffee beans. International Journal
 of Systematic and Evolution Microbiology, (in press).
- Palacios-Cabrera, H., Taniwaki, M.H., Menezes, H.C., and Iamanaka, B.T., 2004. The
 production of ochratoxin A by *Aspergillus ochraceus* in raw coffee at different
 equilibrium relative humidity and under alternating temperatures. Food Control, 15, 531 535.

Patiño, B., González-Salgado, A., González-Jaén, M. T., and Vázquez, C., 2005. PCR detection assays for the ochratoxin-producing *Aspergillus carbonarius* and *Aspergillus ochraceus* species. International Journal of Food Microbiology, 104, 207-214.

- Samson, R. A., Hoekstra, E. H. and Frisvad, J. C., 2004a. Introduction to food and air-borne
 fungi. 7th ed. Centraalbureau voor Schimmelcultures, Utrecht.
- ⁰ 25 Samson, R. A., Houbraken, J. A. M. P., Kuijpers, A. F. A., Frank, J. M. and Frisvad, J. C., 2004b.

2	
3	1
4 5	-
6	2
7 8	3
9 10	1
11	4
12	5
14 15 16	6
10 17 19	7
10 19 20	8
20	0
22 23	9
24 25	10
26 27	11
28 29 20	12
30 31 32	13
33 34	1.0
35	14
36 37	15
38 39 40	16
40 41 42	17
43 44	18
45 46	19
47 48	• •
49	20
50 51	21
52 53	22
54 55	
56	
57 58	
59	
60	

New ochratoxin A or sclerotium producing species in Aspergillus section Nigri. Studies in Mycology, 50, 45-61. Samson, R. A., Noonim, P., Meijer, M., Houbraken, J., Frisvad, J. C. and Varga, J. 2007. Diagnostic tools to identify black Aspergilli. Studies in Mycology, 59, 129-146. Satori, D, Furlaneto, M. C., Martins, M. K., de Paula, M. R. F., Pizzirani-Kleiner, A. A., Taniwaki, M. H. and Fungaro, M. H. P., 2006. PCR method for the detection of potential ochratoxin-producing Aspergillus species in coffee beans. Research in Microbiology, 157, 350-354. Soliman, K. M., 2002. Incidence, level, and behavior of aflatoxins during coffee bean roasting and decaffeination. Journal of Agricultural and Food Chemistry, 50, 7477 -7481. Suarez-Quiroz, M., Gonzalez-Rios, O., Barel, M., Guyot, B., Schorr-Galindo, S., and Guiraud, J.-P., 2005. Effect of the post-harvest processing procedure on OTA occurrence in artificially contaminated coffee. International Journal of Food Microbiology, 103, 339-345. Taniwaki, M. H., Pitt, J. I., Teixeira, A. A., and Iamanaka, B. T., 2003. The source of ochratoxin A in Brazilian coffee and its formation in relation to processing methods. International Journal of Food Microbiology, 82, 173-179. Yoshizawa, T., Yamashita, A., and Luo, Y., 1994. Fumonisin occurrence in corn from high-risk and low-risk areas for human esophageal cancer in China. Applied and Environmental Microbiology, 60, 1626-1629.

1 2		
2 3 4	1	Figure 1. Diversity of fungal population in coffee beans from direct plating of two types of
5 6 7	2	coffee bean samples from 2 different regions of Thailand.
7 8 9	3	A. Arabica green coffee, MEA. B. Arabica green coffee, DG18. C. Robusta green coffee, MEA.
10 11	4	D. Robusta green coffee, DG18. E. Robusta cherries, DG18. F. Robusta cherries, MEA. G.
12 13	5	Arabica parchment coffee DG18. H. Arabica parchment coffee, MEA;
14 15 16	6	
17 18	7	Figure 2 . A. showing the total ion chromatogram (TIC) of a fumonisin B_1 and B_2 standard mix.
19 20 21	8	B. extracted ion chromatogram m/z 706.39-706.41, from plug extract of Aspergillus niger 1F6
21 22 23	9	grown on CYAS for 7 days. C. TIC of same extract, and D. mass spectrum of fumonisin B_2 in the
24 25	10	extract
26 27 28	11	
29 30	12	Figure 3.
31 32	13	Tandem spectra (40 V collision), of fumonisin B_2 peak from A. niger (A) extract and a reference
33 34 35	14	standard (B).
36 37	15	
38 39	16	Figure 4.
40 41 42	17	LC-MS/MS chromatograms of R1 sample, showing the MRM transitions from FB ₂ (A and B) as
43 44	18	well as FB ₄ (C and D).
45 46	19	
47 48 49	20	
50 51	21	
52 53	22	
54 55 56	23	
57 58		
59 60		

Table 1. Distribution and fumonisin producing abilities of *Aspergillus* spp. in section *Nigri* isolated from Thai coffee beans as determined by LC-MS

Arabica	Robusta	Fumonisin Production*		
(Northern Thailand)	(Southern Thailand)	Positive	Fumonisin B ₂	
A. niger (44)	A. niger (28)	13/17	+++	
A. tubingensis (19)	A. tubingensis (17)	0/13	-	
A. foetidus (28)	-	0/15	-	
A. aculeatinus (9)	A. aculeatinus (15)	0/14	-	
-	A. carbonarius (35)	0/18	-	
-	A. sclerotiicarbonarius (5)	0/5	-	

Note: In brackets: percent of black Aspergilli isolates identified from each type of Thai coffee beans as determined on non-surface disinfected coffee beans.

* Other analogues are with accurate masses matching FB₁ and FB₃, however retentions times do not match reference standards.



Strain	Fumonisin Production	Coffee trans			
number	$(\mu g/cm^2)^*$	Conee type, source			
1B6	0.4	Arabica, Northern Thailand			
1 B 7	0.7	Arabica, Northern Thailand			
1B8	0.9	Arabica, Northern Thailand			
1F1	0.9	Arabica, Northern Thailand			
1F6	0.7	Arabica, Northern Thailand			
1F7	0.5	Arabica, Northern Thailand			
3A2	1.3	Arabica, Northern Thailand			
3E2	0.8	Arabica, Northern Thailand			
3G2	ND**	Robusta, Southern Thailand			
3G7	ND	Robusta, Southern Thailand			
3H2	ND	Robusta, Southern Thailand			
3H3	1.3	Arabica, Northern Thailand			
3H4	1.2	Robusta, Southern Thailand			
3H7	2	Robusta, Southern Thailand			
4C4	0.3	Robusta, Southern Thailand			
4D4	ND	Robusta, Southern Thailand			
6E3	2	Arabica, Northern Thailand			

*Relative standard deviation 30%. **Not detected.



-	
3	
4	
5	
ĉ	
ю	
7	
8	
à	
3	
10	
11	
12	
13	
13	
14	
15	
16	
17	
17	
18	
19	
20	
21	
<u>~</u> 1	
22	
23	
24	
25	
25	
26	
27	
28	
20	
29	
30	
31	
32	
52	
33	
34	
35	
36	
30	
37	
38	
39	
10	
40	
41	
42	
43	
ΔΛ	
44	
45	
46	
47	
47 42	
47 48	
47 48 49	
47 48 49 50	
47 48 49 50 51	
47 48 49 50 51	
47 48 49 50 51 52	
47 48 49 50 51 52 53	
47 48 49 50 51 52 53 54	
47 48 49 50 51 52 53 54 55	
47 48 49 50 51 52 53 54 55 56	
47 48 49 50 51 52 53 54 55 56	
47 48 49 50 51 52 53 54 55 56 57	
47 48 49 50 51 52 53 54 55 56 57 58	

1

Strain				Extro	olites			
1B6	AU-NA	FU			PY	TE		
1B7	AU-NA	FU	KO	OR	PY	TE		—
1B8	AU-NA	FU	KO	OR	PY	TE		
1F1	AU-NA	FU	KO	OR	PY	TE	OT A	OT B
1F6	AU-NA	FU	KO	OR	PY	TE	OT A	OT B
1F7	AU-NA	FU	KO	OR	PY	TE		
3E2	AU-NA	FU	KO	OR	PY	TE		
3G2	AU-NA	FU	KO	OR	PY	TE		
3G7	AU-NA	FU	KO	OR	PY	TE		
3H2	AU-NA	FU	KO	OR	PY	TE	—	—
3H3	AU-NA	FU	KO	OR	PY	TE		—
3H4	AU-NA	FU	KO	OR	PY	TE		
3H7	AU-NA	FU	KO	OR	PY	TE		
4C4	AU-NA	FU	KO	OR	PY	TE		
4D4	AU-NA	FU	KO	OR	PY	TE		_

Table 3. Extrolite production other than Fumonisin B_2 by isolates of *Aspergillus niger* from Thai green coffee beans:

AU-NA = aurasperone B and other naphtho- γ -pyrones, FU = funalenone, KO = kotanin, OR = orlandin, PY = pyranonigrin A, TE = tensidol A, OT = ochratoxin.



203x100mm (300 x 300 DPI)









Tandem spectra (40 V collision), of fumonisin B_2 peak from *A. niger* (A) extract and a reference standard (B).





LC-MS/MS chromatograms of R1 sample, showing the MRM transitions from FB_2 (A and B) as well as FB_4 (C and D).