



HAL
open science

Distribution of aflatoxins and fumonisins in dry-milled maize fractions

Amedeo Pietri, Marco Zanetti, Terenzio Bertuzzi

► **To cite this version:**

Amedeo Pietri, Marco Zanetti, Terenzio Bertuzzi. Distribution of aflatoxins and fumonisins in dry-milled maize fractions. *Food Additives and Contaminants*, 2009, 26 (03), pp.372-380. 10.1080/02652030802441513 . hal-00577336

HAL Id: hal-00577336

<https://hal.science/hal-00577336>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distribution of aflatoxins and fumonisins in dry-milled maize fractions

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2008-158.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	20-Aug-2008
Complete List of Authors:	Pietri, Amedeo; Università Cattolica del Sacro Cuore, Scienze degli alimenti e della nutrizione Zanetti, Marco; Università Cattolica del Sacro Cuore, Scienze degli alimenti e della nutrizione Bertuzzi, Terenzio; Università Cattolica del Sacro Cuore, Scienze degli alimenti e della nutrizione
Methods/Techniques:	Chromatography - HPLC, Chromatography - LC/MS, Clean-up - affinity columns
Additives/Contaminants:	Aflatoxins, Fumonisins
Food Types:	Cereals, Processed foods

SCHOLARONE™
Manuscripts

Distribution of aflatoxins and fumonisins in dry-milled maize fractions

A. Pietri, M. Zanetti and T. Bertuzzi

Istituto Scienze degli Alimenti e della Nutrizione, Facoltà di Agraria, Università

Cattolica del Sacro Cuore, Via Emilia Parmense, 84 I-29100 Piacenza, Italy

E-mail: amedeo.pietri@unicatt.it

Abstract

The aim of this study was to evaluate the distribution of aflatoxins and fumonisins in fractions derived from dry-milling of contaminated maize. Two maize lots with different contamination levels were processed and sampled: the first (maize 1) had aflatoxin B₁ (AFB₁) and fumonisin B₁ (FB₁) levels of 3.6 and 5379 µg kg⁻¹, respectively; the second (maize 2) had corresponding levels of 91.1 and 8841 µg kg⁻¹, respectively. The cleaning step reduced AFB₁ and FB₁ levels by 8 and 11% in maize 1 and by 57 and 34% in maize 2. The subsequent removal of bran and germ led to a further decrease in contamination levels in the products destined for human consumption. In the latter, AFB₁ was uniformly distributed, while FB₁ was concentrated in the finer size fractions. Contamination of raw maize 1 (3.6 µg kg⁻¹) was below the EU AFB₁ limit of 5 µg kg⁻¹ for unprocessed maize, but among the final products only coarse flour (1.7 µg kg⁻¹) was within the EU limit of 2 µg kg⁻¹, while grits and fine flour showed higher levels (2.7 and 2.5 µg kg⁻¹, respectively). As regards

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

cleaned maize, a different distribution of the two toxins was observed in the kernels: AFB₁ contamination was more superficial and concentrated in germ, while FB₁ contamination affected the inner layers of the kernels.

Keywords: aflatoxins, fumonisins, maize, dry-milling process, milling fractions

Introduction

Maize is one of the most important agricultural crops in the world, both for its potential yield per hectare and its nutritional value. In Italy, the area dedicated to maize cultivation is about 1.1 million hectares, located mainly in Northern Italy, while maize production is about 9.7 million tonnes (Istat, 2006). Although grown principally for animal feed, about 4% of Italian maize production is destined for the milling industry and is processed into products for human consumption.

The natural occurrence in maize of mycotoxins such as aflatoxins, fumonisins, trichothecenes and zearalenone, has been reported in various studies (Vargas et al. 2001; Adejumo et al. 2007; Abbas et al. 2002; Ali et al. 1998; Domijan et al. 2005; Gonzalez et al. 1999). As regards maize produced in Italy, mycotoxins that deserve special attention are: aflatoxins for their extreme acute and chronic toxicity and because a toxic metabolite (aflatoxin M₁) is excreted into the milk of cows fed contaminated feeds; fumonisins for their toxicity and high and widespread contamination that cause several disease states in animals and could affect human health (Pietri et al. 2004; Logrieco et al. 2002; Battilani et al. 2005). Aflatoxins B₁, B₂, G₁ and G₂ are mainly produced by two fungal species: *Aspergillus flavus* and *A. parasiticus*, mould species that grow in

1
2
3
4 warm conditions in tropical and sub-tropical regions of the world. Aflatoxin B₁, (AFB₁)
5
6 the most toxic aflatoxin, is a potent genotoxic carcinogen in laboratory animals and
7
8 there is strong evidence that it is a liver carcinogen in humans. Fumonisin B₁, B₂ and
9
10 B₃ are a group of closely related compounds mainly produced by *Fusarium*
11
12 *verticillioides* and *F. proliferatum*, fungi that commonly contaminate maize in many
13
14 growing regions of the world; studies demonstrate the importance of insect damage,
15
16 drought and temperature stress on fumonisin synthesis. Fumonisins have been shown to
17
18 be hepatocarcinogenic in male rats and female mice and nephrocarcinogenic in male
19
20 rats (Voss et al. 2001). Fumonisins have also been associated with the high incidence of
21
22 human oesophageal cancer in some regions of South Africa and China (Sydenham et al.
23
24 1990; Sun et al. 2007). Recent findings suggest that they might increase the risk of
25
26 neural tube defects in populations consuming large amounts of fumonisin-contaminated
27
28 maize (Missmer et al. 2006).
29
30
31
32
33
34
35
36

37 Aflatoxin B₁ and *Fusarium verticillioides* toxins have been categorised by the
38
39 International Agency for Research on Cancer (IARC) as class 1 (human carcinogen) and
40
41 2B (possible human carcinogen), respectively (IARC 1993). In a later evaluation in
42
43 2002, the IARC declared fumonisin B₁ (FB₁) to be a group 2B carcinogen (IARC 2002).
44
45 The EU legislation fixed maximum admissible levels for AFB₁, total aflatoxins (sum of
46
47 AFB₁, AFB₂, AFG₁ and AFG₂) and fumonisins (sum of FB₁ and FB₂) in cereals and
48
49 derived products for direct human consumption (Commission of the European
50
51 Communities 2006a, 2007). Current limits relevant to maize are given in Table 1.
52
53
54
55

56 Current legislation being very detailed, for maize millers it is vital to know the extent of
57
58 decontamination that can be obtained by the first step (cleaning) of the process and how
59
60

1
2
3
4 these mycotoxins are distributed in final products. Physical procedures generate little
5
6 heat other than that caused by the operation of the machinery for cleaning, de-husking
7
8 and abrasion so no significant thermal breakdown of mycotoxins would be expected at
9
10 this stage. However, moulds and mycotoxins are often concentrated in dust and broken
11
12 grains or in the outer seed coat of grains. The removal of this material can thus result in
13
14 a considerable reduction of the mean mycotoxin concentration.
15
16
17
18

19
20
21 Fumonisin distribution was studied on a laboratory-scale processing apparatus (Katta et
22
23 al. 1997) or sampling, over 9 months, maize products in a commercial dry-mill (Broggi
24
25 et al. 2002). Brera et al. (2004, 2006) carried out a study, in a commercial dry-milling
26
27 plant, of FB₁, aflatoxins and zearalenone distribution in milled corn fractions from two
28
29 maize lots showing medium-low contamination levels. These studies indicated that
30
31 contamination increased, with respect to unprocessed maize, in products like bran, germ
32
33 and flour of low particle size (animal feed flour), while it decreased in grits and flour of
34
35 higher particle size, destined for human consumption. No data are available on the
36
37 effect of the cleaning step.
38
39
40
41
42

43
44 The aim of our work was to investigate in an industrial dry-mill: (a) aflatoxin and
45
46 fumonisin distribution in milling products deriving from the processing of two maize
47
48 lots, that showed (with respect to the Italian situation) medium levels of aflatoxins
49
50 (AFB₁, AFB₂, AFG₁, AFG₂) and fumonisins (FB₁ and FB₂) in the first and high levels
51
52 in the second (about 5 and 8000 µg kg⁻¹ for the first lot, 120 and 13000 µg kg⁻¹ for the
53
54 second, respectively); (b) the efficacy of maize cleaning steps, carried out at the
55
56 beginning of the process, on aflatoxin and fumonisin contamination levels. Further, a
57
58
59
60

1
2
3
4 mass balance of the mycotoxins was carried out, in order to verify the accuracy of the
5
6 results.
7
8
9

10
11 [insert Table 1 about here]
12
13

14 15 16 **Materials and methods**

17 18 *Dry-Milling Process*

19
20 The study was carried out in an industrial dry-milling plant able to process 10 tonnes
21
22 per hour of maize grain and performed de-germination under partially wet conditions by
23
24 a conical de-germinator. First, the unprocessed maize was cleaned through three
25
26 different steps (Figure 1), which included a separator with aspirator (winnowing), a dry
27
28 de-stoner and an intensive scourer coupled with an aspirator. The cleaned maize was
29
30 soaked in warm water (70°C) for 10 min, to increase moisture to 22%. In these
31
32 conditions, the germ becomes soft and elastic; also, partial detachment of bran occurs.
33
34 After soaking, the maize was decorticated and de-germinated in the conical de-
35
36 germinator, then sifted through a 5-mm sieve. The fraction (large size grits and germ)
37
38 retained on the sieve was directly conveyed to the plan sifter, while the sieved fraction
39
40 (flours, bran, small size grits and fractions of germ) was dried to 14% moisture content
41
42 and conveyed to a turbo-sifter, where the flour of a particle size <2 mm (fine flour), was
43
44 separated and destined to animal feed. The fraction with a particle size >2 mm was
45
46 conveyed to a winnowing, which aspirated the bran (lighter) sending it to animal feed,
47
48 while fractions of germ and small size grits were sent to the plan sifter, together with
49
50 large size grits and germ of particle size >5 mm. The plan sifter separated the products
51
52 according to their size: those of size >5.5 mm, were conveyed back to the de-
53
54
55
56
57
58
59
60

1
2
3
4 germinator, while the smaller ones were separated through a 4 mm sieve. The fractions
5
6 which were obtained were sent to different gravity tables, to separate germ from small
7
8 and large size grits. Fine grits were processed again using a horizontal roller mill and
9
10 transformed into flour, that was conveyed to a plan sifter and divided between coarse
11
12 flour (300-850 μm), fine flour (<300 μm) and residual germ (about 1%). Undersized
13
14 flour, bran and cleaning waste, were processed with a hammer mill for the production of
15
16 animal feed flour.
17
18
19
20
21

22
23 [insert Figure 1 about here]
24
25
26
27

28 *Samples*

29
30 Two maize lots (maize 1 and 2), with different aflatoxin and fumonisin contamination,
31
32 were processed and sampled. Both were FAO 600 class hybrids, with flinty endosperm.
33
34 Samples of unprocessed maize and derived fractions were drawn from twelve opening
35
36 slits of the plant (numbered in Figure 1). In order to obtain a dynamic representative
37
38 sampling (corresponding to the flow of different products), a primary sample of about
39
40 300 g was collected from each slit every 90 seconds for one hour (40 samples),
41
42 according to the Regulation 2006/401/EC (Commission of the European Communities
43
44 2006b). From the cleaning plant were collected: unprocessed maize (1), waste of the
45
46 winnower (2), of the dust filter (3), of the intensive scourer (4) and cleaned maize (5).
47
48 From the de-germination process were collected: bran (6), germ (7), coarse grits (8),
49
50 fine grits (9) and animal feed flour (12). From the refining step were collected: coarse
51
52 (10) and fine flour (11). For each collecting-slit, the 40 primary samples were mixed,
53
54 ground with a hammer mill (1 mm sieve), and homogenized for 10 min using a
55
56
57
58
59
60

1
2
3
4 horizontal mixer. For each product, two sub-samples of 1.5 kg were withdrawn and
5 stored at -20°C until the time of analysis. The percentage yields of the different
6 fractions were calculated taking into account the technical specifications of the plant
7 and from measurements carried out by plant-technicians during processing of the two
8 maize lots, before sampling.
9
10
11
12
13
14
15
16
17

18 *Analytical Standards*

19
20 Aflatoxin and fumonisin standards were obtained from Sigma-Aldrich (St. Louis, MO,
21 USA). For each aflatoxin, a stock solution of $5\text{--}8\ \mu\text{g ml}^{-1}$ was prepared in
22 benzene:acetonitrile (98:2, v/v, 2 ml) and stored at -20°C . The solution was calibrated
23 spectrophotometrically at 350 nm (AOAC International, 2005). The working standard
24 solution was prepared after evaporation under nitrogen of an aliquot (100 μl) of the
25 stock solution and re-dissolution in chloroform (10 ml) by ultrasonication. An aliquot
26 (100 μl) of this solution was evaporated under nitrogen and re-dissolved in the HPLC
27 mobile phase (0.5-5 ml), to obtain calibrant solutions at individual concentrations of
28 AFB₁, AFB₂, AFG₁, AFG₂ between 0.3 and 6 $\mu\text{g l}^{-1}$. FB₁ and FB₂ (1 mg) were
29 separately dissolved in 10 ml acetonitrile:water (1:1, v/v); the concentration was
30 calculated using the weight indicated by the manufacturer. These solutions were diluted
31 to obtain HPLC calibrant solutions in acetonitrile:water (30:70, v/v, acidified with 0.4%
32 acetic acid) at individual concentrations of FB₁ and FB₂ between 2.5 and 50 $\mu\text{g l}^{-1}$.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 *Analysis for aflatoxins and fumonisins*

55
56 Aflatoxins were analysed according to the method of Stroka et al. (1999). Aflatoxins
57 were extracted from 25 g of sample with 250 ml methanol:water (80:20, v/v), using a
58
59
60

1
2
3
4 rotary-shaking stirrer for 45 min. After filtration through a folded filter-paper, an aliquot
5
6 of the filtrate (5 ml) was diluted with distilled water (45 ml) and the solution was
7
8 purified through an immunoaffinity column (R-Biopharm Rhône LTD, Glasgow,
9
10 Scotland, UK). After washing of the column with 5 ml distilled water, aflatoxins were
11
12 eluted into a graduated glass vial with methanol (2.5 ml). The eluate, concentrated to 1.0
13
14 ml under a gentle stream of nitrogen, was brought to 2 ml with acetonitrile:water
15
16 (25:75, v/v), and vortex-mixed for a few seconds; then, the extract was filtered
17
18 (Millipore Corporation, Bedford, Massachusetts, USA, HV 0.45 μm) and injected (30
19
20 μl). Analysis was performed using an HPLC instrument, consisting of two PU-1580
21
22 chromatographic pumps, an AS 1555 sampling system, a FP 1520 fluorescence detector
23
24 and a post-column derivatization system (Jasco Corporation, Tokyo, Japan); the
25
26 instrument was controlled by Borwin 1.5 software (Jasco). A Superspher RP-18
27
28 column (4 μm particle size, 125x4 mm i.d., Merck) was used at ambient temperature,
29
30 with a mobile phase of water:methanol:acetonitrile (64:23:13, v/v/v), at 1.0 ml min^{-1} . A
31
32 solution of pyridinium bromide perbromide (25 mg in 500 ml of HPLC-grade water)
33
34 was used as a derivatizing agent. The flow of the post-column derivatizing solution was
35
36 set at 0.1 ml min^{-1} and the reaction tubing was 500 μl . The detector was set at $\lambda_{\text{ex}}=365$
37
38 nm and $\lambda_{\text{em}}=440$ nm.
39
40
41
42
43
44
45
46
47
48
49

50 FB_1 and FB_2 content was analysed according to the method proposed by Visconti et al.
51
52 (2001). Fumonisin were extracted from 10 g of sample in a plastic centrifuge bottle
53
54 with 50 ml of acetonitrile:methanol:water (25:25:50, v/v/v). After extraction for 45 min
55
56 using a rotary-shaking stirrer and centrifugation at 4500 g for 6 min, the supernatant
57
58 was poured into a flask; another 50 ml of the same solution was added to the residue in
59
60

1
2
3
4 the centrifuge bottle, and a second extraction performed for 30 min. The combined
5
6
7 extracts were filtered through a folded filter-paper. An aliquot of 2 ml was diluted with
8
9
10 20 ml of 0.1 M phosphate buffered saline (PBS, pH=7.4) and purified through an
11
12 immunoaffinity column (R-Biopharm Rhône Ltd, Glasgow, Scotland, UK); after
13
14 washing the column with PBS (2 ml), the fumonisins were slowly eluted (0.5 ml min^{-1})
15
16 with methanol (6 ml) into a graduated glass vial; subsequently, the eluate was
17
18 concentrated to 2 ml under a gentle stream of nitrogen. Analysis was carried out using a
19
20 LC-MS/MS system, consisting of a LC 1.4 Surveyor pump (Thermo-Fisher Scientific,
21
22 San Jose, CA, USA), a PAL 1.3.1 sampling system (CTC Analytcs AG, Zwingen,
23
24 Switzerland) and a Quantum Discovery Max triple-quadrupole mass spectrometer; the
25
26 system was controlled by Excalibur 1.4 software (Thermo-Fisher). After dilution of the
27
28 extract (0.1 ml brought to 1 ml) with acetonitrile:water (30:70 v/v, acidified with 0.4%
29
30 acetic acid), the fumonisins were separated on a Betasil RP-18 column (5 μm particle
31
32 size, 150 x 2.1 mm, Thermo-Fisher) with mobile-phase gradient acetonitrile-water
33
34 (both acidified with 0.4% acetic acid) from 25:75 to 55:45 in 9 min, then isocratic for 3
35
36 min; the flow rate was 0.2 ml min^{-1} . Ionisation was carried out with an ESI interface
37
38 (Thermo-Fisher) in positive mode as follows: spray capillary voltage 4.0 kV, sheath and
39
40 auxiliary gas 35 and 14 psi, respectively, temperature of the heated capillary 270 °C.
41
42 The mass spectrometric analysis was operated in selected reaction monitoring (SRM).
43
44 For fragmentation of $[\text{M}+\text{H}]^+$ ions (722 m/z for FB_1 , 706 m/z for FB_2), the argon
45
46 collision pressure was set to 1.5 mTorr and the collision energy to 36 V. The selected
47
48 fragment ions were: 704, 352 and 334 m/z for FB_1 , 688, 336 and 318 m/z for FB_2 .
49
50 Quantitative determination was performed using LC-Quan 2.0 software (Thermo-
51
52
53
54
55
56
57
58
59
60 Fisher).

Both for aflatoxins and fumonisins, four replicates were analysed for each sample.

Mass balance

Both for aflatoxins and fumonisins, mass balances of the cleaning step $[\sum \text{amount}_{(\text{cleaning waste+cleaned maize})} / \text{amount}_{(\text{unprocessed maize})}] * 100$, of the de-germination process $[\sum \text{amount}_{(\text{animal feed flour-cleaning waste+grits+germ})} / \text{amount}_{(\text{cleaned maize})}] * 100$ and of the total milling process $[\sum \text{amount}_{(\text{animal feed flour+coarse grits+flours+germ})} / \text{amount}_{(\text{unprocessed maize})}] * 100$, were calculated.

Results and discussion

Yields of the milling-corn fractions

Yields of the different fractions, calculated with an approximation of 1%, are reported in Table 2. The two maize lots had different characteristics, maize 2 being visually dustier and with a higher amount of broken kernels with respect to maize 1; as a consequence, the cleaning waste was 6 vs. 3% of the processed product for maize 2 and 1, respectively. Further, maize 2 yielded a lower amount of undersized flour deriving from the turbo sifter, with respect to maize 1 (17 vs. 20%). As a consequence, the yield of animal feed flour was the same for maize 1 and 2 (28%). For both maize lots, the ratio between coarse and fine flour, after refining of fine grits, was 85:15.

[insert Table 2 about here]

Performance of the methods

Recovery percentages were evaluated by spiking known blank samples at levels of 2

1
2
3
4 and 1000 $\mu\text{g kg}^{-1}$ for aflatoxins and fumonisins, respectively. Average recovery values
5
6 were: 97.8 \pm 1.6% for AFB₁ and over 93% for the other aflatoxins, 95.5 \pm 1.9% for FB₁
7
8 and 93.6 \pm 2.1% for FB₂. A certified reference material (ground corn, R-Biopharm
9
10 Rhône LTD), with an AFB₁ contamination of 4.1 \pm 1.0 $\mu\text{g kg}^{-1}$, was also analysed; the
11
12 result was 4.2 \pm 0.2 $\mu\text{g kg}^{-1}$. The results of the analyses were not corrected for recovery.
13
14 The limit of detection (LOD, signal-to-noise ratio of 3:1) and of quantification (LOQ,
15
16 signal-to-noise ratio of 10:1) were respectively 0.02 and 0.05 $\mu\text{g kg}^{-1}$ for aflatoxins, 10
17
18 and 30 $\mu\text{g kg}^{-1}$ for fumonisins.
19
20
21
22
23
24
25

26 *Aflatoxin distribution*

27
28 AFB₁ and AFG₁ were detected in both maize lots, but AFB₂ and AFG₂ only in maize 2
29
30 (Table 3). The AFB₁ level of maize 1 (3.6 $\mu\text{g kg}^{-1}$) was lower than the value fixed by
31
32 EC Regulation 1881/2006 for unprocessed maize (5 $\mu\text{g kg}^{-1}$); however, considering the
33
34 final products, only coarse flour was within the limit for human consumption (2 μg
35
36 kg^{-1}), while coarse and fine grits slightly exceeded this limit. After the cleaning steps,
37
38 maize 1 showed an AFB₁ concentration slightly lower than the unprocessed one (3.3
39
40 against 3.6 $\mu\text{g kg}^{-1}$, -8%). Bran and germ deriving from de-cortication and de-
41
42 germination steps, resulted as more contaminated than cleaned maize (+127 and +197%,
43
44 respectively). Animal feed flour, that included cleaning waste, bran, and undersized
45
46 flour ground together, showed an AFB₁ level close to raw maize. The results obtained
47
48 for maize 1 basically accord with those reported by Brera et al. (2006): processing two
49
50 maize lots with an AFB₁ concentration below 3 $\mu\text{g kg}^{-1}$, they observed an increase of
51
52 contamination in bran, germ and animal feed flour and a consequent decrease in grits
53
54 and flour destined for human consumption, that showed values below 2 $\mu\text{g kg}^{-1}$.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Maize 2 was much more contaminated ($91.1 \mu\text{g kg}^{-1}$); after cleaning it showed an AFB_1 level decidedly lower ($39.5 \mu\text{g kg}^{-1}$) with respect to the unprocessed one (-57%), as cleaning steps removed the most contaminated fractions (broken kernels, dust, etc.). Bran and germ of maize 2 showed much higher AFB_1 concentrations (+379 and +629%, respectively) compared to cleaned maize. As a consequence, unlike maize 1, the contamination level decreased remarkably in grits (-80%) and increased (+141% with respect to raw maize) in animal feed flour; however, the latter could not be used, as the value (0.219 mg kg^{-1}) exceeded the limit of 0.02 mg kg^{-1} in force in the EU (Commission of the European Communities 2002). None of the milling-corn fractions destined for human consumption showed a concentration below the legal limit ($2 \mu\text{g kg}^{-1}$), but they could be used as feeds. Considering the AFB_1 percent distribution (Table 3), it is evident that cleaning steps led to very different results for the two maize lots, removing 11 vs. 59% of the toxin from maize 1 and 2, respectively. Analogously, very different results were observed in de-cortication and de-germination steps; toxin removal was 30 vs. 72% for cleaned maize 1 and 2, respectively. Consequently, the AFB_1 percentage remaining in coarse and fine grits, was 55 vs. 13%, for cleaned maize 1 and 2, respectively. Fine grits were subsequently processed to obtain coarse and fine flour, in the ratio 85:15; for maize 1, 61% of the AFB_1 amount in fine grits (7.2% of the total) was present in coarse flour and 16.1% in fine flour; for maize 2, the corresponding percentages were 64.3 and 9.0%. The lower figure was due to a small amount of remaining germ fraction (about 1% of unprocessed maize), more contaminated, that was removed in this step of the process. The different AFB_1 percentage (26.4% against 67.5% for maize 1 and 2, respectively) collected in animal feed flour, was mainly due to the different contribution made by the waste of the

1
2
3
4 cleaning steps.

5
6
7 A similar trend was observed for the other aflatoxins.

8
9
10
11 [insert Table 3 about here]
12
13

14 15 16 *Fumonisin distribution*

17
18 Both maize lots turned out to be contaminated with FB₁ and FB₂ (Table 4) at levels
19 exceeding the limit (4000 µg kg⁻¹) in force in the EU. Maize grains obtained after
20 cleaning steps showed fumonisin concentrations lower (especially maize 2) than the
21 unprocessed ones (-11 and -34% for FB₁). Bran showed higher contamination levels
22 with respect to cleaned maize (+50 and +167% for FB₁); germ turned out to be less
23 contaminated with respect to bran in both maize lots. Coarse and fine grits, fractions
24 intended for human consumption, showed much lower FB₁ concentrations when
25 compared to cleaned maize (-90 and -88% for maize 1 and -90 and -73% for maize 2).
26 Besides, fine grits and fine flours, showed a level higher than the coarse ones.
27 Therefore, as already found by Brera et al. (2004), contamination increases in lower
28 particle size final products. Coarse grits for both maize lots, fine grits and coarse flour
29 for maize 1, showed a contamination level (FB₁ + FB₂) below 1400 µg kg⁻¹, which is
30 the limit fixed by the EU. Animal feed flour, which collected bran and cleaning waste,
31 was three times more contaminated than unprocessed maize. Contamination levels
32 found in different corn-milling fractions were similar to those found in previous studies
33 (Katta et al., Broggi et al., Brera et al.), in which there was always reported an increase
34 of concentration in bran, germ and animal feed flour and a decrease in products
35 intended for human consumption, like grits and flours. As regards the FB₁ percentage
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 distribution, the cleaning plant removed 14% and 38% of the toxin for maize 1 and 2,
5
6 respectively; regarding cleaned maize, de-cortication and de-germination processes
7
8 removed altogether 12% and 27% of the toxin, while 6.8% and 8.8% remained in coarse
9
10 and fine grits. It is evident that the highest fraction of FB₁ remained in the undersized
11
12 flour separated from the turbo sifter and conveyed into animal feed flour; this product
13
14 gathers 83.3 and 90.9% of the total FB₁ present in uncleaned maize 1 and 2,
15
16 respectively. A similar trend has been observed for FB₂.
17
18
19
20
21
22

23 [insert Table 4 about here]
24
25
26
27

28 *Mass balance*

29
30 Mass balances of aflatoxins and fumonisins for the cleaning step, for the degermination
31
32 process and for the complete process are reported in Table 5. The mass balance of the
33
34 complete process for AFG₁ in maize 1 was relatively low (79.3%), probably because of
35
36 a high analytical error associated with the low level of the mycotoxin; except for this
37
38 result, mass balances of mycotoxins in both processes were in the range 90-110%.
39
40 Thus, it can be concluded that the dynamic sampling was effective and the results were
41
42 altogether accurate.
43
44
45
46
47
48

49 [insert Table 5 about here]
50
51
52
53

54 **Conclusions**

55
56 The extent of decontamination obtained from the cleaning step was decidedly higher for
57
58 a “dirty” maize (maize 2) with respect to a “normal” maize (maize 1). In this step, the
59
60

1
2
3
4 percent toxin reduction in maize 2, if calculated in terms of both concentration and
5 mass, was 57-59% and 34-38% for AFB₁ and FB₁, respectively. For maize 1, the
6
7 corresponding values were 8-11% and 11-14% for AFB₁ and FB₁, respectively.
8
9

10
11 The extent of decontamination obtained from the cleaning step of maize 2, was higher
12 for aflatoxins than for fumonisins. Probably, during harvesting, handling and
13 transportation, a fraction of the aflatoxin contaminated kernels broke and produced
14 powder; therefore their removal contributed significantly to decontamination. For maize
15
16 2, discarding of the cleaning waste would decrease the yield of animal feed flour from
17
18 28% to 22%, but the total amount of AFB₁ and FB₁ residual in the product would
19
20 decrease from 67.5% to 8.2% and from 90.9% to 53.2%, respectively. Generally, it is
21
22 evident that an alternative use of cleaning waste would substantially improve the quality
23
24 of animal feed flour.
25
26
27
28
29
30
31
32
33
34

35 As regards products after the cleaning step, a different distribution of the two toxins was
36 observed. There was a large amount of AFB₁ in the germ fraction, of 18.5% and 46.7%
37
38 of the total for maize 1 and 2, respectively; for bran the corresponding values were
39
40 11.8% and 25.6%; most of the remaining AFB₁ was in coarse grits. As regards FB₁,
41
42 4.3% and 12.7% occurred in germ, 7.7% and 14.1% in bran, for maize 1 and 2,
43
44 respectively; the amount remaining in coarse grits was low (5.5-5.6%), while most was
45
46 collected in undersized flour (81-64%), that was intended for animal feed. It is evident
47
48 that aflatoxin contamination of maize kernels was more superficial and concerned
49
50 mainly the germ, while fumonisin contamination affected the inner layers of kernels.
51
52
53
54
55
56
57
58
59
60

References

- Abbas, H., Williams, W., Windham, G., Pringle, H., Xie, W., Shier, W. 2002. Aflatoxin and fumonisin contamination of commercial corn (*Zea mays*) hybrids in Mississippi. *Journal of Agricultural and Food Chemistry* 50(18): 5246-5254.
- Adejumo, T., Hettwer, U., Karlovsky, P. 2007. Survey of maize from south-western Nigeria for zearalenone, α and β zearalenols, fumonisin B₁ and enniatins produced by *Fusarium* species. *Food Additives and Contaminants* 24(9): 993-1000.
- Ali, N., Sardjono, Yamashita, A., Yoshizawa, T. 1998. Natural co-occurrence of aflatoxins and *Fusarium* mycotoxins (fumonisins, deoxynivalenol, nivalenol and zearalenone) in corn from Indonesia. *Food Additives and Contaminants* 15(4): 377-384.
- AOAC International. 2005. Natural Toxins. Aflatoxins. Official Methods of Analysis of the International Association of Official Analytical Chemists. Gaithersburg, MD: AOAC. p. 4.
- Battilani, P., Scandolaro, A., Barbano, C., Pietri, A., Bertuzzi, T., Marocco, A., Berardo, N., Vannozzi, G.P., Baldini, M., Miele, S., Salera, E., Maggiore, T. 2005. Monitoraggio della contaminazione da micotossine in mais. *L'Informatore Agrario* 6:47-49.
- Brera, C., Debegnach, F., Grossi, S., Miraglia, M. 2004. Effect of industrial processing on the distribution of fumonisin B₁ in dry-milling corn fractions. *Journal of Food Protection* 67 (6): 1261-1266.
- Brera, C., Catano, C., De Santis, B., Debegnach, F., De Giacomo, M., Pannunzi, E., Miraglia, M. 2006. Effect of industrial processing on the distribution of aflatoxins

1
2
3
4 and zearalenone in corn-milling fractions. Journal of Agricultural and Food
5
6 Chemistry 54: 5014-5019.
7

8
9 Broggi, L.E., Resnik, S.L., Pacin, A.M., Gonzalez, H., Cano, G., Taglieri, G. 2002.
10
11 Distribution of fumonisins in dry-milled corn fractions in Argentina. Food
12
13 Additives and Contaminants 19 (5):465-469.
14

15
16 Commission of the European Communities. 2002. Commission Directive 2002/32/EC
17
18 of 7 May 2002 setting maximum levels for undesirable substances in animal feed.
19
20 Official Journal of the European Communities L 140: 10-21.
21

22
23 Commission of the European Communities. 2006a. Commission regulation (EC) No.
24
25 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants
26
27 in foodstuffs. Official Journal of the European Union L 364: 5-24.
28

29
30 Commission of the European Communities. 2006b. Commission regulation (EC) No.
31
32 401/2006 of 23 February 2006 laying down the methods of sampling and analysis
33
34 for the official control of the levels of mycotoxins in foodstuffs. Official Journal
35
36 of the European Union L 70: 12-29.
37

38
39 Commission of the European Communities 2007. Commission regulation (EC) No.
40
41 1126/2007 of 28 September 2007 amending Regulation (EC) No 1881/2006
42
43 setting maximum levels for certain contaminants in foodstuffs as regards
44
45 *Fusarium* toxins in maize and maize products. Official Journal of the European
46
47 Union L 255: 14-17.
48

49
50
51 Domijan, A., Peraica, M., Jurjevic, Z., Ivic, D., Cvjetkovic, B. 2005. Fumonisin B1,
52
53 fumonisin B2, zearalenone and ochratoxin A contamination of maize in Croatia.
54
55 Food Additives and Contaminants 22(7): 677-680.
56
57
58
59
60

1
2
3
4 Gonzalez, H., Martinez, E., Pacin, A., Resnik, S., Sydenham, E. 1999. Natural co-
5
6 occurrence of fumonisins, deoxynivalenol, zearalenone and aflatoxins in field trial
7
8 corn in Argentina. Food Additives and Contaminants 16(12): 565-569.
9

10
11 International Agency for Research on Cancer (IARC). 1993. Some naturally occurring
12
13 substances, food items and constituents, heterocyclic aromatic amines and
14
15 mycotoxins. IARC Monographs on the Evaluation of Carcinogenic Risk to
16
17 Humans No. 56. Lyon: IARC: aflatoxins p.245; Fusarium verticillioides toxins
18
19 p.445.
20
21

22
23 International Agency for Research on Cancer (IARC). 2002. Summaries & Evaluations
24
25 No. 82. Lyon: IARC: fumonisin B₁ p.301.
26
27

28 Istat 2006. Available: www.istat.it/agricoltura/datiagri/coltivazioni/anno2006/corg2006.
29
30 Accessed 02 October 2007.
31
32

33 Katta, S.K., Cagampang, A.E., Jackson, L.S., Bullerman, L.B. 1997. Distribution of
34
35 *Fusarium* molds and fumonisins in dry-milled corn fractions. Cereal Chemistry 74
36
37 (6): 858-863.
38
39

40 Logrieco, A., Mulè, G., Moretti, A., Bottalico, A. 2002. Toxigenic *Fusarium* species
41
42 and mycotoxins associated with maize ear rot in Europe. Mycotoxins in Plant
43
44 Disease. European Journal of Plant Pathology, edited by A. Logrieco, J.A. Bailey,
45
46 L. Corazza & B.M. Cooke (Dordrecht:Kluwer), 597-609.
47
48

49 Missmer, S.A., Suarez, L., Felkner, M., Wang, E., Merrill, A.H., Rothman, K.J.,
50
51 Hendricks, K.A. 2006. Exposure to fumonisins and the occurrence of neural tube
52
53 defects along the Texas-Mexico border. Environmental Health Perspectives
54
55 114(2): 237-241.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Pietri, A., Bertuzzi, T., Pallaroni, L., Piva, G. 2004. Occurrence of mycotoxins and ergosterol in maize harvested over 5 years in Northern Italy. *Food Additives and Contaminants* 21 (5):479-487.
- Stroka, J., Petz, M., Joerissen, U., Anklam, E. 1999. Investigation of various extractants for the analysis of aflatoxin B₁ in different food and feed matrices. *Food Additives and Contaminants* 16 (8): 331-338.
- Sun, G., Wang, S., Hu, X., Su, J., Huang, T., Yu, J., Tang, L., Gao, W., Wang, J. 2007. Fumonisin B₁ contamination of home-grown corn in high risk areas for esophageal and liver cancer in China. *Food Additives and Contaminants* 24 (2): 181-185.
- Sydenham, E.W., Thiel, P.G., Marasas, W.F.O., Shephard, G.S., Van Schalkwyk, D.J., Koch, K.R. 1990. Natural occurrence of some *Fusarium* mycotoxin in corn from low and high esophageal cancer prevalence areas of the Transkei, Southern Africa. *Journal Agriculture Food Chemistry* 38: 1900-1903.
- Vargas, E., Preis, R., Castro, L., Silva, C. 2001. Co-occurrence of aflatoxins B₁, B₂, G₁, G₂, zearalenone and fumonisin B₁ in Brazilian corn. 2001. *Food Additives and Contaminants* 18(11): 981-986.
- Visconti, A., Solfrizzo, M., Girolamo, A. de. 2001. Determination of fumonisins B₁ and B₂ in corn and corn flakes by liquid chromatography with immunoaffinity column cleanup: collaborative study. *Journal of AOAC International* 84 (6): 1828-1837.
- Voss, K., Riley, R., Norred, W.P., Bacon, C.W., Meredith, F., Howard, P., Plattner, R., Collins, T., Hansen, D., Porter, J. 2001. An overview of rodent toxicities: liver and kidney effects of fumonisins and *Fusarium moniliforme*. *Environmental Health Perspectives* 109, suppl.2: 259-266.

Table 1. EU maximum admissible limits for AFB₁, total aflatoxins (sum of AFB₁, AFB₂, AFG₁, AFG₂) and fumonisins (sum of FB₁ and FB₂) in maize and derived products.

	AFB ₁ ($\mu\text{g kg}^{-1}$)	Sum of AFB ₁ , AFB ₂ , AFG ₁ and AFG ₂ ($\mu\text{g kg}^{-1}$)	Sum of FB ₁ and FB ₂ ($\mu\text{g kg}^{-1}$)
Unprocessed maize	5.0	10.0	4000
Maize for direct human consumption	2.0	4.0	1000
Maize-based breakfast	-	-	800
Milling fractions of maize with particle size > 500 micron	-	-	1400
Milling fractions of maize with particle size # 500 micron	-	-	2000

Table 2. Yield of maize milling fractions.

Maize milling fractions	Yield maize 1 (%)	Yield maize 2 (%)
Unprocessed maize	100	100
Waste of aspirator	2	3
Waste of dust filter	0,2	1,5
Waste of intensive scourer	0,8	1,5
Cleaned maize	97	94
Bran	5	5
Germ	6	6
Coarse grits	55	55
Fine grits	11	11
from which:		
coarse flour	85	85
fine flour	15	15
Undersized flour	20	17
Animal feed flour (waste+bran+undersized flour)	28	28

Table 3: Aflatoxin levels (mean of 4 replicates, $\mu\text{g kg}^{-1}$), relative standard deviation (RSD, %) and AFB₁ relative distribution (%) in maize-milling fractions referred to unprocessed maize and (between parentheses) to cleaned maize.

Maize-milling fractions	Maize 1						Maize 2								
	AFB ₁			AFG ₁			AFB ₁			AFB ₂		AFG ₁		AFG ₂	
	$\mu\text{g kg}^{-1}$	RSD %	Distribution %	$\mu\text{g kg}^{-1}$	RSD %	$\mu\text{g kg}^{-1}$	RSD %	Distribution %	$\mu\text{g kg}^{-1}$	RSD %	$\mu\text{g kg}^{-1}$	RSD %	$\mu\text{g kg}^{-1}$	RSD %	
Unprocessed maize	3.6	11.1	100	1.7	5.9	91.1	3.8	100	3.4	8.8	26.3	1.5	1.1	9.1	
Waste of aspirator	3.4	11.8	1.9	1.8	11.1	138.3	3.9	4.6	38.1	3.4	26.3	9.5	1.6	12.5	
Waste of dust filter	17	9.4	1.0	8.3	8.4	1296	2.0	21.3	57.8	2.4	290.3	2.3	9.1	5.5	
Waste of intensive scourer	11.5	12.2	2.6	6.1	8.2	1554	0.8	25.6	54.5	2.9	285.5	2.5	9.4	5.3	
Cleaned maize	3.3	15.1	88.7 (100)	1.1	9.1	39.5	6.1	40.7 (100)	1.8	11.1	13.9	4.3	0.7	14.3	
Bran	7.5	9.3	10.5 (11.8)	3.9	7.7	189.4	4.5	10.4 (25.6)	7.3	10.9	42.7	3.5	1.1	9.1	
Germ	9.8	9.2	16.4 (18.5)	5.2	9.6	287.9	1.0	19.0 (46.7)	16.7	3.0	81.9	1.5	5.0	6.0	
Coarse grits	2.7	11.1	41.7 (47.0)	1.1	9.1	7.6	3.9	4.6 (11.3)	0.4	25.0	1.9	5.3	0.3	16.7	
Fine grits	2.3	8.7	7.2 (8.1)	1.0	10	7.8	5.1	0.9 (2.2)	0.7	14.3	2.4	8.3	0.6	16.7	
from which:															
coarse flour	1.7	11.8	61.0	0.8	12.5	5.9	5.1	64.3	0.6	16.7	2.0	10.0	0.5	10.0	
fine flour	2.5	12.0	16.1	1.2	8.3	4.7	6.4	9.0	0.5	20.0	1.4	14.3	0.4	12.5	
Animal feed flour	3.4	11.8	26.4	1.1	9.1	219.6	2.5	67.5	8.8	3.4	48.5	7.4	1.7	11.8	

Table 4: Fumonisin levels (mean of 4 replicates, $\mu\text{g kg}^{-1}$), relative standard deviation (RSD, %) and FB₁ relative distribution (%) in corn-milling fractions referred to unprocessed maize and (between parentheses) to cleaned maize.

Maize-milling fractions	Maize 1					Maize 2				
	FB ₁			FB ₂		FB ₁			FB ₂	
	$\mu\text{g kg}^{-1}$	RSD %	Distribution %	$\mu\text{g kg}^{-1}$	RSD %	$\mu\text{g kg}^{-1}$	RSD %	Distribution %	$\mu\text{g kg}^{-1}$	RSD %
Unprocessed maize	5379	10.5	100	2576	7.7	8841	7.9	100	4234	8.4
Waste of aspirator	17353	4.6	6.4	6539	5.6	48425	5.5	16.5	11465	6.5
Waste of dust filter	42222	2.9	1.6	31389	2.1	50872	3.7	8.7	34103	4.3
Waste of intensive scourer	43402	2.8	6.4	24653	1.8	48565	3.0	8.3	24746	4.1
Cleaned maize	4770	8.5	86.0 (100)	2360	6.7	5862	3.9	62.3 (100)	2975	5.3
Bran	7154	5.3	6.6 (7.7)	2559	6.5	15647	1.0	8.8 (14.1)	5597	2.2
Germ	3332	7.2	3.7 (4.3)	1237	6.6	11674	8.1	7.9 (12.7)	4334	8.2
Coarse grits	458	11.1	4.7 (5.5)	128	11.7	563	8.0	3.5 (5.6)	157	13.4
Fine grits	556	8.1	1.1 (1.3)	163	11.0	1592	2.8	2.0 (3.2)	500	8.4
from which:										
coarse flour	292	10.6	44.6	96	9.4	1266	7.5	67.6	416	7.9
fine flour	1922	7.7	51.8	635	5.7	3058	5.8	28.8	2975	5.6
Animal feed flour	16011	2.1	83.3	8155	3.7	28712	5.9	90.9	14624	2.4

Table 5: Aflatoxin and fumonisin mass balance (%) for maize 1 and 2.

Mass balance	AFB ₁	AFB ₂	AFG ₁	AFG ₂	FB ₁	FB ₂
Cleaning step						
<i>Maize 1</i>	94.2	-	93.8	-	100.4	104.0
<i>Maize 2</i>	96.3	109.1	96.2	96.0	95.6	102.2
De-germination process						
<i>Maize 1</i>	97.2	-	110.0	-	91.2	89.8
<i>Maize 2</i>	99.5	108.0	99.5	99.5	114.6	106.1
Complete milling process						
<i>Maize 1</i>	90.0	-	79.3	-	92.9	95.0
<i>Maize 2</i>	91.7	110.0	92.1	92.1	104.3	106.2

Figure 1. Layout of the industrial plant showing sampling points (numbered) and final products (bold type).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1.

Figure 1. Layout of the industrial plant showing sampling points (numbered) and final products (bold type).

254x190mm (72 x 72 DPI)