



**HAL**  
open science

## Apyap1 affects aflatoxin biosynthesis during *Aspergillus parasiticus* growth in maize seeds

Massimo Reverberi

► **To cite this version:**

Massimo Reverberi. Apyap1 affects aflatoxin biosynthesis during *Aspergillus parasiticus* growth in maize seeds. *Food Additives and Contaminants*, 2007, 24 (10), pp.1070-1075. 10.1080/02652030701553244 . hal-00577410

**HAL Id: hal-00577410**

**<https://hal.science/hal-00577410v1>**

Submitted on 17 Mar 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Apyap1 affects aflatoxin biosynthesis during *Aspergillus parasiticus* growth in maize seeds**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2007-154.R1
Manuscript Type:	Special Issue
Date Submitted by the Author:	04-Jul-2007
Complete List of Authors:	reverberi, massimo; Università degli Studi Å La Sapienza
Methods/Techniques:	Mycology
Additives/Contaminants:	Mycotoxins
Food Types:	

SCHOLARONE™  
Manuscripts

## **Apyap1 affects aflatoxin biosynthesis during *Aspergillus parasiticus* growth in maize seeds**

M. REVERBERI\*, S. ZJALIC\*, F. PUNELLI\*, A. RICELLI<sup>o</sup>, A.A. FABBRI\* and C. FANELLI\*

\*Università degli Studi “La Sapienza”, Largo Cristina di Svezia 24 00165 Roma, Italy

<sup>o</sup>ISPA-CNR, via G. Amendola 122/O 70126 Bari, Italy

### **Abstract**

In fungal cells grown in synthetic media we have demonstrated that the *Apyap1* gene is implicated in the modulation of aflatoxin biosynthesis following the perturbation of the redox balance. In this study, we suggest that an association between oxidative stress and aflatoxin biosynthesis occurs also in maize seeds. We used  $\Delta Apyap1$ , a strain in which the gene *Apyap1* was disrupted, to verify whether this oxidative stress related transcription factor, by affecting cell redox balance, can have a role in the modulation of aflatoxin synthesis. The amount of hydroperoxides (ROOH) produced by wild type (WT) and  $\Delta Apyap1$ , both grown in potato dextrose broth, was assayed in the filtrate. In maize seeds (30 g), which were inoculated with WT and  $\Delta Apyap1$  conidia, and incubated at 30°C for 15 days, we analysed lipoxygenase activity (LOX), lipoperoxides (LOOH) production, fungal growth and aflatoxin biosynthesis. It was observed that  $\Delta Apyap1$  releases more hydroperoxides in the culture media and more aflatoxins in seeds, likely through a stronger stimulation of LOX, which, in turn led to a greater LOOH production in the seeds. On the basis of our results, we formulated hypothesis regarding strategies to control aflatoxin synthesis.

**Keywords:** *Apyap1*, oxidative stress, *Aspergillus parasiticus*, maize seeds, aflatoxin biosynthesis

## Introduction

A more detailed knowledge of the organization, regulation, and expression of genes linked to aflatoxin biosynthesis is important to detect early these mycotoxins and the mycotoxigenic fungi. Additionally, it allows finding rapid prevention strategies to control their presence in different food sources. However, the molecular approach must advantage of information concerning those events that influence the fungal physiology in relation to aflatoxin production.

In the eighties we have demonstrated how lipoperoxidation can intervene in modulating the aflatoxin biosynthesis. This evidence was brought up both *in vitro* studies and in different seeds (Fabbri *et al.*,1983; Fanelli *et al.*,1989). In those systems, various antioxidants and free radicals scavengers affected the aflatoxin production, in different ways (Fanelli *et al.*,1985). Jayashreee and Subramanyam, (2000) further brought up oxidative stress, in particular the increase of lipid peroxidation and free radical generation, as a prerequisite for aflatoxin biosynthesis by *Aspergillus parasiticus*. Other researches have suggested the relationship between aflatoxin production and sporulation process (Guzman De Pena and Ruiz-Herrera, 1997). The formation of conidia is an important factor for the infection onset and for the fungal colony formation on different synthetic and natural media. Calvo *et al.*, (1999, 2001) pointed out the sporogenic effect of polyunsaturated fatty acid and their lipoperoxides (LOOH), during the development of *Aspergillus* spp. Hydroxylated derivatives of oleic, linoleic and linolenic acids, whose formation is directed by putative dioxygenase, modulate sporulation (Tsitsigiannis *et al.*, 2004). Such factors seemed to regulate the ratio of asexual to sexual development in *A. nidulans* (Champe and El Zayat, 1989). 9-S and 13-S hydroperoxy-linoleic acids (9-S and 13-S HPODE), which are by-products of plant lipoxygenase, affect spore production in *A. nidulans*, *A. parasiticus* and *A. flavus* (Calvo *et al.*, 1999). In particular 9-S-HPODE enhanced spore production and aflatoxin biosynthesis. On the contrary, 13-S-HPODE inhibited both sexual spores and aflatoxin biosynthesis. Furthermore, these

1  
2  
3 compounds were able to play also a role in the formation of cleistothecia (*A. nidulans*) and sclerotia  
4  
5 (*A. flavus*) (Burow *et al.*, 1997). The biosynthesis of aflatoxins, like other compounds in the cell, is  
6  
7 controlled by a signal transduction cascade that involves a G protein (FadA), a protein kinase  
8  
9 (PKA) and other factors regulating the expression of AflR, the aflatoxin cluster regulator.

10  
11 Compounds derived from ligninolytic fungi (*Lentinula edodes* and *Trametes versicolor*) are useful  
12  
13 to inhibit aflatoxin production as reported by Fanelli *et al.*, (2000) and Zjalic *et al.*, (2006). These  
14  
15 findings land basis for innovating biological approaches to control aflatoxin biosynthesis and fungal  
16  
17 growth. Antimicrobials and antioxidants compounds such as thioproline, mannitol,  $\alpha$ - and  $\beta$ -  
18  
19 glucans, caffeic acid and derivatives, have been detected in these basidiomycetes and plants and  
20  
21 might be of significance in modulating oxidative stress and aflatoxin formation (Kurashima *et al.*,  
22  
23 1990; Kim *et al.*, 2004). In *A. parasiticus* we have observed (Reverberi *et al.*, 2006) that the culture  
24  
25 filtrates derived from some basidiomycetes affect transcription factors (such as Apyap1) regulating  
26  
27 the expression of genes involved in the removal of reactive oxygen species (ROS), such as  
28  
29 superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX) enzymatic  
30  
31 activities (Harschmann *et al.*, 1988). In budding yeasts, Yap1, a transcription factor homologue of  
32  
33 AP-1, is involved in the defence response against oxidative stress (Estruch, 2000). Some other  
34  
35 regulatory proteins such as Skn7 and Hsf 1-2 are also involved (He *et al.*, 2003; Moye-Rowley,  
36  
37 2003). In a culture media conducive for aflatoxin biosynthesis, an increase of lipoperoxides appears  
38  
39 in the trophophase followed by an enhancement of Apyap1 mRNA expression. Coincidentally, a  
40  
41 rapid increase of SOD, CAT and GPX enzymatic activities accompanied by a decrease of aflatoxin  
42  
43 biosynthesis, occurred. In the following phase, there is a decline of antioxidant enzymatic activities,  
44  
45 while biosynthesis of lipoperoxides and the aflatoxin occurs (Reverberi *et al.*, 2006). From these  
46  
47 results it emerges that the transcription factor Apyap1 regulates the natural antioxidant defence  
48  
49 mechanisms of the aflatoxigenic fungi modulating the toxin biosynthesis.

50  
51 In this paper the behaviour of two *Aspergillus* strains inoculated in maize seeds has been showed.  
52  
53 The first strain has a functioning *Apyap1* gene, whereas the second one presents a gene inactivated  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 by homologous recombination. The results obtained suggested that the lipoperoxides produced by  
4  
5 maize seeds can affect aflatoxin biosynthesis in both *Aspergillus* strain likely by Apyap1 pathway.  
6  
7  
8  
9

## 10 **Materials and methods**

### 11 *Plant and fungal material and culture conditions*

12  
13 The fungal strains used were *Aspergillus parasiticus* (NRRL 2999) wild type aflatoxins (B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>,  
14  
15 G<sub>2</sub>) producer and a  $\Delta$ Apyap1 mutant obtained from the same strain. The isolate was kept on PDA  
16  
17 (Potato Dextrose Agar, Difco) for 7 days at incubation temperature of 30°C before the use. The  
18  
19 inoculum of *A. parasiticus* was performed with conidia suspension ( $5 \times 10^6$ /ml in sterile distilled  
20  
21 water) in PDB medium. The same number of conidia of the two strains was inoculated in 30 g of  
22  
23 maize seeds, cv Cecilia, moistened at  $a_w$  0.95. To avoid interferences due to the growth of other  
24  
25 fungi, seeds were previously sterilized with  $\gamma$ -rays (8 Kgy, irradiation ratio 2602 Gy·h<sup>-1</sup>. This  
26  
27 procedure allowed obtaining micro flora sterilization and maintaining seed germination (about 97  
28  
29 %; data not shown). The inoculated seeds were kept in 100 ml Erlenmeyer flasks for 15 days at  
30  
31 30°C in the dark.  
32  
33

### 34 *FOX-1 assay for the determination of hydroperoxides present in the culture filtrate*

35  
36 The total hydroperoxides (ROOH) produced by *A. parasiticus* grown in PDB were analyzed by  
37  
38 spectrophotometric assay by monitoring the oxidation of xylenol orange at 560 nm (Ferrous ion  
39  
40 Oxidation Xylenol orange: FOX-1). The sensitivity of the method was increased according to recent  
41  
42 modification by the use of triphenylphosphine and stabilizing the reagent at pH 1.7-1.8 as reported  
43  
44 by Banerjee *et al.*, (2003).  
45  
46  
47  
48  
49  
50  
51  
52

### 53 *Assay of LOX activity in maize seeds*

54  
55 The activity of lipoxygenase (LOX, EC 1.13.11.12) was tested at different time points (0-15 days),  
56  
57 in contaminated and not contaminated maize seeds. The assay was performed monitoring the diene  
58  
59 conjugates formation at 234 nm as previously reported by Reverberi *et al.*, (2005b).  
60

### *Analysis of 9- and 13-HODE present on maize seeds surface*

The seeds infected and not infected with WT and  $\Delta Apyap1$  were washed in distilled sterile water, stirring the specimens, to remove mycelia and conidia. Hydroperoxyoctadecadienoic acids (HPODEs) present on the surface of maize seeds were analysed after extraction of the samples with chloroform:methanol 2:1 v/v (20 ml x 3) in presence of 100  $\mu$ g of butylated hydroxytoluene (BHT) as an antioxidant. The two hydroxyoctadienoic acids (9-HODE and 13-HODE) obtained by reduction of the respective HPODEs, were analyzed by HPLC-APCI-MS as reported by Reverberi *et al.* (2006). Data were collected and analyzed using the Chemstation LC/MSD revision A.09.01 (Agilent Technologies). Authentic 9-HODE and 13-HODE were purchased from Cayman (Ann Arbor, Michigan, USA).

### *Analysis of aflatoxins*

The aflatoxins (B<sub>1</sub>+ B<sub>2</sub>+ G<sub>1</sub>+ G<sub>2</sub>) analyses were performed as previously reported (Fanelli *et al.* 2000) by extracting inoculated maize seeds after homogenisation in chloroform:methanol (2:1 v/v) (60 ml x 3). The extracts were collected after filtration on anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated under a N<sub>2</sub> stream and quantified by HPLC-UV-DAD.

### *Ergosterol analysis for fungal growth determination*

Contaminated and not contaminated maize seeds were extracted with 250 ml chloroform:methanol (2:1 v/v) for 1 h in the dark, in the presence of 100  $\mu$ g of butylated hydroxytoluene as an antioxidant. After extraction, the solvents were filtered on anhydrous Na<sub>2</sub>SO<sub>4</sub>, collected and concentrated under a N<sub>2</sub> stream. Ergosterol was analysed by HPLC as reported by Fabbri *et al.* (1997), using a Supelco LC-18, 5 $\mu$ m (25 cm x 4,6mm) column, methanol:water (98:2 v/v) as the mobile phase, and detecting at 282 nm.

### Statistics

All the experiments were performed in triplicate and values, referred to minimum three replicates per sample, were averaged and standard deviations were determined. In all experiments, mean values were compared using Student's *t*-test.

### Results

ROOH production, measured by FOX-1 assay, of the two *Aspergillus* strains (WT and  $\Delta Apyap1$ ) grown in PDB for 168 h at 30°C is reported in Figure 1.  $\Delta Apyap1$  presents a higher ROOH production compared with WT between 36 and 70 h with an evident peak at 42 h. After 60 h both strains show a similar trend of ROOH production. Figures 2 and 3 report LOX activity (U/mg protein) and LOOH formation (ng/g seeds) of maize seeds inoculated or not (MS) with the WT and  $\Delta Apyap1$  strains and incubated for 15 days at 30°C.  $\Delta Apyap1$  promotes a stimulation of LOX activity in maize seeds up to 11 days (stronger at 5 and 11 days) in comparison with WT. Also LOOH were formed at greater extent until the first 7 days of incubation. LOOH and LOX activity of not inoculated seeds followed a trend similar to inoculated ones.

Production of 9- and 13-HODE during the growth of both strains on maize seeds is reported in table 1. In maize seeds, not inoculated (MS) and inoculated with the two strains, a slight prevalence of 13-HODE amount compared to 9-HODE occurred. These differences in the ratio between the two regioisomers in the contaminated and not contaminated maize seeds are not always significant.

The aflatoxin biosynthesis detected in maize seeds inoculated with WT and  $\Delta Apyap1$  for 15 days at 30°C is reported in Figure 4. It was evident that aflatoxins were produced before by the  $\Delta Apyap1$  strain ( $1.1 \pm 0.1$  µg/g maize seeds at the 2<sup>nd</sup> day of incubation) compared with the WT ( $1.5 \pm 0.2$  µg/g maize seeds at the 5<sup>th</sup> day of incubation). In addition the  $\Delta Apyap1$  produces significantly ( $p < 0.001$ ) more toxins in the intervals 3-6 days and 8-12 days. No significant differences in fungal growth occurred as evident from ergosterol results (Figure 5).



## Discussion

1  
2  
3  
4  
5  
6  
7  
8 Recently we have studied *in vitro* the regulation effect of some transcription factors, in particular  
9  
10 Apyap1, hsf2-like and skn7-like, on the intracellular antioxidant defence mechanisms of some  
11  
12 aflatoxigenic strains of *A. parasiticus* (Reverberi *et al.*, 2006). Notably, there is a direct correlation  
13  
14 between levels of oxidative stressors in the cell, such as lipoperoxides, the activity of antioxidant  
15  
16 enzymes (i.e. SOD, CAT and GPX), and aflatoxin biosynthesis. In maize seeds inoculated with  
17  
18  $\Delta$ Apyap1, the activity of LOX and the LOOH amount were higher compared with the WT strain. It  
19  
20 can be argued that when the seed is infected with  $\Delta$ Apyap1 it produces more LOOH probably due to  
21  
22 the ROOH produced at high concentration by this strain as suggested by FOX-1 assay. The  
23  
24 oxidative stressors present on the surface of the seeds can trigger the activation of LOX and, in turn,  
25  
26 lead to the release of LOOH. Thus, the fungus, unable to control the increase of oxidative stressors,  
27  
28 such as LOOH, which are an important factor in the enhancement of aflatoxin biosynthesis on the  
29  
30 surface of maize seeds, produces a higher amount of toxins and more rapidly. These results suggest  
31  
32 that also in *A. parasiticus* - maize seeds interaction, oxidative stress and aflatoxin biosynthesis are  
33  
34 correlated. It appears that the antioxidant defences that control the redox balance of fungal cell, also  
35  
36 plays a role in the modulation of toxin formation on maize seeds contaminated by  $\Delta$ Apyap1. This  
37  
38 defence mechanism is a natural phenomenon that can be improved and controlled with different  
39  
40 methods. For example the production of toxins can be modulated, as previously described, by the  
41  
42 addition of antioxidants, which, in some cases, show also an intrinsic toxicity against pathogenic  
43  
44 fungi (Fanelli *et al.*, 1985, 1986, 2000; Kim *et al.*, 2004). As previously reported (Zjalic *et al.*,  
45  
46 2006), comparable effects can be accomplished with compounds that protect the plant against  
47  
48 oxidative stress. The ability of organisms to oppose oxidative stress has been exploited also as a  
49  
50 biocontrol mechanism. Some yeasts, used as biocontrol agents (Castoria *et al.*, 2003), successfully  
51  
52 compete for space and nutrients over many pathogenic fungi (Droby and Chalutz, 1994). These  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 agents, by resisting to oxidative stress conditions, colonize wounds and protect the plant from the  
4  
5 attack of different pathogens.  
6

7  
8 The results obtained in this paper suggest that Apyap1 plays a role in the regulation of  
9  
10 antioxidant/oxidant balance of the cell in *A. parasiticus* also during maize seeds colonisation. Thus,  
11  
12 it can be hypothesised that aflatoxin biosynthesis occurring during the interaction seed/pathogen is  
13  
14 affected by their redox status. Some strategies can be advanced on the basis of our results, such as  
15  
16 the use of antioxidants to minimise the formation of oxidants both in the pathogenic fungus and in  
17  
18 the plant.  
19  
20  
21

## 22 23 24 **References**

- 25  
26  
27 Banerjee D, Madhusoodanan UK, Sharanabasappa M, Ghosh S, Jacob J, 2003. Measurement of  
28  
29 plasma hydroperoxide concentration by FOX-1 assay in conjunction with triphenylphosphine.  
30  
31 Clinica Chimica Acta 337, 147-152.  
32  
33  
34 Burow GB, Nesbitt TC, Dunlap J, Keller NP. 1997. Seed lipoxygenase products modulate  
35  
36 *Aspergillus* mycotoxin biosynthesis. Molecular Plant-Microbe Interaction 10(3): 380-387.  
37  
38  
39 Calvo AM, Hinze LL, Gardner HW, Keller NP. 1999. Sporogenic effect of polyunsaturated fatty  
40  
41 acids on development of *Aspergillus* spp. Applied and Environmental Microbiology 65(8): 3668-  
42  
43 3673.  
44  
45  
46 Calvo AM, Gardner HW, Keller NP. 2001. Genetic connection between fatty acid metabolism and  
47  
48 sporulation in *Aspergillus nidulans*. Journal of Biological Chemistry 276(28): 25766-25774.  
49  
50  
51 Castoria R, Caputo L, De Curtis F, De Cicco V. 2003. Resistance of post-harvest biocontrol yeasts  
52  
53 to oxidative stress: a possible new mechanism of action. Phytopathology 93(5): 564-572.  
54  
55  
56 Champe SP, El-Zayat AA. 1989. Isolation of a sexual sporulation hormone from *Aspergillus*  
57  
58 *nidulans*. Journal of Bacteriology 171: 3982-3988.  
59  
60  
61 Droby S, Chalutz E. 1994. Mode of action of biocontrol agents of postharvest diseases. In: Wilson  
62  
63 CL, Wisniewski ME, editors. CRC press, Boca Raton FL. p. 63.

- 1  
2  
3 Estruch F. 2000. Stress-controlled transcription factors, stress-induced genes and stress tolerance in  
4 budding yeast. FEMS Microbiological Review 24: 469-486.  
5  
6  
7  
8 Fabbri AA, Fanelli C, Panfili G, Passi S, Fasella P. 1983. Lipoperoxidation and aflatoxin  
9 biosynthesis by *Aspergillus parasiticus* and *A. flavus*. Journal of General Microbiology 129: 3447-  
10 3452.  
11  
12  
13  
14 Fabbri AA, Ricelli A, Brasini S, and Fanelli C. 1997. Effect of different antifungals on the control  
15 of paper biodeterioration caused by fungi. International Biodeterioration and Biodegradation 39: 61-  
16 65.  
17  
18  
19  
20  
21  
22 Fanelli C, Fabbri AA, Pieretti S, Finotti E, Passi S. 1985. Effect of different antioxidants and free  
23 radical scavengers on aflatoxin production. Mycological Research 1: 65-69.  
24  
25  
26  
27 Fanelli C, Fabbri AA. 1989. Relationship between lipids and aflatoxin biosynthesis.  
28 Mycopathologia 107: 115-120  
29  
30  
31  
32 Fanelli C, Tasca V, Ricelli A, Reverberi M, Zjalic S, Finotti E, Fabbri AA. 2000. Inhibiting effect  
33 of medicinal mushroom *Lentinus edodes* (Berk.) Sing (Agaricomycetidae) on aflatoxin production  
34 by *Aspergillus parasiticus* Speare. International Journal of Medicinal Mushrooms 2: 229-236.  
35  
36  
37  
38  
39 Guzman De Pena D, Ruiz-Herrera J. 1997. Relationship between aflatoxin biosynthesis and  
40 sporulation in *Aspergillus parasiticus*. Fungal Genetics and Biology 21(2): 198-205.  
41  
42  
43  
44 Harschman KD, Moye-Rowley WS, Parker CS. 1988. Transcriptional activation by the SV40 AP-1  
45 recognition element in yeast is mediated by a factor similar to Ap-1 that is distinct from GCN4. Cell  
46 53: 321-330  
47  
48  
49  
50  
51 He H, Soncin F, Grammatikakis N, Li Y, Sigandò A, Gong J, Brown SA, Kingston RE, Calderwood  
52 SK. 2003. Elevated expression of heat shock factor (HSF) 2A stimulates HSF1-induced  
53 transcription during stress. Journal of Biological Chemistry 278: 34465-34475  
54  
55  
56  
57  
58 Jayashree T, Subramanyam C. 2000. Oxidative stress as a prerequisite for aflatoxin production by  
59 *Aspergillus parasiticus*. Free Radical Biology and Medicine 29: 981-985.  
60

- 1  
2  
3 Kim JH, Campbell BC, Yu J, Mahoney N, Chan KL, Molyneux RJ, Bhatnagar D, Cleveland TE.  
4  
5 2004. Examination of fungal stress response genes using *Saccharomyces cerevisiae* as a model  
6  
7 system: targeting genes affecting aflatoxin biosynthesis by *Aspergillus flavus* Link. Applied  
8  
9 Microbiology and Biotechnology 67(6): 807-815.  
10  
11  
12 Moye-Rowley WS. 2003. Regulation of transcriptional response to oxidative stress in fungi:  
13  
14 similarities and differences. Eukaryotic Cell 2: 381-389.  
15  
16  
17 Reverberi M, Fanelli C, Zjalic S, Briganti S, Picardo M, Ricelli A, Fabbri AA. 2005. Relationship  
18  
19 among lipoperoxides, jasmonic acid and indole-3-acetic acid formation in potato tuber after  
20  
21 wounding. Free Radical Research 39: 637-647.  
22  
23  
24 Reverberi M, Zjalic S, Ricelli A, Fabbri AA, Fanelli C. 2006. Oxidant/antioxidant balance in  
25  
26 *Aspergillus parasiticus* affects aflatoxin biosynthesis. Mycological Research 22(1):39-47.  
27  
28  
29 Tsitsigiannis D, Kowiesky TM, Zarnowsky R, Keller N P. 2004. Endogenous lipogenic regulators  
30  
31 of spore balance in *Aspergillus nidulans*. Eukaryotic Cell, 3(6): 1398-1411.  
32  
33  
34 Zjalic S, Reverberi M, Ricelli A, Granito VM, Fanelli C, Fabbri AA. 2006. *Trametes versicolor*: a  
35  
36 possible tool for aflatoxin control. International Journal of Food Microbiology 1;107(3):243-249.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Figure legends

**Figure 1.** ROOH detected by FOX-1 assay in culture media (PDB) inoculated with *A. parasiticus* WT or  $\Delta Apyap1$  incubated at 30°C for different periods (10, 12, 18, 21, 24, 30, 36, 42, 48, 60, 72, 96 and 168 h). The results represent the mean  $\pm$  S.E.M. of three experiments.

**Figure 2.** Lipoxygenase activity of maize seeds not inoculated (MS) or inoculated with *A. parasiticus* WT or  $\Delta Apyap1$  and incubated at 30°C for different periods (0-15 days). The results represent the mean  $\pm$  S.E.M. of three experiments.

**Figure 3.** Total lipoperoxides (LOOH) originated from linoleic acid, extracted from maize seeds surface not inoculated (MS) or inoculated with *A. parasiticus* WT or  $\Delta Apyap1$  and incubated at 30°C for different periods (0-15 days). The results represent the mean  $\pm$  S.E.M. of three experiments.

**Figure 4.** Aflatoxin production in maize seeds inoculated with *A. parasiticus* WT or  $\Delta Apyap1$  and incubated at 30°C for different periods (0-15 days). The results represent the mean  $\pm$  S.E.M. of three experiments.

**Figure 5.** Fungal growth reported as ergosterol content (ng/g seeds) of *A. parasiticus* WT or  $\Delta Apyap1$  inoculated in maize seeds and incubated at 30°C for different time periods (0-15 days). The results represent the mean  $\pm$  S.E.M. of three experiments.

**Table 1.** 9 and 13 regioisomers of hydroxylated linoleic acid (ng/g maize seeds) produced on the surface of maize seeds at different times (0-15 days) not inoculated (MS) or inoculated with *A. parasiticus* WT and  $\Delta Apyap1$ . The results represent the mean  $\pm$  S.E.M. of three separate experiments.

<i>Time (days)</i>	WT		$\Delta Apyap1$		MS	
	13-HODE	9-HODE	13-HODE	9-HODE	13-HODE	9-HODE
0	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	0.1 $\pm$ 0.05	0.1 $\pm$ 0.05
1	1.1 $\pm$ 0.1	1.0 $\pm$ 0.1	1.5 $\pm$ 0.2	1.4 $\pm$ 0.2	0.6 $\pm$ 0.1	0.5 $\pm$ 0.1
2	2.9 $\pm$ 0.3	3.2 $\pm$ 0.2	3.5 $\pm$ 0.3	3.7 $\pm$ 0.5	0.9 $\pm$ 0.1	0.7 $\pm$ 0.1
3	1.8 $\pm$ 0.2	1.9 $\pm$ 0.2	3.2 $\pm$ 0.4	2.9 $\pm$ 0.4	1.0 $\pm$ 0.1	1.2 $\pm$ 0.2
5	0.6 $\pm$ 0.1	0.3 $\pm$ 0.1	2.0 $\pm$ 0.5	2.0 $\pm$ 0.3	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1
7	0.9 $\pm$ 0.2	0.5 $\pm$ 0.1	1.2 $\pm$ 0.2	0.6 $\pm$ 0.1	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1
11	2.8 $\pm$ 0.3	1.9 $\pm$ 0.3	3.0 $\pm$ 0.4	1.9 $\pm$ 0.5	0.8 $\pm$ 0.1	1.0 $\pm$ 0.2
15	4.0 $\pm$ 0.5	3.0 $\pm$ 0.4	2.0 $\pm$ 0.2	1.5 $\pm$ 0.2	0.7 $\pm$ 0.1	0.8 $\pm$ 0.2

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only




Figure 1. Reverberi et al.






Figure 2. Reverberi et al.




Figure 3. Reverberi et al.




Figure 4. Reverberi et al.




Figure 5. Reverberi et al.