



HAL
open science

Phytosterol consumption and the anabolic steroid boldenone in humans: a hypothesis piloted

Martine M Ros, Saskia Sterk, Hans Verhagen, Anton Fh Stalenhoef, Nynke de
Jong

► **To cite this version:**

Martine M Ros, Saskia Sterk, Hans Verhagen, Anton Fh Stalenhoef, Nynke de Jong. Phytosterol consumption and the anabolic steroid boldenone in humans: a hypothesis piloted. *Food Additives and Contaminants*, 2007, 24 (07), pp.679-684. 10.1080/02652030701216727 . hal-00577549

HAL Id: hal-00577549

<https://hal.science/hal-00577549v1>

Submitted on 17 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Phytosterol consumption and the anabolic steroid boldenone in humans: a hypothesis piloted

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2006-306.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	12-Jan-2007
Complete List of Authors:	Ros, Martine; National Institute for Public Health and the Environment (RIVM) Sterk, Saskia; National Institute for Public Health and the Environment (RIVM) Verhagen, Hans; National Institute for Public Health and the Environment (RIVM) Stalenhoef, Anton; Radboud University Nijmegen Medical Centre, Department of Internal Medicine de Jong, Nynke; National Institute for Public Health and the Environment (RIVM)
Methods/Techniques:	Chromatography - GC/MS, Chromatography - HPLC, Chromatography - LC/MS
Additives/Contaminants:	Veterinary drug residues - anabolic steroids
Food Types:	Novel food, Oils and fats

SCHOLARONE™
Manuscripts

1
2
3
4 **Phytosterol consumption and the anabolic steroid boldenone in**
5
6 **humans: a hypothesis piloted**
7
8

9
10
11 Martine M Ros¹, Saskia S Sterk¹, Hans Verhagen¹, Anton FH Stalenhoef², Nynke de
12
13 Jong¹
14

15
16
17
18 ¹ National Institute for Public Health and the Environment (RIVM), PO Box 1, 3720
19
20 BA, Bilthoven, the Netherlands
21

22
23 ² Department of Internal Medicine, 463, Radboud University Nijmegen Medical
24
25 Centre, PO Box 9101, 6500 HB Nijmegen, the Netherlands
26
27

28
29
30 Correspondence to:

31
32 Dr. Nynke de Jong, National Institute for Public Health and the Environment, Centre
33
34 for Nutrition and Health, P.O. Box 1, 3720 BA, Bilthoven, the Netherlands
35

36
37 Tel: +31 30 274 4135 Fax: +31 30 274 4466
38

39
40 E-mail: Nynke.de.Jong@rivm.nl
41

42
43
44 Running title: phytosterols and boldenone in humans
45

46
47 Key words: phytosterol enriched margarine, boldenone, human intervention, anabolic
48
49 steroid, postlaunch monitoring, side effect
50

Abstract

The presence of the anabolic steroid boldenone in animals has become a research topic as its occurrence is proposed to be a marker for illegal hormone administration. However, boldenone can also be formed from β -sitosterol, a phytosterol present in animal feed, as well as from endogenous sources. The observations in animals together with the increased consumption of phytosterol-enriched foods in the western population led us to the hypothesis that consumption of phytosterol-enriched foods might possibly lead to increased boldenone levels in humans. We performed a pilot study among female volunteers (n=10) to investigate whether boldenone concentrations in urine were detectable after consumption of 25 g/day of phytosterol-enriched margarines for one week. Urine samples were collected at day 0, day 3 or 4 and day 7. Urine of a sitosterolemia (a rare autosomal recessively inherited lipid metabolic disorder) patient was collected as a positive control case. No traces of boldenone were detected in either the volunteers or in the patient. In conclusion, there is no evidence of formation of boldenone in women after consumption of the recommended amount of phytosterol-enriched margarines.

Introduction

For a number of years, the presence of the anabolic steroid boldenone (1-dehydrotestosterone or androsta-1,4-diene-17 β -ol-3-one) in various animal species has become a topic of research as the occurrence of this hormone or its metabolites in biological samples is proposed to be a marker for illegal hormone administration (De Brabander and others 2004). Especially urine samples of cattle and veal calves have been subject of research in various EU countries. The investigation into the origin of the boldenone in bovine urine is still ongoing. Boldenone can be formed from β -sitosterol (a phytosterol naturally occurring in plants) present in animal feed, but also endogenous production can still be one of the sources (Poelmans and others 2005). Phytosterols are a normal constituent of the human diet. Normal levels of consumption are approximately 200-400 mg/day, which may increase to 700-800 mg/day in those who consume a large amount of soy-based foods (Andersson and others 2004). The recently introduced phytosterol-enriched foods deliver 2-3 g of phytosterols per day if the recommended amount is consumed. The intestinal absorption of phytosterols is estimated between 0.4-3.5% (Hallikainen and others 2000).

The observations in animals together with the increased consumption of phytosterol-enriched foods in the western population to lower (slightly) elevated serum cholesterol levels lead us to the hypothesis that in humans phytosterol-enriched food may possibly lead to an unwanted amount of the anabolic steroid boldenone as in animals. This would then identify a new potential health hazard in humans.

Phytosterol-enriched margarines are on the market for some years now, and other plant sterol-enriched food products including dairy products have also been introduced. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

phytosterols are thought to displace cholesterol from mixed micelles in the intestine and thereby reduce intestinal cholesterol absorption; the exact mechanism, however, remains still unknown (Katan and others 2003). In animals boldenone works as an anabolic agent and is misused in cattle fattening. It is commonly used to enhance athletic performance and muscular development in humans. Hypertension, heart attacks, strokes, liver and other types of internal organ cancers have been associated with anabolic steroid use (De Brabander and others 2004; Sullivan and others 1998).

Within the framework of Postlaunch Monitoring (PLM) of functional foods in which positive but also potential side effects are investigated (de Jong and others 2004; 2005a; 2005b; Wolfs and others 2006) we studied the hypothesis postulated above. To our knowledge there were no data of biotransformation of the phytosterol β -sitosterol into boldenone in humans. We performed a pilot study in 10 women to investigate whether boldenone concentrations in urine were detectable after consumption of the recommended daily amounts of phytosterol-enriched margarines. In addition, a urine sample of a sitosterolemia patient was analysed as a positive control case. Patients with sitosterolemia are characterized by a > 50-fold elevation in plasma phytosterol levels, which results from an increased absorption and decreased hepatic removal of phytosterols that leads to accumulation of phytosterols in blood and tissues (Ketomaki and others 2005). Because these patients have very high levels of plasma phytosterols we anticipated that especially in these patients the presence of boldenone might be demonstrated.

Materials and methods

Study design

1
2
3 For our pilot 10 healthy female volunteers (aged 22 to 51) were recruited. In daily life,
4 these women did not use phytosterol-enriched margarines. We included only women
5 because they excrete less to none anabolic hormones by nature. During our one week
6 trial we provided phytosterol-enriched margarines. The participants were asked to
7 consume 25 grams per day (i.e. 2 grams of phytosterols or ~1 gram of β -sitosterol) of
8 the phytosterol-enriched margarines, which corresponds to the recommended intake by
9 the manufacturer. At the beginning of the study 7 portions of 25 grams were distributed
10 to the volunteers. Spot urine samples were collected on day 0, on day 3 or 4, and on
11 day 7 at around lunchtime. In this way we collected one voidance from a day at about
12 the same time. Within 30 minutes after collection the urine samples were frozen at
13 minus 20°C. As a normal food product was supplied in physiological quantities and the
14 collection of spot urine samples was not invasive, no authorization from the medical
15 ethical committee was necessary (personal communication with the Medical
16 Committee of Ethics of the University Medical Centre Utrecht, April 2006). The
17 volunteers received a little incentive after participation.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 In the blind urine samples, β -sitosterol, campesterol, stigmasterol, β -boldenone, α -
42 boldenone, α -testosterone and β -testosterone were analysed according to ARO-SOP
43 507 (ARO-SOP 507 RIVM: Bilthoven) a detailed description of which follows below.
44 In addition, one spot urine sample of our positive control case, i.e. a patient with
45 sitosterolemia, was collected and analysed.
46
47
48
49
50
51
52
53
54

55 **Method of analysis**

56 *Chemicals*

57
58
59
60

1
2
3 All chemicals and reagents were of high purity quality. α/β -boldenone, α/β -
4 testosterone, androsterone, etiocholanolon, α -androsterone, testosterone-D2 and β -
5 boldenone-D3 were obtained from the RIVM-CRL, Bank of Reference Standards.
6
7 Cholesterol, coprostanol, campesterol, β -sitosterol, stigmasterol were obtained from
8 Matreya. Cholesterol-D6 was obtained from Isotel. β -glucuronidase from E.Coli K12
9 (Roche). Derivatization reagent for coprostanol, cholesterol, cholesterol-D6,
10 campesterol and sitosterol consists of 25 μ l N-methyl-N-
11 trimethylsilyltrifluoroacetamide (MSTFA) / ammoniumiodide/dithiothreitol (1000:2:4,
12 v/w/w) (Alltech), for α/β -boldenone, α/β -testosterone, androsterone, etiocholanolon, α -
13 androsteron, testosterone-D2 and β -boldenone-D3 the derivatization reagent consisted
14 of 10 μ l heptafluorobutyric Acid Anhydride (HFAA) (Pierce) and 40 μ l of dried
15 acetone. Phosphate buffer pH 7.4 was prepared by dissolving 2.278 g of
16 disodiumhydrogenphosphate and 0.416 g of potassium-dihydrogenphosphate in 800 ml
17 of water, the pH was adjusted to 7.4 ± 0.1 and water was added to a final volume of
18 1000 ml.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 *Apparatus*

42
43 Liquid Chromatography (LC): Waters Chromatography autosampler, two Waters
44 pumps, Pharmacia controller, ThermoQuest multi-channel UV-detector. HPLC-column
45 used was a Superspher RP-18 (L 125 mm, 4 mm ID, 4 μ m) with pre-column (Waters).
46
47 Column temperature 40°C. Fraction-collector (Foxy Jr). Datasystem, PC1000
48 ThermoQuest. The LC mobile phase consists of solution A: 65:35 v/v-%
49 methanol/water, solution B: 100% methanol. Gradient starts at 0% B, after 8 min the
50 percentage B is increased in 8 min to 100% and remains at 100% till 25 min. The
51 gradient then returns in 0.1 min to the initial condition. The flow rate was 0.7 ml/min.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 Gas-Chromatography coupled to a mass-spectrometer (GC-MS) analysis was carried
7
8 out on an Agilent 5973 MSD. GC capillary column, 30 meter VF-17MS (Varian) i.d.
9
10 0.25 mm, 0.15 μm film thickness, constant flow of 1.1 ml helium/minute. Injection,
11
12 splitless mode at 250°C, injection volume 2 μl . The oven temperature was kept
13
14 constant at 80°C for 1 min and was increased, 20°C per min, to 340°C and was kept
15
16 constant at this temperature for 4 min.
17
18
19
20
21

22 *Extraction procedure*

23 *Isolation*

24
25 A sample portion of 5 ml of urine was transferred to a 10 ml glass tube. The samples
26
27 were spiked with 10 ng β -boldenone-D3 , 10 ng testosterone-D2 and 25 cholesterol-
28
29 D6. To the samples 1 ml of phosphate buffer (pH 7.4) and 50 μl of β -glucuronidase
30
31 was added. The mixture was vortexed and hydrolysed for 3 hours at 52°C.
32
33
34
35
36
37
38

39 A SPE C₁₈ column (3 ml) was preconditioned with 3 ml of methanol and 3 ml of water.
40
41 The centrifuged sample was passed through the column. The SPE C₁₈ column was
42
43 washed with 3 ml of water and 3 ml 30:70 v/v-% acetonitrile/water. The anabolic
44
45 steroids were eluted with 4 ml of 80:20 v/v-% methanol/water. The eluate was
46
47 evaporated at 55°C under a gentle stream of nitrogen until dry and further processed as
48
49 described in Preparative HPLC. The sterols were eluted with 3 ml of iso-octane, the
50
51 eluate was collected and evaporated at 55°C under a gentle nitrogen until dryness and
52
53 derivatised as described under derivatisation.
54
55
56
57
58
59
60

Preparative HPLC

1
2
3 The dried extracts were reconstituted in 120 μl of 65:35 v/v-% methanol/water, from
4 which 100 μl was injected. One fraction was collected from 4.9 to 20 minutes. The
5
6 collected fraction was dried under a stream of nitrogen. The dried extracts were
7
8 reconstituted in 300 μl ethanol, transferred to a 2 ml vial and evaporated at 55°C under
9
10 a gentle stream of nitrogen to dryness.
11
12
13
14

15 16 17 *Derivatisation*

18
19 The dried extracts containing cholesterol, coprostanol, campesterol, sitosterol and
20
21 cholesterol-D6 were reconstituted in 20 μl of derivatisation reagent (MSTFA++) and
22
23 incubated for one hour at 60°C. After 1 hour the derivatisation reagent was evaporated.
24
25
26 The dried residue was reconstituted in 75 μl of iso-octane.
27
28
29
30

31
32 The dried extracts containing α/β -boldenone, α/β -testosterone, α/β -androsterone,
33
34 etiocholanolon, testosterone-D2 and β -boldenone-D3 were reconstituted in 50 μl of
35
36 derivatisation reagent (HFBA/acetone) and incubated for one hour at 60°C. After 1 hour
37
38 the derivatisation reagent was evaporated. The dried residue was reconstituted in 50 μl
39
40 of iso-octane.
41
42
43
44

45 46 *Detection*

47
48 Mass spectrometric detection was performed in selected ion monitoring (SIM) mode,
49
50 see Table 1 and 2 for an overview of the m/z monitored and typical retention times of
51
52 the analytes.
53
54
55
56

57 58 **Results**

1
2
3 In Table 3 the results of our pilot study are presented. Unexpectedly and in contrast
4
5 with our hypothesis no traces of boldenone in the urine of any of the volunteers could
6
7 be detected. The levels of β -sitosterol, campesterol and stigmasterol were almost the
8
9 same over the 7 days of consumption of phytosterol enriched margarines. Also the
10
11 patient with sitosterolemia showed no traces of urinary boldenone. As expected, the
12
13 patient levels of β -sitosterol, campesterol and stigmasterol were elevated compared to
14
15 the mean levels of the female volunteers. Table 4 presents preliminary data on urinary
16
17 levels of α -testosterone and β -testosterone, which showed a slight increase in
18
19 testosterone compared to baseline levels, after a few days of phytosterol-enriched
20
21 margarine consumption.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

From our pilot we conclude that there is no evidence of formation of the anabolic steroid boldenone in women volunteers after consumption of the recommended amount (i.e. 25 grams per day) of phytosterol-enriched margarines. In addition, although the urinary sitosterol and stigmasterol levels in our sitosterolemia patient are elevated as expected, also no urinary boldenone in this positive control case could be detected. Therefore, within the PLM perspective of this study there seems no health hazard for women forming boldenone associated with this type of functional food.

In contrast to the clear results on the absence of urinary boldenone in both our human volunteers and in our sitosterolemia patient, the results on urinary testosterone are rather uncertain. At day 3 or 4 the ratio of β -testosterone and α -testosterone seems to be high, but this value was influenced by one low concentration of α -testosterone near the detection limit. The variation in these data is large and it should be demonstrated in future larger trials whether any shift in concentration is caused by true differences in human metabolic systems or by measurement errors.

Several (mechanistic) issues remain to be clarified in the future, especially in cattle consuming a large amount of phytosterol -rich feed. There is still debate about the implications of the detection of boldenone in cattle and veal calves' urine. Questions that are raised focus among others on whether boldenone is endogenously synthesized or indeed illegally administered. Also, detection methods might have been improved over time. Another theory that has been proposed is that due to bacteria in feed or a different feed composition (more sterols), the side chain at the c17 position in the D-ring can be broken and the sterols can be formed into anabolic steroids such as

1
2
3 boldenone or testosterone. One should bear in mind that in human food we do not
4
5 expect this type of bacteria, but the implications for humans in the light of beef
6
7 consumption needs more research.
8
9

10
11
12 To our knowledge, this study is the first to investigate the association between
13
14 phytosterol (β -sitosterol) consumption and urinary boldenone concentration in humans.
15
16 And although our pilot study only aimed at hypothesis generation, we very well realize
17
18 that there are limitations to our pilot study. We had a limited number of participants
19
20 who only consumed the recommended dose of physterol-enriched margarine, during a
21
22 limited amount of time. But as the metabolic effects of physiological amounts of
23
24 phytosterol-enriched margarines are normally observed within one day (Ellegard and
25
26 others 2005) we are of the opinion that the duration of our pilot had been sufficiently
27
28 long. Within the scope of the hypothetical aim of this pilot, it was not necessary to
29
30 collect representative 24h urine samples to take into account variations in urine
31
32 volume. Therefore, spot urine samples were appropriate to investigate whether
33
34 boldenone is detectable in urine. Although no levels of boldenone were found, we are
35
36 of the opinion that in theory it would have been possible to detect sterols and
37
38 boldenone in human urine samples. Given a mean urine excretion of 1,5 l/day, a
39
40 phytosterol intake of 2 g/day, and a mean intestinal absorption of 2% (see introduction
41
42 section), the theoretical maximal concentration of sterols in urine is a factor 300 higher
43
44 (2% of 2 g sterols = $40\text{ mg}/1,5\text{ l} = \sim 30\text{mg/l}$) than the detection limit of $0.1\ \mu\text{g/l}$. We did
45
46 not have any compliance measure of the phytosterol-enriched margarine consumption,
47
48 but we do not have any indications that among our 10 female volunteers who were
49
50 contacted regularly during the study period there were any non-compliant subjects.
51
52
53
54
55
56
57
58
59
60

1
2
3 Our main focus was the detection of urinary boldenone. Despite the limitations
4 described above we conclude that within our setting consumption of the recommended
5 amounts of phytosterol -enriched margarines, does not lead to detectable levels of the
6 anabolic steroid boldenone in women. Further research on boldenone formation is
7 dependent on the observations in cattle and veal calves in relation to beef and veal
8 consumption. Also, the differences in the exact metabolic mechanisms between
9 humans and cattle, and among humans themselves with regard to testosterone are
10 topics of further research.
11
12
13
14
15
16
17
18
19
20
21

22 **Acknowledgements**

23
24
25
26 The authors would like to thank the volunteers as well as the sitosterolemia patient for
27 their willingness to participate. We also would like to thank H.J. van Rossum and M.H.
28 Blokland for their technical assistance. We acknowledge the Netherlands Organisation
29 for Health Research and Development (ZonMW) for supporting the project on
30 PostLaunch Monitoring (Grant Number 014-12-010).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- ARO-SOP 507: Analysis of free and conjugated boldenone in bovine urine by GC-MS. Standard Operating Procedure nr 507, Laboratory of Food and Residue Analyses. RIVM: Bilthoven.
- Andersson SW, Skinner J, Ellegard L, Welch AA, Bingham S, Mulligan A, Andersson H, Khaw KT. 2004. Intake of dietary plant sterols is inversely related to serum cholesterol concentration in men and women in the EPIC Norfolk population: a cross-sectional study. *Eur J Clin Nutr* 58(10):1378-85.
- De Brabander HF, Poelmans S, Schilt R, Stephany RW, Le Bizec B, Draisci R, Sterk SS, van Ginkel LA, Courtheyn D, Van Hoof N. 2004. Presence and metabolism of the anabolic steroid boldenone in various animal species: a review. *Food Addit Contam* 21(6):515-25.
- de Jong N, Ocké MC. 2004. Postlaunch monitoring on functional foods. Methodology development (I). Bilthoven: RIVM reportnr. 35003001/2004.
- de Jong N, Buurma-Rethans EJM, Fransen HP, Ocké MC. 2005a. Postlaunch monitoring of functional foods. Methodology development (II). Bilthoven: RIVM reportnr. 35003005/2005.
- de Jong N, Fransen HP, van den Berg SW, Ocké MC. 2005b. Postlaunch monitoring of functional foods. Methodology development (III). Bilthoven: RIVM reportnr. 35003006/2005.
- Ellegard L, Andersson H, Bosaeus I. 2005. Rapeseed oil, olive oil, plant sterols, and cholesterol metabolism: an ileostomy study. *Eur J Clin Nutr* 59(12):1374-8.

- 1
2
3 Hallikainen MA, Sarkkinen ES, Gylling H, Erkkila AT, Uusitupa MI. 2000.
4
5 Comparison of the effects of plant sterol ester and plant stanol ester-enriched
6
7 margarines in lowering serum cholesterol concentrations in
8
9 hypercholesterolaemic subjects on a low-fat diet. *Eur J Clin Nutr* 54(9):715-25.
10
11
12
13 Katan MB, Grundy SM, Jones P, Law M, Miettinen T, Paoletti R. 2003. Efficacy and
14
15 safety of plant stanols and sterols in the management of blood cholesterol
16
17 levels. *Mayo Clin Proc* 78(8):965-78.
18
19
20
21 Ketomaki A, Gylling H, Miettinen TA. 2005. Non-cholesterol sterols in serum,
22
23 lipoproteins, and red cells in statin-treated FH subjects off and on plant stanol
24
25 and sterol ester spreads. *Clin Chim Acta* 353(1-2):75-86.
26
27
28
29
30 Poelmans S, De Wasch K, Noppe H, Van Hoof N, Van Cruchten S, Le Bizec B,
31
32 Deceuninck Y, Sterk S, Van Rossum HJ, Hoffman MK. 2005. Endogenous
33
34 occurrence of some anabolic steroids in swine matrices. *Food Addit Contam*
35
36 22(9):808-15.
37
38
39
40 Sullivan ML, Martinez CM, Gennis P, Gallagher EJ. 1998. The cardiac toxicity of
41
42 anabolic steroids. *Prog Cardiovasc Dis* 41(1):1-15.
43
44
45
46 Wolfs M, de Jong N, Ocke MC, Verhagen H, Verschuren WM. 2006. Effectiveness of
47
48 customary use of phytosterol/-stanol enriched margarines on blood cholesterol
49
50 lowering. *Food Chem Toxicol* 44(10):1682-8.
51
52
53
54
55
56
57
58
59
60

Table 1. Overview of the m/z of the different TMS-compounds measured.

Compound	m/z	Retention time (min.)
Cholesterol	368	13.08
Campesterol	382	13.43
Sitosterol	396	13.68
Stigmasterol	394	13.50
Coprostanol	370	12.66
Cholesterol-D6	374	13.08

Table 2. Overview of the m/z of the different (HFB) compounds measured.

Compound	m/z	Retention time (min.)
β-Boldenone	678	12.97
Boldenone-D3	681	12.95
α-Boldenone	678	13.30
α-testosterone	680	12.81
β-testosterone	680	13.29
androsterone	486	15.02
etiocholanolone	486	15.11
α-androsteron	486	15.83
testosterone-D2	682	13.28

Table 3. Urine levels of boldenone and phytosterols in 10 healthy human volunteers and a sitosterolemia patient

	Volunteers			Patient with sitosterolemia ¹
	Baseline	Day 3-4	Day 7	
β-boldenone ($\mu\text{g/l}$)	nd	nd	nd	nd ²
α-boldenone ($\mu\text{g/l}$)	nd	nd	nd	nd
Campesterol ($\mu\text{g/l}$)				
mean \pm sd	0.8 \pm 0.3	0.9 \pm 0.4	0.9 \pm 0.3	7.1
median (min-max)	0.7 (0.5-1.5)	0.8 (0.4-1.5)	1.0 (0.5-1.2)	
Stigmasterol ($\mu\text{g/l}$)				
mean \pm sd	0.2 \pm 0.2	0.1 \pm 0.2	0.3 \pm 0.3	1.5
median ³ (min-max)	-(0.0-0.5)	-(0.0-0.3)	-(0.0-0.5)	
β-sitosterol ($\mu\text{g/l}$)				
mean \pm sd	1.1 \pm 0.6	0.9 \pm 0.5	1.2 \pm 0.4	13.5
median (min-max)	1.1 (0.5-2.4)	0.8 (0.5-1.9)	1.1 (0.6-1.9)	

¹ In this patient levels of boldenone and phytosterols were measured once

² nd: not detected below limit of detection: i.e. for campesterol, stigmasterol and β -sitosterol and boldenone $< 0.1\mu\text{g/l}$

³ not calculated due to statistical limitations



Table 4. Urine levels of testosterone in 10 healthy human volunteers and a sitosterolemia patient

	Volunteers			Patient with sitosterolemia ¹
	Baseline	Day 3-4	Day 7	
α-testosterone ($\mu\text{g/l}$)				
mean \pm sd	1.5 \pm 1.3	3.5 \pm 4.1	3.3 \pm 1.8	2.8
median (min-max)	0.9 (0.5-3.6)	3.2 (0.0-13.3)	3.5 (0.8-5.2)	
β- testosterone ($\mu\text{g/l}$)				
mean \pm sd	1.3 \pm 0.4	3.2 \pm 2.1	1.8 \pm 0.7	0.1
median (min-max)	1.2 (0.8-1.9)	2.8 (0.8-7.0)	1.9 (1.0-2.7)	
Ratio				
β- testosterone/ α- testosterone				
mean \pm sd	1.2 \pm 0.6	1.4 \pm 1.0 ²	0.9 \pm 0.8	-
median (min-max)	1.0 (0.5-2.1)	1.2 (0.3-2.7)	0.5 (0.2-2.3)	


¹ In this patient levels of testosterone were measured once

² Mean ratio is 23.2 if one person with values near detection limit is included.


a. Testosterone

b. β -sitosterol

c. Campesterol



d. Stigmasterol



e. Boldenone




Figure 1. Structures of a. testosterone, b. β -sitosterol, c. campesterol, d. stigmasterol and e. boldenone