



HAL
open science

Semi-classical trace formulas and heat expansions

Yves Colin de Verdière

► **To cite this version:**

| Yves Colin de Verdière. Semi-classical trace formulas and heat expansions. 2011. hal-00578731v3

HAL Id: hal-00578731

<https://hal.science/hal-00578731v3>

Preprint submitted on 29 Mar 2011 (v3), last revised 20 Jul 2011 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Semi-classical trace formulas and heat expansions

Yves Colin de Verdière *

March 30, 2011

Introduction

There is a strong similarity between the expansions of the heat kernel as worked out by people in Riemannian geometry in the seventies (starting with the famous “Can one hear the shape of a drum” by Mark Kac [Kac], the Berger paper [Berger] and the Mc-Kean-Singer paper [McK-Si]) and the so-called semi-classical trace formulas developed by people in semi-classical analysis (starting with Helffer-Robert [He-Ro]). In fact, this is not only a similarity, but, as we will prove, each of these expansions, even if they differ when expressed numerically for some example, can be deduced from the other one as formal expressions of the fields.

Let us look first at the *heat expansion* on a smooth closed Riemannian manifold of dimension d , (X, g) , with the (negative) Laplacian Δ_g ¹. The heat kernel $e(t, x, y)$, with $t > 0$ and $x, y \in X$, is the Schwartz kernel of $\exp(t\Delta_g)$: the solution of the heat equation $u_t - \Delta_g u = 0$ with initial datum u_0 is given by

$$u(t, x) = \int_X e(t, x, y) u_0(y) |dy|_g .$$

The function $e(t, x, x)$ admits, as $t \rightarrow 0^+$, the following asymptotic expansion:

$$e(t, x, x) \sim (4\pi t)^{-d/2} (1 + a_1(x)t + \dots + a_l(x)t^l + \dots) .$$

The a_l 's are given explicitly in [Gilkey2], page 201, for $l \leq 3$ and are known for $l \leq 5$ [Avramidi, vdV]. See also the related works by Hitrik and Polterovich [Hi, Hi-Po-1, Hi-Po-2, Po]. They are universal polynomials in the components of the curvature tensor and its co-variant derivatives. For example $a_0 = 1$, $a_1 = \tau_g/6$ where τ_g is the scalar curvature.

*Institut Fourier, Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France); yves.colin-de-verdiere@ujf-grenoble.fr

¹In this note, we will not follow the usual sign convention of geometers, but the convention of analysts

The previous asymptotic expansion gives the expansion of the trace by integration over X and has been used as an important tool in spectral geometry:

$$\text{trace}(e^{t\Delta_g}) = \int_X e(t, x, x) |dx|_g = \sum_{k=1}^{\infty} e^{\lambda_k t} ,$$

where $-\lambda_1 = 0 \leq -\lambda_2 \leq \dots \leq -\lambda_k \leq \dots$ is the sequence of eigenvalues of $-\Delta_g$ with the usual convention about multiplicities. If $d = 2$, this gives

$$\text{trace}(e^{t\Delta_g}) = \frac{1}{4\pi t} \left(\text{Area}(X) + \frac{2\pi\chi(X)}{6}t + o(t^2) \right) ,$$

where $\chi(X)$ is the Euler characteristic of X .

There is an extension of the previous expansion in the case of Laplace type operators on fiber bundles: the coefficients of the expansion are then polynomials in the co-variant derivatives of the curvature of the metric and of the connection on the fiber bundle. The heat expansion can be re-interpreted as an expansion of the Schwartz kernel of $f(-\hbar^2\Delta_g)$ on the diagonal $x = y$ in powers of \hbar with $f(u) = \exp(-u)$ and $t = \hbar^2$. This is a particular case of the semi-classical trace.

Let us describe the semi-classical setting in the flat case: \widehat{H}_\hbar is a self-adjoint \hbar -pseudo-differential operator with Weyl symbol $H(x, \xi)$ in some open domain X in \mathbb{R}^d , or more generally on a Riemannian manifold. Let $f \in \mathcal{S}(\mathbb{R})$ and look at $f(\widehat{H}_\hbar)$. Under some suitable assumptions (ellipticity at infinity in ξ) on H , $f(\widehat{H}_\hbar)$ is a pseudo-differential operator whose Weyl symbol $f^\star(H)$ is a formal power series in \hbar , given using the Moyal product denoted by \star , by the following formula (see [Gracia] for explicit formulas) at the point $z_0 \in T^\star X$:

$$f^\star(H)(z_0) = (2\pi\hbar)^{-d} \left(\sum_{l=0}^{\infty} \frac{1}{l!} f^{(l)}(H(z_0)) (H - H(z_0))^\star{}^l(z_0) \right) .$$

From the previous formula, we see that the symbol of $f(\widehat{H}_\hbar)$ at the point z depends only of the Taylor expansions of H at the point z and of f at the point $H(z)$. In the paper [H-P], the authors study the case of the magnetic Schrödinger operator whose Weyl symbol is $H_{a,V}(x, \xi) = \sum_{j=1}^d (\xi_j - a_j(x))^2 + V(x)$ and show that the Schwartz kernel of $f(\widehat{H}_{\hbar,a,V})$ at the point (x, x) admits an asymptotic expansion of the form

$$[f(\widehat{H}_{\hbar,a,V})](x, x) = (2\pi\hbar)^{-d} \left[\sum_{j=0}^{\infty} \hbar^{2j} \left(\sum_{l=0}^{k_j} \int_{\mathbb{R}^d} f^{(l)}(\|\xi\|^2 + V(x)) Q_{j,l}^{a,V}(x, \xi) |d\xi| \right) \right]$$

where the $Q_{j,l}^{a,V}(x, \xi)$'s are polynomials in ξ calculated from the Taylor expansions of the magnetic field $B = da$ and V at the point x . The proof in [H-P] uses a pseudo-differential calculus adapted to the magnetic field.

We will give a simplified version of the expansion replacing the (non unique) $Q_{j,l}^{a,V}(x, \xi)$'s by functions $P_{j,l}^{B,V}(x)$'s which are uniquely defined and are given by universal $O(d)$ -invariant polynomials of the Taylor expansions of B and V at the point x . We present then two ways to compute the $P_{j,l}^{B,V}$'s:

- we can first use Weyl's invariant theory (see Gilkey's book [Gilkey2]) in order to reduce the problem to the determination of a finite number of numerical coefficients; then simple examples, like harmonic oscillator and constant magnetic field, allow to determine (part of) these coefficients.
- The $P_{j,l}^{B,V}$'s are related in a very simple way to the coefficients of the heat expansion; it is possible to compute the $P_{j,l}^{B,V}$'s from the knowledge of the a_l 's for $j + 1 \leq l \leq 3j$. This is enough to re-compute the coefficient of \hbar^2 and also, in principle, the coefficients of \hbar^4 in the expansion, because the a_l 's are known up to $l = 6$ in the case of a flat metric (see [vdV]).

In this note, we will first describe precisely the semi-classical expansion for Schrödinger operators (in the case of an Euclidian metric) and the properties of the functions $P_{j,l}^{B,V}(x)$'s. Then, we will show how to compute the $P_{j,l}^{B,V}(x)$'s using an adaptation of the method used for the heat kernel (Weyl's Theorem on invariants and explicit examples). Finally, we will explain how the a_l 's are related to the $P_{j,l}^{B,V}(x)$'s. This gives us two proofs of the main formula given by Helffer and Purice in [H-P]; this paper was the initial motivation to this work.

1 Semi-classical trace for Schrödinger operators

Let us give a Schrödinger operator, with magnetic field $B = \sum_{1 \leq i < j \leq d} b_{ij} dx_i \wedge dx_j$ (a closed real 2-form) and electric potential V (a real valued smooth function) in some open domain $X \subset \mathbb{R}^d$. Locally, the 2-form B is exact and can be written $B = da$ and

$$\widehat{H_{h,a,V}} = \sum_{j=1}^d \left(\frac{\hbar}{i} \frac{\partial}{\partial x_j} - a_j(x) \right)^2 + V(x) .$$

The Weyl symbol of $\widehat{H_{h,a,V}}$ is $H_{a,V}(x, \xi) = \|\xi - a(x)\|^2 + V(x)$. Let us give $f \in \mathcal{S}(\mathbb{R})$ and $\phi \in C_0^\infty(X)$ and consider the trace of $\phi f(\widehat{H_{h,a,V}})$ as a distribution on $X \times \mathbb{R}$ (the density of states):

$$\text{Trace}(\phi f(\widehat{H_{h,a,V}})) = \int_X Z_{h,a,V}(g)(x) \phi(x) |dx| ,$$

where $Z_{h,a,V}(g)(x)$ is the value at the point (x, x) of the Schwartz kernel of $f(\widehat{H_{h,a,V}})$.

Theorem 1 *We have the following asymptotic expansion in power of \hbar :*

$$Z_{\hbar,a,V}(g)(x) \sim (2\pi\hbar)^{-d} \left[\int_{\mathbb{R}^d} f(\|\xi\|^2 + V(x)) |d\xi| + \sum_{j=1}^{\infty} \hbar^{2j} \left(\sum_{l=j+1}^{l=3j} P_{j,l}^{B,V}(x) \int_{\mathbb{R}^d} f^{(l)}(\|\xi\|^2 + V(x)) |d\xi| \right) \right]$$

We have the explicit formulas

$$\begin{aligned} P_{1,2}^{B,V} &= -\frac{1}{6} (\Delta V + \|B\|^2) , \\ P_{1,3}^{B,V} &= -\frac{1}{12} \|\nabla V\|^2 , \\ P_{2,3}^{B,V} &= -\frac{1}{180} (8\|\nabla B\|^2 + \|d^*B\|^2 + 12\langle \Delta B|B \rangle + 3\Delta^2 V) . \end{aligned}$$

Here $\|B\|^2 = \sum_{1 \leq i < j \leq d} b_{ij}^2$. If $d = 3$, $\|B\|$ is the Euclidean norm of the vector field associated to B .

The $P_{j,l}^{B,V}(x)$ are polynomials of the derivatives of B and V at the point x . Moreover, if λ, μ, c are constants and we define $\lambda^*(f)(x) = f(\lambda x)$, we have the following scaling properties:

1. $P_{j,l}^{\lambda, \lambda^*(B), \lambda^*(V)}(x) = \lambda^{2j} P_{j,l}^{B,V}(\lambda x)$. This will be used with $x = 0$.
2. $P_{j,l}^{\mu B, \mu^2 V}(x) = \mu^{2(l-j)} P_{j,l}^{B,V}(x)$
3. $P_{j,l}^{B, V+c}(x) = P_{j,l}^{B,V}(x)$
4. $P_{j,l}^{-B, V}(x) = P_{j,l}^{B,V}(x)$.
5. The $P_{j,l}^{B,V}$'s are invariant by the natural action of the orthogonal group $O(d)$ on the Taylor expansions of B and V at the point x .

As a consequence, we can get the following full trace expansion under some more assumptions:

Corollary 1 *Let us assume that $E_0 = \inf V < E_\infty = \liminf_{x \rightarrow \partial X} V(x)$ and $f \in C_0^\infty(-\infty, E_\infty]$, then the trace of $f(\widehat{H_{\hbar,a,V}})$ admits the following asymptotic expansion*

$$\begin{aligned} \text{Trace}(f(\widehat{H_{\hbar,a,V}})) &\sim (2\pi\hbar)^{-d} \int_X \left(\int_{\mathbb{R}^d} f(\|\xi\|^2 + V(x)) |d\xi| + \dots \right. \\ &\quad \left. \dots \sum_{j=1}^{\infty} \hbar^{2j} \sum_{l=j+1}^{l=3j} P_{j,l}^{B,V}(x) \int_{\mathbb{R}^d} f^{(l)}(\|\xi\|^2 + V(x)) |d\xi| \right) |dx| , \end{aligned}$$

The coefficient of \hbar^2 can be written as

$$-\frac{1}{12} \int_{X \times \mathbb{R}^d} f^{(2)}(\|\xi\|^2 + V(x)) (\Delta V(x) + 2\|B(x)\|^2) |dx d\xi| .$$

The expansion follows from [He-Ro]. An integration by part in x gives

$$\int_X f^{(3)}(\|\xi\|^2 + V(x))\|\nabla V(x)\|^2|dx| = - \int_X f^{(2)}(\|\xi\|^2 + V(x))\Delta V(x)|dx| .$$

2 Existence of the \hbar -expansion of $Z_{\hbar,a,V}$

The existence of the expansion is known in general from [He-Ro] and the calculus of the symbol of $f(\widehat{H_{\hbar,a,V}})$. We get

$$\int_X Z_{\hbar,a,V}(f)(x)\phi(x)|dx| = (2\pi\hbar)^{-d} \left(\sum_{j=0}^{\infty} \hbar^{2j} \sum_{l=0}^{k_j} \int \phi(x)f^{(l)}(H_{a,V}(x,\xi)Q_{j,l}(x,\xi)|dx d\xi| \right)$$

where the $Q_{j,l}(x,\xi)$'s are polynomials in the Taylor expansion of $H_{a,V}$ at the point (x,ξ) . The previous expansion is valid for any (admissible) pseudo-differential operator. In the case of Schrödinger operators we can make integrations by part in the integrals $\int f^{(l)}(H_{a,V}(x,\xi)Q_{j,l}(x,\xi)|d\xi|$ which reduces to a similar formula where we can replace the $Q_{j,l}(x,\xi)$'s by $P_{j,l}(x)$'s. This is based on the expansion of $Q_{j,l}$ as a polynomial in ξ in powers of $(\xi - a)$ and

$$d_\xi((\xi_j - a_j)f^{(l)}(H_{a,V})\iota(\partial_{\xi_j})d\xi) = 2\|\xi_j - a_j\|^2 f^{(l+1)}(H_{a,V})d\xi + f^{(l)}(H_{a,V})d\xi .$$

We have only to check that the powers of ξ in $Q_{j,l}(x,\xi)$ is less than l : this is due to the fact that the l -th Moyal power of $H_{a,V}(z) - H_{a,V}(z_0)$ use monomials of degree l in the derivatives of order ≥ 1 of $H_{a,V}(z)$ which are all of degree ≤ 1 in ξ . Using Gauge invariance at the point x (Section 3), we can assume that $a(x) = 0$.

3 Gauge invariance

We have

$$\text{Trace}(\phi e^{-iS(x)/\hbar} f(\widehat{H_{\hbar,a,V}}) e^{iS(x)/\hbar}) = \text{Trace}(\phi f(\widehat{H_{\hbar,a,V}}))$$

and

$$e^{-iS(x)/\hbar} f(\widehat{H_{\hbar,a,V}}) e^{iS(x)/\hbar} = f(\widehat{H_{a+dS,V}}) .$$

Hence, we can chose any local gauge a in order to compute the expansion: using the synchronous gauge (see Section 7), we get the individual terms

$$\int f^{(l)}(H_{0,V})P_{j,l}^{B,V}(x)|d\xi|$$

for the expansion, where the $P_{j,l}^{B,V}(x)$ depend only of the Taylor expansions of B and V at the point x .

4 Properties of the $P_{j,l}$'s

4.1 Range of l for j fixed

From the scaling properties, we deduce that, in a monomial

$$D^{\alpha_1} B_{i_1, j_1} \cdots D^{\alpha_k} B_{i_k, j_k} D^{\beta_1} V \cdots D^{\beta_m} V ,$$

belonging to $P_{j,l}$, we have $k + 2m = 2(l - j)$ and $k + |\alpha_1| + \cdots + |\alpha_k| + |\beta_1| + \cdots + |\beta_m| = 2j$. Moreover, for $j \geq 1$, $k + m \geq 1$ and $|\beta_p| \geq 1$. Hence $j + 1 \leq l \leq 3j$. The previous bounds are sharp: take the monomials $\Delta^j V$ and $\|\nabla V\|^{2j}$ which give $l = j + 1$ and $l = 3j$.

4.2 Invariance properties

1. Let us assume that we look at the point $x = 0$ and consider the operator $D_\mu(f)(x) = f(\mu x)$. We have

$$D_\mu \circ \hat{H}_{\hbar, A, V} \circ D_{1/\mu} = \hat{H}_{\hbar/\mu, A \circ D_\mu, V \circ D_\mu} .$$

The same relation is true for any function $f(\hat{H}_{\hbar, A, V})$ and then we have, looking at the Schwartz kernels and using the Jacobian μ^d of D_μ :

$$P_{j,l}^{B,V}(0) \int_{\mathbb{R}^d} f^{(l)}(\|\xi\|^2 + V(0)) |d\xi| = \mu^{-2j} P_{j,l}^{\mu^* B, \mu^* V}(0) \int_{\mathbb{R}^d} f^{(l)}(\|\xi\|^2 + V(0)) |d\xi| .$$

2. We have

$$\hat{H}_{\hbar, \mu a, \mu^2 V} = \mu^2 \hat{H}_{\frac{\hbar}{\mu}, a, V} .$$

3. Changing V into $V + c$ gives a translation by c in the function f but does not change the $P_{j,l}^{B,V}$'s.
4. Changing B into $-B$ gives a complex conjugation in the computations. The final result is real valued.
5. Orthogonal invariance is clear: an orthogonal change of coordinates around the point x preserves the density of states.

4.3 The case $d = 2$

We deduce from the scaling properties and invariance by the orthogonal group, that there exists constants a_d , b_d , c_d so that $P_{1,2}^{B,V}(x) = a_d \Delta V + b_d \|B\|^2$, $P_{1,3}(x) = c_d \|\nabla V\|^2$.

5 Explicit examples

The calculation for the harmonic oscillators and the constant magnetic fields allows to determine the constants a_d, b_d, c_d .

5.1 Harmonic oscillators

Let us consider $\Omega = -\hbar^2 \frac{d^2}{dx^2} + x^2$ with $d = 1$. The kernel of $P(t, x, y)$ of $\exp(-t\Omega)$ is given by the Mehler formula:

$$P(t, x, y) = (2\pi\hbar \sinh(2t\hbar))^{-\frac{1}{2}} \exp\left(-\frac{1}{2\hbar \sinh(2t\hbar)} (\cosh(2t\hbar)(x^2 + y^2) - 2xy)\right).$$

Hence

$$P(t, x, x) \sim (2\pi\hbar)^{-1} e^{-tx^2} \left(\int_{\mathbb{R}} e^{-t\xi^2} d\xi \right) (1 - \hbar^2(t^2 - t^3x^2)/3 + 0(\hbar^4)).$$

Hence $P_{1,2}(x) = -V''(x)/6$, $P_{1,3}(x) = -V'(x)^2/12$.

Similarly, in dimension $d > 1$, we get $P_{1,2}(x) = -\Delta V(x)/6$, $P_{1,3}(x) = -\|\nabla V\|^2/12$.

5.2 Constant magnetic field

Let us consider the case of a constant magnetic field B in the plane and denote by $Q(t, x, y)$ the kernel of $\exp(-tH_{B,0})$. We have (see [A-H-S])

$$Q(t, x, x) = \frac{B\hbar}{4\pi \sinh Bt\hbar}.$$

Hence the asymptotic expansion

$$Q(t, x, x) = (2\pi\hbar)^{-2} \int \exp(-t\|\xi\|^2) |d\xi| (1 - t^2\hbar^2 B^2/6 + 0(\hbar^4)),$$

hence $P_{1,2}(x) = -B^2/6$, $P_{1,3}(x) = 0$.

Using the normal form $B = b_{12}dx_1 \wedge dx_2 + b_{34}dx_3 \wedge dx_4 + \dots$, we get in dimension $d > 2$, $P_{1,2}(x) = -\|B\|^2/6$, $P_{1,3}(x) = 0$.

6 Heat expansion from the semi-classical expansions

We have $t\widehat{H_{1,a,V}} = \widehat{H_{\sqrt{t}, \sqrt{ta}, tV}}$. Using the expansion of Theorem 1 with $f(E) = e^{-E}$, we get easily the pointwise expansion of the heat kernel on the diagonal as

$t \rightarrow 0^+$:

$$[\exp(-t\widehat{H}_{1,a,V})](x, x) \sim \frac{1}{(4\pi t)^{d/2}} e^{-tV(x)} \left[\sum_{l=0}^{\infty} \left(\sum_{l/3 \leq j \leq l-1} P_{j,l}^{B,V}(x) \right) (-t)^l \right].$$

In particular, $a_1(x) = -V(x)$ and the coefficient $a_2(x)$ is given by

$$a_2(x) = \frac{1}{2}V(x)^2 - \frac{1}{6}\Delta V(x) - \frac{1}{6}\|B(x)\|^2.$$

This formula agrees with Equation (3) of Theorem 3.3.1 in [Gilkey2].

This give another way to compute the $P_{j,l}$'s: if, as power series in t ,

$$\sum_{l=0}^{\infty} (-1)^l b_l(x) t^l = e^{tV(x)} \left(\sum_{l=0}^{\infty} a_l(x) t^l \right),$$

we have

$$\sum_{l/3 \leq j \leq l-1} P_{j,l}^{B,V}(x) = b_l(x).$$

$P_{j,l}^{B,V}$ is the sum of monomials homogeneous of degree $2(l-j)$ in b_l where B and its derivatives have weights 1 while V and its derivatives have weights 2.

The heat coefficients a_l on flat spaces are known for $l \leq 6$ from [vdV]. This is enough to check the term in \hbar^2 (uses a_2 and a_3) in [H-P] and to compute the term in \hbar^4 in the semi-classical expansion (uses the a_l 's for $3 \leq l \leq 6$).

We have also a mixt expansion writing $t\widehat{H}_{h,a,V} = \widehat{H}_{\sqrt{t\hbar}, \sqrt{t}a, tV}$, we get a power series expansion in powers of \hbar and t valid in the domain $\hbar^2 t \rightarrow 0$ and $0 < t \leq t_0$ for the pointwise trace of $\exp(-t\widehat{H}_{h,a,V})$:

$$Z_{t,\hbar}(x) \sim \frac{1}{(4\pi t)^{d/2}} e^{-tV(x)} \left(1 + \sum_{j \geq 1, j+1 \leq l \leq 3j} \hbar^{2j} (-t)^l P_{j,l}^{B,V}(x) \right).$$

This shows that the integrals $\int_X V(x)^k |dx|$ and $\int_X P_{j,l}^{B,V}(x) |dx|$ are recoverable from the semi-classical spectrum.

7 The synchronous gauge

The main idea is to find an appropriate gauge a adapted to the point x_0 where we want to make the symbolic computation. We use the trivialization of the bundle by parallel transportation along the rays: the potential a vanishes on the radial vector field ².

²This gauge is sometimes called the Fock-Schwinger gauge; in [A-B-P], it is called the synchronous framing

We will do that for the Taylor expansions degree by degree. In what follows we will use a decomposition for 1-forms, but it works also for k -forms.

Let us denote by Ω_N^k the finite dimensional vector space of k -differential forms on \mathbb{R}^d whose coefficients are homogeneous polynomial of degree N and by $W = \sum_{j=1}^d x_j \frac{\partial}{\partial x_j}$ the radial vector field. The exterior differential induces a linear map from Ω_N^k into Ω_{N-1}^{k+1} and the inner product $\iota(W)$ a map from Ω_N^k into Ω_{N+1}^{k-1} . They define complexes which are exact except at $k = N = 0$. Moreover, we have a situation similar to Hodge theory:

$$\Omega_N^k = d\Omega_{N+1}^{k-1} \oplus \iota(W)\Omega_{N-1}^{k+1} .$$

This is due to Cartan's formula: the Lie derivative of a form $\omega \in \Omega_N^k$ satisfies, from the direct calculation, $\mathcal{L}_W\omega = (k + N)\omega$, and, by Cartan's formula, $\mathcal{L}_W\omega = d(\iota(W)\omega) + \iota(W)d\omega$. So

$$\omega = \frac{1}{k + N} (d(\iota(W)\omega) + \iota(W)d\omega) .$$

It remains to show that this is a direct sum: if $\omega = d\alpha = \iota(W)\gamma$, we have $\iota(W)\omega = 0$ and $d\omega = 0$; from the previous decomposition, we see that $\omega = 0$. Let us denote by $J^N\omega$, where ω is a differential form of degree k , the form in Ω_N^k which appears in the Taylor expansion of ω .

We get

Proposition 1 *If $P(J^0a, J^1a, \dots, J^Na)$ is a polynomial in the Taylor expansion of the 1-form a at some order N which is invariant by $a \rightarrow a + dS$, P is independent of J^0a and*

$$P(J^1a, \dots, J^Na) = P\left(\frac{1}{2}J^1\iota(W)B, \dots, \frac{1}{N+1}J^N\iota(W)B\right)$$

is a polynomial of the Taylor expansion of B to the order $N - 1$.

References

- [A-B-P] M. Atiyah, R. Bott & Patodi. On the heat equation and the index theorem. *Inv. Math.* **19**:279–330 (1973).
- [Avramidi] I.G. Avramidi. The covariant technique for the calculation of the heat kernel asymptotic expansion. *Phys. Lett. B.* **238**:92–97 (1990).
- [A-H-S] J. Avron, I. Herbst & B. Simon. Schrödinger operators with magnetic fields. I. General interactions. *Duke Math. J.*, **45**:847–883 (1978).
- [Berger] M. Berger. Sur le spectre d'une variété riemannienne. *C. R. Acad. Sci. Paris Sér. A-B*, **263**:A13–A16 (1966).

- [B-G-M] M. Berger, P. Gauduchon & E. Mazet. Le spectre d'une variété riemannienne compacte. *Lecture Notes in Math, Springer* **194** (1971).
- [Gilkey1] P. B. Gilkey. The spectral geometry of a Riemannian manifold. *J. Diff. Geom.*, **10**:601–618 (1975).
- [Gilkey2] P. B. Gilkey. Asymptotic formulas in spectral geometry. *Chapman and Hall/CRC* (2004).
- [Gracia] A. Gracia-Saz. The symbol of a function of a pseudo-differential operator. *Ann. Inst. Fourier* **55**:2257–2284 (2005).
- [H-P] B. Helffer & R. Purice. Magnetic calculus and semi-classical trace formulas. *Journal of Physics A*, 43474-0288 (2011) and *arXiv*:1005.1795v1 (2010).
- [He-Ro] B. Helffer & D. Robert. Calcul fonctionnel par la transformée de Mellin et applications. *Jour. of Funct. Anal.* **53**:246–268 (1983).
- [Hi] M. Hitrik, Existence of resonances in magnetic scattering. On the occasion of the 65th birthday of Professor Michael Eastham. *J. Comput. Appl. Math.* **148(1)**:91–97 (2002).
- [Hi-Po-1] M. Hitrik & I. Polterovich, Regularized traces and Taylor expansions for the heat semigroup. *J. London Math. Soc.* **68(2)**: 402–418 (2003).
- [Hi-Po-2] M. Hitrik and I. Polterovich, Resolvent expansions and trace regularizations for Schrödinger operators. *Advances in differential equations and mathematical physics (Birmingham, AL)* **161** (2002).
- [Kac] M. Kac. Can one hear the shape of a drum?. *Amer. Math. Monthly*, **73**: 1–23 (1966).
- [McK-Si] H. P. Mc Kean & I. Singer. Curvature and the eigenvalues of the Laplacian. *Jour. Diff. Geometry* **1**:43–69 (1967).
- [Po] I. Polterovich, Heat invariants of Riemannian manifolds, *Israel J. Math.* **119**: 239–252 (2000).
- [vdV] A. van de Ven. Index-free heat kernel coefficients. *Class. Quantum Gravity*. **15**:2311–2344 (1998).